



C4.3 Functional Analytic Methods for PDEs

Lecture 5

Luc Nguyen
luc.nguyen@maths

University of Oxford

MT 2021

In the last 4 lectures

- L^p spaces and their properties.

This lecture

- Pre-compactness criterion in $L^p(\Omega)$.
- Divergence theorem and Integration by parts formula.
- Weak derivatives.
- Sobolev spaces $W^{k,p}(\Omega)$ and $W_0^{k,p}(\Omega)$ as Banach spaces.
- Differentiation rule for convolution of Sobolev functions.

Theorem (Ascoli-Arzelà's theorem)

Let K be a compact subset of \mathbb{R}^n . Suppose that (f_i) is a sequence of functions of $C(K)$ such that

- ① (Boundedness) $\sup_i \|f_i\|_{C(K)} < \infty$,
- ② (Equi-continuity) For every $\varepsilon > 0$, there exists $\delta > 0$ such that $|f_i(x) - f_i(y)| < \varepsilon$ for all i and all $x, y \in K$ with $|x - y| < \delta$.

Then there exists a subsequence (f_{i_j}) which converges uniformly on K .

Theorem (Kolmogorov-Riesz-Fréchet's theorem)

Let $1 \leq p < \infty$ and Ω be an open bounded subset of \mathbb{R}^n . Suppose that a sequence (f_i) of $L^p(\Omega)$ satisfies

- ① (Boundedness) $\sup_i \|f_i\|_{L^p(\Omega)} < \infty$,
- ② (Equi-continuity in L^p) For every $\varepsilon > 0$, there exists $\delta > 0$ such that $\|\tau_y \tilde{f}_i - \tilde{f}_i\|_{L^p(\Omega)} < \varepsilon$ for all $|y| < \delta$, where \tilde{f}_i is the extension by zero of f_i to the whole of \mathbb{R}^n .

Then, there exists a subsequence (f_{i_j}) which converges in $L^p(\Omega)$.

By definition $\tilde{f}_i : \mathbb{R}^n \rightarrow \mathbb{R}$ is given by $\tilde{f}_i = f_i$ in Ω and $\tilde{f}_i = 0$ in $\mathbb{R}^n \setminus \Omega$.

Pre-compactness criterion in $L^p(\Omega)$

Proof

- As in the proof of Ascoli-Arzelà's theorem, it suffices to show that, for every given $\varepsilon > 0$, there exists a subsequence $(f_{i_j}^\varepsilon)$ of (f_i) such that

$$\|f_{i_j}^\varepsilon - f_{i_k}^\varepsilon\|_{L^p(\Omega)} \leq 3\varepsilon \text{ for large } j, k. \quad (***)$$

- Claim: For every fixed $\varphi \in C_c^\infty(\mathbb{R}^n)$, the sequence $(\tilde{f}_i * \varphi|_{\bar{\Omega}})$ satisfies the condition of Ascoli-Arzelà's theorem.
 - ★ First, by Hölder's inequality, we have

$$\|\tilde{f}_i\|_{L^1(\mathbb{R}^n)} = \|f_i\|_{L^1(\Omega)} \leq \|f_i\|_{L^p(\Omega)} |\Omega|^{1/p'}.$$

Thus, by the boundedness of (f_i) in $L^p(\Omega)$, we have that (\tilde{f}_i) is bounded in $L^1(\mathbb{R}^n)$.

Pre-compactness criterion in $L^p(\Omega)$

Proof

- Claim: $(\tilde{f}_i * \varphi|_{\bar{\Omega}})$ satisfies the condition of Ascoli-Arzelà's theorem.

- ★ $\sup_i \|\tilde{f}_i\|_{L^1(\mathbb{R}^n)} < \infty$.
- ★ By Young's convolution inequality

$$\|\tilde{f}_i * \varphi\|_{L^\infty(\mathbb{R}^n)} \leq \|\tilde{f}_i\|_{L^1(\mathbb{R}^n)} \|\varphi\|_{L^\infty(\mathbb{R}^n)}.$$

So $\sup_i \|\tilde{f}_i * \varphi\|_{C(\bar{\Omega})} < \infty$.

- ★ Next,

$$\begin{aligned} |\tilde{f}_i * \varphi(x) - \tilde{f}_i * \varphi(y)| &\leq \int_{\mathbb{R}^n} |\varphi(x-z) - \varphi(y-z)| |\tilde{f}_i(z)| dz \\ &\leq \|\varphi\|_{Lip(\mathbb{R}^n)} |x-y| \|\tilde{f}_i\|_{L^1(\mathbb{R}^n)}. \end{aligned}$$

So by squeezing $|x-y|$, we can make $\sup_i |\tilde{f}_i * \varphi(x) - \tilde{f}_i * \varphi(y)|$ as small as we want.

Pre-compactness criterion in $L^p(\Omega)$

Proof

- $(\tilde{f}_i * \varphi|_{\bar{\Omega}})$ satisfies the condition of Ascoli-Arzelà's theorem.
- Now, take a non-negative function $\varrho \in C_c^\infty(B_1)$ with $\int_{\mathbb{R}^n} \varrho = 1$ and, for $\eta > 0$, let $\varrho_\eta(x) = \frac{1}{\eta^n} \varrho(x/\eta)$ be the standard mollifiers. Recall that we have the estimate

$$\begin{aligned} \|\tilde{f}_i * \varrho_\eta - \tilde{f}_i\|_{L^p}^p &\leq \int_{\mathbb{R}^n} |\varrho_\eta(y)| \|\tau_{-y}\tilde{f}_i - \tilde{f}_i\|_{L^p}^p dy \\ &\leq \sup_{|y| \leq \eta} \|\tau_y \tilde{f}_i - \tilde{f}_i\|_{L^p}^p \int_{\mathbb{R}^n} |\varrho_\eta(y)| dy \\ &= \sup_{|y| \leq \eta} \|\tau_y \tilde{f}_i - \tilde{f}_i\|_{L^p}^p. \end{aligned}$$

Pre-compactness criterion in $L^p(\Omega)$

Proof

- $(\tilde{f}_i * \varphi|_{\bar{\Omega}})$ satisfies the condition of Ascoli-Arzelà's theorem.
- $\|\tilde{f}_i * \varrho_\eta - \tilde{f}_i\|_{L^p} \leq \sup_{|y| \leq \eta} \|\tau_y \tilde{f}_i - \tilde{f}_i\|_{L^p}$.
- We are now ready to prove (***):
 - ★ By the equi-continuity, there exists a small $\eta > 0$ such that $\|\tilde{f}_i * \varrho_\eta - \tilde{f}_i\|_{L^p} \leq \varepsilon$ for all i .
 - ★ Using Ascoli-Arzelà's theorem, select a subsequence $(f_{i_j}^\varepsilon)$ of (f_i) such that $(\tilde{f}_{i_j}^\varepsilon * \varrho_\eta|_{\bar{\Omega}})$ is convergent in $C(\bar{\Omega})$.
 - ★ It follows that $\|\tilde{f}_{i_j}^\varepsilon * \varrho_\eta - \tilde{f}_{i_k}^\varepsilon * \varrho_\eta\|_{L^p(\Omega)} \leq \varepsilon$ for large j, k .
 - ★ Consequently, by triangle inequality,

$$\begin{aligned} \|f_{i_j}^\varepsilon - f_{i_k}^\varepsilon\|_{L^p(\Omega)} &\leq \|f_{i_j}^\varepsilon * \varrho_\eta - f_{i_k}^\varepsilon * \varrho_\eta\|_{L^p(\Omega)} + \|f_{i_j}^\varepsilon * \varrho_\eta - f_{i_j}^\varepsilon\|_{L^p(\Omega)} \\ &\quad + \|\tilde{f}_{i_k}^\varepsilon * \varrho_\eta - f_{i_k}^\varepsilon\|_{L^p(\Omega)} \leq 3\varepsilon \text{ for large } j, k, \end{aligned}$$

which is (***) .

Frequently used terminologies/notations


- Ω denotes a domain in \mathbb{R}^n .
- $C^k(\Omega)$ denotes the space of functions which are k -times continuously differentiable in Ω .
- $C^k(\bar{\Omega})$ denotes the subspace of $C^k(\Omega)$ consisting of functions which can be extended to a k -times continuously differentiable functions on some open set containing $\bar{\Omega}$.
- $C_c^k(\Omega)$ denotes the subspace of $C^k(\Omega)$ consisting of functions f such that $Supp(f) = \overline{\{f \neq 0\}}$ is a bounded closed subset of Ω .

Frequently used terminologies/notations

- Ω is said to be a Lipschitz (resp. C^k) domain, or equivalently, $\partial\Omega$ is said to be Lipschitz (resp. C^k), if for every $x_0 \in \partial\Omega$ there exists a radius $r_0 > 0$ such that, after a relabeling of coordinate axes if necessary,

$$\Omega \cap B_{r_0}(x_0) = \{x \in B_{r_0}(x_0) : x_n > \gamma(x_1, \dots, x_{n-1})\}$$

for some Lipschitz (resp. C^k) function γ .



Divergence theorem

Let Ω be a bounded Lipschitz domain in \mathbb{R}^n . Fact: $\partial\Omega$ admits an 'outward pointing' unit normal n .

Theorem (Divergence theorem)

Let $F \in C^1(\bar{\Omega}; \mathbb{R}^n)$. Then

$$\int_{\Omega} \operatorname{div} F \, dx = \int_{\partial\Omega} F \cdot n \, dS.$$

In particular, if $F \in C_c^1(\Omega; \mathbb{R}^n)$, then

$$\int_{\Omega} \operatorname{div} F \, dx = 0.$$

IBP formula

Let Ω be a bounded Lipschitz domain in \mathbb{R}^n .

Theorem (Integration by parts formula)

Let $f, g \in C^1(\bar{\Omega})$. Then

$$\int_{\Omega} f \partial_i g \, dx = \int_{\partial\Omega} f g n_i \, dS - \int_{\Omega} \partial_i f g \, dx.$$

In particular, if f or g has compact support in Ω , then

$$\int_{\Omega} f \partial_i g \, dx = - \int_{\Omega} \partial_i f g \, dx.$$

Weak derivatives

Let Ω be a domain in \mathbb{R}^n .

Definition

Let $f \in L^1_{loc}(\Omega)$ and $\alpha = (\alpha_1, \dots, \alpha_n)$ be a multi-index. A function $g \in L^1_{loc}(\Omega)$ is said to be a weak α -derivative of f if

$$\int_{\Omega} f \partial^{\alpha} \varphi \, dx = (-1)^{|\alpha|} \int_{\Omega} g \varphi \, dx \text{ for all } \varphi \in C_c^{\infty}(\Omega). \quad (1)$$

We write $g = \partial^{\alpha} f$ in the weak sense.

The function φ is called a *test function*.

Example of weak derivatives

- If $f \in C^1(\bar{\Omega})$ and Ω is a bounded Lipschitz domain, then its classical derivatives are also its weak derivatives.
- Suppose $\Omega = (-1, 1)$ and $f(x) = |x|$. Then, if $\varphi \in C_c^\infty(-1, 1)$, we have by IBP that

$$\begin{aligned}\int_{-1}^1 f(x) \varphi'(x) dx &= \int_{-1}^0 (-x) \varphi'(x) dx + \int_0^1 x \varphi'(x) dx \\ &= -x\varphi(x) \Big|_{-1}^0 - \int_{-1}^0 (-1) \varphi(x) dx \\ &\quad + x\varphi(x) \Big|_0^1 - \int_0^1 (1) \varphi(x) dx \\ &= - \int_{-1}^1 \text{sign}(x) \varphi(x) dx.\end{aligned}$$

So $f'(x) = \text{sign}(x)$ in the weak sense.

Uniqueness of weak derivatives

Lemma

Let $f \in L^1_{loc}(\Omega)$ and $\alpha = (\alpha_1, \dots, \alpha_n)$ be a multi-index. The weak α -derivative of f , if exists, is uniquely defined up to a set of measure zero.

This follows from the definition of weak derivative and the following:

Lemma (Fundamental lemma of the Calculus of Variations)

Let $g \in L^1_{loc}(\Omega)$. If $\int_{\Omega} g\varphi = 0$ for all $\varphi \in C_c^{\infty}(\Omega)$, then $g = 0$ a.e. in Ω .

Uniqueness of weak derivatives

Proof

- We will only consider the case Ω is a bounded domain and $g \in L^1(\Omega)$. The general case is left as an exercise.
- In Sheet 1, you showed that $C_c^\infty(\Omega)$ is dense in $L^1(\Omega)$. Thus, for any $\varepsilon > 0$, we can select $h \in C_c^\infty(\Omega)$ such that $\|g - h\|_{L^1} \leq \varepsilon$. Furthermore, by triangle inequality $\|h\|_{L^1} \geq \|g\|_{L^1} - \varepsilon$.
- For $\delta > 0$, let $h_\delta = \frac{h}{\sqrt{\delta^2 + h^2}}$ so that $h_\delta \in C_c^\infty(\Omega)$ and $|h_\delta| \leq 1$.
- By hypotheses, $\int_\Omega gh_\delta dx = 0$.
- By construction, $\left| \int_\Omega (g - h)h_\delta dx \right| \leq \|g - h\|_{L^1} \|h_\delta\|_{L^\infty} \leq \varepsilon$.
- It follows that

$$\varepsilon \geq \int_\Omega gh_\delta dx - \int_\Omega (g - h)h_\delta dx = \int_\Omega hh_\delta dx.$$

Uniqueness of weak derivatives

Proof

- Recalling the expression of h_δ , we have

$$\varepsilon \geq \int_{\Omega} \frac{h^2}{\sqrt{\delta^2 + h^2}} dx.$$

- The integrand on the right hand side converges monotonically increasingly to $|h|$. Thus, by Lebesgue's monotone convergence theorem,

$$\varepsilon \geq \int_{\Omega} |h| dx = \|h\|_{L^1}.$$

- Recall that $\|h\|_{L^1} \geq \|g\|_{L^1} - \varepsilon$, we obtain that $2\varepsilon \geq \|g\|_{L^1}$. Sending $\varepsilon \rightarrow 0$, we obtain $\|g\|_{L^1} = 0$, i.e. $g = 0$ a.e. in Ω .

Remark

Suppose that

- (i) $f \in L^1(\Omega)$ is weakly differentiable with weak derivatives $\partial_1^w f, \dots, \partial_n^w f$,
- (ii) and, for some subdomain $\omega \subset \Omega$, f is classically differentiable in ω with classical derivatives $\partial_1^c f, \dots, \partial_n^c f$.

Then

$$\partial_i^w f = \partial_i^c f \text{ a.e. in } \omega \text{ for all } i = 1, \dots, n.$$

A relation between classical and weak derivatives

Sketch of proof

- Using the definition of weak derivatives, $f|_{\omega}$ is weakly differentiable with weak derivatives $\partial_1^w f|_{\omega}, \dots, \partial_n^w f|_{\omega}$.
- As f is classically differentiable in ω , its classical derivatives are also weak derivatives of $f|_{\omega}$.
- By the uniqueness of weak derivatives, the conclusion follows.

Example of non-existence of weak derivatives

If $\Omega = (-1, 1)$ and $u(x) = \text{sign}(x)$, then u has no weak derivative.

Proof

- Suppose otherwise that $u' = g \in L^1_{loc}(-1, 1)$. Then, for $\varphi \in C_c^\infty(-1, 1)$,

$$\begin{aligned}\int_{-1}^1 g(x)\varphi(x) dx &= \int_{-1}^0 \varphi'(x) dx - \int_0^1 \varphi'(x) dx \\ &= [\varphi(0) - \varphi(-1)] - [\varphi(1) - \varphi(0)] \\ &= 2\varphi(0).\end{aligned}$$

- In particular, if we take $\varphi \in C_c^\infty(-1, 0)$, we have

$\int_{-1}^0 g(x)\varphi(x) dx = 0$. So $g = 0$ a.e. in $(-1, 0)$. Likewise, $g = 0$ a.e. in $(0, 1)$. So $g = 0$ a.e. in $(-1, 1)$.

- We thus have $0 = \int_{-1}^1 g(x)\varphi(x) dx = 2\varphi(0)$ for all $\varphi \in C_c^\infty(-1, 1)$, which is impossible.

The Sobolev spaces $W^{k,p}(\Omega)$

- Ω : a domain of \mathbb{R}^n .
- For $k \geq 0$ and $1 \leq p \leq \infty$, define

$$W^{k,p}(\Omega) = \left\{ f \in L^p(\Omega) \mid \forall |\alpha| \leq k, \text{ the weak derivative } \partial^\alpha f \text{ exists and belongs to } L^p(\Omega) \right\}.$$

We equip $W^{k,p}(\Omega)$ with the norm

$$\|u\|_{W^{k,p}(\Omega)} = \left[\sum_{|\alpha| \leq k} \|\partial^\alpha u\|_{L^p(\Omega)}^p \right]^{\frac{1}{p}}$$

so that $W^{k,p}(\Omega)$ is a normed vector space (check this!).

- For $p = 2$, we also write $H^k(\Omega)$ for $W^{k,2}(\Omega)$. These are inner product spaces (check this!) with inner product

$$\langle u, v \rangle_{W^{k,2}(\Omega)} = \sum_{|\alpha| \leq k} \langle \partial^\alpha u, \partial^\alpha v \rangle_{L^2(\Omega)}.$$

Examples of Sobolev functions

Let $\Omega = (-1, 1)$ and $f(x) = |x|$.

- We have that $f'(x) = \text{sign}(x)$ and so $f \in W^{1,p}(-1, 1)$ for every $p \in [1, \infty]$.
- The function $f'(x) = \text{sign}(x)$ has no weak derivatives, and so $f \notin W^{2,p}(-1, 1)$ for any $p \in [1, \infty]$.

Completeness of $W^{k,p}(\Omega)$

Theorem

For $k \geq 0$ and $1 \leq p \leq \infty$, $W^{k,p}(\Omega)$ is a Banach space. When $p = 2$, $W^{k,2}(\Omega)$ is a Hilbert space.

Proof

- We have seen that $W^{k,p}$ is a normed vector space and $W^{k,2}$ is an inner product space. It remains to show that $W^{k,p}$ is complete.
- Suppose that (u_m) is a Cauchy sequence in $W^{k,p}$. We need to show that there exists $u \in W^{k,p}$ such that $\|u_m - u\|_{W^{k,p}} \rightarrow 0$.
- For $|\alpha| \leq k$, $(\partial^\alpha u_m)$ is Cauchy in L^p , as

$$\|\partial^\alpha u_m - \partial^\alpha u_j\|_{L^p} \leq \|u_m - u_j\|_{W^{k,p}}.$$

By Riesz-Fischer's theorem, we have that $(\partial^\alpha u_m)$ converges in L^p to some $v_\alpha \in L^p$.

Completeness of $W^{k,p}(\Omega)$

Proof

- (u_m) is Cauchy in $W^{k,p}$.
- For $|\alpha| \leq k$, $(\partial^\alpha u_m)$ converges in L^p to some $v_\alpha \in L^p$.
- To conclude, we show that $u := v_{(0,\dots,0)}$ belongs to $W^{k,p}$ and $u_m \rightarrow u$ in $W^{k,p}$.

★ By definition of weak derivatives, we have for $|\alpha| \leq k$ that

$$\int_{\Omega} u_m \partial^\alpha \varphi \, dx = (-1)^{|\alpha|} \int_{\Omega} \partial^\alpha u_m \varphi \, dx \text{ for all } \varphi \in C_c^\infty(\Omega),$$

★ Now we would like to pass $m \rightarrow \infty$. By Hölder's inequality

$$\left| \int_{\Omega} (u_m - u) \partial^\alpha \varphi \, dx \right| \leq \|u_m - u\|_{L^p} \|\partial^\alpha \varphi\|_{L^{p'}} \rightarrow 0.$$

So $\int_{\Omega} u_m \partial^\alpha \varphi \, dx \rightarrow \int_{\Omega} u \partial^\alpha \varphi \, dx$.

★ Similarly, $\int_{\Omega} \partial^\alpha u_m \varphi \, dx \rightarrow \int_{\Omega} v_\alpha \varphi \, dx$.

Completeness of $W^{k,p}(\Omega)$

Proof

- (u_m) is Cauchy in $W^{k,p}$.
- For $|\alpha| \leq k$, $(\partial^\alpha u_m)$ converges in L^p to some $v_\alpha \in L^p$.
- ★ We thus have

$$\int_{\Omega} u \partial^\alpha \varphi = (-1)^{|\alpha|} \int_{\Omega} v_\alpha \varphi \text{ for all } \varphi \in C_c^\infty(\Omega).$$

So v_α is the weak α -derivative of u . So $u \in W^{k,p}$.

★ Now

$$\begin{aligned} \|u_m - u\|_{W^{k,p}}^p &= \sum_{|\alpha| \leq k} \|\partial^\alpha u_m - \partial^\alpha u\|_{L^p}^p \\ &= \sum_{|\alpha| \leq k} \|\partial^\alpha u_m - v_\alpha\|_{L^p}^p \xrightarrow{m \rightarrow \infty} 0. \end{aligned}$$

So $u_m \rightarrow u$ in $W^{k,p}$.

- We conclude that $W^{k,p}$ is complete.