

BO1 History of Mathematics
Lecture XI
19th-century rigour in real analysis

MT 2021 Week 6

Summary

Part 1

- ▶ New difficulties emerge
- ▶ Continuity and convergence

Part 2

- ▶ Integration
- ▶ The Fundamental Theorem of Calculus
- ▶ New ideas about integration

Recall from lecture VII: Fourier series, 1822

Joseph Fourier, *Théorie analytique de la chaleur* [*Analytic theory of heat*] (1822):

Suppose that $\phi(x) = a \sin x + b \sin 2x + c \sin 3x + \dots$

and also that $\phi(x) = x\phi'(0) + \frac{1}{6}x^3\phi'''(0) + \dots$

After many pages of calculations, multiplying and comparing power series, Fourier found that the coefficient of $\sin nx$ must be

$$\frac{2}{\pi} \int_0^{\pi} \phi(x) \sin nx \, dx$$

Fourier's derivation was based on 'naive' manipulations of infinite series. It was ingenious but non-rigorous, shaky.

BUT it led to profound results

New doubts in the early 19th century

Fourier's work converged with more philosophical investigation to stimulate questions concerning:

New doubts in the early 19th century

Fourier's work converged with more philosophical investigation to stimulate questions concerning:

- ▶ functions — what exactly should they be?

New doubts in the early 19th century

Fourier's work converged with more philosophical investigation to stimulate questions concerning:

- ▶ functions — what exactly should they be?
- ▶ convergence — what exactly should it be?

New doubts in the early 19th century

Fourier's work converged with more philosophical investigation to stimulate questions concerning:

- ▶ functions — what exactly should they be?
- ▶ convergence — what exactly should it be?
- ▶ convergence of functions — what properties are preserved?

New doubts in the early 19th century

Fourier's work converged with more philosophical investigation to stimulate questions concerning:

- ▶ functions — what exactly should they be?
- ▶ convergence — what exactly should it be?
- ▶ convergence of functions — what properties are preserved?
- ▶ integration — what exactly should it be?

New doubts in the early 19th century

Fourier's work converged with more philosophical investigation to stimulate questions concerning:

- ▶ functions — what exactly should they be?
- ▶ convergence — what exactly should it be?
- ▶ convergence of functions — what properties are preserved?
- ▶ integration — what exactly should it be?
- ▶ existence of limits — what are the essential properties of real numbers? [Lecture XII]

Recall from Lecture VII: Cauchy sequences, 1821

Augustin-Louis Cauchy, *Cours d'analyse* (1821), Ch. VI, pp. 124, 125:

In order for the series u_0, u_1, u_2, \dots [that is, $\sum u_i$] to be convergent ... it is necessary and sufficient that the partial sums

$$s_n = u_0 + u_1 + u_2 + \&c. \dots + u_{n-1}$$

converge to a fixed limit s : in other words, it is necessary and sufficient that for infinitely large values of the number n , the sums

$$s_n, s_{n+1}, s_{n+2}, \&c. \dots$$

differ from the limit s , and consequently from each other, by infinitely small quantities.

Cauchy and continuity revisited

In *Cours d'analyse*, p. 34, Cauchy defined a function f to be **continuous** between certain limits if, for each x between those limits, the value of $f(x)$ is unique and finite, and $|f(x + \alpha) - f(x)|$, where α is indefinitely small, decreases indefinitely with α .

Cauchy and continuity revisited

In *Cours d'analyse*, p. 34, Cauchy defined a function f to be **continuous** between certain limits if, for each x between those limits, the value of $f(x)$ is unique and finite, and $|f(x + \alpha) - f(x)|$, where α is indefinitely small, decreases indefinitely with α .

In other words (p. 35): for x between the given limits, an infinitely small increase in x produces an infinitely small increase in $f(x)$.

Cauchy and continuity revisited

In *Cours d'analyse*, p. 34, Cauchy defined a function f to be **continuous** between certain limits if, for each x between those limits, the value of $f(x)$ is unique and finite, and $|f(x + \alpha) - f(x)|$, where α is indefinitely small, decreases indefinitely with α .

In other words (p. 35): for x between the given limits, an infinitely small increase in x produces an infinitely small increase in $f(x)$.

So Cauchy defined continuity **on an interval**, rather than at a point.

Cauchy and continuity revisited

In *Cours d'analyse*, p. 34, Cauchy defined a function f to be **continuous** between certain limits if, for each x between those limits, the value of $f(x)$ is unique and finite, and $|f(x + \alpha) - f(x)|$, where α is indefinitely small, decreases indefinitely with α .

In other words (p. 35): for x between the given limits, an infinitely small increase in x produces an infinitely small increase in $f(x)$.

So Cauchy defined continuity **on an interval**, rather than at a point.

He went on to derive basic results concerning continuous functions: that the composition of two continuous functions is continuous, the Intermediate Value Theorem, etc.

A theorem of Cauchy (1821)

Cauchy, *Cours d'analyse*, pp. 131–132:

When the various terms of a series are functions of a variable x , continuous with respect to this variable in the neighbourhood of a particular value for which the series is convergent, the sum s of the series is also, in the neighbourhood of this value, a continuous function of x .

A theorem of Cauchy (1821)

Cauchy, *Cours d'analyse*, pp. 131–132:

When the various terms of a series are functions of a variable x , continuous with respect to this variable in the neighbourhood of a particular value for which the series is convergent, the sum s of the series is also, in the neighbourhood of this value, a continuous function of x .

In other words: a convergent series of continuous functions converges to a continuous function.

A theorem of Cauchy (1821)

Cauchy, *Cours d'analyse*, pp. 131–132:

When the various terms of a series are functions of a variable x , continuous with respect to this variable in the neighbourhood of a particular value for which the series is convergent, the sum s of the series is also, in the neighbourhood of this value, a continuous function of x .

In other words: a convergent series of continuous functions converges to a continuous function.

Not true!

Cauchy's argument

Cauchy considered a sequence of continuous functions $u_0(x), u_1(x), u_2(x), \dots$ on a given interval.

Cauchy's argument

Cauchy considered a sequence of continuous functions $u_0(x), u_1(x), u_2(x), \dots$ on a given interval. He supposed that the corresponding series converges to a function $s(x)$.

Cauchy's argument

Cauchy considered a sequence of continuous functions $u_0(x), u_1(x), u_2(x), \dots$ on a given interval. He supposed that the corresponding series converges to a function $s(x)$. Partial sums are denoted by $s_n(x) = \sum_{j=0}^{n-1} u_j(x)$.

Cauchy's argument

Cauchy considered a sequence of continuous functions $u_0(x), u_1(x), u_2(x), \dots$ on a given interval. He supposed that the corresponding series converges to a function $s(x)$. Partial sums are denoted by $s_n(x) = \sum_{j=0}^{n-1} u_j(x)$. The n th remainder term $r_n(x)$ is defined by $s(x) = s_n(x) + r_n(x)$.

Cauchy's argument

Cauchy considered a sequence of continuous functions $u_0(x), u_1(x), u_2(x), \dots$ on a given interval. He supposed that the corresponding series converges to a function $s(x)$. Partial sums are denoted by $s_n(x) = \sum_{j=0}^{n-1} u_j(x)$. The n th remainder term $r_n(x)$ is defined by $s(x) = s_n(x) + r_n(x)$.

Cauchy noted that each s_n is evidently continuous for values of x in the given interval.

Cauchy's argument

Cauchy considered a sequence of continuous functions $u_0(x), u_1(x), u_2(x), \dots$ on a given interval. He supposed that the corresponding series converges to a function $s(x)$. Partial sums are denoted by $s_n(x) = \sum_{j=0}^{n-1} u_j(x)$. The n th remainder term $r_n(x)$ is defined by $s(x) = s_n(x) + r_n(x)$.

Cauchy noted that each s_n is evidently continuous for values of x in the given interval. Suppose that we increase x by an infinitely small quantity α .

Cauchy's argument

Cauchy considered a sequence of continuous functions $u_0(x), u_1(x), u_2(x), \dots$ on a given interval. He supposed that the corresponding series converges to a function $s(x)$. Partial sums are denoted by $s_n(x) = \sum_{j=0}^{n-1} u_j(x)$. The n th remainder term $r_n(x)$ is defined by $s(x) = s_n(x) + r_n(x)$.

Cauchy noted that each s_n is evidently continuous for values of x in the given interval. Suppose that we increase x by an infinitely small quantity α . For all values of n , the corresponding increase in $s_n(x)$ will also be infinitely small.

Cauchy's argument

Cauchy considered a sequence of continuous functions $u_0(x), u_1(x), u_2(x), \dots$ on a given interval. He supposed that the corresponding series converges to a function $s(x)$. Partial sums are denoted by $s_n(x) = \sum_{j=0}^{n-1} u_j(x)$. The n th remainder term $r_n(x)$ is defined by $s(x) = s_n(x) + r_n(x)$.

Cauchy noted that each s_n is evidently continuous for values of x in the given interval. Suppose that we increase x by an infinitely small quantity α . For all values of n , the corresponding increase in $s_n(x)$ will also be infinitely small. For n very large ('très-considérable'), the increase in $r_n(x)$ becomes 'insensible'.

Cauchy's argument

Cauchy considered a sequence of continuous functions $u_0(x), u_1(x), u_2(x), \dots$ on a given interval. He supposed that the corresponding series converges to a function $s(x)$. Partial sums are denoted by $s_n(x) = \sum_{j=0}^{n-1} u_j(x)$. The n th remainder term $r_n(x)$ is defined by $s(x) = s_n(x) + r_n(x)$.

Cauchy noted that each s_n is evidently continuous for values of x in the given interval. Suppose that we increase x by an infinitely small quantity α . For all values of n , the corresponding increase in $s_n(x)$ will also be infinitely small. For n very large ('très-considérable'), the increase in $r_n(x)$ becomes 'insensible'. Therefore, the increase in $s(x)$ can only be an infinitely small quantity.

NB. All notation except ' \sum ' is Cauchy's.

Cauchy's argument

est convergente, la somme de cette série est représentée par

$$u_0 + u_1 + u_2 + u_3 + \&c. \dots$$

En vertu de cette convention, la valeur du nombre ϵ se trouvera déterminée par l'équation

$$(6) \quad \epsilon = 1 + \frac{1}{1} + \frac{1}{1,2} + \frac{1}{1,2,3} + \frac{1}{1,2,3,4} + \&c. \dots;$$

et, si l'on considère la progression géométrique

$$1, x, x^2, x^3, \&c. \dots,$$

on aura, pour des valeurs numériques de x inférieures à l'unité,

$$(7) \quad 1 + x + x^2 + x^3 + \&c. \dots = \frac{1}{1-x}.$$

La série

$$u_0, u_1, u_2, u_3, \&c. \dots$$

étant supposée convergente, si l'on désigne sa somme par s , et par s_n la somme de ses n premiers termes, on trouvera

$$\begin{aligned} s &= u_0 + u_1 + u_2 + \dots + u_{n-1} + u_n + u_{n+1} + \&c. \dots \\ &= s_n + u_n + u_{n+1} + \&c. \dots, \end{aligned}$$

et par suite

$$s - s_n = u_n + u_{n+1} + \&c. \dots$$

De cette dernière équation il résulte que les quantités

$$u_n, u_{n+1}, u_{n+2}, \&c. \dots$$

formeront une nouvelle série convergente dont la somme sera équivalente à $s - s_n$. Si l'on représente cette même somme par r_n , on aura

$$s = s_n + r_n;$$

et r_n sera ce qu'on appelle le *reste* de la série (1) à partir du $n.$ ^{me} terme.

Lorsque, les termes de la série (1) renfermant une même variable x , cette série est convergente, et ses différens termes fonctions continues de x , dans le voisinage d'une valeur particulière attribuée à cette variable ;

$$s_n, r_n \text{ et } s$$

sont encore trois fonctions de la variable x , dont la première est évidemment continue par rapport à x dans le voisinage de la valeur particulière dont il s'agit. Cela posé, considérons les accroissemens que reçoivent ces trois fonctions, lorsqu'on fait croître x d'une quantité infiniment petite α . L'accroissement de s_n sera, pour toutes les valeurs possibles de n , une quantité infiniment petite; et celui de r_n deviendra insensible en même temps que r_n , si l'on attribue à n une valeur très-considérable. Par suite, l'accroissement de la fonction s ne pourra être qu'une quantité infiniment petite. De cette remarque on déduit immédiatement la proposition suivante.

1.^{er} THÉORÈME. *Lorsque les différens termes de la série (1) sont des fonctions d'une même variable x ,*

A modern counterexample

For each $n \in \mathbb{N}$, define continuous functions f_n by

$$f_n(x) = \begin{cases} -1 & \text{if } x \leq -\frac{1}{n}; \\ nx & \text{if } -\frac{1}{n} \leq x \leq \frac{1}{n}; \\ +1 & \text{if } x \geq \frac{1}{n}. \end{cases}$$

A modern counterexample

For each $n \in \mathbb{N}$, define continuous functions f_n by

$$f_n(x) = \begin{cases} -1 & \text{if } x \leq -\frac{1}{n}; \\ nx & \text{if } -\frac{1}{n} \leq x \leq \frac{1}{n}; \\ +1 & \text{if } x \geq \frac{1}{n}. \end{cases}$$

Now set $u_1(x) = f_1(x)$, and define new functions u_n recursively by

$$u_n(x) = f_n(x) - f_{n-1}(x).$$

A modern counterexample

For each $n \in \mathbb{N}$, define continuous functions f_n by

$$f_n(x) = \begin{cases} -1 & \text{if } x \leq -\frac{1}{n}; \\ nx & \text{if } -\frac{1}{n} \leq x \leq \frac{1}{n}; \\ +1 & \text{if } x \geq \frac{1}{n}. \end{cases}$$

Now set $u_1(x) = f_1(x)$, and define new functions u_n recursively by

$$u_n(x) = f_n(x) - f_{n-1}(x).$$

Notice then that

$$s_n(x) = \sum_{j=1}^n u_j(x) = f_n(x).$$

A modern counterexample

For each $n \in \mathbb{N}$, define continuous functions f_n by

$$f_n(x) = \begin{cases} -1 & \text{if } x \leq -\frac{1}{n}; \\ nx & \text{if } -\frac{1}{n} \leq x \leq \frac{1}{n}; \\ +1 & \text{if } x \geq \frac{1}{n}. \end{cases}$$

Now set $u_1(x) = f_1(x)$, and define new functions u_n recursively by

$$u_n(x) = f_n(x) - f_{n-1}(x).$$

Notice then that

$$s_n(x) = \sum_{j=1}^n u_j(x) = f_n(x).$$

But we see that $s_n \rightarrow s$ as $n \rightarrow \infty$, where

$$s(x) = \begin{cases} -1 & \text{if } x < 0; \\ 0 & \text{if } x = 0; \\ +1 & \text{if } x > 0, \end{cases}$$

which is discontinuous at $x = 0$.

A modern counterexample

What happens to the remainders $r_n(x) = s(x) - s_n(x)$?

A modern counterexample

What happens to the remainders $r_n(x) = s(x) - s_n(x)$?

Outside the range $-\frac{1}{n} \leq x \leq \frac{1}{n}$, $r_n(x) = 0$,

A modern counterexample

What happens to the remainders $r_n(x) = s(x) - s_n(x)$?

Outside the range $-\frac{1}{n} \leq x \leq \frac{1}{n}$, $r_n(x) = 0$, but inside:

$$r_n(x) = \begin{cases} -1 - nx & \text{if } -\frac{1}{n} \leq x < 0; \\ 0 & \text{if } x = 0; \\ 1 - nx & \text{if } 0 < x \leq \frac{1}{n}. \end{cases}$$

A modern counterexample

What happens to the remainders $r_n(x) = s(x) - s_n(x)$?

Outside the range $-\frac{1}{n} \leq x \leq \frac{1}{n}$, $r_n(x) = 0$, but inside:

$$r_n(x) = \begin{cases} -1 - nx & \text{if } -\frac{1}{n} \leq x < 0; \\ 0 & \text{if } x = 0; \\ 1 - nx & \text{if } 0 < x \leq \frac{1}{n}. \end{cases}$$

For each x , $r_n(x) \rightarrow 0$ as $n \rightarrow \infty$, but this does not happen **simultaneously** for all values of x .

Cauchy's remainders

Cauchy: For n very large, the increase in $r_n(x)$ becomes 'insensible'.

Cauchy's remainders

Cauchy: For n very large, the increase in $r_n(x)$ becomes 'insensible'. But what does this mean?

Cauchy's remainders

Cauchy: For n very large, the increase in $r_n(x)$ becomes 'insensible'. But what does this mean?

One of the following modern statements? (Denoting Cauchy's interval by I .)

Cauchy's remainders

Cauchy: For n very large, the increase in $r_n(x)$ becomes 'insensible'. But what does this mean?

One of the following modern statements? (Denoting Cauchy's interval by I .)

$$\forall \varepsilon > 0 : \exists N : \forall x \in I : n > N \Rightarrow |r_n(x)| < \varepsilon$$

Cauchy's remainders

Cauchy: For n very large, the increase in $r_n(x)$ becomes 'insensible'. But what does this mean?

One of the following modern statements? (Denoting Cauchy's interval by I .)

$$\forall \varepsilon > 0 : \exists N : \forall x \in I : n > N \Rightarrow |r_n(x)| < \varepsilon$$

$$\forall \varepsilon > 0 : \forall x \in I : \exists N : n > N \Rightarrow |r_n(x)| < \varepsilon$$

Cauchy's remainders

Cauchy: For n very large, the increase in $r_n(x)$ becomes 'insensible'. But what does this mean?

One of the following modern statements? (Denoting Cauchy's interval by I .)

$$\forall \varepsilon > 0 : \exists N : \forall x \in I : n > N \Rightarrow |r_n(x)| < \varepsilon$$

$$\forall \varepsilon > 0 : \forall x \in I : \exists N : n > N \Rightarrow |r_n(x)| < \varepsilon$$

The second is true for our modern counterexample, but the first is not

Cauchy's remainders

Cauchy: For n very large, the increase in $r_n(x)$ becomes 'insensible'. But what does this mean?

One of the following modern statements? (Denoting Cauchy's interval by I .)

$$\forall \varepsilon > 0 : \exists N : \forall x \in I : n > N \Rightarrow |r_n(x)| < \varepsilon$$

$$\forall \varepsilon > 0 : \forall x \in I : \exists N : n > N \Rightarrow |r_n(x)| < \varepsilon$$

The second is true for our modern counterexample, but the first is not — so there really is a distinction between the two.

Cauchy's remainders

Cauchy: For n very large, the increase in $r_n(x)$ becomes 'insensible'. But what does this mean?

One of the following modern statements? (Denoting Cauchy's interval by I .)

$$\forall \varepsilon > 0 : \exists N : \forall x \in I : n > N \Rightarrow |r_n(x)| < \varepsilon$$

$$\forall \varepsilon > 0 : \forall x \in I : \exists N : n > N \Rightarrow |r_n(x)| < \varepsilon$$

The second is true for our modern counterexample, but the first is not — so there really is a distinction between the two.

Cauchy clearly didn't make this distinction

Cauchy's remainders

Cauchy: For n very large, the increase in $r_n(x)$ becomes 'insensible'. But what does this mean?

One of the following modern statements? (Denoting Cauchy's interval by I .)

$$\forall \varepsilon > 0 : \exists N : \forall x \in I : n > N \Rightarrow |r_n(x)| < \varepsilon$$

$$\forall \varepsilon > 0 : \forall x \in I : \exists N : n > N \Rightarrow |r_n(x)| < \varepsilon$$

The second is true for our modern counterexample, but the first is not — so there really is a distinction between the two.

Cauchy clearly didn't make this distinction — but should this really be regarded as a 'mistake'?

Reactions to Cauchy's 'mistake'

Abel (Crelle's Journal, 1826) on the *Cours d'analyse*:
the excellent work of M. Cauchy . . . which must be read by every analyst who aims at rigour in mathematical research

Reactions to Cauchy's 'mistake'

Abel (Crelle's Journal, 1826) on the *Cours d'analyse*:
the excellent work of M. Cauchy . . . which must be read by every analyst who aims at rigour in mathematical research

Four pages later, on Cauchy's 'theorem' on sums of continuous functions:


it seems to me that the theorem admits exceptions. For example, the series

$$\sin x - \frac{1}{2} \sin 2x + \frac{1}{3} \sin 3x - \dots$$

is discontinuous for every value $(2m + 1)\pi$ of x , m being a whole number. There are, as one knows, many series of this kind.

Abel's counterexample

$$\sin x - \frac{1}{2} \sin 2x + \frac{1}{3} \sin 3x - \dots$$



Abel's counterexample elsewhere

Abel to Holmboe, January 1826:

One applies all operations to infinite series as if they were finite, but is this permissible? I think not. — Where is it proved that one gets the differential of an infinite series by differentiating each term? It is easy to give an example for which this is not true, e.g.

$$\frac{1}{2}x = \sin x - \frac{1}{2} \sin 2x + \frac{1}{3} \sin 3x - \dots$$

Differentiation gives

$$\frac{1}{2} = \cos x - \cos 2x + \cos 3x - \dots \text{etc.}$$

a result which is quite false because this series is divergent.

Can theorems have exceptions?

Abel was not the first to study this example:

Can theorems have exceptions?

Abel was not the first to study this example:

- ▶ Euler had discussed it in 1783;

Can theorems have exceptions?

Abel was not the first to study this example:

- ▶ Euler had discussed it in 1783;
- ▶ Lacroix had included it in his *Traité du Calcul Différentiel et du Calcul Intégral* of 1810;

Can theorems have exceptions?

Abel was not the first to study this example:

- ▶ Euler had discussed it in 1783;
- ▶ Lacroix had included it in his *Traité du Calcul Différentiel et du Calcul Intégral* of 1810;
- ▶ it had appeared in Fourier's work — which is probably where Abel found it.

Can theorems have exceptions?

Abel was not the first to study this example:

- ▶ Euler had discussed it in 1783;
- ▶ Lacroix had included it in his *Traité du Calcul Différentiel et du Calcul Intégral* of 1810;
- ▶ it had appeared in Fourier's work — which is probably where Abel found it.

Had Cauchy seen it?

Can theorems have exceptions?

Abel was not the first to study this example:

- ▶ Euler had discussed it in 1783;
- ▶ Lacroix had included it in his *Traité du Calcul Différentiel et du Calcul Intégral* of 1810;
- ▶ it had appeared in Fourier's work — which is probably where Abel found it.

Had Cauchy seen it?

Abel: Cauchy's argument is essentially correct, only failing in certain anomalous situations.

Can theorems have exceptions?

Abel was not the first to study this example:

- ▶ Euler had discussed it in 1783;
- ▶ Lacroix had included it in his *Traité du Calcul Différentiel et du Calcul Intégral* of 1810;
- ▶ it had appeared in Fourier's work — which is probably where Abel found it.

Had Cauchy seen it?

Abel: Cauchy's argument is essentially correct, only failing in certain anomalous situations.

Cauchy: no (immediate) reaction

Can theorems have exceptions?

Abel was not the first to study this example:

- ▶ Euler had discussed it in 1783;
- ▶ Lacroix had included it in his *Traité du Calcul Différentiel et du Calcul Intégral* of 1810;
- ▶ it had appeared in Fourier's work — which is probably where Abel found it.

Had Cauchy seen it?

Abel: Cauchy's argument is essentially correct, only failing in certain anomalous situations.

Cauchy: no (immediate) reaction — so deemed the example irrelevant?

Can theorems have exceptions?

Abel was not the first to study this example:

- ▶ Euler had discussed it in 1783;
- ▶ Lacroix had included it in his *Traité du Calcul Différentiel et du Calcul Intégral* of 1810;
- ▶ it had appeared in Fourier's work — which is probably where Abel found it.

Had Cauchy seen it?

Abel: Cauchy's argument is essentially correct, only failing in certain anomalous situations.

Cauchy: no (immediate) reaction — so deemed the example irrelevant?

(See: [Henrik Kragh Sørensen, Exceptions and counterexamples: Understanding Abel's comment on *Cauchy's Theorem*, *Historia Mathematica* **32** \(2005\) 453–480](#))

Abel's interpretation

Abel read Cauchy as: if the series is convergent at a point x_0 and the individual terms of the series are continuous on a neighbourhood of x_0 , then the series is also continuous on that neighbourhood.

Abel's interpretation

Abel read Cauchy as: if the series is convergent at a point x_0 and the individual terms of the series are continuous on a neighbourhood of x_0 , then the series is also continuous on that neighbourhood.

But what did Cauchy mean by 'continuity' and 'convergence'?

Abel's interpretation

Abel read Cauchy as: if the series is convergent at a point x_0 and the individual terms of the series are continuous on a neighbourhood of x_0 , then the series is also continuous on that neighbourhood.

But what did Cauchy mean by 'continuity' and 'convergence'?

Note that he only ever defined continuity on an **interval**

Abel's interpretation

Abel read Cauchy as: if the series is convergent at a point x_0 and the individual terms of the series are continuous on a neighbourhood of x_0 , then the series is also continuous on that neighbourhood.

But what did Cauchy mean by 'continuity' and 'convergence'?

Note that he only ever defined continuity on an **interval** — so was it in fact **uniform continuity**?

Abel's interpretation

Abel read Cauchy as: if the series is convergent at a point x_0 and the individual terms of the series are continuous on a neighbourhood of x_0 , then the series is also continuous on that neighbourhood.

But what did Cauchy mean by 'continuity' and 'convergence'?

Note that he only ever defined continuity on an **interval** — so was it in fact **uniform continuity**? If, in addition, we regard his notion of convergence on an interval as being **uniform convergence**, then the theorem holds in all cases.

Abel's interpretation

Abel read Cauchy as: if the series is convergent at a point x_0 and the individual terms of the series are continuous on a neighbourhood of x_0 , then the series is also continuous on that neighbourhood.

But what did Cauchy mean by 'continuity' and 'convergence'?

Note that he only ever defined continuity on an **interval** — so was it in fact **uniform continuity**? If, in addition, we regard his notion of convergence on an interval as being **uniform convergence**, then the theorem holds in all cases.

Similarly, what did Abel mean by 'continuity' and 'convergence'? The same as Cauchy? Or did he use a similar form of words but with a different meaning?

Conditions for Cauchy's theorem to work

Dirichlet (1829): Fourier series can represent discontinuous functions.

Conditions for Cauchy's theorem to work

Dirichlet (1829): Fourier series can represent discontinuous functions.

But the reason why Cauchy's theorem fails (if indeed it does) remained unclear.

Conditions for Cauchy's theorem to work

Dirichlet (1829): Fourier series can represent discontinuous functions.

But the reason why Cauchy's theorem fails (if indeed it does) remained unclear.

The need for **uniform convergence** was gradually recognised:

Conditions for Cauchy's theorem to work

Dirichlet (1829): Fourier series can represent discontinuous functions.

But the reason why Cauchy's theorem fails (if indeed it does) remained unclear.

The need for **uniform convergence** was gradually recognised:

- ▶ Karl Weierstrass (lectures in Berlin), 1841;

Conditions for Cauchy's theorem to work

Dirichlet (1829): Fourier series can represent discontinuous functions.

But the reason why Cauchy's theorem fails (if indeed it does) remained unclear.

The need for **uniform convergence** was gradually recognised:

- ▶ Karl Weierstrass (lectures in Berlin), 1841;
- ▶ Emmanuel Björling (Uppsala), 1846 (or not?);

Conditions for Cauchy's theorem to work

Dirichlet (1829): Fourier series can represent discontinuous functions.

But the reason why Cauchy's theorem fails (if indeed it does) remained unclear.

The need for **uniform convergence** was gradually recognised:

- ▶ Karl Weierstrass (lectures in Berlin), 1841;
- ▶ Emmanuel Björling (Uppsala), 1846 (or not?);
- ▶ Gabriel Stokes (Cambridge), 1847;

Conditions for Cauchy's theorem to work

Dirichlet (1829): Fourier series can represent discontinuous functions.

But the reason why Cauchy's theorem fails (if indeed it does) remained unclear.

The need for **uniform convergence** was gradually recognised:

- ▶ Karl Weierstrass (lectures in Berlin), 1841;
- ▶ Emmanuel Björling (Uppsala), 1846 (or not?);
- ▶ Gabriel Stokes (Cambridge), 1847;
- ▶ Phillip Seidel (Berlin), 1848.

Conditions for Cauchy's theorem to work

Dirichlet (1829): Fourier series can represent discontinuous functions.

But the reason why Cauchy's theorem fails (if indeed it does) remained unclear.

The need for **uniform convergence** was gradually recognised:

- ▶ Karl Weierstrass (lectures in Berlin), 1841;
- ▶ Emmanuel Björling (Uppsala), 1846 (or not?);
- ▶ Gabriel Stokes (Cambridge), 1847;
- ▶ Phillip Seidel (Berlin), 1848.

See: G. H. Hardy, 'Sir George Stokes and the concept of uniform convergence', *Proc. Camb. Phil. Soc.* **19** (1918) 148–156 (also: *Collected Papers of G. H. Hardy*, vol. VII, 505–513)

Cauchy revisits his theorem (1853)

it is easy to see how one can modify the statement of the theorem so that it will no longer have any exception. This is what I am going to explain in a few words.

Cauchy revisits his theorem (1853)

it is easy to see how one can modify the statement of the theorem so that it will no longer have any exception. This is what I am going to explain in a few words.

Theorem. If the different terms of the series

$$u_0, u_1, u_2, \dots, u_n, u_{n+1}, \dots$$

are functions of a real variable x , continuous with respect to this variable within the given limits; and if, in addition, the sum

$$u_n + u_{n+1} + \dots + u_{n'}$$

always becomes infinitely small for infinitely large values of the whole numbers n and $n' > n$, then the series will be convergent and the sum of the series will be, within the given limits, a continuous function of the variable x .

Cauchy revisits his theorem (1853)

it is easy to see how one can modify the statement of the theorem so that it will no longer have any exception. This is what I am going to explain in a few words.

Theorem. If the different terms of the series

$$u_0, u_1, u_2, \dots, u_n, u_{n+1}, \dots$$

are functions of a real variable x , continuous with respect to this variable within the given limits; and if, in addition, the sum

$$u_n + u_{n+1} + \dots + u_{n'}$$

always becomes infinitely small for infinitely large values of the whole numbers n and $n' > n$, then the series will be convergent and the sum of the series will be, within the given limits, a continuous function of the variable x .

But it was becoming clear that the language of infinities and infinitesimals was inadequate for expressing the problems at hand.

Part 2: Integration

Integration


Integration

- ▶ Recall that in the 17th century, 'integration' was designed for 'quadrature', for measuring space or calculating area.

Integration


- ▶ Recall that in the 17th century, 'integration' was designed for 'quadrature', for measuring space or calculating area.
- ▶ In the 18th century, 'integration' was essentially regarded as the inverse of differentiation.

Integration in the 18th century (1)



Leonhard Euler, *Foundations of
integral calculus* (1768):

Integration in the 18th century (1)




Leonhard Euler, *Foundations of integral calculus* (1768):

Definition 1: Integral calculus is the method of finding, from a given relationship between differentials, a relationship between the quantities themselves: and the operation by which this is carried out is usually called integration.


(See *Mathematics emerging*, §14.2.1.)

Integration in the 18th century (2)



Corollary 1: Therefore where differential calculus teaches us to investigate the relationship between differentials from a given relationship between variable quantities, integral calculus supplies the inverse method.

Integration in the 18th century (2)



Corollary 1: Therefore where differential calculus teaches us to investigate the relationship between differentials from a given relationship between variable quantities, integral calculus supplies the inverse method.

Corollary 2: Clearly just as in Analysis two operations are always contrary to each other, as subtraction to addition, division to multiplication, extraction of roots to raising of powers, so also by similar reasoning integral calculus is contrary to differential calculus.

Integration in the 18th century (3)

4 DE CALCULO INTEGRALI

ratione conscripti prodierint, huiusmodi conciliatio nullum usum esset habitura.

Definitio 2.

7. Cum functionis cuiuscunque ipsius x differentiale huiusmodi habeat formam Xdx , proposita tali forma differentiali Xdx , in qua X sit functio quaecunque ipsius x , illa functio, cuius differentiale est $=Xdx$, huius vocatur integrale, et praefixo signo \int indicari solet, ita ut $\int Xdx$ eam denotet quantitatem variabilem, cuius differentiale est $=Xdx$.

Coroll. 1.

8. Quemadmodum ergo propositae formulae differentialis Xdx integrale, seu ea functio ipsius x , cuius differentiale est $=Xdx$, quae hac scriptura $\int Xdx$ indicatur, inuestigari debeat, in calculo integrali est explicandum.

Coroll. 2.

9. Vti ergo littera d signum est differentiationis, ita littera \int pro signo integrationis utimur, sicque haec duo signa sibi mutuo opponuntur, et quasi se destruunt scilicet $\int dX$ erit $=X$, quia ea quantitas denotatur cuius differentiale est dX , quae utique est X .

Coroll. 3.

10. Cum igitur harum ipsius x functionum x^a , x^b , $\sqrt{(aa-xx)}$ differentialia sint $a x^{a-1} dx$, $b x^{b-1} dx$, $\frac{-x dx}{\sqrt{(a^2-xx)}}$ signo integrationis \int adhibendo patet fore

$\int 2x$

Definition 2: Since the differentiation of any function of x has a form of this kind: $X dx$, when such a differential form $X dx$ is proposed, in which X is any function of x , that function whose differential $= X dx$ is called its integral, and is usually indicated by the prefix \int , so that $\int X dx$ denotes that variable quantity whose differential $= X dx$.

Integration in the 18th century (3)

4 DE CALCULO INTEGRALI

ratione conscripti prodierint, huiusmodi conciliatio nullum usum esset habitura.

Definitio 2.

7. Cum functionis cuiuscunque ipsius x differentiale huiusmodi habeat formam Xdx , proposita tali forma differentiali Xdx , in qua X sit functio quaecunque ipsius x , illa functio, cuius differentiale est $=Xdx$, huius vocatur integrale, et praefixo signo \int indicari solet, ita ut $\int Xdx$ eam denotet quantitatem variabilem, cuius differentiale est $=Xdx$.

Coroll. 1.

8. Quemadmodum ergo propositae formulae differentialis Xdx integrale, seu ea functio ipsius x , cuius differentiale est $=Xdx$, quae hac scriptura $\int Xdx$ indicatur, inuestigari debeat, in calculo integrali est explicandum.

Coroll. 2.

9. Vt ergo littera d signum est differentiationis, ita littera \int pro signo integrationis utimur, sicque haec duo signa sibi mutuo opponuntur, et quasi se destruant scilicet $\int dX$ erit $=X$, quia ea quantitas denotatur cuius differentiale est dX , quae utique est X .

Coroll. 3.

10. Cum igitur harum ipsius x functionum x^a , x^b , $\sqrt{(aa-xx)}$ differentialia sint $axdx$, $bx^{b-1}dx$, $\frac{-x dx}{\sqrt{(aa-xx)}}$ signo integrationis \int adhibendo patet fore $\int ax = \frac{a}{2} x^2$

Definition 2: Since the differentiation of any function of x has a form of this kind: $X dx$, when such a differential form $X dx$ is proposed, in which X is any function of x , that function whose differential $= X dx$ is called its integral, and is usually indicated by the prefix \int , so that $\int X dx$ denotes that variable quantity whose differential $= X dx$.

Corollary 2: Therefore just as the letter d is the sign of differentiation, so we use the letter \int as the sign of integration, and thus these two signs are mutually contrary to each other, as though they destroy each other: certainly $\int dX = X, \dots$

Coroll. 3.

10. Cum igitur harum ipsius x functionum
 x^2 , x^n , $\sqrt{aa-xx}$ differentialia sint $2x dx$, $nx^{n-1} dx$,
 $\frac{-x dx}{\sqrt{aa-xx}}$ signo integrationis \int adhibendo patet fore
 $\int 2x dx = xx$; $\int nx^{n-1} dx = x^n$; $\int \frac{-x dx}{\sqrt{aa-xx}} = \sqrt{aa-xx}$
vnde vsus huius signi clarius perspicitur.

Some 19th-century ideas

Recall that Fourier coefficients are given by $\frac{2}{\pi} \int_0^\pi \phi(x) \sin nx \, dx$.

Some 19th-century ideas

Recall that Fourier coefficients are given by $\frac{2}{\pi} \int_0^\pi \phi(x) \sin nx \, dx$.

It is not always possible to solve such an integral algebraically.

Some 19th-century ideas

Recall that Fourier coefficients are given by $\frac{2}{\pi} \int_0^\pi \phi(x) \sin nx \, dx$.

It is not always possible to solve such an integral algebraically.

Fourier (1822): but we can draw the curve of $\phi(x)$, and hence that of $\phi(x) \sin nx$, under which there is clearly an **area**.

Some 19th-century ideas

Recall that Fourier coefficients are given by $\frac{2}{\pi} \int_0^\pi \phi(x) \sin nx \, dx$.

It is not always possible to solve such an integral algebraically.

Fourier (1822): but we can draw the curve of $\phi(x)$, and hence that of $\phi(x) \sin nx$, under which there is clearly an **area**.

Fourier thus returned to the idea of integral as area and influenced Cauchy almost immediately...

A theory of definite integrals (1823)

Cauchy's *Résumé*, 1823, Lesson 21:

A theory of definite integrals (1823)

Cauchy's *Résumé*, 1823, Lesson 21:

Suppose $f(x)$ continuous between $x = x_0$ and $x = X$. Choose x_1, x_2, \dots, x_{n-1} between these limits. Define

$$S = (x_1 - x_0)f(x_0) + (x_2 - x_1)f(x_1) + \cdots + (X - x_{n-1})f(x_{n-1})$$

A theory of definite integrals (1823)

Cauchy's *Résumé*, 1823, Lesson 21:

Suppose $f(x)$ continuous between $x = x_0$ and $x = X$. Choose x_1, x_2, \dots, x_{n-1} between these limits. Define

$$S = (x_1 - x_0)f(x_0) + (x_2 - x_1)f(x_1) + \cdots + (X - x_{n-1})f(x_{n-1})$$

[much discussion of dependence on partition followed by]

A theory of definite integrals (1823)

Cauchy's *Résumé*, 1823, Lesson 21:

Suppose $f(x)$ continuous between $x = x_0$ and $x = X$. Choose x_1, x_2, \dots, x_{n-1} between these limits. Define

$$S = (x_1 - x_0)f(x_0) + (x_2 - x_1)f(x_1) + \dots + (X - x_{n-1})f(x_{n-1})$$

[much discussion of dependence on partition followed by]

If the numerical values of the elements are made to decrease indefinitely by increasing their number, the value of S will become essentially constant, or in other words, it will finish by attaining a certain limit which will depend only on the form of the function $f(x)$ and the boundary values $x = x_0, x = X$ given to the variable x . This limit is what one calls a definite integral.

A theory of definite integrals (1823)

Cauchy's *Résumé*, 1823, Lesson 21:

Suppose $f(x)$ continuous between $x = x_0$ and $x = X$. Choose x_1, x_2, \dots, x_{n-1} between these limits. Define

$$S = (x_1 - x_0)f(x_0) + (x_2 - x_1)f(x_1) + \dots + (X - x_{n-1})f(x_{n-1})$$

[much discussion of dependence on partition followed by]

If the numerical values of the elements are made to decrease indefinitely by increasing their number, the value of S will become essentially constant, or in other words, it will finish by attaining a certain limit which will depend only on the form of the function $f(x)$ and the boundary values $x = x_0, x = X$ given to the variable x . This limit is what one calls a definite integral.

[further issues connected with uniform convergence]

Cauchy and integrals

84

COURS D'ANALYSE.

Observons maintenant que, si l'on désigne par $\Delta x = h = dx$ un accroissement fini attribué à la variable x , les différens termes dont se compose la valeur de S , tels que les produits $(x, -x_0)f(x_0)$, $(x, -x_1)f(x_1)$, &c... seront tous compris dans la formule générale

$$(8) \quad hf(x) = f(x) dx$$

de laquelle on les déduira l'un après l'autre, en posant d'abord $x = x_0$, et $h = x_1 - x_0$, puis $x = x_1$, et $h = x_2 - x_1$, &c... On peut donc énoncer que la quantité S est une somme de produits semblables à l'expression (8); ce qu'on exprime quelquefois à l'aide de la caractéristique Σ en écrivant

$$(9) \quad S = \Sigma hf(x) = \Sigma f(x) \Delta x.$$

Quant à l'intégrale définie vers laquelle converge la quantité S , tandis que les élémens de la différence $X - x_0$ deviennent infiniment petits, on est convenu de la représenter par la notation $\int hf(x)$ ou $\int f(x) dx$, dans laquelle la lettre f substituée à la lettre Σ indique, non plus une somme de produits semblables à l'expression (8), mais la limite d'une somme de cette espèce. De plus, comme la valeur de l'intégrale définie que l'on considère dépend des valeurs extrêmes x_0, X attribuées à la variable x , on est convenu de placer ces deux valeurs, la première au-dessous, la seconde au-dessus de la lettre f , ou de les écrire à côté de l'intégrale, que l'on désigne en conséquence par l'une des notations

$$(10) \quad \int_{x_0}^X f(x) dx, \quad \int_{x_0}^X f(x) dx \left[\begin{matrix} x_0 \\ X \end{matrix} \right], \quad \int_{x_0}^X f(x) dx \left[\begin{matrix} x_0 \\ x=X \end{matrix} \right].$$

La première de ces notations, imaginée par M. *Fourier*, est la plus simple. Dans le cas particulier où la fonction $f(x)$ est remplacée par une quantité constante a , on trouve, quel que soit le mode de division de la différence $X - x_0$,

$$S = a(X - x_0), \quad \text{et l'on en conclut}$$

$$(11) \quad \int_{x_0}^X a dx = a(X - x_0),$$

Si, dans cette dernière formule on pose $a = 1$, on en tirera

$$(12) \quad \int_{x_0}^X dx = X - x_0.$$

Cauchy and integrals

84

COURS D'ANALYSE.

Observons maintenant que, si l'on désigne par $\Delta x = h = dx$ un accroissement fini attribué à la variable x , les différens termes dont se compose la valeur de S , tels que les produits $(x, -x_0)f(x_0)$, $(x, -x_1)f(x_1)$, &c... seront tous compris dans la formule générale

$$(8) \quad hf(x) = f(x) dx$$

de laquelle on les déduira l'un après l'autre, en posant d'abord $x = x_0$, et $h = x_1 - x_0$, puis $x = x_1$, et $h = x_2 - x_1$, &c... On peut donc énoncer que la quantité S est une somme de produits semblables à l'expression (8); ce qu'on exprime quelquefois à l'aide de la caractéristique Σ en écrivant

$$(9) \quad S = \Sigma hf(x) = \Sigma f(x) \Delta x.$$

Quant à l'intégrale définie vers laquelle converge la quantité S , tandis que les élémens de la différence $X - x_0$ deviennent infiniment petits, on est convenu de la représenter par la notation $\int hf(x)$ ou $\int f(x) dx$, dans laquelle la lettre f substituée à la lettre Σ indique, non plus une somme de produits semblables à l'expression (8), mais la limite d'une somme de cette espèce. De plus, comme la valeur de l'intégrale définie que l'on considère dépend des valeurs extrêmes x_0, X attribuées à la variable x , on est convenu de placer ces deux valeurs, la première au-dessous, la seconde au-dessus de la lettre f , ou de les écrire à côté de l'intégrale, que l'on désigne en conséquence par l'une des notations

$$(10) \quad \int_{x_0}^X f(x) dx, \quad \int_{x_0}^{x_1} f(x) dx, \quad \int_{x=X}^{x=x_0} f(x) dx.$$

La première de ces notations, imaginée par M. *Fourier*, est la plus simple. Dans le cas particulier où la fonction $f(x)$ est remplacée par une quantité constante a , on trouve, quel que soit le mode de division de la différence $X - x_0$,

$$S = a(X - x_0), \quad \text{et l'on en conclut}$$

$$(11) \quad \int_{x_0}^X a dx = a(X - x_0),$$

Si, dans cette dernière formule on pose $a = 1$, on en tirera

$$(12) \quad \int_{x_0}^X dx = X - x_0.$$

Is it valid to use the symbol \int here?

Cauchy and the Fundamental Theorem of Calculus

VINGT-SIXIÈME LEÇON.

Intégrales indéfinies.

Si, dans l'intégrale définie $\int_{x_0}^X f(x) dx$, on fait varier l'une des deux limites, par exemple, la quantité X , l'intégrale variera elle-même avec cette quantité; et, si l'on remplace la limite X devenue variable par x , on obtiendra pour résultat une nouvelle fonction de x , qui sera ce qu'on appelle une intégrale prise à partir de l'origine $x = x_0$. Soit

$$(1) \quad \mathcal{F}(x) = \int_{x_0}^x f(x) dx$$

cette fonction nouvelle. On tirera de la formule (19) [22.^e leçon]

$$(2) \quad \mathcal{F}(x) = (x - x_0) f[x_0 + \theta(x - x_0)], \quad \mathcal{F}(x_0) = 0,$$

θ étant un nombre inférieur à l'unité; et de la formule (7) [23.^e leçon]

$$\int_{x_0}^{x+a} f(x) dx - \int_{x_0}^x f(x) dx = \int_x^{x+a} f(x) dx = a f(x + \theta a), \quad \text{ou}$$

$$(3) \quad \mathcal{F}(x+a) - \mathcal{F}(x) = a f(x + \theta a).$$

Il suit des équations (2) et (3) que, si la fonction $f(x)$ est finie et continue dans le voisinage d'une valeur particulière attribuée à la variable x , la nouvelle fonction $\mathcal{F}(x)$ sera non-seulement finie, mais encore continue dans le voisinage de cette valeur, puisqu'à un accroissement infiniment petit de x correspondra un accroissement infiniment petit de $\mathcal{F}(x)$. Donc, si la fonction $f(x)$ reste finie et continue depuis $x = x_0$ jusqu'à $x = X$, il en sera de même de la fonction $\mathcal{F}(x)$. Ajoutons que, si l'on divise par a les deux membres de la formule (3), on en conclura, en passant aux limites,

$$(4) \quad \mathcal{F}'(x) = f(x).$$

Donc l'intégrale (1), considérée comme fonction de x , a pour dérivée la fonction $f(x)$ renfermée sous le signe \int dans cette intégrale. On prouverait de la même manière que l'intégrale $\int_x^X f(x) dx = -\int_X^x f(x) dx$;

If in the definite integral $\int_{x_0}^X f(x) dx$ one makes one of the two limits vary, for example the quantity X , the integral itself will vary with this quantity; and if one replaces the variable limit X by x , there results a new function of x , ...

Cauchy and the Fundamental Theorem of Calculus

VINGT-SIXIÈME LEÇON.

Intégrales indéfinies.

Si, dans l'intégrale définie $\int_{x_0}^X f(x) dx$, on fait varier l'une des deux limites, par exemple, la quantité X , l'intégrale variera elle-même avec cette quantité; et, si l'on remplace la limite X devenue variable par x , on obtiendra pour résultat une nouvelle fonction de x , qui sera ce qu'on appelle une intégrale prise à partir de l'origine $x = x_0$. Soit

$$(1) \quad \mathcal{F}(x) = \int_{x_0}^x f(x) dx$$

cette fonction nouvelle. On tirera de la formule (19) [22.^e leçon]

$$(2) \quad \mathcal{F}(x) = (x - x_0) f[x_0 + \theta(x - x_0)], \quad \mathcal{F}(x_0) = 0,$$

θ étant un nombre inférieur à l'unité; et de la formule (7) [23.^e leçon]

$$\int_{x_0}^{x_0+a} f(x) dx - \int_{x_0}^{x_0} f(x) dx = \int_{x_0}^{x_0+a} f(x) dx = a f(x_0 + \theta a), \quad \text{ou}$$

$$(3) \quad \mathcal{F}(x_0 + a) - \mathcal{F}(x_0) = a f(x_0 + \theta a).$$

Il suit des équations (2) et (3) que, si la fonction $f(x)$ est finie et continue dans le voisinage d'une valeur particulière attribuée à la variable x , la nouvelle fonction $\mathcal{F}(x)$ sera non-seulement finie, mais encore continue dans le voisinage de cette valeur, puisqu'à un accroissement infiniment petit de x correspondra un accroissement infiniment petit de $\mathcal{F}(x)$. Donc, si la fonction $f(x)$ reste finie et continue depuis $x = x_0$ jusqu'à $x = X$, il en sera de même de la fonction $\mathcal{F}(x)$. Ajoutons que, si l'on divise par a les deux membres de la formule (3), on en conclura, en passant aux limites,

$$(4) \quad \mathcal{F}'(x) = f(x).$$

Donc l'intégrale (1), considérée comme fonction de x , a pour dérivée la fonction $f(x)$ renfermée sous le signe \int dans cette intégrale. On prouverait de la même manière que l'intégrale $\int_a^X f(x) dx = -\int_X^a f(x) dx$;

Let

$$\mathcal{F}(x) = \int_{x_0}^x f(x) dx$$

be this new function.

Cauchy and the Fundamental Theorem of Calculus

VINGT-SIXIÈME LEÇON.

Intégrales indéfinies.

Si, dans l'intégrale définie $\int_{x_0}^X f(x) dx$, on fait varier l'une des deux limites, par exemple, la quantité X , l'intégrale variera elle-même avec cette quantité; et, si l'on remplace la limite X devenue variable par x , on obtiendra pour résultat une nouvelle fonction de x , qui sera ce qu'on appelle une intégrale prise à partir de l'origine $x = x_0$. Soit

$$(1) \quad \mathcal{F}(x) = \int_{x_0}^x f(x) dx$$

cette fonction nouvelle. On tirera de la formule (19) [22.^e leçon]

$$(2) \quad \mathcal{F}(x) = (x - x_0) f[x_0 + \theta(x - x_0)], \quad \mathcal{F}(x_0) = 0,$$

θ étant un nombre inférieur à l'unité; et de la formule (7) [23.^e leçon]

$$\int_{x_0}^{x_0+a} f(x) dx - \int_{x_0}^{x_0} f(x) dx = \int_{x_0}^{x_0+a} f(x) dx = a f(x_0 + \theta a), \quad \text{ou}$$

$$(3) \quad \mathcal{F}(x_0 + a) - \mathcal{F}(x_0) = a f(x_0 + \theta a).$$

Il suit des équations (2) et (3) que, si la fonction $f(x)$ est finie et continue dans le voisinage d'une valeur particulière attribuée à la variable x , la nouvelle fonction $\mathcal{F}(x)$ sera non-seulement finie, mais encore continue dans le voisinage de cette valeur, puisqu'à un accroissement infiniment petit de x correspondra un accroissement infiniment petit de $\mathcal{F}(x)$. Donc, si la fonction $f(x)$ reste finie et continue depuis $x = x_0$ jusqu'à $x = X$, il en sera de même de la fonction $\mathcal{F}(x)$. Ajoutons que, si l'on divise par a les deux membres de la formule (3), on en conclura, en passant aux limites,

$$(4) \quad \mathcal{F}'(x) = f(x).$$

Donc l'intégrale (1), considérée comme fonction de x , a pour dérivée la fonction $f(x)$ renfermée sous le signe \int dans cette intégrale. On prouverait de la même manière que l'intégrale $\int_x^X f(x) dx = -\int_x^X f(x) dx$;

Let

$$\mathcal{F}(x) = \int_{x_0}^x f(x) dx$$

be this new function.

Proved that $\mathcal{F}'(x) = f(x)$,

Cauchy and the Fundamental Theorem of Calculus

VINGT-SIXIÈME LEÇON.

Intégrales indéfinies.

Si, dans l'intégrale définie $\int_{x_0}^X f(x) dx$, on fait varier l'une des deux limites, par exemple, la quantité X , l'intégrale variera elle-même avec cette quantité; et, si l'on remplace la limite X devenue variable par x , on obtiendra pour résultat une nouvelle fonction de x , qui sera ce qu'on appelle une intégrale prise à partir de l'origine $x = x_0$. Soit

$$(1) \quad \mathcal{F}(x) = \int_{x_0}^x f(x) dx$$

cette fonction nouvelle. On tirera de la formule (19) [22.^e leçon]

$$(2) \quad \mathcal{F}(x) = (x - x_0) f[x_0 + \theta(x - x_0)], \quad \mathcal{F}(x_0) = 0,$$

θ étant un nombre inférieur à l'unité; et de la formule (7) [23.^e leçon]

$$\int_{x_0}^{x_0+a} f(x) dx - \int_{x_0}^{x_0} f(x) dx = \int_{x_0}^{x_0+a} f(x) dx = a f(x_0 + \theta a), \quad \text{ou}$$

$$(3) \quad \mathcal{F}(x_0 + a) - \mathcal{F}(x_0) = a f(x_0 + \theta a).$$

Il suit des équations (2) et (3) que, si la fonction $f(x)$ est finie et continue dans le voisinage d'une valeur particulière attribuée à la variable x , la nouvelle fonction $\mathcal{F}(x)$ sera non-seulement finie, mais encore continue dans le voisinage de cette valeur, puisqu'à un accroissement infiniment petit de x correspondra un accroissement infiniment petit de $\mathcal{F}(x)$. Donc, si la fonction $f(x)$ reste finie et continue depuis $x = x_0$ jusqu'à $x = X$, il en sera de même de la fonction $\mathcal{F}(x)$. Ajoutons que, si l'on divise par a les deux membres de la formule (3), on en conclura, en passant aux limites,

$$(4) \quad \mathcal{F}'(x) = f(x).$$

Donc l'intégrale (1), considérée comme fonction de x , a pour dérivée la fonction $f(x)$ renfermée sous le signe \int dans cette intégrale. On prouverait de la même manière que l'intégrale $\int_x^X f(x) dx = -\int_X^x f(x) dx$;

Let

$$\mathcal{F}(x) = \int_{x_0}^x f(x) dx$$

be this new function.

Proved that $\mathcal{F}'(x) = f(x)$, and also that

$$\varpi'(x) = 0 \Rightarrow \varpi(x) = \text{const},$$

Cauchy and the Fundamental Theorem of Calculus

VINGT-SIXIÈME LEÇON.

Intégrales indéfinies.

Si, dans l'intégrale définie $\int_{x_0}^X f(x) dx$, on fait varier l'une des deux limites, par exemple, la quantité X , l'intégrale variera elle-même avec cette quantité; et, si l'on remplace la limite X devenue variable par x , on obtiendra pour résultat une nouvelle fonction de x , qui sera ce qu'on appelle une intégrale prise à partir de l'origine $x = x_0$. Soit

$$(1) \quad \mathcal{F}(x) = \int_{x_0}^x f(x) dx$$

cette fonction nouvelle. On tirera de la formule (19) [22.^e leçon]

$$(2) \quad \mathcal{F}(x) = (x - x_0) f[x_0 + \theta(x - x_0)], \quad \mathcal{F}(x_0) = 0,$$

θ étant un nombre inférieur à l'unité; et de la formule (7) [23.^e leçon]

$$\int_{x_0}^{x_0+a} f(x) dx - \int_{x_0}^{x_0+a} f(x) dx = \int_{x_0}^{x_0+a} f(x) dx = a f(x_0 + \theta a), \quad \text{ou}$$

$$(3) \quad \mathcal{F}(x_0 + a) - \mathcal{F}(x_0) = a f(x_0 + \theta a).$$

Il suit des équations (2) et (3) que, si la fonction $f(x)$ est finie et continue dans le voisinage d'une valeur particulière attribuée à la variable x , la nouvelle fonction $\mathcal{F}(x)$ sera non-seulement finie, mais encore continue dans le voisinage de cette valeur, puisqu'à un accroissement infiniment petit de x correspondra un accroissement infiniment petit de $\mathcal{F}(x)$. Donc, si la fonction $f(x)$ reste finie et continue depuis $x = x_0$ jusqu'à $x = X$, il en sera de même de la fonction $\mathcal{F}(x)$. Ajoutons que, si l'on divise par a les deux membres de la formule (3), on en conclura, en passant aux limites,

$$(4) \quad \mathcal{F}'(x) = f(x).$$

Donc l'intégrale (1), considérée comme fonction de x , a pour dérivée la fonction $f(x)$ renfermée sous le signe \int dans cette intégrale. On prouverait de la même manière que l'intégrale $\int_a^X f(x) dx = -\int_X^a f(x) dx$;

Let

$$\mathcal{F}(x) = \int_{x_0}^x f(x) dx$$

be this new function.

Proved that $\mathcal{F}'(x) = f(x)$, and also that

$$\varpi'(x) = 0 \Rightarrow \varpi(x) = \text{const},$$

which may be used to show that if $F'(x) = f(x)$, then

$$\int_{x_0}^X f(x) dx = F(X) - F(x_0).$$

The Fundamental Theorem of Calculus

What is the Fundamental Theorem of Calculus?

The Fundamental Theorem of Calculus

What is the Fundamental Theorem of Calculus?

- ▶ integration is the inverse of differentiation?

The Fundamental Theorem of Calculus

What is the Fundamental Theorem of Calculus?

- ▶ integration is the inverse of differentiation?
- ▶ integration 'as a sum' is the same as integration 'by rule'?

The Fundamental Theorem of Calculus

What is the Fundamental Theorem of Calculus?

- ▶ integration is the inverse of differentiation?
- ▶ integration 'as a sum' is the same as integration 'by rule'?
- ▶ Cauchy's integration is the same as Euler's integration?

The Fundamental Theorem of Calculus

What is the Fundamental Theorem of Calculus?


- ▶ integration is the inverse of differentiation?
- ▶ integration 'as a sum' is the same as integration 'by rule'?
- ▶ Cauchy's integration is the same as Euler's integration?
- ▶ 19th-century integration is the same as 18th-century integration?

The Fundamental Theorem of Calculus

What is the Fundamental Theorem of Calculus?

- ▶ integration is the inverse of differentiation?
- ▶ integration 'as a sum' is the same as integration 'by rule'?
- ▶ Cauchy's integration is the same as Euler's integration?
- ▶ 19th-century integration is the same as 18th-century integration?
- ▶ ...

Bernhard Riemann (1826–1866)



Riemann's integral (1853)

Function $f(x)$ no longer required to be continuous on $[a, b]$.

Riemann's integral (1853)

Function $f(x)$ no longer required to be continuous on $[a, b]$. Take $x_1 < x_2 < \dots < x_{n-1}$. Define $\delta_1 := x_1 - a$, $\delta_2 := x_2 - x_1$, ..., $\delta_n := b - x_{n-1}$.

Riemann's integral (1853)

Function $f(x)$ no longer required to be continuous on $[a, b]$. Take $x_1 < x_2 < \cdots < x_{n-1}$. Define $\delta_1 := x_1 - a$, $\delta_2 := x_2 - x_1$, ..., $\delta_n := b - x_{n-1}$. Choose numbers ε_i between 0 and 1. Then define

$$S := \delta_1 f(a + \varepsilon_1 \delta_1) + \delta_2 f(x_1 + \varepsilon_2 \delta_2) \\ + \delta_3 f(x_2 + \varepsilon_3 \delta_3) + \cdots + \delta_n f(x_{n-1} + \varepsilon_n \delta_n)$$

Riemann's integral (1853)

Function $f(x)$ no longer required to be continuous on $[a, b]$. Take $x_1 < x_2 < \dots < x_{n-1}$. Define $\delta_1 := x_1 - a$, $\delta_2 := x_2 - x_1$, ..., $\delta_n := b - x_{n-1}$. Choose numbers ε_i between 0 and 1. Then define

$$S := \delta_1 f(a + \varepsilon_1 \delta_1) + \delta_2 f(x_1 + \varepsilon_2 \delta_2) \\ + \delta_3 f(x_2 + \varepsilon_3 \delta_3) + \dots + \delta_n f(a_{n-1} + \varepsilon_n \delta_n)$$

If this has the property that it comes infinitely close to a fixed value A when all the δ_i become infinitely small, then this is the value of $\int_a^b f(x) dx$.

Riemann's integral (1853)

Function $f(x)$ no longer required to be continuous on $[a, b]$. Take $x_1 < x_2 < \dots < x_{n-1}$. Define $\delta_1 := x_1 - a$, $\delta_2 := x_2 - x_1$, ..., $\delta_n := b - x_{n-1}$. Choose numbers ε_i between 0 and 1. Then define

$$S := \delta_1 f(a + \varepsilon_1 \delta_1) + \delta_2 f(x_1 + \varepsilon_2 \delta_2) \\ + \delta_3 f(x_2 + \varepsilon_3 \delta_3) + \dots + \delta_n f(a_{n-1} + \varepsilon_n \delta_n)$$

If this has the property that it comes infinitely close to a fixed value A when all the δ_i become infinitely small, then this is the value of $\int_a^b f(x) dx$.

Many variants over the years, all called **Riemann integral**.

Lebesgue's integral (1901)

Considers step functions on subsets that are not necessarily intervals, thus requiring the notion of a **measure** (Borel, 1894).

Lebesgue's integral (1901)

Considers step functions on subsets that are not necessarily intervals, thus requiring the notion of a **measure** (Borel, 1894).

Results in a notion of integral of wider applicability than Riemann's;

Lebesgue's integral (1901)

Considers step functions on subsets that are not necessarily intervals, thus requiring the notion of a **measure** (Borel, 1894).

Results in a notion of integral of wider applicability than Riemann's; for example:

can integrate highly discontinuous functions, such as the Dirichlet function:

$$f(x) := \begin{cases} 1 & \text{if } x \text{ is rational;} \\ 0 & \text{if } x \text{ is irrational.} \end{cases}$$