

## Dynamics: Problem Sheet 5 (of 8)

1. A particle moves on the inside surface of a smooth cone with its axis vertical, defined by the equation  $z = r$  in cylindrical polar coordinates  $(r, \theta, z)$ . Initially the particle is at height  $z = a$ , and its velocity is horizontal, speed  $v$ , in the  $\mathbf{e}_\theta$  direction. Starting from Newton's second law show that  $r^2\dot{\theta}$  is constant. Explain why the total energy is conserved, and deduce that

$$\dot{z}^2 + \frac{1}{2} \frac{a^2 v^2}{z^2} + gz = \frac{1}{2} v^2 + ga .$$

Hence show that the particle remains at all times between two heights, which should be determined.

2. A particle of mass  $m$ , moving under gravity, is disturbed from rest at the highest point on the outside of a smooth sphere of radius  $a$ .
- (a) Explain why the particle subsequently moves on a great circle.
- (b) By introducing plane polar coordinates in the vertical plane of this circle (or otherwise), show that

$$\ddot{\theta} = \frac{g}{a} \sin \theta , \quad N = mg \cos \theta - ma\dot{\theta}^2 .$$

Here  $\theta(t)$  denotes the angle between the *upward* vertical axis of the sphere and the straight line from the particle to the centre of the sphere (the usual polar angle for a sphere), and  $N$  is the magnitude of the normal reaction.

- (c) Show that the normal reaction is given by

$$N = mg \left( \frac{3z}{a} - 2 \right) ,$$

where  $z$  is the height of the particle above the centre of the sphere. At what height does the particle lose contact with the sphere?

3. A bead of mass  $m$  is free to slide on a smooth wire that is made to rotate at constant angular velocity  $\omega$  about the vertical axis through a fixed point  $O$  on the wire. The wire is bent into the shape of a parabola,  $z = r^2/2a$ , where  $z$  is measured vertically upwards from  $O$ , and  $r$  is the horizontal distance from  $O$ .

- (a) Show that if  $z(t)$  and  $r(t)$  are the vertical and horizontal distances of the bead from  $O$ , then

$$m [(\ddot{r} - r\omega^2)\mathbf{e}_r + 2\omega\dot{r}\mathbf{e}_\theta + \ddot{z}\mathbf{e}_z] = -mg\mathbf{e}_z + \mathbf{N} ,$$

where  $\mathbf{N}$  is the normal reaction.

- (b) Hence deduce that

$$(a^2 + r^2)\ddot{r} + r\dot{r}^2 = (a^2\omega^2 - ga)r \quad (*) .$$

- (c) Show that  $r = 0$  is an equilibrium point. The *linearized equation of motion* about  $r = 0$  is

$$a^2\ddot{\xi} = (a^2\omega^2 - ga)\xi ,$$

where we have written  $r = \xi$ , and kept only the linear terms in  $\xi, \dot{\xi}$  in equation (\*), in a Taylor expansion around  $\xi = 0$ . Discuss the stability of the equilibrium point.

4. (*Optional*) A particle of mass  $m$  is released from rest at a very large height  $z = z_0$  above the Earth. The Newtonian gravitational potential energy of the particle is

$$V(z) = -\frac{G_N M m}{z},$$

where  $M$  is the mass of the Earth, and  $G_N$  is Newton's gravitational constant.

- (a) Using conservation of energy show that the trajectory  $z(t)$  satisfies

$$\sqrt{2G_N M} t = -\int_{z_0}^z \left( \frac{1}{s} - \frac{1}{z_0} \right)^{-1/2} ds.$$

- (b) Using the substitution  $s = z_0 \sin^2 \theta$ , show  $z(t)$  satisfies the unlikely looking equation

$$\frac{\pi}{2} - \sin^{-1} \left( \sqrt{\frac{z(t)}{z_0}} \right) + \frac{1}{2} \sin \left[ 2 \sin^{-1} \left( \sqrt{\frac{z(t)}{z_0}} \right) \right] = \sqrt{\frac{2G_N M}{z_0^3}} t.$$

This is a *radial Kepler trajectory* (c.f. section 6.2 of the lecture notes).

Please send comments and corrections to [gaffney@maths.ox.ac.uk](mailto:gaffney@maths.ox.ac.uk).