

## Linear Algebra II Problem Sheet 1, HT 2019

1. Calculate the determinants of the five matrices

$$A = \begin{pmatrix} 2 & 0 & -1 \\ 1 & 2 & 2 \\ 3 & 2 & 4 \end{pmatrix}, B = \begin{pmatrix} 2 & 1 & -2 \\ -1 & 1 & 5 \\ 4 & 2 & 3 \end{pmatrix}, AB^2, A + B, AB + A^2.$$

2. Solve the equation

$$\det \begin{pmatrix} a-x & b-x & c \\ a-x & c & b-x \\ a & b-x & c-x \end{pmatrix} = 0$$

for the variable  $x$ .

3. Let  $A$  be an  $n \times n$  matrix. Suppose that  $A$  has the form  $\begin{pmatrix} U & V \\ W & X \end{pmatrix}$  in which  $U, V, W$  and  $X$  are  $n_1 \times n_1, n_1 \times n_2, n_2 \times n_1$  and  $n_2 \times n_2$  matrices respectively, with  $n_1 + n_2 = n$ . Show that if  $W = 0$  then  $\det(A) = \det(U) \det(X)$ .

4. Show that

$$\det \begin{pmatrix} 0 & 1 & 1 & 1 \\ 1 & 0 & \alpha + \beta & \alpha + \gamma \\ 1 & \beta + \alpha & 0 & \beta + \gamma \\ 1 & \gamma + \alpha & \gamma + \beta & 0 \end{pmatrix} = -4(\alpha\beta + \beta\gamma + \gamma\alpha).$$

What is the value of this when  $\alpha, \beta, \gamma$  are the three roots of the equation  $x^3 - 1 = 0$ ?

5. Here we derive an explicit formula for the inverse of a matrix with non-zero determinant.

- (i) (Cramer's rule) Let  $\mathbf{a}_1, \dots, \mathbf{a}_n \in \mathbb{R}^n$  (column vectors) and  $x_1, \dots, x_n \in \mathbb{R}$  with

$$x_1 \mathbf{a}_1 + \dots + x_n \mathbf{a}_n = \mathbf{b}$$

for some  $\mathbf{b} \in \mathbb{R}^n$ . Show that for each  $i$  we have

$$x_i \det[\mathbf{a}_1, \dots, \mathbf{a}_n] = \det[\mathbf{a}_1, \dots, \mathbf{b}, \dots, \mathbf{a}_n]$$

where the  $\mathbf{b}$  occurs in the  $i$ th place. [Hint: Use the properties of  $\det$  from Definition 1.1 and Proposition 1.2 of the lecture notes.]

- (ii) Now let  $A = [\mathbf{a}_1, \dots, \mathbf{a}_n]$  and assume  $\det(A) \neq 0$ , and so there exists  $B = (b_{ij}) \in M_n(\mathbb{R})$  such that  $AB = I_n$ . Write  $\mathbf{e}_j \in \mathbb{R}^n$  for the column vector with 1 in the  $j$ th place and zeros elsewhere. Show using (i) that

$$b_{ij} = \frac{\det[\mathbf{a}_1, \dots, \mathbf{e}_j, \dots, \mathbf{a}_n]}{\det(A)}$$

where the  $\mathbf{e}_j$  occurs in the  $i$ th position.

- (iii) By expanding  $\det[\mathbf{a}_1, \dots, \mathbf{e}_j, \dots, \mathbf{a}_n]$  down the  $i$ th column using the Laplace expansion, show that

$$b_{ij} = \frac{(-1)^{i+j} \det(A_{ji})}{\det(A)}$$

where  $A_{ji}$  is the matrix obtained from  $A$  by deleting the  $j$ th row and  $i$ th column.