

Part A Linear Algebra MT 2018, Sheet 1 of 4 <sup>1</sup>

1. Show that the vector space of polynomials  $\mathbb{R}[x]$  is isomorphic to a proper subspace of itself.
2. (Harder) Show that the space of functions  $f : \mathbb{N} \rightarrow \mathbb{R}$  does not have a countable basis.
3. Let  $\mathbb{F}$  be a field and  $f(x)$  be an irreducible polynomial in  $\mathbb{F}[x]$ . Show that the set of polynomials modulo  $f(x)$  form a field.
4. (a) Show that the set  $M_n(R)$  of  $(n \times n)$ -matrices with entries in a ring  $R$  is a ring with the usual matrix addition and multiplication.  
(b) Show that the canonical surjection  $R \rightarrow R/I$  induces a surjective ring homomorphism  $M_n(R) \rightarrow M_n(R/I)$ . What is the kernel?  
(c) Describe, with justification, the ideals of  $M_n(R)$  for a ring  $R$  with multiplicative unit 1.
5. Prove that a linear transformation  $P : V \rightarrow V$  of a finite dimensional vector space satisfies  $P^2 = P$  if and only if there exists a basis such that the matrix of  $P$  with respect to that basis is a block matrix

$$\begin{pmatrix} I & 0 \\ 0 & 0 \end{pmatrix}.$$

Hence determine the minimal and characteristic polynomials of  $P$ .

6. Let  $T : V \rightarrow V$  be a linear transformation of a finite dimensional vector space over a field  $\mathbb{F}$  to itself. Prove that  $T$  is invertible if and only if  $x$  does not divide the minimal polynomial  $m_T(x)$ .
7. Let  $T : V \rightarrow V$  be a linear transformation of a finite dimensional vector space over a field  $\mathbb{F}$  to itself. Assume that  $\{v, Tv, T^2v, \dots\}$  span  $V$  for some  $v \in V$ . Show that  
(i) there exists a  $k$  such that  $v, Tv, \dots, T^{k-1}v$  are linearly independent and for some  $\alpha_i \in \mathbb{F}$

$$T^k v = \alpha_0 v + \alpha_1 Tv + \dots + \alpha_{k-1} T^{k-1} v;$$

- (ii) the set  $\{v, Tv, \dots, T^{k-1}v\}$  forms a basis for  $V$ ;
  - (iii) its minimal polynomial is given by  $m_T(x) = x^k - \alpha_{k-1}x^{k-1} - \dots - \alpha_0$ .
- What is the characteristic polynomial  $\chi_T(x)$ ?

8. Suppose  $U$  is a subspace of  $V$  invariant under a linear transformation  $T : V \rightarrow V$ . Prove that  $T$  induces a linear map  $\bar{T} : V/U \rightarrow V/U$  of quotients given by  $\bar{T}(v + U) = T(v) + U$ . Show further that when  $V$  is finite dimensional, the minimal polynomial of  $\bar{T}$  divides the minimal polynomial of  $T$ .

---

<sup>1</sup>Many of the problems on these sheets can be found in Kaye & Wilson.

9. Let  $\mathcal{P} = \mathbb{F}[x]$  be the vector space of polynomials over the field  $\mathbb{F}$ . Determine whether or not  $\mathcal{P}/\mathcal{M}$  is finite dimensional when  $\mathcal{M}$  is
- (i) the subspace  $\mathcal{P}_n$  of polynomial of degree less or equal  $n$ ;
  - (ii) the subspace  $\mathcal{E}$  of even polynomials;
  - (iii) the subspace  $x^n\mathcal{P}$  of all polynomials divisible by  $x^n$ .
10. Let  $\mathcal{P}$  be as above and  $L : \mathcal{P} \rightarrow \mathcal{P}$  be given by  $L(f(x)) = x^2f(x)$ . Prove that  $L$  is linear. In the examples above, determine whether  $L$  induces a map of quotients  $\bar{L} : \mathcal{P}/\mathcal{M} \rightarrow \mathcal{P}/\mathcal{M}$ . When it does, choose a convenient basis for the quotient space and find a matrix representation of  $\bar{L}$ .