

## Problem Sheet 4

- Suppose that  $X_1, \dots, X_n$  each have a geometric distribution with probability mass function  $f(x|\theta) = (1-\theta)^x \theta$  for  $x = 0, 1, \dots$ . Suppose that the prior for  $\theta$  is a Beta( $a, b$ ) density. Find the posterior distribution of  $\theta$ .
- Let  $\theta > 0$  be an unknown parameter and let  $c > 0$  be a known constant. Conditional on  $\theta$ , suppose  $X_1, \dots, X_n$  are independent each with probability density function

$$f(x|\theta) = \theta c^\theta x^{-(\theta+1)}, \quad x \geq c$$

and suppose the prior for  $\theta$  is a Gamma( $\alpha, \beta$ ) density. Find the posterior distribution of  $\theta$ .

- Let  $r \geq 1$  be a known integer and let  $\theta \in [0, 1]$  be an unknown parameter. The negative binomial distribution with index  $r$  and parameter  $\theta$  has probability mass function

$$f(x|\theta) = \binom{x+r-1}{x} (1-\theta)^x \theta^r \quad \text{for } x = 0, 1, \dots$$

Let  $\theta$  have a Beta( $a, b$ ) prior density and suppose, given  $\theta$ , that  $X_1, \dots, X_n$  are independent each with the above negative binomial distribution.

- Show that the posterior density is also a Beta density.
  - Explain how to construct a  $100(1-\alpha)\%$  equal-tailed credible interval for  $\theta$ . Will this interval be a highest posterior density interval?
- Suppose that  $X$  has a  $N(\theta, \phi)$  distribution, where  $\phi$  is known, Suppose also that the prior distribution for  $\theta$  is  $N(\theta_0, \phi_0)$ , where  $\theta_0$  and  $\phi_0$  are known.
    - Find the posterior distribution of  $\theta$  given  $X = x$ .
    - Show that the posterior mean of  $\theta$  always lies between the prior mean and the observed value  $x$ .
    - Construct a  $100(1-\alpha)\%$  highest posterior density interval for  $\theta$ .
    - Let  $\phi = 2$ ,  $\theta_0 = 0$  and  $\phi_0 = 2$ .
      - Suppose the observed value is  $x = 4$ . What are the mean and variance of the resulting posterior distribution? Sketch the prior, likelihood, and posterior on a single set of coordinate axes.
      - Repeat (i) assuming  $\phi_0 = 18$ . Explain any resulting differences. Which of these two priors would likely have more appeal for a frequentist statistician?

- Let  $X$  be the number of heads when a coin with probability  $\theta$  of heads is flipped  $n$  times.

- When the prior is  $\pi(\theta)$ , the prior predictive distribution for  $X$  (the predictive distribution before observing any data) is given by

$$P(X = k) = \int_0^1 P(X = k|\theta)\pi(\theta) d\theta, \quad k = 0, 1, \dots, n.$$

Find the prior predictive distribution when  $\pi(\theta)$  is uniform on  $(0, 1)$ .

- Suppose you assign a Beta( $a, b$ ) prior for  $\theta$ , and then you observe  $x$  heads out of  $n$  flips. Show that the posterior mean of  $\theta$  is always lies between your prior mean,  $a/(a+b)$ , and the observed relative frequency of heads,  $x/n$ .

- (c) Show that, if the prior distribution on  $\theta$  is uniform, then the posterior variance is always less than the prior variance.
- (d) Give an example of a  $\text{Beta}(a, b)$  prior distribution and values of  $x, n$  for which the posterior variance is larger than the prior variance. (Try  $x = n = 1$ .)
6. A coin, with probability  $\theta$  of heads, is flipped  $n$  times and  $r$  heads are observed.
- (a) If the prior for  $\theta$  is a uniform distribution on  $(0, 1)$ , what is the probability that the next flip is a head?
- (b) Can you generalise to the case where  $\theta$  has a  $\text{Beta}(a, b)$  prior and where we wish to find the probability of getting  $k$  heads from  $m$  further flips?
7. (a) Let  $X \sim N(\theta, \sigma_0^2)$ , where  $\sigma_0^2$  is known. Find the Jeffreys' prior for  $\theta$ .
- (b) Let  $X \sim N(\mu_0, \sigma^2)$ , where  $\mu_0$  is known. Find the Jeffreys' prior for  $\sigma$ .
- (c) Let  $X$  be Poisson with parameter  $\lambda$ . Find the Jeffreys' prior for  $\lambda$ . Check that the posterior distribution of  $\theta$  given  $X = x$  is proper, but that the Jeffreys' prior is not.
8. Suppose  $X$  is the number of successes in a binomial experiment with  $n$  trials and probability of success  $\theta$ . Either  $H_0 : \theta = \frac{1}{2}$  or  $H_1 : \theta = \frac{3}{4}$  is true. Show that the posterior probability that  $H_0$  is true is greater than the prior probability for  $H_0$  if and only if

$$x \log 3 < n \log 2.$$

9. Let  $X \sim \text{Binomial}(n, \theta)$ , where the prior for  $\theta$  is uniform on  $(0, 1)$ . Suppose that we wish to compare the hypotheses  $H_0 : \theta \leq \frac{1}{2}$  and  $H_1 : \theta > \frac{1}{2}$ .
- What are the prior odds of  $H_0$  relative to  $H_1$ ?
- Find an expression for the posterior odds of  $H_0$  relative to  $H_1$ .
- If we observe  $X = n$ , find the Bayes factor  $B$  of  $H_0$  relative to  $H_1$ .
- Check that  $B \rightarrow 0$  as  $n \rightarrow \infty$ . Why is this expected?
10. Suppose we have a random sample  $X_1, \dots, X_n$  from a Poisson distribution with mean  $\theta$ . Suppose we wish to test the hypothesis  $H_0 : \theta = \theta_0$  against  $H_1 : \theta \neq \theta_0$  and that, under  $H_1$ , the prior distribution  $\pi(\theta|H_1)$  for  $\theta$  is given by

$$\pi(\theta|H_1) = \frac{\beta^\alpha}{\Gamma(\alpha)} \theta^{\alpha-1} e^{-\beta\theta}, \quad \theta > 0.$$

Calculate the Bayes factor of  $H_0$  relative to  $H_1$ .

When  $n = 6$ ,  $\sum x_i = 19$ ,  $\theta_0 = 2$ , find the numerical value of the Bayes factor (i) when  $\alpha = 4$  and  $\beta = \frac{2}{3}$ , and (ii) when  $\alpha = 36$  and  $\beta = 6$ . Compare and interpret the values of the Bayes factor in cases (i) and (ii).