

Part A Integration: HT 2019

Problem Sheet 2: Measurable functions, Lebesgue integral

1. Let (ω_n) be a sequence of non-negative real numbers. For a subset E of \mathbb{N} , let

$$\mu_\omega(E) = \sum_{n \in E} \omega_n.$$

Show that μ_ω is a measure on $(\mathbb{N}, \mathcal{P}(\mathbb{N}))$.

Now let ν be any measure on $(\mathbb{N}, \mathcal{P}(\mathbb{N}))$, and define $\omega_n = \nu(\{n\})$. Show that $\nu(E) = \mu_\omega(E)$ for all subsets E of \mathbb{N} .

2. Let $(\Omega, \mathcal{F}, \mu)$ be any measure space. If (A_n) is a decreasing sequence of sets in \mathcal{F} and $\mu(A_1) < \infty$, prove that

$$\mu\left(\bigcap_{n=1}^{\infty} A_n\right) = \lim_{n \rightarrow \infty} \mu(A_n).$$

Is this still true if $\mu(A_1) = \infty$?

3. Let $b \in \mathbb{R}$. Show that $(-\infty, b) = \bigcup_{n=1}^{\infty} (\mathbb{R} \setminus (b - \frac{1}{n}, \infty))$.

Deduce that if \mathcal{F} is a σ -algebra on \mathbb{R} containing the intervals (a, ∞) for each $a \in \mathbb{R}$, then \mathcal{F} contains all open intervals, and hence all open subsets of \mathbb{R} .

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a function. Show that

$$\mathcal{G} := \{G \subseteq \mathbb{R} : f^{-1}(G) \in \mathcal{M}_{\text{Leb}}\}$$

is a σ -algebra. Deduce that if $f^{-1}(a, \infty) \in \mathcal{M}_{\text{Leb}}$ for every a , then $f^{-1}(G) \in \mathcal{M}_{\text{Leb}}$ for every $G \in \mathcal{M}_{\text{Bor}}$.

4. Let $(\mathcal{F}_\lambda)_{\lambda \in \Lambda}$ be a non-empty family of σ -algebras on the same set Ω . Show that $\bigcap_{\lambda \in \Lambda} \mathcal{F}_\lambda$ is a σ -algebra.

By considering the family of all σ -algebras containing \mathcal{B} , deduce that if \mathcal{B} is any subset of $\mathcal{P}(\Omega)$, there is a unique σ -algebra $\mathcal{F}_{\mathcal{B}}$ such that

(i) $\mathcal{B} \subseteq \mathcal{F}_{\mathcal{B}}$;

(ii) If \mathcal{G} is a σ -algebra on Ω and $\mathcal{B} \subseteq \mathcal{G}$, then $\mathcal{F}_{\mathcal{B}} \subseteq \mathcal{G}$.

5. Let $(\Omega, \mathcal{F}, \mu)$ be a measure space, Ω_* be a set, and $f : \Omega \rightarrow \Omega_*$ be a function. Let

$$f_*(\mathcal{F}) = \{G \subseteq \Omega_* : f^{-1}(G) \in \mathcal{F}\}, \quad (f_*\mu)(G) = \mu(f^{-1}(G)).$$

Show that $(\Omega_*, f_*(\mathcal{F}), f_*\mu)$ is a measure space.

Now let $(\Omega, \mathcal{F}, \mu) = (\mathbb{R}, \mathcal{M}_{\text{Bor}}, m)$, and $\Omega_* = \mathbb{R}$. Determine $f_*(\mathcal{M}_{\text{Bor}})$ and f_*m when

(i) $f(x) = \tan x$ if $\cos x \neq 0$, and $f(x) = 0$ if $\cos x = 0$,

* (ii) $f(x) = \arctan x$ (taking values in $(-\pi/2, \pi/2)$).

6. *(a) Let I be an interval of positive length, let $a \in I$, $f, g : I \rightarrow \mathbb{R}$ be functions such that $f(t) = g(t)$ a.e., and suppose that f and g are continuous at a . Show that $f(a) = g(a)$.
- (b) Is $\chi_{\mathbb{Q}}$ continuous a.e.? Does there exist a continuous function g such that $\chi_{\mathbb{Q}} = g$ a.e.?
- (c) Is $\chi_{(0,\infty)}$ continuous a.e.? Does there exist a continuous function g such that $\chi_{(0,\infty)} = g$ a.e.? [Use (a).]
7. Let f, g be measurable functions from \mathbb{R} to \mathbb{R} , and $h : \mathbb{R} \rightarrow \mathbb{R}$ be continuous. Recall from lectures that $f + g$ and $h \circ f$ are measurable. Prove that the following functions are measurable. [Complicated constructions are not required. Everything can be quickly deduced from the information from lectures recalled above, plus a couple of simple formulae.]
- $f^2 : x \mapsto f(x)^2$,
 - $fg : x \mapsto f(x)g(x)$,
 - $|f| : x \mapsto |f(x)|$,
 - $\max(f, g) : x \mapsto \max(f(x), g(x))$.
- *8. Suppose that g is a measurable function and $f = g$ a.e. Show that f is measurable.
- Suppose that f is continuous a.e. Show that there is a sequence of step functions (ϕ_n) such that $f = \lim_{n \rightarrow \infty} \phi_n$ a.e. Deduce that f is measurable.
9. Let $f : \mathbb{R} \rightarrow [-\infty, \infty]$ be an integrable function, and let $\alpha > 0$. Show that
- $$m(\{x : |f(x)| \geq \alpha\}) \leq \frac{1}{\alpha} \int |f|.$$
- Deduce that
- $f(x) \in \mathbb{R}$ a.e.
 - If $\int |f| = 0$, then $f(x) = 0$ a.e.
10. In each of the following cases, state whether the function f is Lebesgue integrable over the interval I . Justify your answers, *and calculate $\int_I f$ in those cases where this is feasible.
- $I = \mathbb{R}$, $f(x) = x$ if x is rational, $f(x) = 0$ if x is irrational,
 - $I = (0, \pi/2)$, $f(x) = \tan x$,
 - $I = [1, \infty)$, $f(x) = (-1)^n/n$ if $n \leq x < n + 1$, $n = 1, 2, 3, \dots$,
 - $I = (0, 1]$, $f(x) = \sin(1/x)$,
 - $I = [0, \infty)$, $f(x) = x^n e^{-x}$ where n is a positive integer,
 - $I = (0, \infty)$, $f(x) = (\log x)e^{-x}$,
 - * $I = [1, \infty)$, $f(x) = x^\alpha \log x$ where $\alpha \in \mathbb{R}$,
 - * $I = (0, \pi)$, $f(x) = (\operatorname{cosec} x)^{1/2}$,
 - * $I = (0, \infty)$, $f(x) = (1 + x)^{-1} \cos x$,
 - * $I = [1, \infty)$, $f(x) = \sin(1/x)$.