

# Introduction to Manifolds

Trinity 2019

szendroi@maths.ox.ac.uk

## Example sheet 1

1. The function  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  is defined by

$$f(x, y) = \begin{cases} \frac{|xy|^\alpha}{x^2 + y^2} & \text{for } (x, y) \neq (0, 0), \\ 0 & \text{for } (x, y) = (0, 0); \end{cases}$$

where  $\alpha > 0$ . Find the values of  $\alpha$  for which  $f$  is

- (a) continuous at  $(0, 0)$ ;
  - (b) differentiable at  $(0, 0)$ .
2. A function is called *homogeneous of degree  $k$*  if  $f(\lambda x) = \lambda^k f(x)$  for all  $\lambda > 0$  and all  $x \in \mathbb{R}^n$ .
- (a) Show that if  $f$  is homogeneous of degree  $k$ , then

$$\langle \nabla f(x), x \rangle = kf(x).$$

- (b) Show conversely that if  $f$  satisfies this equation, then  $f$  is homogeneous of degree  $k$ .
3. The function  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  is given by

$$f(x, y) = \begin{cases} \frac{xy^2}{x^2 + y^4} & \text{for } (x, y) \neq (0, 0), \\ 0 & \text{for } (x, y) = (0, 0). \end{cases}$$

Show that all the directional derivatives of  $f$  exist at the origin, but  $f$  is not differentiable at the origin.

4. In this question we use the Hilbert-Schmidt matrix norm

$$\| A \| = \left( \sum_{i,j} A_{ij}^2 \right)^{\frac{1}{2}}.$$

Show that if  $H$  has Hilbert-Schmidt norm less than 1, then  $I - H$  is invertible (you may assume that  $\| AB \| \leq \| A \| \| B \|$ ).

5. Let  $M_{n \times n}(\mathbb{R})$  denote the vector space of  $n \times n$  real matrices. Show that the derivative at the identity of the determinant function

$$\det : M_{n \times n}(\mathbb{R}) \rightarrow \mathbb{R}$$

is

$$d(\det)_I : h \mapsto \text{trace } h$$

Deduce that the derivative at an arbitrary invertible matrix  $A$  is

$$d(\det)_A : h \mapsto \det A \cdot \text{trace } (A^{-1}h).$$

6. (a) Show that the set  $GL(n, \mathbb{R})$  of invertible matrices is an open set in  $M_{n \times n}(\mathbb{R})$ .  
(b) Show that the derivative of the inversion map  $\text{Inv} : GL(n, \mathbb{R}) \rightarrow GL(n, \mathbb{R})$  is

$$d(\text{Inv})_A : h \mapsto -A^{-1}hA^{-1}$$

(Hint: look at the case where  $A$  is the identity first).