

FURTHER MATHEMATICAL BIOLOGY: PROBLEM SHEET 3
MICHAELMAS TERM 2018

PATTERN FORMATION.

Question 1.

Consider the reaction-diffusion system

$$\begin{aligned}\frac{\partial u}{\partial t} &= f(u, v) + D_1 \frac{\partial^2 u}{\partial x^2}, \\ \frac{\partial v}{\partial t} &= g(u, v) + D_2 \frac{\partial^2 v}{\partial x^2},\end{aligned}$$

where f and g describe the reaction kinetics, and D_1 and D_2 are positive constants.

- (a) State the conditions for diffusion-driven instability.
- (b) Show that, when these conditions hold, bifurcation to solutions oscillating in time (and space) cannot occur.

Question 2.

Consider the Gierer-Meinhardt reaction-diffusion system in one dimension:

$$\begin{aligned}\frac{\partial A}{\partial t} &= \frac{\rho A^2}{(1 + K A^2)H} - \mu A + D_A \frac{\partial^2 A}{\partial x^2}, \\ \frac{\partial H}{\partial t} &= \rho' A^2 - \nu H + D_H \frac{\partial^2 H}{\partial x^2},\end{aligned}$$

where A and H are the reactants and ρ , K , μ , ν , ρ' , D_A and D_H are positive constants.

- (a) Draw a phase portrait of the system in the absence of diffusion and show that diffusion-driven instability may be possible if the nullclines intersect in a certain way.
- (b) Write down the conditions for diffusion-driven instability.

[In (a) and (b) consider only the non-zero steady states.]

Question 3.

The amoebae of the slime mould *Dictyostelium discoideum* secrete a chemical attractant, cyclic-AMP, and spatial aggregations of the amoebae start to form. This process can be modelled by the following system of dimensional equations:

$$\begin{aligned}\frac{\partial \tilde{n}}{\partial \tilde{t}} &= \tilde{D}_n \frac{\partial^2 \tilde{n}}{\partial \tilde{x}^2} - \tilde{\chi} \frac{\partial}{\partial \tilde{x}} \left(\tilde{n} \frac{\partial \tilde{a}}{\partial \tilde{x}} \right), \\ \frac{\partial \tilde{a}}{\partial \tilde{t}} &= h \tilde{n} - k \tilde{a} + \tilde{D}_a \frac{\partial^2 \tilde{a}}{\partial \tilde{x}^2},\end{aligned}$$

where $\tilde{n}(\tilde{x}, \tilde{t})$ and $\tilde{a}(\tilde{x}, \tilde{t})$ are the cell density of the amoebae and the attractant concentration. The parameters h , k , $\tilde{\chi}$ and the diffusion coefficients, \tilde{D}_n and \tilde{D}_a , are positive constants.

(a) Non-dimensionalise the system to obtain

$$\begin{aligned}\frac{\partial n}{\partial t} &= D_n \frac{\partial^2 n}{\partial x^2} - \chi \frac{\partial}{\partial x} \left(n \frac{\partial a}{\partial x} \right), \\ \frac{\partial a}{\partial t} &= n - a + D_a \frac{\partial^2 a}{\partial x^2},\end{aligned}$$

where the variables and parameters are now dimensionless.

- (b) Suppose that the amoebae and chemical occupy an infinite domain. Examine the linear stability about the steady state (which introduces a further parameter), and derive the dispersion relation. Obtain the conditions on the parameters for the mechanism to initiate spatially heterogeneous solutions.
- (c) Suppose now that the amoebae and the chemical attractant are confined within a finite domain ($0 \leq \tilde{x} \leq L$ in dimensional variables), with zero flux boundary conditions imposed on both n and a at the ends of the domain. Determine the minimum domain size for which spatially structured solutions arise.
- (d) Briefly describe the physical processes operating and explain intuitively how spatial aggregation takes place.

Question 4.

Consider a tissue containing cells of density $n(x, t)$ which produce a chemical $c(x, t)$ according to the model equations

$$\frac{\partial n}{\partial t} = \mu \frac{\partial^2 n}{\partial x^2} - \frac{\partial}{\partial x} \left(n \frac{\partial c}{\partial x} \right), \quad \frac{\partial c}{\partial t} = D \frac{\partial^2 c}{\partial x^2} + \frac{n}{(1+n)^2} - c,$$

where μ and D are positive constants.

(a) If n_0 is the initial, spatially-uniform distribution of cells, what is the corresponding initial distribution of chemoattractant (*i.e.* what is the steady state for c when $n = n_0$)?

(b) Carry out a linear stability analysis about the positive steady state, by seeking solutions of the form

$$(n, c) \sim (n_0, c_0) + e^{ikx + \sigma t} (N, K), \quad 0 < |N|, |K| \ll n_0,$$

obtaining a dispersion relation for σ in terms of the wavenumber, k .

(c) Let $\mu_{max} = \max_x [x(1-x)/(1+x)^3]$. Show that when $0 < \mu < \mu_{max}$, no spatial patterns can be obtained if the initial cell density n_0 is sufficiently large or sufficiently small.