
Topology & Groups Michaelmas 2016 Question Sheet 4

- 1. Let K be a simplicial complex, and let α_1 and α_2 be edge paths. Suppose that α_1 and α_2 are homotopic relative to their endpoints. Show that α_1 and α_2 are equivalent as edge paths. [You should adapt the proof of Theorem III.27.]
- 2. Triangulate the torus as shown below



Let x and y be the loops (1, 2, 3, 1) and (1, 4, 7, 1), and let K be the union of these two loops (ie. K comes from the boundary of the square).

- (i) Show that any edge path that starts and ends on K but with the remainder of the path missing K is equivalent to an edge path lying entirely in K.
- (ii) Prove that any edge loop based at 1 is equivalent to an edge loop lying entirely in K.
- (iii) Deduce that any edge loop based at 1 is equivalent to a word in the alphabet $\{x,y\}.$
- (iv) Show that the edge loops xy and yx are equivalent.
- (v) Deduce that any edge loop based at 1 is equivalent to $x^m y^n$, for $m, n \in \mathbb{Z}$.
- (vi) Prove that if $x^m y^n \sim x^M y^N$, then m = M and n = N. [Hint: define 'winding numbers' as in the proof of Theorem III.32.]
- (vii) Deduce that the fundamental group of the torus is isomorphic to $\mathbb{Z} \times \mathbb{Z}$.

- 2. Prove that every non-trivial element of a free group has infinite order.
- 3. The centre Z(G) of a group G is $\{g \in G : gh = hg \ \forall h \in G\}$. Let S be a set with more than one element. Prove that the centre of F(S) is the identity element.
- 4. (i) Let F be the free group on the three generators x, y and z. For non-zero integers r, s and t, show that the subgroup of F generated by x^r , y^s and z^t is freely generated by these elements.
 - (ii) Let H be the subgroup of $F(\{x,y\})$ generated by x^2 , y^2 , xy and yx. Show that H is not freely generated by these elements.
- 5. Compute an explicit free generating set for the fundamental group of the following graph:

