

B2.1 Introduction to Representation Theory  
Problem Sheet 3, MT 2017

The groups below are assumed to be finite and the representations finite-dimensional, unless stated otherwise.

1. Let  $V, W$  be two  $G$ -representations over  $\mathbb{C}$ . Prove that:
  - (a)  $\chi_{V \otimes W}(g) = \chi_V(g)\chi_W(g)$  for all  $g \in G$ ;
  - (b)  $\chi_{V^*}(g) = \chi_V(g^{-1}) = \overline{\chi_V(g)}$  for all  $g \in G$ , where  $V^*$  denotes the representation contragredient to  $V$ .
  - (c) Suppose  $W$  is a one-dimensional representation. Prove that  $V \otimes W$  is irreducible if and only if  $V$  is irreducible.
  - (d) Prove that  $V$  is irreducible if and only if  $V^*$  is irreducible.
  - (e) Let  $\text{St}$  be the standard 3-dimensional representation of  $S_4$ . Decompose  $\text{St} \otimes \text{St}$  into a direct sum of irreducible representations.
2. Let  $\chi$  be the character of a  $\mathbb{C}G$ -module  $M$ . Show that  $N = \{g \in G \mid \chi(g) = \chi(1)\}$  is a normal subgroup of  $G$   
Deduce that a finite group  $G$  is simple if and only if  $\chi(g) \neq \chi(1)$  for every  $g \in G \setminus \{1\}$  and every nontrivial character of  $G$ .
3. Show if two  $\mathbb{C}G$ -modules  $M_1$  and  $M_2$  have the same characters then they are isomorphic.
4. Let  $G$  act on a finite set  $\Omega$  and let  $M$  be the permutation module with basis  $\{e_w \mid w \in \Omega\}$  defined in lectures. Let  $\chi = \chi_M$  be the character of  $M$ . Show that  $\sum_{g \in G} \chi(g) = r|G|$  where  $r$  is the number of orbits of  $G$  on  $\Omega$ .  
Suppose now that  $G$  is 2-transitive, that is  $G$  has two orbits acting on  $\Omega \times \Omega$  in the action defined by  $g \cdot (w_1, w_2) := (g \cdot w_1, g \cdot w_2)$   
Show that  $\sum_{g \in G} \chi(g)^2 = 2|G|$  and deduce that  $M$  is a sum of two irreducible submodules  $V_1 \oplus V_2$  where  $V_1$  is the trivial module.
5. Find the character tables of  $Q_8$  and  $D_8$ . Does the character table determine the group?
6. For a group  $G$  we denote by  $[G, G]$  the subgroup generated by all elements  $x^{-1}y^{-1}xy$  for all  $x, y \in G$ . The subgroup  $[G, G]$  is the smallest normal subgroup  $N$  of  $G$  such that  $G/N$  is abelian.  
Let now  $G_1$  and  $G_2$  be two groups with the same character table. Show that  $|G_1 : [G_1, G_1]| = |G_2 : [G_2, G_2]|$ . Show further that the centre of  $G_1$  has the same size as the centre of  $G_2$ .
7. Show that an element  $g$  of a finite group is conjugate to its inverse if and only if  $\chi(g) \in \mathbb{R}$  for all characters of  $G$ .

[Optional for those who are taking Galois theory]: More generally show that  $\chi(g) \in \mathbb{Z}$  if and only if  $g$  is conjugate to  $g^n$  for each integer  $n$  coprime to the order of  $g$  in  $G$ . What does this tell us about the character table of  $S_n$ ?