

## 6. A deductive system for propositional calculus

- We have introduced '*logical consequence*':  
 $\Gamma \models \phi$  – whenever (each formula of)  $\Gamma$  is true so is  $\phi$
- But we don't know yet how to give an actual **proof** of  $\phi$  from the **hypotheses**  $\Gamma$ .
- A **proof** should be a finite sequence  $\phi_1, \phi_2, \dots, \phi_n$  of statements such that
  - either  $\phi_i \in \Gamma$
  - or  $\phi_i$  is some **axiom** (which should *clearly* be true)
  - or  $\phi_i$  should follow from previous  $\phi_j$ 's by some **rule of inference**
  - AND  $\phi = \phi_n$

## 6.1 Definition

Let  $\mathcal{L}_0 := \mathcal{L}[\{\neg, \rightarrow\}]$  (which is an adequate language). Then the **system**  $L_0$  consists of the following axioms and rules:

### Axioms

An **axiom** of  $L_0$  is any formula of the following form ( $\alpha, \beta, \gamma \in \text{Form}(\mathcal{L}_0)$ ):

$$\mathbf{A1} \quad (\alpha \rightarrow (\beta \rightarrow \alpha))$$

$$\mathbf{A2} \quad ((\alpha \rightarrow (\beta \rightarrow \gamma)) \rightarrow ((\alpha \rightarrow \beta) \rightarrow (\alpha \rightarrow \gamma)))$$

$$\mathbf{A3} \quad ((\neg\beta \rightarrow \neg\alpha) \rightarrow (\alpha \rightarrow \beta))$$

### Rules of inference

Only one: **modus ponens**

(for any  $\alpha, \beta \in \text{Form}(\mathcal{L}_0)$ )

**MP** From  $\alpha$  and  $(\alpha \rightarrow \beta)$  infer  $\beta$ .

## 6.2 Definition

For any  $\Gamma \subseteq \text{Form}(\mathcal{L}_0)$  we say that  $\alpha$  is **deducible** (or **provable**) from the hypotheses  $\Gamma$  if there is a finite sequence  $\alpha_1, \dots, \alpha_m \in \text{Form}(\mathcal{L}_0)$  such that for each  $i = 1, \dots, m$  either

- (a)  $\alpha_i$  is an axiom, or
- (b)  $\alpha_i \in \Gamma$ , or
- (c) there are  $j < k < i$  such that  $\alpha_i$  follows from  $\alpha_j, \alpha_k$  by MP,  
i.e.  $\alpha_j = (\alpha_k \rightarrow \alpha_i)$  or  $\alpha_k = (\alpha_j \rightarrow \alpha_i)$

AND

- (d)  $\alpha_m = \alpha$ .

The sequence  $\alpha_1, \dots, \alpha_m$  is then called a **proof** or **deduction** or **derivation** of  $\alpha$  from  $\Gamma$ .

Write  $\Gamma \vdash \alpha$ .

If  $\Gamma = \emptyset$  write  $\vdash \alpha$  and say that  $\alpha$  is a **theorem** (of the system  $L_0$ ).

### 6.3 Example For any $\phi \in \text{Form}(\mathcal{L}_0)$

$$(\phi \rightarrow \phi)$$

is a theorem of  $L_0$ .

*Proof:*

$$\alpha_1 \quad (\phi \rightarrow (\phi \rightarrow \phi))$$

[A1 with  $\alpha = \beta = \phi$ ]

$$\alpha_2 \quad (\phi \rightarrow ((\phi \rightarrow \phi) \rightarrow \phi))$$

[A1 with  $\alpha = \phi$ ,  $\beta = (\phi \rightarrow \phi)$ ]

$$\alpha_3 \quad ((\phi \rightarrow ((\phi \rightarrow \phi) \rightarrow \phi)) \rightarrow \\ \rightarrow ((\phi \rightarrow (\phi \rightarrow \phi)) \rightarrow (\phi \rightarrow \phi)))$$

[A2 with  $\alpha = \phi$ ,  $\beta = (\phi \rightarrow \phi)$ ,  $\gamma = \phi$ ]

$$\alpha_4 \quad ((\phi \rightarrow (\phi \rightarrow \phi)) \rightarrow (\phi \rightarrow \phi))$$

[MP  $\alpha_2, \alpha_3$ ]

$$\alpha_5 \quad (\phi \rightarrow \phi)$$

[MP  $\alpha_1, \alpha_4$ ]

Thus,  $\alpha_1, \alpha_2, \dots, \alpha_5$  is a deduction of  $(\phi \rightarrow \phi)$  in  $L_0$ .

□

## 6.4 Example

For any  $\phi, \psi \in \text{Form}(\mathcal{L}_0)$ :

$$\{\phi, \neg\phi\} \vdash \psi$$

*Proof:*

$$\alpha_1 (\neg\phi \rightarrow (\neg\psi \rightarrow \neg\phi))$$

[A1 with  $\alpha = \neg\phi, \beta = \neg\psi$ ]

$$\alpha_2 \neg\phi [\in \Gamma]$$

$$\alpha_3 (\neg\psi \rightarrow \neg\phi) [\text{MP } \alpha_1, \alpha_2]$$

$$\alpha_4 ((\neg\psi \rightarrow \neg\phi) \rightarrow (\phi \rightarrow \psi))$$

[A3 with  $\alpha = \phi, \beta = \psi$ ]

$$\alpha_5 (\phi \rightarrow \psi) [\text{MP } \alpha_3, \alpha_4]$$

$$\alpha_6 \phi [\in \Gamma]$$

$$\alpha_7 \psi [\text{MP } \alpha_5, \alpha_6]$$

□

## 6.5 The Soundness Theorem for $L_0$

$L_0$  is **sound**, i.e. for any  $\Gamma \subseteq \text{Form}(\mathcal{L}_0)$  and for any  $\alpha \in \text{Form}(\mathcal{L}_0)$ :

if  $\Gamma \vdash \alpha$  then  $\Gamma \models \alpha$ .

*In particular, any theorem of  $L_0$  is a tautology.*

*Proof:*

Assume  $\Gamma \vdash \alpha$  and let  $\alpha_1, \alpha_2, \dots, \alpha_m = \alpha$  be a deduction of  $\alpha$  in  $L_0$ .

Let  $v$  be any valuation such that  $\tilde{v}(\phi) = T$  for all  $\phi \in \Gamma$ .

We have to show that  $\tilde{v}(\alpha) = T$ .

We show by induction on  $i \leq m$  that

$$\tilde{v}(\alpha_1) = \dots = \tilde{v}(\alpha_i) = T \quad (\star)$$

**$i = 1$**

either  $\alpha_1$  is an axiom, so  $\tilde{v}(\alpha_1) = T$  or  $\alpha_1 \in \Gamma$ ,  
so, by hypothesis,  $\tilde{v}(\alpha_1) = T$ .

### **Induction step**

Suppose  $(\star)$  is true for some  $i < m$ .  
Consider  $\alpha_{i+1}$ .

Either  $\alpha_{i+1}$  is an axiom or  $\alpha_{i+1} \in \Gamma$ ,  
so  $\tilde{v}(\alpha_{i+1}) = T$  as above,

or else there are  $j \neq k < i + 1$  such that  
 $\alpha_j = (\alpha_k \rightarrow \alpha_{i+1})$ .

By induction hypothesis

$$\tilde{v}(\alpha_k) = \tilde{v}(\alpha_j) = \tilde{v}((\alpha_k \rightarrow \alpha_{i+1})) = T.$$

But then, by tt  $\rightarrow$ ,  $\tilde{v}(\alpha_{i+1}) = T$   
(since  $T \rightarrow F$  is  $F$ ).

□

For the proof of the converse

## Completeness Theorem

*If  $\Gamma \models \alpha$  then  $\Gamma \vdash \alpha$ .*

we first prove

### 6.6 The Deduction Theorem for $L_0$

*For any  $\Gamma \subseteq \text{Form}(\mathcal{L}_0)$  and  
for any  $\alpha, \beta \in \text{Form}(\mathcal{L}_0)$ :*

*if  $\Gamma \cup \{\alpha\} \vdash \beta$  then  $\Gamma \vdash (\alpha \rightarrow \beta)$ .*