

## Mathematical physiology

### PROBLEM SHEET 4.

1. The delayed logistic equation is

$$\dot{x} = \alpha x(1 - x_1), \quad x_1 = x(t - 1).$$

Show that the steady solution  $x = 1$  is oscillatorily unstable if  $\alpha > \frac{1}{2}\pi$ .

When  $\alpha$  is large (a value  $\alpha > 3$  is sufficient), the resulting oscillations take on the appearance of periodic isolated pulses. Find an approximation to this solution by assuming  $x \approx e^{\alpha t}$  for  $t < 0$ , and integrating forward using the method of steps to find the solution in  $0 < t < 1$ , then in  $1 < t < 2$  and finally in  $2 < t < 3$ . [You should write the solution in terms of  $\phi = \ln x$ .]

Show that, approximately,

$$\phi = \alpha t - e^{\alpha(t-1)}, \quad 0 < t < 1;$$

using this, show that

$$\phi \approx \alpha t + e^{\alpha} [e^{-e^{\alpha(t-2)}} - 1], \quad 1 < t < 2,$$

and using this, show that

$$\phi \approx \alpha t + e^{\alpha} [e^{-e^{\alpha(t-2)}} - 1], \quad t > 2,$$

where the last approximation is valid for  $t < 2.5$ .

Show that for  $t \gtrsim 2$ ,

$$\phi \approx \alpha(t - P), \quad P = \frac{e^{\alpha}}{\alpha},$$

and deduce that the solution is periodic with approximate period  $P$ .

Show that the maximum and minimum values of  $x$  are

$$x_{max} \approx e^{\alpha-1}, \quad x_{min} \approx \alpha e^{2\alpha-e^{\alpha}}.$$

Sketch the form of the solution for  $x(t)$ . Why is the approximate solution already good for  $\alpha = 3$ ?

2. In respiratory physiology, what is meant by the *minute ventilation*? Describe the way in which respiration is controlled by the blood gas concentrations at the central and peripheral chemoreceptors.

The Mackey-Glass model is a one compartment model of respiratory control, and can be represented by the equations

$$K\dot{p} = M - p\dot{V},$$

$$\dot{V} = \dot{V}(p_\tau);$$

explain what the various terms represent, and their physiological interpretation.

Suppose that

$$\dot{V} = G[p - p_0]_+,$$

and that  $M = 200 \text{ mm Hg l(BTPS) min}^{-1}$ ,  $p_0 = 35 \text{ mm Hg}$ ,  $K = 40 \text{ l(BTPS)}$ ,  $G = 2 \text{ l(BTPS) min}^{-1} \text{ mm Hg}^{-1}$ ,  $\tau = 0.2 \text{ min}$ . Show how to non-dimensionalise the equations to obtain the dimensionless form

$$\dot{p} = \alpha[1 - (1 + \mu p)v],$$

$$v = [p_1]_+,$$

and give the definitions of  $\alpha$  and  $\mu$ . Check that they are dimensionless, and find their values.

3. A simplified version of the Grodins model describes  $\text{CO}_2$  partial pressures in arteries, veins, brain and tissues by the equations

$$\begin{aligned} K_L \dot{P}_{\text{aCO}_2} &= -\dot{V} P_{\text{aCO}_2} + 863 K_{\text{CO}_2} Q [P_{\text{vCO}_2} - P_{\text{aCO}_2}], \\ K_{\text{CO}_2} K_B \dot{P}_{\text{BCO}_2} &= MR_{\text{BCO}_2} + K_{\text{CO}_2} Q_B [P_{\text{aCO}_2} (t - \tau_{\text{aB}}) - P_{\text{BCO}_2}] \\ K_{\text{CO}_2} K_T \dot{P}_{\text{TCO}_2} &= MR_{\text{TCO}_2} + (Q - Q_B) K_{\text{CO}_2} [P_{\text{aCO}_2} (t - \tau_{\text{aT}}) - P_{\text{TCO}_2}], \end{aligned}$$

with the venous pressure being determined by

$$Q P_{\text{vCO}_2} = Q_B P_{\text{BCO}_2} (t - \tau_{\text{vB}}) + (Q - Q_B) P_{\text{TCO}_2} (t - \tau_{\text{vT}}).$$

Explain the meaning of the equations and their constituent terms.

Use values  $K_L = 3 \text{ l}$ ,  $\dot{V} \sim V^* = 5 \text{ l min}^{-1}$ ,  $863 K_{\text{CO}_2} Q = 26 \text{ l min}^{-1}$ ,  $K_B = 1 \text{ l}$ ,  $Q = 6 \text{ l min}^{-1}$ ,  $Q_B = 0.75 \text{ l min}^{-1}$ ,  $K_T = 39 \text{ l}$ , to evaluate response time scales for arterial, brain and tissue  $\text{CO}_2$  partial pressures.

Deduce that for oscillations on a time scale of a minute, one can assume that the arterial pressure is in quasi-equilibrium, and that the tissue (and thus also venous) partial pressures are approximately constant.

Hence derive an approximate expression for  $P_{\text{aCO}_2}$  in terms of the ventilation  $\dot{V}$ .

4. Red blood cell precursors are produced from pluripotential stem cells in the bone marrow at a rate  $F$ . They mature for a period of  $\tau$  days before being released into the blood, where they circulate for a further  $A$  days. If the apoptotic rates in bone marrow and blood are  $\delta$  and  $\gamma$ , respectively, show that the developing cell density  $p$  and circulating RBC density  $e$  satisfy the equations

$$\frac{\partial p}{\partial t} + \frac{\partial p}{\partial m} = -\delta p,$$

$$\frac{\partial e}{\partial t} + \frac{\partial e}{\partial a} = -\gamma e,$$

for  $0 < m < \tau$  and  $0 < a < A$ , where

$$p(t, 0) = F[E(t)], \quad e(t, 0) = p(t, \tau),$$

and we assume  $F$  depends on the total circulating blood cell population,

$$E = \int_0^A e \, da.$$

Solve the equations using the method of characteristics, and hence show that for  $t > \tau + A$ ,  $E$  satisfies

$$\dot{E} = F[E_\tau]e^{-\delta\tau} - F[E_{A+\tau}]e^{-\delta\tau-\gamma A} - \gamma E, \quad t > \tau + A.$$

Compare this model to that which assumes no age limit to the circulating RBC. Under what circumstances does the model reduce to the no age limit model?

Suppose that  $F = F_0 f$ , where  $f$  is  $O(1)$  and is a positive monotone decreasing function. Show how to non-dimensionalise the model to the form

$$\dot{E} = \mu[f(E_1) - f(E_{\Lambda+1})e^{-\mu\Lambda} - E],$$

where  $\mu = \gamma\tau$  and  $\Lambda = A/\tau$ . Supposing that  $A = 120$  days and  $\tau = 6$  days, explain why you might expect  $\mu$  to be small.

Write down an equation for the exponent  $\sigma$  in solutions  $\propto \exp(\sigma t)$  describing small perturbations about the steady state, and show that the steady state is stable if  $|f'| < \frac{1}{2}$ .

5. The Hopf bifurcation curve for the equation

$$\sigma = -\alpha - \Gamma e^{-\sigma}$$

is defined parametrically by

$$\Gamma_0(\alpha) = \frac{\Omega}{\sin \Omega}, \quad \alpha = -\frac{\Omega}{\tan \Omega},$$

for  $\Omega \in (0, \pi)$ . By expanding for small  $\Omega$ , show that

$$\Gamma_0 = 1 + \frac{1}{6}\Omega^2 + \frac{7}{360}\Omega^4 + \frac{31}{15120}\Omega^6 + O(\Omega^8),$$

and that

$$\alpha + 1 = \frac{1}{3}\Omega^2 + \frac{1}{45}\Omega^4 + \frac{2}{945}\Omega^6 + O(\Omega^8).$$

Deduce that

$$\Gamma_0 = 1 + \frac{1}{2}(\alpha + 1) + \frac{3}{40}(\alpha + 1)^2 - \frac{9}{2800}(\alpha + 1)^3 + O[(\alpha + 1)^4],$$

i. e.,

$$\Gamma_0 \approx 1 + 0.5(\alpha + 1) + 0.075(\alpha + 1)^2 - 0.0032(\alpha + 1)^3 + O[(\alpha + 1)^4].$$

Plot  $\Gamma_0$  versus  $\alpha$  using suitable graphical software, and show that an accurate quadratic approximation for  $\alpha \in (-1, 2)$  is

$$\Gamma_0 \approx 1 + 0.5(\alpha + 1) + 0.058(\alpha + 1)^2,$$

and that an accurate cubic approximation for  $\alpha \in (-1, 5)$  is

$$\Gamma_0 \approx 1 + 0.5(\alpha + 1) + 0.075(\alpha + 1)^2 - 0.005(\alpha + 1)^3.$$

Can you find a value of  $c$  for which

$$\Gamma_0 \approx 1 + 0.5(\alpha + 1) + 0.075(\alpha + 1)^2 - 0.0032(\alpha + 1)^3 + c(\alpha + 1)^4$$

provides an accurate approximation for larger values of  $\alpha$ ?

Show (plot it and compare with the quadratic approximation) that a better two coefficient approximation is given by

$$\Gamma_0 = \frac{1 + b(\alpha + 1) + c(\alpha + 1)^2}{1 + c(\alpha + 1)},$$

if  $b = 0.65$  and  $c = 0.15$ . Why would these values of  $b$  and  $c$  be chosen? Show (graphically) that an even better approximation is obtained if  $b = 0.69$  and  $c = 0.3$ . (The maximum error for  $\alpha < 100$  is less than 0.05.) How could you extend this type of approximation to larger values of  $\alpha$ ?

6. In a model for the evolution of the resting phase cells in a blood cell maturation model, the cell density  $M$  is given by

$$\frac{\partial M}{\partial t} + \frac{\partial M}{\partial \xi} = -RM + Q,$$

where  $\xi$  is the maturation variable,

$$Q = \begin{cases} 2e^{-\gamma\tau}R[t - \tau, \xi - \tau]M[t - \tau, \xi - \tau], & \xi > \tau, \quad t > \tau, \\ 2e^{-(\gamma_0 + V_0)\tau}e^{(\gamma_0 + V_0 - \gamma)\xi}V_0R_0(t - \tau)N_0(t - \tau), & \xi < \tau, \quad t > \xi, \end{cases}$$

If  $R = (1 + \lambda)R_0$ ,  $\lambda = 2e^{-\gamma\tau} - 1$ ,  $\gamma_0 + V_0 = \gamma$ , and all these quantities and also  $N_0$  are constant, then the equation for  $M$  can be written as

$$\frac{\partial M}{\partial t} + \frac{\partial M}{\partial \xi} = -RM + (1 + \lambda)RM_{\tau, \tau},$$

with initial data being

$$M = M_0 = N_0V_0 \quad \text{on} \quad \xi = 0 \quad \text{and} \quad t > \xi.$$

By careful consideration of how the characteristic equations are solved, show that for  $t > \xi$ ,  $M = M(\xi) \equiv M_0 u[(\xi - \tau)/\tau]$ , where  $u(\eta)$  satisfies

$$\frac{du}{d\eta} = -\alpha u - \Gamma u_1,$$

and  $\alpha = R\tau$ ,  $\Gamma = -(1 + \lambda)R\tau$ , and  $u = 1$  for  $\eta \in [-1, 0)$ .

By taking the Laplace transform of the equation (exercising due care with the delayed term), show that the Laplace transform  $U(p)$  of  $u$  is given by

$$U(p) = \frac{h(p)}{f(p)},$$

where

$$f(p) = p + \alpha + \Gamma e^{-p}$$

and

$$h(p) = 1 - \Gamma \left( \frac{1 - e^{-p}}{p} \right).$$

Deduce that  $U$  can also be written as

$$U(p) = \frac{\Lambda e^p}{p[(p + \alpha)e^p + \Gamma]} + \frac{1}{p},$$

where

$$\Lambda = \lambda R\tau.$$

Hence show that if the inversion contour for  $U$  is completed as a square with upper and lower sides at  $\text{Im } p = \pm(n + \frac{1}{2})\pi$ , with  $n$  even for  $\Gamma < 0$ , as here, then by taking the limit as  $n \rightarrow \infty$ ,  $u$  can be found as

$$u = \sum_{j=-\infty}^{\infty} c_j \exp(p_j \eta),$$

where  $p_j$  are the zeros of  $f(p)$  ( $p_0$  is the real root, and  $p_{-j}$  is the complex conjugate of  $p_j$ ). Write down the definition of the constants  $c_j$  in terms of  $p_j$ , and show that they can be expressed as

$$c_j = \frac{\Lambda}{p_j (1 + \alpha + p_j)},$$

so that  $c_j = O(1/j^2)$  for  $j \gg 1$ .