

**Problem Sheet 1**

**QUESTION 1. Uniqueness of Solutions to ODEs.** Let  $H$  be a real Hilbert space endowed with the scalar product  $(\cdot, \cdot)$ . Show that the initial value problem for  $y: \mathbb{R} \rightarrow H$ , given by

$$y'(t) = f(t, y(t)) \text{ for } t > 0, \quad y(0) = y_0,$$

has at most one continuously differentiable solution on the interval  $[0, T]$ , provided that  $f: \mathbb{R} \times H \rightarrow H$  is continuous and satisfies for some  $L > 0$

$$(1) \quad (f(t, y) - f(t, z), y - z) \leq L\|y - z\|^2 \text{ for all } y, z \in H.$$

[Hint: Use the product rule  $\frac{d}{dt}(y(t), z(t)) = (y'(t), z(t)) + (z'(t), y(t))$  for functions  $y, x: \mathbb{R} \rightarrow H$  and Gronwall's Lemma.]

Give furthermore an example of a function  $f$  for which (1) is satisfied but for which the Lipschitz-condition of Picard's theorem does not hold.

**QUESTION 2. Euler-Lagrange Equations.**

(i) Let  $p \in (1, \infty)$  and  $\Omega \subset \mathbb{R}^n$  be a domain. Derive the Euler-Lagrange equation for the functional

$$I(v) = \int_{\Omega} \frac{1}{p} |\nabla v|^p - \frac{1}{4} v^4 dx$$

where  $v: \Omega \rightarrow \mathbb{R}$  and  $|\nabla v| = \sqrt{(\partial_1 v)^2 + \dots + (\partial_n v)^2}$  once by using the formula derived in the lecture and once by direct computation of  $\frac{d}{dt} I(v + t\phi)$ ,  $\phi \in C_c^\infty(\Omega)$ .

(ii) Let  $\Omega \subset \subset \mathbb{R}^3$  and  $1 \leq p \leq 6$ . Show that the functional

$$E(u) := \frac{\|\nabla u\|_{L^2}^2}{\|u\|_{L^p}^2}$$

is well defined for all  $u \in H_0^1(\Omega)$ ,  $u \neq 0$  and satisfies  $\inf\{E(u) : u \in H_0^1(\Omega)\} > 0$ . Derive furthermore its Euler-Lagrange equation.

Then consider

$$E_0(u) := \int |\nabla u|^2 dx$$

and explain what condition has to be satisfied for a function  $u \in H_0^1(\Omega)$  which minimises  $E_0$  in the set  $M := \{v : \|v\|_{L^p} = 1\}$

**QUESTION 3. Counter-example to Brouwer's Fixed Point Theorem in an infinite dimensional space.** Consider the real Hilbert Space

$$l^2 = \left\{ (x_i)_{i \in \mathbb{N}} \text{ such that } \sum_{i=0}^{\infty} x_i^2 < \infty \right\} \text{ with the norm } \|x\|_{l^2} = \sqrt{\sum_{i=0}^{\infty} x_i^2}.$$

Let  $B$  be its closed unit ball.

- Consider the map

$$T: B \rightarrow B \text{ given by } T(x) = (\sqrt{1 - \|x\|_{l^2}^2}, x_0, x_1, x_2, \dots).$$

Show that  $T$  is continuous and does not have a fixed point.

- Construct a continuous retraction from  $B$  to  $\partial B$ .

**QUESTION 4. Application of Brouwer's FPT.** Given a map  $f \in C(\mathbb{R}^n : \mathbb{R}^n)$  such that  $|f(x)| \leq a + b|x|$ , with  $a \geq 0$  and  $0 < b < 1$ , show that  $f$  has a fixed point.