

Elliptic Curves. HT 2018/19. Sheet 2.

1. Let K be a field with non-Archimedean valuation $|\cdot|$.
 - (a). For any $x, y \in K$ show that, if $|x| \neq |y|$ then $|x \pm y| = \max(|x|, |y|)$.
 - (b). If $x_1, \dots, x_n \in K$ and if there exists ℓ such that $|x_\ell| > |x_i|$ for all $i \neq \ell$, then show that $|x_1 + \dots + x_n| = |x_\ell|$.
 - (c). Suppose that $s_n \rightarrow s$ in $K, |\cdot|$. Show that $|s_n| \rightarrow |s|$ in $\mathbb{R}, |\cdot|_\infty$. When $s \neq 0$, show that there exists N such that, for all $n > N$, $|s_n| = |s|$.
- 2(a). Find: $|3/50|_5, |3/50|_3, |3/50|_7, d_5(2/3, 1/5), d_7(2/3, 1/5), d_{11}(2/3, 1/5)$.
 - (b). Describe $|3/7|_p$ for all p . What is the product $\prod |3/7|_i$, taken over $i = p$, for all primes p , and $i = \infty$? Given any $x \in \mathbb{Q} (x \neq 0)$, what is $\prod |x|_i$?
3. Which of the following are convergent in \mathbb{Q}_5 ?
 - (a). $1/5^n$. (b). n . (c). $n!$ (d). $3 + 10^n$. (e). $\sum_0^\infty 10^n$. (f). $\sum_0^\infty 7^n$.
4. For each p, m, r , either find an $x \in \mathbb{Z}$ such that $|x^2 - r|_p \leq p^{-m}$ or show that no such x exists.
 - (a). $p = 5, r = -1, m = 4$. (b). $p = 3, r = 7/8, m = 7$. (c). $p = 5, r = 5/4, m = 4$.
5. Find the 7-adic expansion of each of: 200 and $3/14$. Determine the member of \mathbb{Q} expressed by the 5-adic expansion $2, \overline{34}$.
6. Let $x \in \mathbb{Q}$. Show that $x \in \mathbb{Z} \iff (x \in \mathbb{Z}_p \text{ for all } p)$.
7. Decide whether there exists $x \in \mathbb{Q}_p$ such that $x^2 = -28$ for each of: $p = 2, 3, 5, 7, 11$.
8. Show that $(X^2 - 2)(X^2 - 17)(X^2 - 34)$ has a root in \mathbb{R} and in every \mathbb{Q}_p , but not in \mathbb{Q} .
9. Is 4 a cube in \mathbb{Q}_3 ? Is 28 a cube in \mathbb{Q}_3 ? Is 13 a cube in \mathbb{Q}_7 ?