## Prelims Introductory Calculus MT 2019: Sheet 6

1. Evaluate

(a) 
$$\int_{x=0}^{x=1} \int_{y=0}^{y=3-3x} dy dx$$
, (b)  $\int_{y=0}^{y=1} \int_{x=0}^{x=\sqrt{1-y^2}} dx dy$ , (c)  $\int_{x=0}^{x=4} \int_{y=0}^{y=\sqrt{x}} dy dx$ .

In each case, sketch the area of integration.

- 2. Use a double integral to find the area of the region bounded by the curves  $y^2 = 4x$  and 2x + y = 4.
- 3. Let a > 0. Sketch the curve

$$x^{2/3} + y^{2/3} = a^{2/3}$$

in the quadrant x, y > 0.

The variables u and v are given in terms of x and y by

$$x = u\cos^3 v, \quad y = u\sin^3 v.$$

What is the equation of the curve in terms of the new coordinates u and v?

Calculate the Jacobian  $\frac{\partial(x,y)}{\partial(u,v)}$  and hence find the area of the region bounded by the curve and the positive x- and y-axes.

4. Let a > 0. The curve with polar equation

$$r = a(1 + \cos\theta), \quad 0 \le \theta < 2\pi$$

is called a cardioid. Sketch the curve and, using a double integral, show that the area bounded by it equals  $\frac{3}{2}\pi a^2$ .

5. Calculate the scalar line integral of the vector field

$$\mathbf{F}(\mathbf{r}) = (2xy^2 - 3yz + 1, 2yx^2 - 3xz, -3xy)$$

along the path consisting of the straight-line segment joining the origin to the point (a, b, c).

6. Find the length of the curve given by

$$x = t$$
,  $y = -\ln(\cos t)$ ,  $0 \le t \le \pi/4$ .