

# INTEGRATION: H.T. 2020, 16 LECTURES

## Reading

- Z. Qian, *Part A: Integration*, Maths Institute Website  
M. Capinski and E. Kopp, *Measure, Integration and Probability*, Springer SUMS (2nd edition, 2004)  
H.A. Priestley, *Introduction to Integration*, OUP, 1997  
E. M. Stein & R. Shakarchi, *Real Analysis: Measure Theory, Integration and Hilbert Spaces*, Princeton Lectures in Analysis III, Princeton University Press, 2005  
D.J.H. Garling, *A Course in Mathematical Analysis, III (Part 6)*, CUP, 2014.

Qian's notes were written for the course as he gave it in 2014-17, based on previous versions of the course given by Alison Etheridge and myself. I will cover more or less the same material, but I will not follow his notes exactly.

Capinski and Kopp is the most basic of the books, giving the theory in a basic style, but with not many worked examples; I shall follow rather closely their approach to the theory. Priestley adopts a very different approach to the construction of the integral, so early parts of her book look quite different from what I shall do, but about the 8th lecture onward everything comes together; she has lots of worked examples.

Stein and Shakarchi, and Garling, are a little more sophisticated in the theory. Garling's book is based on lectures given in Cambridge, and it has a good number of worked examples.

Numerous other useful books may be found in libraries. Some may adopt different approaches to the construction of the integral, but when they talk about Lebesgue integration they all mean the same class of integrable functions and the same theorems.

## INTRODUCTION

In Prelims, you saw how to define  $\int_a^b f(x) dx$  for a continuous function  $f : [a, b] \rightarrow \mathbb{R}$  or more generally for Riemann integrable  $f$ . It had some good properties: the Fundamental Theorem of Calculus shows that it is more or less an inverse of differentiation, leading to rigorous statements concerning A level calculus. Moreover you saw that

$$(*) \quad \lim_{n \rightarrow \infty} \int_a^b f_n(x) dx = \int_a^b f(x) dx$$

if  $(f_n)$  converges to  $f$  uniformly on  $[a, b]$ . This was useful (a) for integrating power series term-by-term, (b) for finding  $\lim_{n \rightarrow \infty} \int_\gamma f_n(z) dz$ , where  $\gamma$  is a contour of finite length, in complex analysis last term. However, the Riemann integral has various deficiencies:

- (a) There are still functions which one feels one should be able to integrate, for which the Prelims definition fails to work. For example, let  $f = \chi_{\mathbb{Q} \cap [0,1]}$  be the characteristic function of  $\mathbb{Q} \cap [0,1]$ . Then

$$\int_0^1 f(x) dx = 0, \quad \overline{\int_0^1 f(x) dx} = 1$$

so the definition of the integral fails.

In particular, if we want to define the length of a subset  $E$  of  $\mathbb{R}$  by

$$m(E) = \int \chi_E(x) dx,$$

we need to extend the definition of integrals in some way beyond Riemann integration.

- (b) There is a lack of theorems saying that

$$f_n \rightarrow f \implies \int f_n(x) dx \rightarrow \int f(x) dx$$

particularly for integrals over  $\mathbb{R}$  or unbounded subsets of  $\mathbb{R}$ . To some extent, this is unavoidable because of the following example:

**Example 0.1.** Let  $f_n(x) = n^2 x^n (1-x)$  ( $0 \leq x \leq 1$ ). Then  $\lim_{n \rightarrow \infty} f_n(x) = 0$  for all  $x \in [0,1]$ , but  $\lim_{n \rightarrow \infty} \int_0^1 f_n(x) dx = 1$ .

This example is going to arise in any reasonable theory. But we would like some more theorems of the form

*Suppose  $(f_n)$  is a sequence of integrable functions,  $f_n(x) \rightarrow f(x)$  for each  $x$ , and [supplementary assumptions to be inserted]. Then  $f$  is integrable and  $\int f(x) dx = \lim_{n \rightarrow \infty} \int f_n(x) dx$ .*

Lebesgue's integration theory provides two very powerful theorems of this form (Monotone Convergence Theorem, Dominated Convergence Theorem). The theorems are less good in Riemann integration, because one has to assume that the limiting function is integrable.

- (c) Riemann's integration theory does not generalise to include various other contexts such as:
- probability theory, taking expectations of arbitrary random variables (continuous, discrete, hybrid, singular);
  - summing infinite series.

Lebesgue's theory resolves these difficulties, except where there is an unavoidable obstruction. In a sense the passage from Riemann integration to Lebesgue integration resembles the passage from rational numbers to real numbers—it completes the space of integrable functions, or it fills in the gaps.

The crucial ideas of the Lebesgue's construction are:

- (i) Instead of using integrals to define lengths of sets, define the length of a set directly; then define integrals.
- (ii) Instead of partitioning the  $x$ -axis into intervals and using step functions, partition the  $y$ -axis into intervals and considering corresponding “simple” functions.

As I said, there are other ways of constructing Lebesgue's integral on  $\mathbb{R}$ , including ways which use step functions (see Priestley), but they don't generalise so easily to probability (for example). Once one gets the Monotone Convergence Theorem, then everything is the same, however you got there. We then get a whole host of theorems about:

- passing limits through integrals,
- passing infinite sums through integrals,
- differentiating through integrals,
- interchanging two integrals (Fubini's Theorem)
- changing variables.

Note that these processes do not always work—there are simple counterexamples for the first 4! So all these theorems have conditions which must be checked before using in applications. In this course, we do not take the position that you can just assume all these processes work. On the other hand, we shall not go pedantically through all details of the construction of the integral and the proofs of the theorems. I shall present the construction in a way which generalises easily, but the proofs are often not interesting. The construction up to the MCT will take some time - around 8 lectures - and then useful theorems and applications will come thick and fast.

Please be aware that all the Prelims theory remains valid in this context. Lebesgue integration theory extends Riemann's theory by enabling you to integrate more functions. In particular, the Fundamental Theorem of Calculus (both versions), Integration by Parts and Substitution remain valid under the assumptions given in Prelims.

## 1. EXTENDED REAL NUMBER SYSTEM

In this course, we shall often take infinite series of non-negative terms and limits of (monotone) sequences. In order to avoid complications concerning divergence, it will be convenient to work in the extended real numbers including  $-\infty$  and  $\infty$ , and to use the notions of  $\limsup$  and  $\liminf$ .

Thus we consider the set  $[-\infty, \infty] = \mathbb{R} \cup \{-\infty, \infty\}$ . Addition and multiplication by  $\infty$  are defined as follows (for  $x \in \mathbb{R}$ ):

$$\begin{aligned} x + \infty &= \infty + x = \infty, \\ x - \infty &= -\infty + x = -\infty, \\ x \cdot \infty &= \infty \cdot x = (-x) \cdot (-\infty) = \begin{cases} \infty & (x > 0), \\ -\infty & (x < 0), \\ 0 & (x = 0). \end{cases} \end{aligned}$$

Note that

- $\infty - \infty$  is undefined;
- the usual laws (commutativity, associativity and distributivity) apply, provided that the relevant expressions are defined;
- the above are uncontroversial, except for  $0 \cdot \infty = 0$  which is convenient for our particular context but might be inappropriate in other mathematical contexts.

The ordering on  $[-\infty, \infty]$  is the obvious one, and  $\lim_{n \rightarrow \infty} a_n = \infty$  has the same meaning as in Prelims Analysis.

In this system, any subset  $E$  has a supremum and an infimum in  $[-\infty, \infty]$ . Note that  $\sup \emptyset = -\infty$ . If  $E \subseteq \mathbb{R}$ ,  $\sup E = \infty$  if and only if  $E$  is not bounded above. For an increasing sequence  $(a_n)$ ,  $\lim_{n \rightarrow \infty} a_n = \sup\{a_n\}$ . If  $a_n \geq 0$  for all  $n$ , then  $\sum a_n = \infty$  if and only if the series diverges.

**Proposition 1.1.** 1. Let  $(a_n)$  be a sequence of non-negative terms. Then

$$\sum_{n=1}^{\infty} a_n = \sup \left\{ \sum_{n \in J} a_n : J \text{ finite subset of } \mathbb{N} \right\}.$$

2. Let  $(b_{mn})_{m,n \geq 1}$  be a double sequence of non-negative terms, and  $\{(m_k, n_k) : k \geq 1\}$  be any enumeration of  $\mathbb{N} \times \mathbb{N}$ . Then

$$\sum_{m=1}^{\infty} \sum_{n=1}^{\infty} b_{mn} = \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} b_{mn} = \sum_{k=1}^{\infty} b_{m_k, n_k} = \sup \left\{ \sum_{(m,n) \in J} b_{mn} : J \text{ finite subset of } \mathbb{N} \times \mathbb{N} \right\}.$$

In particular, Proposition 1.1 implies that  $\sum a_n$  is independent of the order of the terms, and similarly  $\sum \sum b_{mn}$  can be arbitrarily rearranged.

A bounded sequence  $(a_n)$  in  $\mathbb{R}$  may not have a limit. It has a supremum and infimum, but for some large values of  $n$ ,  $a_n$  may not be close to them. Think for example about  $a_n = (1 + 1/n) \sin n$ . Asymptotically the values oscillate between  $-1$  and  $1$ , but there are infinitely many values bigger than  $1$  and infinitely many smaller than  $-1$ .

For a sequence  $(a_n)$  in  $[-\infty, \infty]$ , define

$$\begin{aligned} \limsup_{n \rightarrow \infty} a_n &= \lim_{m \rightarrow \infty} \left( \sup_{n \geq m} a_n \right), \\ \liminf_{n \rightarrow \infty} a_n &= \lim_{m \rightarrow \infty} \left( \inf_{n \geq m} a_n \right). \end{aligned}$$

The limits exist, because  $(\sup_{n \geq m} a_n)_{m \geq 1}$  is a decreasing sequence

So,  $\limsup_{n \rightarrow \infty} a_n$  is the largest number  $\ell$  such that there is a subsequence of  $(a_n)$  converging to  $\ell$ .

**Examples 1.2.** 1. Let  $a_n = (1 + 1/n) \sin n$ . Then

$$\limsup_{n \rightarrow \infty} a_n = 1, \quad \liminf_{n \rightarrow \infty} a_n = -1.$$

2. Let  $a_n = (-1)^n$ . Then

$$\limsup_{n \rightarrow \infty} a_n = 1, \quad \liminf_{n \rightarrow \infty} a_n = -1.$$

3. Let  $a_n = n(-1)^n$ . Then

$$\limsup_{n \rightarrow \infty} a_n = \infty, \quad \liminf_{n \rightarrow \infty} a_n = -\infty.$$

4. Let  $a_n = \begin{cases} 1 + 2^{-n} & (n \text{ prime}), \\ 0 & \text{otherwise.} \end{cases}$  Then

$$\limsup_{n \rightarrow \infty} a_n = 1, \quad \liminf_{n \rightarrow \infty} a_n = 0.$$

- Proposition 1.3.** 1.  $\liminf_{n \rightarrow \infty} a_n = -\limsup_{n \rightarrow \infty} (-a_n)$ ;  
 2.  $\liminf_{n \rightarrow \infty} a_n \leq \limsup_{n \rightarrow \infty} a_n$ ;  
 3.  $\lim_{n \rightarrow \infty} a_n$  exists if and only if  $\liminf_{n \rightarrow \infty} a_n = \limsup_{n \rightarrow \infty} a_n$ ; then all are equal;  
 4. If  $a_n \leq b_n$  for all  $n$ , then  $\limsup_{n \rightarrow \infty} a_n \leq \limsup_{n \rightarrow \infty} b_n$ ;  
 5.  $\limsup_{n \rightarrow \infty} (a_n + b_n) \leq \limsup_{n \rightarrow \infty} a_n + \limsup_{n \rightarrow \infty} b_n$  (if all sums exist).

$\limsup$  and  $\liminf$  are useful for avoiding epsilontics. For example, consider the Sandwich Rule, i.e., suppose that  $a_n \leq b_n \leq c_n$  for all  $n$  and  $\lim a_n = \lim c_n$ . Then

$$\begin{aligned} \limsup b_n &\leq \limsup c_n && \text{(Proposition 1.3(4))} \\ &= \lim c_n && \text{(Proposition 1.3(3))} \\ &= \lim a_n && \text{(assumption)} \\ &= \liminf a_n && \text{(Proposition 1.3(3))} \\ &\leq \liminf b_n && \text{(Proposition 1.3(4))} \\ &\leq \limsup b_n && \text{(Proposition 1.3(2)).} \end{aligned}$$

Hence equality holds throughout, so  $\lim b_n = \lim a_n$ , by Proposition 1.3(3).

## 2. LEBESGUE MEASURE

A measure of length for (all) subsets of  $\mathbb{R}$  should be a function  $m : \mathcal{P}(\mathbb{R}) \rightarrow [0, \infty]$  satisfying:

- (i)  $m(\emptyset) = 0$ ,  $m(\{x\}) = 0$ ;
- (ii)  $m(I) = b - a$  if  $I$  is an interval with endpoints  $a, b$ , where  $a < b$ ;
- (iii)  $m(A + x) = m(A)$ ;
- (iv)  $m(\alpha A) = |\alpha| m(A)$ ;
- (v)  $m(A) \leq m(B)$  if  $A \subseteq B$ ; ( $m$  is *monotone*);
- (vi)  $m(A \cup B) = m(A) + m(B)$  if  $A \cap B = \emptyset$  ( $m$  is *finitely additive*);
- (vi)'  $m(\bigcup_{n=1}^{\infty} A_n) = \sum_{n=1}^{\infty} m(A_n)$  if  $A_n \cap A_k = \emptyset$  for  $k \neq n$  ( $m$  is *countably additive*);
- (vii)  $m(\bigcup_{n=1}^{\infty} A_n) = \lim_{n \rightarrow \infty} m(A_n)$  if  $(A_n)$  is an increasing sequence of sets.

In fact, there is very considerable redundancy here. For example, (v), (vi) and (vii) follow from (i) and (vi)'.

The status of (vi)' is perhaps debatable, but it is usually assumed. It is equivalent to (vi) and (vii) together, and (vii) is essential to have a Monotone Convergence Theorem.

Let us attempt to construct such an  $m$ . For  $A \subseteq \mathbb{R}$ , suppose that  $A \subseteq \bigcup_{n=1}^{\infty} I_n$  for intervals  $I_n$ . Letting  $I'_n = I_n \setminus (I_1 \cup \dots \cup I_{n-1})$ , we have

$$m(A) \leq m\left(\bigcup I'_n\right) = \sum m(I'_n) \leq \sum m(I_n).$$

So we attempt to define  $m$  as follows. First, for any interval  $I$  with endpoints  $a$  and  $b$ , define

$$m(I) = b - a.$$

For  $A \subseteq \mathbb{R}$ , we define the *outer measure* of  $A$  to be

$$m^*(A) = \inf \left\{ \sum_{n=1}^{\infty} m(I_n) : I_n \text{ intervals, } A \subseteq \bigcup_{n=1}^{\infty} I_n \right\}.$$

We can always take  $I_n = [-n, n]$ , so the infimum is not over the empty set (but  $m^*(A)$  may be infinite). It makes no difference if we restrict  $I_n$  to being closed intervals, or open intervals.

- Proposition 2.1.** 1.  $m^*(\emptyset) = 0$ ,  $m^*({x}) = 0$ ;  
 2.  $m^*(I) = b - a$  if  $I$  is any interval with endpoints  $a, b$ ;  
 3.  $m^*(A + x) = m^*(x)$ ;  
 4.  $m^*(\alpha A) = |\alpha| m^*(A)$ ;  
 5.  $m^*(A) \leq m^*(B)$  if  $A \subseteq B$ ;  
 6.  $m^*(A \cup B) \leq m^*(A) + m^*(B)$ ;  
 6'.  $m^*(\bigcup_{n=1}^{\infty} A_n) \leq \sum_{n=1}^{\infty} m^*(A_n)$ .

*Proof.* (1), (3), (4), (5) are easy; (6) and (6)' are moderately tricky exercises. Let us prove (2); we will do it for  $I = [a, b]$ ; then the other cases follow using (1), (5) and (6).

Firstly,  $m^*[a, b] \leq b - a$ , because we may take  $I_1 = [a, b]$  and  $I_n = \{0\}$  for  $n \geq 2$ .

Now suppose that  $[a, b] \subseteq \bigcup_{n=1}^{\infty} I_n$  where  $I_n$  is an interval with endpoints  $a_n, b_n$ . Take  $\varepsilon > 0$ . Let

$$J_n = (a_n - \varepsilon 2^{-n}, b_n + \varepsilon 2^{-n}) =: (c_n, d_n).$$

Then  $J_n$  is open and  $[a, b] \subseteq \bigcup_{n=1}^{\infty} J_n$ . By the Heine-Borel Theorem,  $[a, b]$  is compact, so  $[a, b] \subseteq \bigcup_{n=1}^N J_n$  for some  $N$ .

Now it is almost obvious that  $b - a \leq \sum_{n=1}^N m(J_n)$ . Enumerate  $\{c_n, d_n : n = 1, \dots, N\}$  in increasing order:

$$x_1 < x_2 < \dots < x_k.$$

Then  $x_1 < a < b < x_k$ , each interval  $(x_i, x_{i+1})$  is contained in some  $J_n$ , and  $J_n$  has endpoints  $c_n = x_{k_n}$ ,  $d_n = x_{\ell_n}$ , say. Hence

$$b - a < x_k - x_1 = \sum_{i=1}^{k-1} (x_{i+1} - x_i) \leq \sum_{n=1}^N \sum_{i=k_n}^{\ell_n-1} (x_{i+1} - x_i) = \sum_{n=1}^N m(J_n).$$

Now  $\sum_{n=1}^{\infty} m(I_n) = \sum_{n=1}^{\infty} (m(J_n) - 2^{-(n-1)}\varepsilon) > b - a - 2\varepsilon$ . This holds for every  $\varepsilon > 0$ , so  $\sum_{n=1}^{\infty} m(I_n) \geq b - a$ . Hence  $m^*[a, b] \geq b - a$ .  $\square$

A subset  $E$  of  $\mathbb{R}$  is said to be *null* if  $m^*(E) = 0$ .

- Corollary 2.2.** 1. Any subset of a null set is null.  
 2. If  $E_n$  is a null set for  $n = 1, 2, \dots$ , then  $\bigcup_{n=1}^{\infty} E_n$  is null.  
 3. Any countable subset of  $\mathbb{R}$  is null.

*Proof.* **[Direct proof of (2)]** Let  $\varepsilon > 0$ . There exist intervals  $I_{rn}$  such that  $E_n \subseteq \bigcup_{r=1}^{\infty} I_{rn}$  and  $\sum_r m(I_{rn}) < \varepsilon 2^{-n}$ . Now  $\{I_{rn} : r, n = 1, 2, \dots\}$  is a countable family of intervals covering  $\bigcup E_n$ , and  $\sum_n \sum_r m(I_{rn}) < \sum_n \varepsilon 2^{-n} = \varepsilon$ . Hence  $m^*(\bigcup_n E_n) = 0$ .  $\square$

**Example 2.3.** Let  $C_0 = [0, 1]$ ,  $C_1 = [0, \frac{1}{3}] \cup [\frac{2}{3}, 1]$ ,  $C_2 = [0, \frac{1}{9}] \cup [\frac{2}{9}, \frac{1}{3}] \cup [\frac{2}{3}, \frac{7}{9}] \cup [\frac{8}{9}, 1]$ , etc. In general,  $C_n$  is the union of  $2^n$  disjoint closed intervals, each of length  $3^{-n}$ , and  $C_{n+1}$  is obtained from  $C_n$  by deleting the open middle third of each of those intervals.

Let  $C = \bigcap_{n=1}^{\infty} C_n$ . Then  $C$  is a closed subset of  $\mathbb{R}$ , known as the *Cantor set*. Clearly,  $m^*(C) \leq 2^n 3^{-n}$  for each  $n$ . Letting  $n \rightarrow \infty$  shows that  $C$  is null.

Let  $x \in [0, 1]$ . Then  $x \in C$  if and only if  $x$  has a ternary expansion  $x = \sum_{n=1}^{\infty} a_n 3^{-n}$ , where each  $a_n = 0$  or  $2$ . Then a variation of Cantor's proof shows that  $C$  is uncountable.

A property  $Q$  of real numbers is said to hold *almost everywhere* (a.e.) if the set of real numbers for which  $Q$  does not hold is a null set. For example,  $\chi_C = 0$  a.e., i.e.,  $\chi_C(x) = 0$  for almost all  $x$ , because  $C$  is null.

Now let us consider the question whether  $m^*$  is countably additive.

**Example 2.4.** Let  $A$  be a subset of  $[0, 1]$  with the following properties;

- (i)  $x, y \in A, x \neq y \implies x - y \notin \mathbb{Q}$ ;
- (ii) For any  $x \in [0, 1]$ , there exists  $q \in \mathbb{Q}$  such that  $x + q \in A$ .

Then

$$[0, 1] \subseteq \bigcup_{q \in \mathbb{Q} \cap [-1, 1]} (A - q) \subseteq [-1, 2].$$

Moreover, the sets  $A - q$  are disjoint (as  $q$  varies), and there are countably many of them. If  $m^*$  is countably additive, then

$$1 = m^*[0, 1] \leq \sum_{q \in \mathbb{Q} \cap [-1, 1]} m^*(A - q) = \sum_{q \in \mathbb{Q} \cap [-1, 1]} m^*(A) \leq 3.$$

This is impossible.

Thus  $m^*$  is not countably additive, provided that such a set  $A$  exists. The additive group  $\mathbb{R}$  is partitioned into the cosets of its additive subgroup  $\mathbb{Q}$ , and (i) and (ii) say that  $A$  contains exactly one member of each coset of  $\mathbb{Q}$ . The existence of such a set follows from the Axiom of Choice, an axiom of set theory beyond the basic axioms. This shows that it is impossible to prove that  $m^*$  is countably additive without using some weird axiom which contradicts the Axiom of Choice. On the other hand, it can be proved that it is impossible to show that  $m^*$  is not countably additive, using only the basic axioms of set theory.

This is bad news, but it is not so very bad because the badness occurs only with sets which cannot be explicitly described. So we can rescue things by restricting attention to a class of sets with good behaviour.

A subset  $E$  of  $\mathbb{R}$  is said to be (*Lebesgue*) *measurable* if

$$m^*(A) = m^*(A \cap E) + m^*(A \setminus E)$$

for all subsets  $A$  of  $\mathbb{R}$ . Here,  $A \setminus E = A \cap (\mathbb{R} \setminus E)$ —it is not assumed that  $E \subseteq A$ . [NB: I am using the same definition as Capinski & Kopp and Zhongminh Qian’s lecture notes (2017). Etheridge had a different definition, Stein & Shakarchi have another, Garling has another; and Priestley has yet another. All these definitions are equivalent, but this is not obvious.]

Let  $\mathcal{M}_{\text{Leb}}$  be the set of all Lebesgue measurable subsets of  $\mathbb{R}$ .

**Proposition 2.5.** 1. If  $E$  is null then  $E \in \mathcal{M}_{\text{Leb}}$ .

2. If  $I$  is any interval, then  $I \in \mathcal{M}_{\text{Leb}}$ .

3. If  $E \in \mathcal{M}_{\text{Leb}}$ , then  $\mathbb{R} \setminus E \in \mathcal{M}_{\text{Leb}}$ .

4. If  $E_n \in \mathcal{M}_{\text{Leb}}$  for  $n = 1, 2, \dots$ , then  $\bigcup_{n=1}^{\infty} E_n \in \mathcal{M}_{\text{Leb}}$ .

5. If  $E_n \in \mathcal{M}_{\text{Leb}}$  for  $n = 1, 2, \dots$  and  $E_n \cap E_k = \emptyset$  whenever  $n \neq k$ , then  $m^*(\bigcup_{n=1}^{\infty} E_n) = \sum_{n=1}^{\infty} m^*(E_n)$ .

The proofs are exercises, or can be found in books such as Capinski & Kopp. (3) is almost trivial.

Note  $\bigcap_{n=1}^{\infty} E_n = \mathbb{R} \setminus (\bigcup_{n=1}^{\infty} \mathbb{R} \setminus E_n)$ ,  $\mathcal{M}_{\text{Leb}}$  is also closed under (finite or countable) intersections. The set  $A$  of Example 2.4 is not Lebesgue measurable.

**Corollary 2.6.** All open subsets, and all closed subsets of  $\mathbb{R}$ , are Lebesgue measurable.

*Proof.* Any open subset of  $\mathbb{R}$  is a countable union of intervals (Exercise).  $\square$

For  $E \in \mathcal{M}_{\text{Leb}}$ , we shall write  $m(E)$  for  $m^*(E)$ . Then  $m : \mathcal{M}_{\text{Leb}} \rightarrow [0, \infty]$  is countably additive.

### 3. MEASURE SPACES AND MEASURABLE FUNCTIONS

Let  $\Omega$  be any set, and  $\mathcal{F} \subseteq \mathcal{P}(\Omega)$ . We say that  $\mathcal{F}$  is a  $\sigma$ -algebra (or  $\sigma$ -field) on  $\Omega$  if:

- (i)  $\emptyset \in \mathcal{F}$ ,
- (ii) If  $E \in \mathcal{F}$ , then  $\Omega \setminus E \in \mathcal{F}$ ,
- (iii) If  $E_n \in \mathcal{F}$  for  $n = 1, 2, \dots$ , then  $\bigcup_{n=1}^{\infty} E_n \in \mathcal{F}$ .

Then  $(\Omega, \mathcal{F})$  is a *measurable space*, and sets in  $\mathcal{F}$  are  $\mathcal{F}$ -measurable. As before,  $\bigcap E_n \in \mathcal{F}$  if  $E_n \in \mathcal{F}$  for  $n = 1, 2, \dots$ .

A *measure* on  $(\Omega, \mathcal{F})$  is a function  $\mu : \mathcal{F} \rightarrow [0, \infty]$  such that

- (i)  $\mu(\emptyset) = 0$ ,
- (ii)  $\mu(\bigcup_{n=1}^{\infty} E_n) = \sum_{n=1}^{\infty} \mu(E_n)$  whenever  $E_n$  are disjoint sets in  $\mathcal{F}$ .

Then  $\mu$  is *finite* if  $\mu(\Omega) < \infty$ ;  $\mu$  is a *probability measure* if  $\mu(\Omega) = 1$ .

**Examples 3.1.** 1.  $(\mathbb{R}, \mathcal{M}_{\text{Leb}}, m)$  is a measure space. Also,  $([0, 1], \mathcal{M}_{\text{Leb}}|_{[0,1]}, m)$  is a probability space, where  $\mathcal{M}_{\text{Leb}}|_{[0,1]}$  is the set of all Lebesgue measurable subsets of  $[0, 1]$ .

2. Let  $\Omega$  be any set,  $\mathcal{F} = \mathcal{P}(\Omega)$  and  $\mu(E) = |E|$  (the number of elements of  $E$ ). This is a measure space;  $\mu$  is *counting measure* on  $\Omega$ .



3. In probability theory, let  $\Omega$  be a sample space of all possible outcomes,  $\mathcal{F}$  be the collection of all events  $E$ , and  $\mathbb{P}(E)$  be the probability that event  $E$  occurs. Then  $\mathbb{P}$  is a probability measure on  $(\Omega, \mathcal{F})$ , and  $(\Omega, \mathcal{F}, \mathbb{P})$  is a probability space.
4. Let  $F : \mathbb{R} \rightarrow \mathbb{R}$  be an increasing function. Note that  $F$  may be discontinuous, but its left and right limits exist at each point. We assume that  $F(x) = \lim_{y \rightarrow x^+} F(y)$  for all  $x$  (without essential loss). Define

$$m_F(a, b] = F(b) - F(a),$$

$$m_F^*(E) = \inf \left\{ \sum_{n=1}^{\infty} m_F(J_n) : J_n = (a_n, b_n], E \subseteq \bigcup_{n=1}^{\infty} J_n \right\}.$$

Then  $m_F^*$  has similar properties to  $m^*$ , but one has to be aware that  $m_F^*(a, b) = F(b-) - F(a)$ ,  $m_F^*([a, b]) = F(b) - F(a-)$ ; and  $m_F^*({x}) = 0$  if and only if  $F$  is continuous at  $x$ . One can then define a  $\sigma$ -algebra  $\mathcal{M}_F$ , containing all intervals, in the same way as  $\mathcal{M}_{\text{Leb}}$ , and  $m_F^*$  is a measure, written  $m_F$  on  $\mathcal{M}_F$ . This is the *Lebesgue-Stieltjes measure associated with  $F$* .

**Proposition 3.2.** *Let  $(\Omega, \mathcal{F}, \mu)$  be a measure space.*

1. *If  $A, B \in \mathcal{F}$  and  $A \subseteq B$ , then  $\mu(A) \leq \mu(B)$ .*
2. *If  $A_n \in \mathcal{F}$  and  $A_n \subseteq A_{n+1}$  for all  $n$ , then  $\mu(\bigcup_n A_n) = \lim_{n \rightarrow \infty} \mu(A_n)$ .*
3. *If  $A_n \in \mathcal{F}$  and  $A_n \supseteq A_{n+1}$  for all  $n$  and  $\mu(A_1) < \infty$ , then  $\mu(\bigcap_n A_n) = \lim_{n \rightarrow \infty} \mu(A_n)$ .*

*Proof.* (1) Since  $B = A \cup (B \setminus A)$  (disjoint union),  $\mu(B) = \mu(A) + \mu(B \setminus A) \geq \mu(A)$ .

(2) Let  $A'_1 = A_1$  and  $A'_r = A_r \setminus A_{r-1}$  for  $r \geq 2$ . Then  $A_n = \bigcup_{r=1}^n A'_r$ ,  $\bigcup_{n=1}^{\infty} A_n = \bigcup_{r=1}^{\infty} A'_r$  (disjoint unions), so

$$\mu\left(\bigcup_n A_n\right) = \sum_{r=1}^{\infty} \mu(A'_r) = \lim_{n \rightarrow \infty} \sum_{r=1}^n \mu(A'_r) = \lim_{n \rightarrow \infty} \mu(A_n).$$

(3) is an exercise. □

Let  $\mathcal{B} \subseteq \mathcal{P}(\Omega)$ . There is a (unique)  $\sigma$ -algebra  $\mathcal{F}_{\mathcal{B}}$  on  $\Omega$  which is generated by  $\mathcal{B}$  in the following sense:

- (i)  $\mathcal{F}_{\mathcal{B}}$  is a  $\sigma$ -algebra and  $\mathcal{B} \subseteq \mathcal{F}_{\mathcal{B}}$ ,
- (ii) If  $\mathcal{F}$  is  $\sigma$ -algebra on  $\Omega$  and  $\mathcal{B} \subseteq \mathcal{F}$  then  $\mathcal{F}_{\mathcal{B}} \subseteq \mathcal{F}$ .

The  $\sigma$ -algebra  $\mathcal{M}_{\text{Bor}}$  generated by the intervals is the *Borel  $\sigma$ -algebra* on  $\mathbb{R}$ . It can be described as the class of all subsets of  $\mathbb{R}$  which can be obtained from intervals in a countable number of steps, each of which is one of taking the complement of a set, taking a countable union of sets, or a countable intersection of sets. However this has to be treated with caution, because it is not necessarily possible to obtain a given Borel set by performing the countable number of steps in a single sequence.

**Proposition 3.3.** 1. *Let  $\mathcal{B}$  be any one of the following classes of subsets of  $\mathbb{R}$ .*

- (i) *All intervals*
- (ii) *All intervals of the form  $(a, \infty)$*
- (iii) *All intervals of the form  $[a, b]$*
- (iv) *All open sets.*

Then  $\mathcal{M}_{\text{Bor}}$  is the smallest  $\sigma$ -algebra containing  $\mathcal{B}$ .

2.  $\mathcal{M}_{\text{Bor}} \neq \mathcal{M}_{\text{Leb}}$ .
3. If  $E \in \mathcal{M}_{\text{Leb}}$  there exist  $A, B \in \mathcal{M}_{\text{Bor}}$  such that  $A \subseteq E \subseteq B$  and  $B \setminus A$  is null (so  $E \setminus A$  and  $B \setminus E$  are null).

*Proof.* (1) is an exercise involving showing each interval can be obtained from members of  $\mathcal{B}$ , and each member of  $\mathcal{B}$  can be obtained from intervals. (2) and (3) are quite deep results; (2) is discussed in a document on the course Webpage; (3) is Theorem 2.28 in Capinski & Kopp.  $\square$

Let  $(\Omega, \mathcal{F})$  be a measurable space. A function  $f : \Omega \rightarrow \mathbb{R}$  is  $\mathcal{F}$ -measurable if  $f^{-1}(I) \in \mathcal{F}$  for each interval  $I$ .

**Proposition 3.4.** *Let  $\mathcal{B}$  be any one of the classes of subsets of  $\mathbb{R}$  listed in Proposition 3.3. Let  $f : \Omega \rightarrow \mathbb{R}$ . Then  $f$  is  $\mathcal{F}$ -measurable if and only if  $f^{-1}(G) \in \mathcal{F}$  for all  $G \in \mathcal{M}_{\text{Bor}}$  or for all  $G \in \mathcal{B}$ .*

*Proof.* It is easily verified that  $f_*(\mathcal{F}) := \{G \subseteq \mathbb{R} : f^{-1}(G) \in \mathcal{F}\}$  is a  $\sigma$ -algebra. Hence if  $\mathcal{B}$  generates the  $\sigma$ -algebra  $\mathcal{M}_{\text{Bor}}$ , then the result holds.  $\square$

In this course, we shall usually take  $(\Omega, \mathcal{F})$  to be  $(\mathbb{R}, \mathcal{M}_{\text{Leb}})$  or minor variants, but much of this section will apply to the general case as well. We may refer to  $\mathcal{M}_{\text{Leb}}$ -measurable functions simply as measurable functions, for simplicity; or as Lebesgue measurable functions. We shall also be interested in cases where  $\Omega$  is an interval (or a Lebesgue measurable subset) and  $\mathcal{F} = \mathcal{M}_{\text{Leb}}|_{\Omega} = \{E \in \mathcal{M}_{\text{Leb}} : E \subseteq \Omega\}$ . However,  $f : \Omega \rightarrow \mathbb{R}$  is  $\mathcal{M}_{\text{Leb}}|_{\Omega}$ -measurable if and only if  $\tilde{f} : \mathbb{R} \rightarrow \mathbb{R}$  is measurable, where  $\tilde{f}(x) = f(x)$  for  $x \in \Omega$ , and  $\tilde{f}(x) = 0$  otherwise. So we may state results just for functions defined on  $\mathbb{R}$ .

Recall from the Analysis courses that  $f : \mathbb{R} \rightarrow \mathbb{R}$  is continuous if and only if  $f^{-1}(G)$  is open for every open set (or open interval)  $G$ . By Proposition 3.4, we have that  $f$  is (Lebesgue) measurable if and only if  $f^{-1}(G)$  is (Lebesgue) measurable for every open set (or open interval)  $G$ .

**Examples 3.5.** 1. Constant functions are measurable.

2. The characteristic function  $\chi_A$  of a subset  $A$  of  $\mathbb{R}$  is measurable if and only if  $A$  is a measurable set. In particular, if  $A$  is as in Example 2.4, then  $\chi_A$  is not (Lebesgue) measurable.
3. Continuous functions  $f : \mathbb{R} \rightarrow \mathbb{R}$  are measurable.
4. Monotone functions  $f : \mathbb{R} \rightarrow \mathbb{R}$  are measurable.
5. If  $f$  is continuous a.e., then  $f$  is measurable.
6. If  $f : \mathbb{R} \rightarrow \mathbb{R}$  is (Lebesgue) measurable and  $g = f$  a.e., then  $g$  is (Lebesgue) measurable.
7. In probability theory, measurable functions are called random variables.

It follows from the definition of measurable functions and Example 3.5(2) that the existence of a non-measurable function is equivalent to the existence of a non-measurable set. So their existence depends on the Axiom of Choice. Thus, we have the following:

**Fact of Life.** ALL FUNCTIONS  $f : \mathbb{R} \rightarrow \mathbb{R}$  THAT CAN BE EXPLICITLY DEFINED ARE LEBESGUE MEASURABLE.

This is not exactly a mathematical theorem—it becomes one if one interprets “explicitly defined” in the right technical way. It is a true statement about the real world: a non-measurable function involves some non-explicit choice process. Priestley compares the existence of non-measurable functions to the existence of yetis.

Nevertheless, measurability is a real issue in some more advanced mathematics, because:

- (a) One may be interested not in Lebesgue measurability of functions  $f$  on  $\mathbb{R}$ , but in measurability on some other measurable space  $(\Omega, \mathcal{F})$ . This occurs frequently in time-dependent probability theory, where  $\mathcal{F}_t$  is the class of all events depending only on past history up to time  $t$ , not on the future (cf. Part B courses on martingales and stochastic calculus).
- (b) One may be interested in functions  $f$  which are not real-valued, but take values in an infinite-dimensional space. Then measurability is a real issue in many areas of analysis, although you probably won't see this in your undergraduate course.

So it is useful to accumulate general results about measurable functions, even if we only state them for functions  $f : (\mathbb{R}, \mathcal{M}_{\text{Leb}}) \rightarrow \mathbb{R}$ .

**Proposition 3.6.** *Let  $f$  and  $g$  be measurable functions from  $\mathbb{R}$  to  $\mathbb{R}$ . The following functions are measurable:*

$$f + g, fg, \max(f, g), \quad h \circ f \text{ for any continuous function } h.$$

For example,  $\alpha f$  is measurable, where  $\alpha \in \mathbb{R}$ .

*Proof.* For example,

$$(f + g)^{-1}(a, \infty) = \bigcup_{q \in \mathbb{Q}} f^{-1}(q, \infty) \cap g^{-1}(a - q, \infty).$$

If  $G$  is open in  $\mathbb{R}$ , then  $h^{-1}(G)$  is open. Since  $f$  is measurable,  $f^{-1}(h^{-1}(G))$  is measurable, i.e.,  $(h \circ f)^{-1}(G)$  is measurable.  $\square$

In fact, it suffices in Proposition 3.6 that  $h$  should be Borel measurable.

Now we want to consider limits and suprema of sequences of functions  $(f_n)$ . Even if each  $f_n$  is real-valued, the resulting functions may take the values  $\infty$  and  $-\infty$ .

A function  $f : \mathbb{R} \rightarrow [-\infty, \infty]$  is *measurable* if  $f^{-1}(a, \infty] \in \mathcal{M}_{\text{Leb}}$  for all  $a \in \mathbb{R}$ ; equivalently  $f^{-1}(B) \in \mathcal{M}_{\text{Leb}}$  for all  $B \in \mathcal{M}_{\text{Bor}}$  and  $f^{-1}(\{\infty\}) \in \mathcal{M}_{\text{Leb}}$ ; equivalently,  $\arctan \circ f$  is measurable, where  $\arctan : [-\infty, \infty] \rightarrow [-\pi/2, \pi/2]$  is the inverse tan function.

**Proposition 3.7.** *Let  $(f_n)$  be a sequence of measurable functions from  $\mathbb{R} \rightarrow [-\infty, \infty]$ . Then the following functions are measurable:*

$$\sup_n f_n, \inf_n f_n, \limsup_{n \rightarrow \infty} f_n, \liminf_{n \rightarrow \infty} f_n.$$

Hence, if  $f(x) = \lim_{n \rightarrow \infty} f_n(x)$  a.e., then  $f$  is measurable.

*Proof.* First,

$$(\sup f_n)^{-1}(a, \infty] = \bigcup_n f_n^{-1}(a, \infty] \in \mathcal{M}_{\text{Leb}}.$$

Then

$$\begin{aligned} \inf f_n &= -\sup(-f_n), \\ \limsup f_n &= \inf g_m, \text{ where } g_m = \sup_{n \geq m} f_n. \end{aligned}$$

□

A function  $\phi : \mathbb{R} \rightarrow \mathbb{R}$  is *simple* if it is measurable and it takes only finitely many real values. So  $\chi_E$  is simple if  $E \in \mathcal{M}_{\text{Leb}}$ . If  $\phi, \psi$  are simple, then so are  $\phi + \psi$ ,  $\phi \cdot \psi$ ,  $\alpha\phi$ ,  $\max(\phi, \psi)$ ,  $h \circ \phi$  for any function  $h$ .

Any function of the form  $\sum_{j=1}^n \beta_j \chi_{E_j}$ , where  $\beta_j \in \mathbb{R}$  and  $E_j \in \mathcal{M}_{\text{Leb}}$  is simple. On the other hand, if  $\phi$  is simple with non-zero values  $\alpha_1, \dots, \alpha_k$ , and  $B_i = \phi^{-1}(\{\alpha_i\})$ , then  $B_i$  is measurable, and

$$(*) \quad \phi = \sum_{i=1}^k \alpha_i \chi_{B_i}.$$

In this form, we have

- (i)  $\alpha_i$  are distinct and non-zero,
- (ii)  $B_i$  are disjoint.

If these additional properties hold, then (\*) is unique (up to reordering of the terms). We shall then say that  $\phi$  is in *standard*, or *canonical*, form. For example, the standard form of  $\chi_{(0,2)} + \chi_{[1,3]}$  is  $1\chi_{(0,1) \cup [2,3]} + 2\chi_{[1,2]}$ .

In defining simple functions, some authors insist that the sets  $B_i$ , corresponding to non-zero  $\alpha_i$ , must be bounded [Etheridge] or of finite measure [Stein & Shakarchi]. [Garling and Priestley avoid introducing simple functions.]

**Examples 3.8.** 1. Any step function is a simple function—for a step function, the sets  $B_i$  in the standard representation must be finite unions of bounded intervals (or single points).

2. The function  $\chi_{\mathbb{Q} \cap [0,1]}$  is a simple function but it is not a step function.

**Proposition 3.9.** *Let  $f : \mathbb{R} \rightarrow [0, \infty]$  be measurable. There is an increasing sequence  $(\phi_n)$  of non-negative simple functions  $\phi_n$  such that*

$$f(x) = \lim_{n \rightarrow \infty} \phi_n(x)$$

for all  $x \in \mathbb{R}$ .

*Proof.* For  $n = 1, 2, \dots$  and  $k = 0, 1, 2, \dots, 4^n - 1$ , let

$$B_{kn} = \{x : k2^{-n} \leq f(x) < (k+1)2^{-n}\}.$$

Let

$$\phi_n(x) = \begin{cases} k2^{-n} & \text{if } x \in B_{kn} \text{ for some (unique) } k, \\ 2^n & \text{if } f(x) > 2^n. \end{cases}$$

Then  $\phi_n \leq \phi_{n+1}$ ,  $\phi_n \leq f$ ,  $\phi_n(x) > f(x) - 2^{-n}$  for all sufficiently large  $n$  if  $f(x) < \infty$ , and  $\phi_n(x) = 2^n$  for all  $n$  if  $f(x) = \infty$ .  $\square$

Notice here that the approximating simple functions are constructed by taking horizontal strips, unlike Prelims where vertical strips were used.

**Theorem 3.10.** *A function  $f : \mathbb{R} \rightarrow \mathbb{R}$  is measurable if and only if there is a sequence of step functions  $\psi_n$  such that  $f = \lim \psi_n$  a.e.*

*Proof.* Stein & Shakarchi, Theorem 4.3, p.32.  $\square$

#### 4. THE LEBESGUE INTEGRAL

For a non-negative simple function  $\phi$  with standard form  $\sum_{i=1}^k \alpha_i \chi_{B_i}$  (so  $\alpha_i > 0$ ), the *integral* of  $\phi$  is defined to be:

$$\int_{\mathbb{R}} \phi = \int_{-\infty}^{\infty} \phi(x) dx = \sum_{i=1}^k \alpha_i m(B_i).$$

Note that  $\int \phi < \infty$  if and only if  $m(B_i) < \infty$  for each  $i$ .

**Proposition 4.1.** *Let  $\phi, \psi$  be non-negative simple functions,  $\alpha \in [0, \infty)$ .*

1. *If  $\phi = \sum_{j=1}^n \beta_j \chi_{E_j}$  where  $\beta_j \geq 0$  and  $E_j$  are measurable (but not necessarily in standard form), then  $\int \phi = \sum_j \beta_j m(E_j)$ .*
2.  *$\int(\phi + \psi) = \int \phi + \int \psi$ ,  $\int \alpha \phi = \alpha \int \phi$ .*
3. *If  $\phi \leq \psi$  then  $\int \phi \leq \int \psi$ .*

The first statement of Proposition 4.1 is not completely obvious, but fortunately it is true! [Capinski & Kopp define  $\int \phi$  to be  $\sum_j \beta_j m(E_j)$ , ignoring the question whether this is well-defined.]

For a non-negative measurable function  $f : \mathbb{R} \rightarrow [0, \infty]$ , we define the *integral* of  $f$  to be

$$\int_{\mathbb{R}} f = \sup \left\{ \int_{\mathbb{R}} \phi : \phi \text{ simple, } 0 \leq \phi \leq f \right\}.$$

For a measurable subset  $E$  of  $\mathbb{R}$ , we define

$$\int_E f = \int_{\mathbb{R}} f \chi_E.$$

For a measurable function  $f : E \rightarrow [0, \infty)$ , we define  $\int_E f = \int_{\mathbb{R}} \tilde{f}$ , where  $\tilde{f}$  agrees with  $f$  on  $E$  and is 0 on  $\mathbb{R} \setminus E$ .

In either case, we say that  $f$  is *integrable over  $E$*  if  $\int_E f < \infty$ .

This definition of integral corresponds to the lower integral in Prelims, but with simple functions replacing step functions. If the Monotone Convergence Theorem is to be true, then Proposition 3.9 shows that the integral must equal this supremum, but it is still necessary to show that our definition has good properties.

It is clear from the definition of integral that

- (i)  $\int \alpha f = \alpha \int f$  ( $\alpha \geq 0$ );
- (ii) If  $f \leq g$ , then  $\int f \leq \int g$ ,

The first things to establish are

- (iii)  $\int(f + g) = \int f + \int g$ ,
- (iv) The Monotone Convergence Theorem.

**Theorem 4.2. [Monotone Convergence Theorem, Version 1]** *If  $(f_n)$  is an increasing sequence of non-negative measurable functions and  $f = \lim_{n \rightarrow \infty} f_n$ , then  $\int f = \lim_{n \rightarrow \infty} \int f_n$ .*

*Proof.* Since  $f_n \leq f$ , it is immediate that  $\sup_n \int f_n \leq \int f$ .

For the reverse inequality, we consider a simple function  $\phi$  such that  $0 \leq \phi \leq f$ . We have to show that  $\int \phi \leq \lim_{n \rightarrow \infty} \int f_n$ . It then follows from the definition of  $\int f$  that  $\int f \leq \lim_{n \rightarrow \infty} \int f_n$ .

Take  $\alpha \in (0, 1)$ , and let

$$B_n = \{x : f_n(x) \geq \alpha\phi(x)\}.$$

Then  $B_n$  is measurable (since  $f_n - \alpha\phi$  is measurable),  $B_n \subseteq B_{n+1}$  and  $\bigcup_{n=1}^{\infty} B_n = \mathbb{R}$  (for each  $x$ , either  $\phi(x) = 0$  or  $f(x) > \alpha\phi(x)$ ). Since  $\alpha\phi\chi_{B_n} \leq f_n\chi_{B_n} \leq f_n$ ,

$$(*) \quad \alpha \int_{B_n} \phi \leq \int_{\mathbb{R}} f_n.$$

If  $\phi = \sum_{i=1}^k \beta_i \chi_{E_i}$ , then

$$\int_{B_n} \phi = \sum_{i=1}^k \beta_i m(E_i \cap B_n) \rightarrow \sum_{i=1}^k \beta_i m(E_i) = \int_{\mathbb{R}} \phi$$

as  $n \rightarrow \infty$ , by Proposition 3.2(2). Taking limits in (\*),

$$\alpha \int_{\mathbb{R}} \phi \leq \lim_{n \rightarrow \infty} \int_{\mathbb{R}} f_n.$$

Letting  $\alpha \rightarrow 1-$  gives the required inequality. □

**Corollary 4.3. [Baby MCT]** *Let  $f$  be a non-negative measurable function,  $(E_n)$  be an increasing sequence of measurable sets, and  $E = \bigcup_{n=1}^{\infty} E_n$ . Then  $f$  is integrable over  $E$  if and only if  $\sup_n \int_{E_n} f < \infty$ . Then  $\int_E f = \sup_n \int_{E_n} f = \lim_{n \rightarrow \infty} \int_{E_n} f$ .*

*Proof.* Apply Theorem 4.2 with  $f_n = f\chi_{E_n}$ , noting that  $\chi_{E_n} \leq \chi_{E_{n+1}}$  and  $f \geq 0$ , so  $f_n \leq f_{n+1}$  and  $\chi_E(x) = \lim_{n \rightarrow \infty} \chi_{E_n}(x)$ . □

**Corollary 4.4.** *For non-negative measurable functions  $f$  and  $g$ ,*

$$\int (f + g) = \int f + \int g.$$

*Proof.* Let  $(\phi_n)$  and  $\psi_n$  be increasing sequences of simple functions, converging pointwise to  $f$  and  $g$  respectively (Proposition 3.9). Then  $(\phi_n + \psi_n)$  is an increasing sequence, converging to  $f + g$ . By MCT and Proposition 4.1(2),

$$\int (f+g) = \lim_{n \rightarrow \infty} \int (\phi_n + \psi_n) = \lim_{n \rightarrow \infty} \left( \int \phi_n + \int \psi_n \right) = \lim_{n \rightarrow \infty} \int \phi_n + \lim_{n \rightarrow \infty} \int \psi_n = \int f + \int g.$$

□

**Corollary 4.5. [MCT for Series]** *Let  $f_n$  be non-negative measurable functions and  $f = \sum_{n=1}^{\infty} f_n$ . Then  $\int f = \sum_{n=1}^{\infty} \int f_n$ . In particular,  $f$  is integrable if and only if  $\sum_n \int f_n < \infty$ .*

*Proof.* Let  $g_n = \sum_{r=1}^n f_r$ , and apply MCT. □

Before giving an example, we note that our integral agrees with the Riemann integral in a fundamental case.

**Corollary 4.6.** *Let  $f : [a, b] \rightarrow [0, \infty)$  be continuous. Then the Lebesgue integral  $\int_{[a,b]}^{\mathcal{L}} f$  as defined above equals the Riemann integral  $\int_{[a,b]}^{\mathcal{R}} f$  as defined in first-year Integration.*

*Proof.* As shown in first-year, there is an increasing sequence  $(\phi_n)$  of step functions such that  $\lim_{n \rightarrow \infty} \phi_n(x) = f(x)$  for all  $x \in [a, b]$  and  $\lim_{n \rightarrow \infty} \int_a^b \phi_n = \int_{[a,b]}^{\mathcal{R}} f$ . By MCT (Theorem 4.2),  $\lim_{n \rightarrow \infty} \int_a^b \phi_n = \int_{[a,b]}^{\mathcal{L}} f$ . □

**Example 4.7.** Consider  $f(x) = (1 - x)^{-1/2}$  on  $(0, 1)$ . By Baby MCT (Corollary 4.3) and FTC (from Prelims),

$$\int_0^1 (1 - x)^{-1/2} dx = \lim_{n \rightarrow \infty} \int_0^{1 - \frac{1}{n}} (1 - x)^{-1/2} dx = \lim_{n \rightarrow \infty} 2(1 - n^{-1/2}) = 2.$$

For  $0 \leq x < 1$ , the Binomial Theorem with exponent  $-1/2$  or Taylor's Theorem in complex analysis gives

$$(1 - x)^{-1/2} = \sum_{n=0}^{\infty} \frac{(2n)!}{4^n (n!)^2} x^n.$$

By Corollary 4.5 and FTC,

$$\int_0^1 (1 - x)^{-1/2} dx = \sum_{n=0}^{\infty} \frac{(2n)!}{4^n (n!)^2} \int_0^1 x^n dx = \sum_{n=0}^{\infty} \frac{(2n)!}{4^n n!(n+1)!}.$$

The fact that the series above converges to 2 can be obtained directly from the Binomial Expansion of  $(1 - x)^{1/2}$ , via Abel's continuity theorem (A2, Theorem 13.24).

Now we turn to integrability of functions which are not necessarily non-negative.

Let  $f : \mathbb{R} \rightarrow [-\infty, \infty]$  be measurable. Let

$$f^+ = \max(f, 0), \quad f^- = \max(-f, 0).$$

Note that  $f^+$  and  $f^-$  are measurable and non-negative, and

$$f = f^+ - f^-, \quad |f| = f^+ + f^-.$$

We say that  $f$  is *integrable* if  $f$  is measurable and  $\int f^+$  and  $\int f^-$  are both finite. Then the *integral* of  $f$  is

$$\int f = \int f^+ - \int f^-.$$

Moreover,  $f$  is *integrable over a measurable subset  $E$*  if  $f\chi_E$  is integrable. If  $f : E \rightarrow [-\infty, \infty]$ , then  $f$  is *integrable over  $E$*  if  $\tilde{f}$  is integrable over  $\mathbb{R}$ .

- Proposition 4.8.**
1. If  $f$  is integrable, then  $|f|$  is integrable.
  2. If  $f$  is measurable and  $|f|$  is integrable, then  $f$  is integrable.
  3. [**Comparison Test**] If  $f$  is measurable and  $|f| \leq g$  for some integrable function  $g$ , then  $f$  is integrable. If  $|f| \geq g \geq 0$  for some measurable function  $g$  which is not integrable, then  $f$  is not integrable.
  4. If  $f, g$  are both integrable and  $f+g$  is defined, then  $f+g$  is integrable and  $\int(f+g) = \int f + \int g$ . For  $\alpha \in \mathbb{R}$ ,  $\alpha f$  is integrable and  $\int \alpha f = \alpha \int f$ . If  $f \leq g$ , then  $\int f \leq \int g$ .
  5. If  $f$  is integrable and  $g = f$  a.e., then  $g$  is integrable and  $\int g = \int f$ .
  6. If  $f$  is integrable then  $f(x) \in \mathbb{R}$  a.e.
  7. If  $f$  is integrable and  $\int |f| = 0$  then  $f(x) = 0$  a.e.
  8. If  $f$  is integrable over a measurable set  $E$  and  $(E_n)$  is an increasing sequence of measurable sets with  $\bigcup_{n=1}^{\infty} E_n = E$  then  $\int_E f = \lim_{n \rightarrow \infty} \int_{E_n} f$ .

*Proof.* (omitted) (1) and (2) follow from  $\int f^\pm \leq \int |f| = \int f^+ + \int f^-$ . (3) follows from  $|f| \leq g \implies \int |f| \leq \int g$ . (4) follows from  $(f+g)^\pm \leq f^\pm + g^\pm$  and  $(f+g)^+ + f^- + g^- = (f+g)^- + f^+ + g^+$ . (5): Since  $|g-f| = 0$  a.e., any simple function  $\phi$  with  $0 \leq \phi \leq |g-f|$  is a.e. 0, so its integral is 0. Hence  $\int |g-f| = 0$ . (6), (7): Exercises. (8): Apply Baby MCT to  $f^+$  and  $f^-$ .  $\square$

By (5), changing a function on a null set does not affect integrability. So if we have a function defined a.e., we can talk about it being integrable by considering any extension of  $f$ —for example, the extension by 0. Also, integrability over  $[a, b]$  is the same as integrability over  $(a, b)$ .

The following are corollaries of the Comparison Test.

- Corollary 4.9.**
1. If  $g$  is integrable and  $h$  is bounded and measurable, then  $hg$  is integrable.
  2. If  $g$  is integrable over  $\mathbb{R}$ , then  $g$  is integrable over any measurable subset of  $\mathbb{R}$ .
  3. If  $h$  is a bounded measurable function, then  $h$  is integrable over any measurable subset of finite measure.

*Proof.* These follow from the Comparison Test, using

$$|g \cdot h| \leq c|g|, \quad |g\chi_E| \leq |g|, \quad |h\chi_E| \leq c\chi_E. \quad \square$$

Apart from Corollary 4.6, almost all the theory in Section 4 up to this point applies to general measure spaces. Now we make some comments which are specific to the case of Lebesgue measure.



Firstly, the Lebesgue integral is more general than the Riemann (Prelims) integral. In fact,  $f : [a, b] \rightarrow \mathbb{R}$  is Riemann integrable if and only if  $f$  is bounded and continuous a.e. Any such  $f$  is measurable and bounded, hence Lebesgue integrable. Moreover,

$$\begin{aligned} \int_{[a,b]}^{\mathcal{R}} f &= \sup \left\{ \int_a^b \phi : \phi \text{ step, } \phi \leq f \right\} \leq \sup \left\{ \int_a^b \phi : \phi \text{ simple, } \phi \leq f \right\} \\ &\leq \int_{[a,b]}^{\mathcal{L}} f \leq \inf \left\{ \int_a^b \psi : \psi \text{ step, } f \leq \psi \right\} = \int_{[a,b]}^{\mathcal{R}} f \end{aligned}$$

Hence equality holds throughout, so the Lebesgue integral equals the Riemann integral.

The following is an argument showing that every Riemann integrable function on  $[a, b]$  is measurable, without showing that it is continuous a.e. (that is trickier; it is shown in Garling, Section 29.2)

*(this paragraph not included in lectures)* If  $f$  is Riemann integrable, then  $f$  is bounded and there are sequences  $(\phi_n)$  and  $(\psi_n)$  of step functions such that  $\phi_n \leq f \leq \psi_n$  and  $\lim_{n \rightarrow \infty} \int_a^b \phi_n = \int_{[a,b]}^{\mathcal{R}} f = \lim_{n \rightarrow \infty} \int_a^b \psi_n$ . Let  $g = \sup_n \phi_n$  and  $h = \inf_n \psi_n$ . Then  $g$  and  $h$  are measurable,  $g \leq f \leq h$  and  $\int_{[a,b]}^{\mathcal{L}} (h - g) \leq \lim_{n \rightarrow \infty} \int_a^b (\psi_n - \phi_n) = 0$ . By Proposition 4.8(7),  $g = h$  a.e. Then  $f = g$  a.e., so  $f$  is (Lebesgue) measurable. By Corollary 4.9(3),  $f$  is Lebesgue integrable.

Given a function  $f : I \rightarrow \mathbb{R}$ , where  $I$  is an interval in  $\mathbb{R}$ , how does one test whether  $f$  is integrable over  $I$ ? We can do the following:

- Note that  $f$  is measurable (for example, using Examples 3.5).
- Replace  $f$  by  $|f|$ : we can assume that  $f$  is non-negative. (Proposition 4.8(1),(2))
- If  $I$  is bounded and  $f$  is bounded, then  $f$  is integrable over  $I$ . (Corollary 4.9(3))
- If  $I$  or  $f$  is unbounded, we can probably consider an increasing sequence of bounded subintervals  $I_n$ , with union  $I$ , such that  $f$  is bounded on each  $I_n$ .
- We may be able to evaluate  $\int_{I_n} f$  by means of the FTC, Integration by Parts, or Substitution from Prelims theory. Then we can use Baby MCT (Corollary 4.3).
- If we cannot easily evaluate the integral of  $f$ , use the Comparison Test—we look for a simpler measurable function  $g$  such that  $g$  is known to be integrable and  $0 \leq f \leq g$  (if we think  $f$  is going to be integrable), or  $g$  is known not to be integrable and  $0 \leq g \leq f$  (if we think  $f$  will not be integrable).

**Examples 4.10.** 1. Consider  $x^\alpha$  over  $(0, 1)$ , where  $\alpha \in \mathbb{R}$ . Note first that  $x^\alpha$  is continuous, hence measurable, and non-negative. If  $\alpha \geq 0$ , then  $x^\alpha$  is bounded (by 1) on  $(0, 1)$ , hence integrable. If  $\alpha < 0$ ,  $x^\alpha$  has a singularity at  $x = 0$ , so we use Baby MCT with  $I_n = [1/n, 1]$ . By FTC,

$$\int_{1/n}^1 x^\alpha dx = \begin{cases} \frac{1-n^{-(\alpha+1)}}{\alpha+1} & (\alpha \neq -1) \\ \log n & (\alpha = -1) \end{cases} \rightarrow \begin{cases} \infty & (\alpha \leq -1) \\ -\frac{1}{\alpha+1} & (\alpha > -1). \end{cases}$$

By Baby MCT,  $x^\alpha$  is integrable over  $(0, 1)$  if and only if  $\alpha > -1$ , and then  $\int_0^1 x^\alpha dx = -(\alpha + 1)^{-1}$ .

2. Consider  $x^\alpha$  over  $[1, \infty)$ . This is similar, but with  $I_n = [1, n]$ . Now

$$\int_1^n x^\alpha dx = \begin{cases} \frac{n^{\alpha+1}-1}{\alpha+1} & (\alpha \neq -1) \\ \log n & (\alpha = -1) \end{cases} \rightarrow \begin{cases} \infty & (\alpha \geq -1) \\ \frac{1}{\alpha+1} & (\alpha < -1). \end{cases}$$

By Baby MCT,  $x^\alpha$  is integrable over  $(1, \infty)$  if and only if  $\alpha < -1$ , and then  $\int_0^1 x^\alpha dx = (\alpha + 1)^{-1}$ .

3. Consider  $f(x) = x^\alpha/(1+x^\beta)$  over  $(0, \infty)$ , where  $\alpha \in \mathbb{R}$  and  $\beta \geq 0$ . For  $0 < x \leq 1$ ,  $x^\alpha/2 \leq f(x) \leq x^\alpha$ . By comparison,  $f$  is integrable over  $(0, 1)$  if and only if  $x^\alpha$  is, i.e.,  $\alpha > -1$ . For  $x > 1$ ,  $x^{\alpha-\beta}/2 < f(x) < x^{\alpha-\beta}$ , so, by comparison,  $f$  is integrable over  $(1, \infty)$  if and only if  $x^{\alpha-\beta}$  is, i.e.,  $\alpha - \beta < -1$ . Hence  $f$  is integrable over  $(0, \infty)$  if and only if  $-1 < \alpha < \beta - 1$ . [The case when  $\beta < 0$  can be reduced to the previous case because  $f(x) = x^{\alpha-\beta}/(1+x^{-\beta})$ .]
4. Consider  $f(x) = (\sin x)/x$  over  $(0, 2\pi)$ . This function is continuous on  $(0, 2\pi]$ , hence measurable. If we define  $f(0) = 1$ , it becomes continuous, hence bounded on  $[0, 2\pi]$ —in fact it is bounded above by 1 and below by  $-1/\pi$ . So it is integrable over  $(0, 2\pi)$ .
5. Consider  $f(x) = (\sin x)/x$  over  $(0, \infty)$ . Now

$$\int_{r\pi}^{(r+1)\pi} \left| \frac{\sin x}{x} \right| dx \geq \int_{r\pi}^{(r+1)\pi} \frac{|\sin x|}{(r+1)\pi} dx = \frac{2}{(r+1)\pi}.$$

Hence,

$$\lim_{n \rightarrow \infty} \int_0^{n\pi} |f(x)| dx \geq \lim_{n \rightarrow \infty} \sum_{r=0}^{n-1} \frac{2}{(r+1)\pi} = \infty.$$

So  $|f|$  is not integrable, and hence  $f$  is not integrable, over  $(0, \infty)$ .

Let us discuss the first-year theorems a little more carefully.

**Theorem 4.11. (Fundamental Theorem of Calculus)** *Let  $g$  be a function with a continuous derivative on a closed bounded interval  $[a, b]$ . Then  $g'$  is integrable over  $[a, b]$ , and*

$$\int_a^b g'(x) dx = g(b) - g(a).$$

The FTC should be treated with care, if the range of integration is unbounded (as already discussed), or if the derivative does not exist at some points as the following examples show.

**Examples 4.12.** 1. Let  $f(x) = x \sin(\frac{1}{x})$  ( $x \in (0, 1]$ );  $f(0) = 0$ . Then  $f$  is continuous on  $[0, 1]$  and differentiable on  $(0, 1]$  but  $f'(x) = \sin(\frac{1}{x}) - \frac{1}{x} \cos(\frac{1}{x}) \notin \mathcal{L}^1(0, 1)$ .

2. We define a function  $\Phi : [0, 1] \rightarrow [0, 1]$  as follows. On the Cantor set  $C$ ,

$$\Phi \left( \sum_{n=1}^{\infty} a_n 3^{-n} \right) = \sum_{n=1}^{\infty} \frac{a_n}{2} 2^{-n} \quad (a_n = 0 \text{ or } 2).$$

Then put  $\Phi = \frac{1}{2}$  on  $[\frac{1}{3}, \frac{2}{3}]$ ,  $\frac{1}{4}$  on  $[\frac{1}{9}, \frac{2}{9}]$ , etc. Then  $\Phi$  is continuous, monotonic, differentiable at each point of  $[0, 1] \setminus C$  with  $\Phi'(x) = 0$ . So

$$\int_0^1 \Phi'(x) dx = 0 \neq \Phi(1) - \Phi(0).$$

This function  $\Phi$  is called the *Cantor-Lebesgue function*, or the *devil's staircase*.

**Theorem 4.13. (Integration by Parts)** *Let  $f$  and  $g$  be continuously differentiable functions on a closed bounded interval  $[a, b]$ . Then*

$$\int_a^b f(x)g'(x) dx = f(b)g(b) - f(a)g(a) - \int_a^b f'(x)g(x) dx.$$

Integration by parts must be treated with great care if the interval of integration is an unbounded interval or the integrand has a singularity and you do not know whether the integrals exist. In those circumstances you cannot infer the existence of one integral from the existence of the other.

**Example 4.14.** Consider  $\int_0^a \frac{\sin x}{x} dx$ . Integration by parts gives

$$\int_1^a \frac{\sin x}{x} dx = \cos 1 - \frac{\cos a}{a} - \int_1^a \frac{\cos x}{x^2} dx.$$

But  $|\frac{\cos x}{x^2}| \leq \frac{1}{x^2}$ , so  $\frac{\cos x}{x^2}$  is integrable over  $[1, \infty)$ , by Example 4.10(2) and the Comparison Test. It follows from Proposition 4.8(8) that

$$\lim_{a \rightarrow \infty} \int_0^a \frac{\sin x}{x} dx = \int_0^1 \frac{\sin x}{x} dx + \cos 1 - \int_1^{\infty} \frac{\cos x}{x^2} dx.$$

Nevertheless,  $\sin x/x$  is not integrable over  $(0, \infty)$ , by Example 4.10(5).

In the case of substitution, one can infer the existence of one integral from the other. [Note: Priestley's comment near the bottom of p.133 is misleading.]

**Theorem 4.15. (Substitution)** *Let  $g : I \rightarrow \mathbb{R}$  be a monotonic function with a continuous derivative on an interval  $I$ , and let  $J$  be the interval  $g(I)$ . A (measurable) function  $f : J \rightarrow \mathbb{R}$  is integrable over  $J$  if and only if  $(f \circ g) \cdot g'$  is integrable over  $I$ . Then*

$$\int_J f(x) dx = \int_I f(g(y))|g'(y)| dy.$$

This theorem is not contained in the one in the first-year course, because  $f$  is not required to be continuous or Riemann integrable. FTC gives the result when  $f = \chi_{J'}$  for a bounded interval  $J' \subseteq J$ . One has to extend this to  $f = \chi_E$  when  $E \in \mathcal{M}_{\text{Leb}}, E \subseteq J$ , i.e., one needs  $m(E) = \int_{g^{-1}(E)} g'$ . After that, the rest follows fairly easily. See Theorem 7.4 in Qian's notes.

**Example 4.16.** Let  $I = (0, 1)$ ,  $g(y) = 1/y$ , so  $J = (1, \infty)$ . Let  $f(x) = x^\alpha$ . Then  $x^\alpha \in \mathcal{L}^1(1, \infty)$  if and only if  $y^{-\alpha-2} \in \mathcal{L}^1(0, 1)$ . This provides a passage between Example 4.10, (1) and (2).

**Other measures.** We make some comments about integration with respect to measures other than Lebesgue.

A function  $f : \mathbb{N} \rightarrow \mathbb{R}$  is integrable with respect to counting measure  $\mu$  if and only if  $\sum f(n)$  is absolutely convergent, and then  $\int f d\mu = \sum_{n=1}^{\infty} f(n)$ . Thus the general theorems that follow will provide theorems about summing absolutely convergent series.

Next, consider a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ . A measurable function is now just a random variable  $X$  on this space, and the integral of  $X$  with respect to  $\mathbb{P}$  is just the expectation  $\mathbb{E}(X)$ ;  $X$  is integrable if and only if  $|X|$  has finite expectation. The theory that follows applies to all random variables simultaneously—discrete, continuous, hybrid, singular.

## 5. THE CONVERGENCE THEOREMS

The feature of Lebesgue integration theory which distinguishes it from other theories, and makes it much more manageable, is the group of theorems known as convergence theorems. These are the theorems, mentioned in the introduction, which enable one to pass limits or infinite sums through integrals, under certain conditions. We have already seen the MCT, but we restate it here in a slightly more general form.

**Theorem 5.1. [Monotone Convergence Theorem, Version 2]** *Let  $(f_n)$  be a sequence of integrable functions such that:*

- (1) for each  $n$ ,  $f_n \leq f_{n+1}$  a.e.,
- (2)  $\sup_n \int f_n < \infty$ .

*Then  $(f_n)$  converges a.e. to an integrable function  $f$ , and  $\int f = \lim_{n \rightarrow \infty} \int f_n$ .*

*Proof.* Redefining  $f_n$  on the union of countably many null sets, we may assume that  $f_n(x) \leq f_{n+1}(x)$  and  $f_n(x) \in \mathbb{R}$  for all  $x$ . Apply Theorem 4.2 applied to  $f_n - f_1$ . One obtains that  $\int(f - f_1)$  is finite, which implies that  $f$  is finite a.e., by Proposition 4.8.  $\square$

We have not specified the range of integration. It could be  $\mathbb{R}$ , or it could be a fixed interval  $I$ . We can also apply the MCT when the the range of integration depends on  $n$ , by taking  $f_n$  to be 0 elsewhere.

**Example 5.2.** Consider  $\int_0^{n\pi} \left(\cos \frac{x}{2n}\right) x^2 e^{-x^3} dx$ . It is not obvious how to evaluate the integral for a given value of  $n$ , but we can use the MCT to find the limit of the integrals, as  $n \rightarrow \infty$ , as follows.

Let

$$f_n(x) = \begin{cases} \left(\cos \frac{x}{2n}\right) x^2 e^{-x^3} & \text{if } 0 \leq x \leq n\pi \\ 0 & \text{otherwise.} \end{cases}$$

Fix  $n$  for a moment. We wish to show that  $f_n(x) \leq f_{n+1}(x)$  for all  $x$ . If  $0 \leq x \leq n\pi$ , then  $\cos \frac{x}{2n} \leq \cos \frac{x}{2(n+1)}$ , so  $f_n(x) \leq f_{n+1}(x)$ . If  $n\pi < x \leq (n+1)\pi$ , then  $f_n(x) = 0 \leq f_{n+1}(x)$ . If  $x > (n+1)\pi$  (or if  $x < 0$ ), then  $f_n(x) = 0 = f_{n+1}(x)$ . Thus we have established our claim that  $f_n(x) \leq f_{n+1}(x)$  for all  $x$ .

Now

$$\int_0^\infty f_n(x) dx = \int_0^{n\pi} \left(\cos \frac{x}{2n}\right) x^2 e^{-x^3} dx \leq \int_0^{n\pi} x^2 e^{-x^3} dx = \frac{1 - e^{-n^3\pi^3}}{3} \nearrow \frac{1}{3}.$$

Thus the conditions of the MCT Theorem 5.1 are satisfied, and we conclude that  $f_n(x) \rightarrow f(x)$  a.e. for some integrable function  $f$ , and  $\int f_n \rightarrow \int f$ . In this case,  $f(x)$  is obvious. If we fix  $x \geq 0$ , then  $f_n(x) = \left(\cos \frac{x}{2n}\right) x^2 e^{-x^3}$  whenever  $n \geq x/\pi$ , so

$$f(x) = \lim_{n \rightarrow \infty} \left(\cos \frac{x}{2n}\right) x^2 e^{-x^3} = x^2 e^{-x^3}.$$

Thus the MCT tells us that

$$\lim_{n \rightarrow \infty} \int_0^{n\pi} \left(\cos \frac{x}{2n}\right) x^2 e^{-x^3} dx = \int_0^\infty x^2 e^{-x^3} dx = \frac{1}{3}.$$

**Theorem 5.3. [Fatou's Lemma]** *Let  $(f_n)$  be a sequence of non-negative measurable functions. Then*

$$\int \liminf_{n \rightarrow \infty} f_n \leq \liminf_{n \rightarrow \infty} \int f_n.$$

*Proof.* Let  $g_r := \inf_{n \geq r} f_n$ . Then  $(g_r)$  increases to  $\liminf_{n \rightarrow \infty} f_n$  and  $g_r \leq f_r$ , so, by MCT,  $\int \liminf_{n \rightarrow \infty} f_n = \lim_{r \rightarrow \infty} \int g_r \leq \liminf_{r \rightarrow \infty} \int f_r$ .  $\square$

Note that in Example 0.1 with  $f_n(x) = n^2 x^n (1-x)$  on  $(0, 1)$ ,  $f_n \geq 0$ ,  $\lim_{n \rightarrow \infty} f_n = 0$ ,  $\lim_{n \rightarrow \infty} \int f_n = 1$ . So one can have  $\int \limsup_{n \rightarrow \infty} f_n < \liminf_{n \rightarrow \infty} \int f_n$ . However if  $f_n \leq g$  for all  $n$  where  $g$  is integrable, then  $\int \limsup_{n \rightarrow \infty} f_n \geq \limsup_{n \rightarrow \infty} \int f_n$  (apply Fatou to  $g - f_n$ ).

One can also have  $\int \limsup_{n \rightarrow \infty} f_n > \limsup_{n \rightarrow \infty} \int f_n$ —for example,  $f_n(x) = \sin^2(x+n)$  on  $(0, \pi)$ .

**Theorem 5.4. [Dominated Convergence Theorem]** *Let  $(f_n)$  be a sequence of integrable functions such that:*

- (1)  $(f_n(x))$  converges a.e. to a limit  $f(x)$ ,
- (2) there is an integrable function  $g$  such that, for each  $n$ ,  $|f_n(x)| \leq g(x)$  a.e.

*Then  $f$  is integrable, and  $\int f = \lim_{n \rightarrow \infty} \int f_n$ .*

*Proof.* Since  $f$  is measurable (Proposition 3.7) and  $|f(x)| \leq g(x)$  a.e.,  $f$  is integrable by comparison. Apply Fatou's Lemma to  $g + f_n$  and  $g - f_n$ , to obtain  $\int(g + f) \leq \int g + \liminf_{n \rightarrow \infty} \int f_n$  and  $\int(g - f) \leq \int g - \limsup_{n \rightarrow \infty} \int f_n$ .  $\square$

**Example 5.5.** Consider  $\int_0^1 \frac{n^{3/2} x e^x}{1 + n^2 x^2} dx$ . It is difficult (impossible?) to evaluate the integrals themselves, but we can find the limit of the integrals, with the help of the DCT (Theorem 5.4). Let

$$f_n(x) = \frac{n^{3/2} x e^x}{1 + n^2 x^2} = \frac{(nx)^{3/2} e^x}{1 + n^2 x^2 x^{1/2}}.$$

The function  $\frac{y^{3/2}}{1 + y^2}$  tends to 0 as  $y \rightarrow \infty$ , so it is bounded for  $y > 0$ . It follows that  $f_n(x) \rightarrow 0$  as  $n \rightarrow \infty$ , and there is a constant  $c$  such that

$$0 \leq f_n(x) \leq \frac{ce^x}{x^{1/2}} \leq \frac{ce}{x^{1/2}} \quad (0 < x < 1).$$

Now let  $g(x) = \frac{ce}{x^{1/2}}$ . Then  $g$  is integrable over  $(0, 1)$  (Example 4.10(1)), so we have verified the conditions of the DCT (with  $f = 0$ ). We can therefore conclude that

$$\lim_{n \rightarrow \infty} \int_0^1 \frac{n^{3/2} x e^x}{1 + n^2 x^2} dx = 0.$$

**Corollary 5.6. [Bounded Convergence Theorem]** *Let  $I$  be a bounded interval,  $(f_n)$  be a sequence in  $\mathcal{L}^1(I)$  converging a.e. to  $f$ , and suppose that there is a constant  $c$  such that  $|f_n(x)| \leq c$  a.e., for all  $n$ . Then  $f \in \mathcal{L}^1(I)$ , and  $\int_I f = \lim_{n \rightarrow \infty} \int_I f_n$ .*

The next example involves, for the first time in this course, integration of a complex-valued function. A function  $f : \mathbb{R} \rightarrow \mathbb{C}$  is *integrable* if  $\operatorname{Re} f$  and  $\operatorname{Im} f$  are both integrable. Results which hold for real-valued integrable functions and which make sense for complex-valued functions are almost invariably true in the complex case, and can easily be deduced by applying the result to the real and imaginary parts separately. This is the case, for example, with the Comparison Test, FTC, Integration by Parts and the DCT. Note, however, that in Theorem 4.15 (Substitution), the function  $f$  may be complex-valued, but the substitution  $g(t)$  is assumed to be real-valued.

**Example 5.7.** Let  $\gamma_r$  be the semi-circular contour  $\{re^{i\theta} : 0 \leq \theta \leq \pi\}$ , and consider

$$\int_{\gamma_r} \frac{e^{iz}}{z} dz = i \int_0^\pi e^{ir \cos \theta} e^{-r \sin \theta} d\theta.$$

Since

$$\begin{aligned} \left| e^{ir \cos \theta} e^{-r \sin \theta} \right| &\leq 1 \quad \text{for all } r > 0, 0 \leq \theta \leq \pi \\ e^{ir \cos \theta} e^{-r \sin \theta} &\rightarrow \begin{cases} 0 & \text{as } r \rightarrow \infty, \text{ if } 0 < \theta < \pi, \\ 1 & \text{as } r \rightarrow 0 \end{cases} \end{aligned}$$

the Bounded Convergence Theorem gives

$$\int_{\gamma_{R_n}} \frac{e^{iz}}{z} dz \rightarrow 0 \quad (R_n \rightarrow \infty), \quad \int_{\gamma_{\varepsilon_n}} \frac{e^{iz}}{z} dz \rightarrow \pi i \quad (\varepsilon_n \rightarrow 0).$$

By Cauchy's Theorem,

$$0 = \int_{\gamma_{R_n}} \frac{e^{iz}}{z} dz - \int_{\gamma_{\varepsilon_n}} \frac{e^{iz}}{z} dz + \int_{\varepsilon_n}^{R_n} \frac{e^{ix} - e^{-ix}}{x} dx.$$

Letting  $n \rightarrow \infty$ , we obtain

$$\lim_{n \rightarrow \infty} \int_{\varepsilon_n}^{R_n} \frac{\sin x}{x} dx = \frac{\pi}{2}.$$

Hence  $\lim_{a \rightarrow \infty} \int_0^a \frac{\sin x}{x} dx = \pi/2$  (see Example 4.14, and Part A Complex Analysis, Example 22.9 in MT2020 notes).

Next we will apply the results above to term-by-term integration of series. We start by recalling the MCT for Series (Corollary 4.5 above).

**Theorem 5.8. [Monotone Convergence Theorem for Series]** *Let  $(g_n)$  be a sequence of integrable functions such that:*

- (1) for each  $n$ ,  $g_n \geq 0$  a.e.,
- (2)  $\sum_n \int g_n < \infty$ .

Then  $\sum_{n=1}^{\infty} g_n$  converges a.e. to an integrable function, and  $\int \sum_{n=1}^{\infty} g_n = \sum_{n=1}^{\infty} \int g_n$ .

**Theorem 5.9. [Lebesgue's Series Theorem; Beppo Levi Theorem, ...]** Let  $(g_n)$  be a sequence of integrable functions such that  $\sum_n \int |g_n| < \infty$ . Then  $\sum_{n=1}^{\infty} g_n$  converges a.e. to an integrable function, and  $\int \sum_{n=1}^{\infty} g_n = \sum_{n=1}^{\infty} \int g_n$ .

*Proof.* Apply MCT for Series to  $g_n^+$  and  $g_n^-$ . Alternatively, apply MCT for Series to  $|g_n|$  and use the fact that absolute convergence implies convergence.  $\square$

**Theorem 5.10.** Let  $(g_n)$  be a sequence of integrable functions such that  $\sum_n |g_n|$  is integrable. Then  $\sum_{n=1}^{\infty} g_n$  converges a.e. to an integrable function, and  $\int \sum_{n=1}^{\infty} g_n = \sum_{n=1}^{\infty} \int g_n$ .

*Proof.* Clearly  $\sum_{n=1}^k \int |g_n| \leq \int \sum_{n=1}^{\infty} |g_n|$  for all  $k$ , so  $\sum_{n=1}^{\infty} \int |g_n| \leq \int \sum_{n=1}^{\infty} |g_n|$ . Apply Theorem 5.9.  $\square$

**Example 5.11.** Let  $\alpha > 0$ , and consider  $\int_0^1 x^{\alpha-1} e^{-x} dx$ . Let  $g_n(x) = (-1)^n x^{\alpha+n-1}/n!$ , so that  $\sum_{n=0}^{\infty} g_n(x) = x^{\alpha-1} e^{-x}$ . Now

$$\int_0^1 |g_n(x)| dx = \frac{1}{(\alpha+n)n!},$$

so  $\sum_n \int_0^1 |g_n(x)| dx < \infty$ . Thus Lebesgue's Series Theorem tells us that our integral exists (we could have established this directly, by comparing the integrand with  $x^{\alpha-1}$ ), and that

$$\int_0^1 x^{\alpha-1} e^{-x} dx = \sum_{n=0}^{\infty} \int_0^1 (-1)^n x^{\alpha+n-1}/n! dx = \sum_{n=0}^{\infty} \frac{(-1)^n}{(\alpha+n)n!}.$$

**Example 5.12.** Let  $s \in \mathbb{R}$ , and consider  $\int_{-\infty}^{\infty} e^{-isx} e^{-x^2} dx$ . The integrand is continuous,  $|e^{-isx} e^{-x^2}| = e^{-x^2} \leq e^{-|x|} \in \mathcal{L}^1$  (exercise). If  $g_n(x) = \frac{(-isx)^n}{n!} e^{-x^2}$ , then  $\sum_{n=0}^{\infty} g_n(x) = e^{-isx} e^{-x^2}$ , and

$$\sum_{n=0}^{\infty} |g_n(x)| = e^{|sx|-x^2} \leq e^{s^2/2} e^{-x^2/2} \in \mathcal{L}^1.$$

It follows that  $\sum_n |g_n| \in \mathcal{L}^1$ , so Theorem 5.10 shows that term-by-term integration is permissible, and

$$\int_{-\infty}^{\infty} e^{-isx} e^{-x^2} dx = \sum_{n=0}^{\infty} \int_{-\infty}^{\infty} \frac{(-isx)^n}{n!} e^{-x^2} dx.$$

Now

$$\int_{-\infty}^{\infty} x^n e^{-x^2} dx = \begin{cases} 0 & \text{if } n \text{ is odd} \\ \frac{(2m)! \sqrt{\pi}}{4^m m!} & \text{if } n = 2m, \end{cases}$$

(for  $m = 0$  this is a standard trick, and one can use integration by parts and induction on  $m$ ). Thus

$$\int_{-\infty}^{\infty} e^{-isx} e^{-x^2} dx = \sum_{m=0}^{\infty} \frac{(-is)^{2m} \sqrt{\pi}}{4^m m!} = \sqrt{\pi} e^{-s^2/4}.$$

The integral which we have just evaluated is very important—for example, apart from a few constants, it is the characteristic function of the normal distribution (as in Part A Probability); in analysts' language, it is the Fourier transform of the function  $e^{-x^2}$  (as in DEs). There are other methods of evaluating the integral; one is given in Priestley (*Complex Analysis*, 22.12) and Part A Integral Transforms (Example 77 in HT2020 notes), and another will be given in Example 6.6.

All theorems in this Section hold in general measure spaces. Corollary 5.6 holds in finite measure spaces.

## 6. INTEGRALS DEPENDING ON A PARAMETER

Let  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  be a function of two variables. In a while, we shall discuss the (double) integral, and the repeated integrals, of  $f$ . First, we merely consider the partial integral of  $f$ , obtained by integration with respect to one of the variables. Thus we suppose that for each fixed  $y$ , the function  $x \mapsto f(x, y)$  is integrable. We can then define a function  $F$  by:

$$F(y) = \int f(x, y) dx.$$

A natural, and important, question is whether  $F$  is continuous, or differentiable, assuming that  $f$  has corresponding properties. In general, the answer is negative (see Example 6.1), but if we impose some mild conditions of the type that appear in the DCT, then the answer is positive.

**Example 6.1.** Let  $f(x, y) = ye^{-x^2 y^2}$ . Since  $f(x, 0) = 0$  for all  $x$ ,  $F(0) = 0$ . For fixed  $y \neq 0$ , we can make the substitution  $t = yx$  and deduce that  $F(y) = \int_{-\infty}^{\infty} e^{-t^2} dt (= \sqrt{\pi}) (y \neq 0)$ . Thus  $F$  is discontinuous, even though  $f$  is differentiable.

**Theorem 6.2. [Continuous-parameter DCT]** *Let  $I$  and  $J$  be intervals in  $\mathbb{R}$ , and  $f : I \times J \rightarrow \mathbb{R}$  be a function such that:*

- (1) *for each  $y$  in  $J$ ,  $x \mapsto f(x, y)$  is integrable over  $I$ ,*
- (2) *for each  $y$  in  $J$ ,  $\lim_{y' \rightarrow y} f(x, y') = f(x, y)$  a.e.( $x$ ),*
- (3) *there exists  $g \in \mathcal{L}^1(I)$  such that for each  $y$  in  $J$ ,  $|f(x, y)| \leq g(x)$  a.e.( $x$ ).*

*Define  $F(y) = \int_I f(x, y) dx$  ( $y \in J$ ). Then  $F$  is continuous on  $J$ .*

*Remark.* In condition (3) of Theorem 6.2, the function  $g$  does not depend on  $y$ .

*Proof.* Let  $(y_n)$  be any sequence in  $J$  converging to  $y \in J$ . Let  $f_n(x) = f(x, y_n)$ . Then  $|f_n(x)| \leq g(x)$  a.e., for all  $n$ , and  $\lim_{n \rightarrow \infty} f_n(x) = f(x, y)$  a.e., so the conditions of the



DCT are satisfied. The DCT implies that:

$$F(y_n) = \int_I f(x, y_n) dx \rightarrow \int_I f(x, y) dx = F(y).$$

Thus  $F$  is continuous. □

**Example 6.3.** The Gamma function  $\Gamma$  is defined by:

$$\Gamma(y) = \int_0^\infty e^{-x} x^{y-1} dx \quad (y > 0).$$

We wish to show that  $\Gamma$  is continuous, firstly for  $y \in [1, 2]$ . In order to apply Theorem 6.2, we take  $I = (0, \infty)$ ,  $J = [1, 2]$ , and  $f(x, y) = e^{-x} x^{y-1}$ . Condition (1) of Theorem 6.2 is an exercise, and (2) is more or less trivial. For condition (3), we need to ensure that

$$(6.1) \quad g(x) \geq \sup_{1 \leq y \leq 2} f(x, y) = \begin{cases} e^{-x} & (0 < x \leq 1) \\ x e^{-x} & (x > 1). \end{cases}$$

We choose to take  $g$  equal to the right-hand side of (6.1). Then  $g$  is integrable over  $(0, \infty)$  (exercise), so condition (3) of Theorem 6.2 is satisfied. Thus, Theorem 6.2 shows that  $\Gamma$  is continuous on  $[1, 2]$ .

In fact,  $\Gamma$  is continuous on  $(0, \infty)$ . However, it is impossible to establish this by applying Theorem 6.2 with  $J = (0, \infty)$ , for in condition (3), it would be necessary that

$$g(x) \geq \sup_{y > 0} f(x, y) = \begin{cases} x^{-1} e^{-x} & (0 < x \leq 1) \\ \infty & (x > 1). \end{cases}$$

Such a function  $g$  cannot possibly be integrable over  $(0, \infty)$ , so it is impossible to satisfy condition (3) of Theorem 6.2. Instead, we proceed as follows. For each  $b > 0$ , let  $J_b = (a, c)$ , where  $a$  and  $c$  are chosen so that  $0 < a < b < c$ , for example,  $a = b/2$  and  $c = 2b$ . Then let

$$g_b(x) = \sup_{a < y < c} f(x, y) = \begin{cases} x^{a-1} e^{-x} & (0 < x \leq 1) \\ x^{c-1} e^{-x} & (x > 1). \end{cases}$$

Then  $g_b$  is integrable over  $(0, \infty)$ . Thus, Theorem 6.2 shows that  $\Gamma$  is continuous on  $(a, c)$ , and in particular at  $b$ . But  $b$  is arbitrary, so  $\Gamma$  is continuous on  $(0, \infty)$ .

We abstract this method to obtain the following version of Theorem 6.2, where the dominating function  $g$  depends on the parameter to some extent.

**Corollary 6.4.** *Let  $I$  and  $J$  be intervals in  $\mathbb{R}$ , and  $f : I \times J \rightarrow \mathbb{R}$  be a function such that (1) and (2) of Theorem 6.2 hold, and*

- (3') *for each  $b \in J$ , there is an open subinterval  $J_b$  of  $J$  containing  $b$  and  $g_b \in \mathcal{L}^1(I)$  such that, for each  $y \in J_b$ ,  $|f(x, y)| \leq g_b(x)$  a.e.( $x$ ).*

*Then  $F$  is continuous on  $J$ , where  $F$  is as in Theorem 6.2.*

*Remark.* The method of Theorem 6.2 can also be used to cover cases where  $y \rightarrow y_0$  for a single point  $y_0$  or  $y \rightarrow \infty$ . For example, suppose that there exists  $a$  in  $\mathbb{R}$  and a function  $h : I \rightarrow \mathbb{R}$  such that

- (1) for each  $y > a$ ,  $x \mapsto f(x, y)$  is integrable over  $I$ ,
- (2)  $\lim_{y \rightarrow \infty} f(x, y) = h(x)$  a.e.( $x$ ),
- (3) there exists  $g \in \mathcal{L}^1(I)$  such that for each  $y > a$ ,  $|f(x, y)| \leq g(x)$  a.e.( $x$ ).

Then  $F(y) \rightarrow \int_I h(x) dx$  as  $y \rightarrow \infty$ .

Now we turn to the question of differentiability of  $F$ . The sort of result which we hope to have is that if  $\frac{\partial f}{\partial y}$  exists, and some supplementary conditions are satisfied, then  $F$  is differentiable and

$$F'(y) = \int \frac{\partial f}{\partial y}(x, y) dx$$

(differentiation through, or under, the integral sign). The standard supplementary condition is that  $\frac{\partial f}{\partial y}$  should be dominated by an integrable function, *independent of  $y$* .

**Theorem 6.5.** *Let  $I$  and  $J$  be intervals in  $\mathbb{R}$ , and  $f : I \times J \rightarrow \mathbb{R}$  be a function such that:*

- (1) for each  $y$  in  $J$ ,  $x \mapsto f(x, y)$  is integrable over  $I$ ,
- (2) for each  $x$  in  $I$  and  $y$  in  $J$ ,  $\frac{\partial f}{\partial y}(x, y)$  exists,
- (3) there is an integrable function  $g : I \rightarrow \mathbb{R}$  such that for each  $y$  in  $J$ ,  $\left| \frac{\partial f}{\partial y}(x, y) \right| \leq g(x)$  a.e.( $x$ ).

Define  $F(y) = \int_I f(x, y) dx$  ( $y \in J$ ). Then  $F$  is differentiable on  $J$  and

$$F'(y) = \int_I \frac{\partial f}{\partial y}(x, y) dx.$$

*Proof.* Fix  $y$  in  $J$ , and let  $(y_n)$  be any sequence in  $J$  converging to  $y$  (with  $y_n \neq y$ ). Let

$$g_n(x) = \frac{f(x, y_n) - f(x, y)}{y_n - y}.$$

Then  $g_n$  is integrable over  $I$ ,  $g_n(x) \rightarrow \frac{\partial f}{\partial y}(x, y)$  as  $n \rightarrow \infty$ . Moreover, the Mean Value Theorem says that there exists a point  $\xi$  (depending on  $x$  and  $n$ ) between  $y_n$  and  $y$  such that  $g_n(x) = \frac{\partial f}{\partial y}(x, \xi)$ . It follows from (3) that  $|g_n(x)| \leq g(x)$  a.e.( $x$ ). This shows that the Dominated Convergence Theorem is applicable, so

$$\frac{F(y_n) - F(y)}{y_n - y} = \int_I g_n(x) dx \rightarrow \int_I \frac{\partial f}{\partial y}(x, y) dx \quad \text{as } n \rightarrow \infty.$$

Since  $(y_n)$  is an arbitrary sequence tending to  $y$ , and the right-hand side is independent of the choice of sequence, it follows that

$$\frac{F(y') - F(y)}{y' - y} \rightarrow \int_I \frac{\partial f}{\partial y}(x, y) dx \quad \text{as } y' \rightarrow y,$$

which completes the proof. □

**Example 6.6.** Let  $f(x, s) = e^{-isx}e^{-x^2}$ , and  $F(s) = \int_{-\infty}^{\infty} f(x, s) dx$  (compare Example 5.12). This integral exists for all  $s$ . Moreover,

$$\frac{\partial f}{\partial s}(x, s) = -ixe^{-ixs}e^{-x^2},$$

so

$$\left| \frac{\partial f}{\partial s}(x, s) \right| = |x|e^{-x^2}.$$

Since

$$\int_{-n}^n |x|e^{-x^2} dx = 2 \int_0^n xe^{-x^2} dx = 1 - e^{-n^2} \rightarrow 1$$

as  $n \rightarrow \infty$ ,  $|x|e^{-x^2} \in \mathcal{L}^1(\mathbb{R})$  (Baby MCT). Thus Theorem 6.5 is applicable, with  $I = J = \mathbb{R}$  and  $g(x) = |x|e^{-x^2}$ . It follows that  $F$  is differentiable on  $\mathbb{R}$ , and

$$F'(s) = -i \int_{-\infty}^{\infty} xe^{-isx} e^{-x^2} dx.$$

By integration by parts,

$$F'(s) = -\frac{s}{2}F(s).$$

Hence  $F(s) = Ae^{-s^2/4}$  for some constant  $A$ . But  $F(0) = \int_{-\infty}^{\infty} e^{-x^2} dx = \sqrt{\pi}$ , so  $A = \sqrt{\pi}$ .

**Corollary 6.7.** *Let  $I$  and  $J$  be intervals in  $\mathbb{R}$ , and  $f : I \times J \rightarrow \mathbb{R}$  be a function such that (1) and (2) of Theorem 6.5 hold, and*

(3') *for each  $b$  in  $J$ , there is an open subinterval  $J_b$  of  $J$  containing  $b$  and an integrable function  $g_b : I \rightarrow \mathbb{R}$  such that, for each  $y \in J_b$ ,  $\left| \frac{\partial f}{\partial y}(x, y) \right| \leq g_b(x)$  a.e.( $x$ ).*

*Then the conclusions of Theorem 6.5 hold.*

**Example 6.8.** Let  $f(x, y) = e^{-xy}(1+x^3)^{-1}$  ( $x \geq 0, y \geq 0$ ). Since  $0 \leq f(x, y) \leq (1+x^3)^{-1}$ ,  $x \mapsto f(x, y)$  is integrable over  $[0, \infty)$  for each  $y \geq 0$ . Moreover,

$$\frac{\partial f}{\partial y}(x, y) = -\frac{xe^{-xy}}{1+x^3},$$

so

$$\left| \frac{\partial f}{\partial y}(x, y) \right| \leq \frac{x}{1+x^3} \quad (x \geq 0, y \geq 0).$$

Since  $x(1+x^3)^{-1}$  is integrable over  $[0, \infty)$  (by comparison with  $x^{-2}$  for  $x \geq 1$ ), Theorem 6.5 is applicable, and shows that  $F$  is differentiable on  $[0, \infty)$  and

$$F'(y) = - \int_0^{\infty} \frac{xe^{-xy}}{1+x^3} dx.$$

We would like to repeat this argument to show that  $F''(y)$  exists (at least for  $y > 0$ ), but this is more complicated. Indeed,

$$\frac{\partial^2 f}{\partial y^2}(x, y) = \frac{x^2 e^{-xy}}{1+x^3}.$$

For  $y = 0$ , this function is not integrable (by comparison with  $x^{-1}$ ), so we should only consider  $y > 0$ . However, it is not possible to apply Theorem 6.5 with  $f$  replaced by

$\frac{\partial f}{\partial y}$  and with  $J = (0, \infty)$ , because

$$\sup_{y>0} \frac{\partial^2 f}{\partial y^2}(x, y) = \frac{x^2}{1+x^3},$$

which is not integrable over  $[0, \infty)$ . Instead, we must apply Corollary 6.7. Thus we take  $b > 0$ , let  $J_b = (b/2, \infty)$ , and

$$g_b(x) = \sup_{y>b/2} \frac{\partial^2 f}{\partial y^2}(x, y) = \frac{x^2 e^{-xb/2}}{1+x^3} \leq x^2 e^{-xb/2}.$$

This function is integrable on  $[0, \infty)$ , and we conclude from Corollary 6.7, with  $f$  replaced by  $\frac{\partial f}{\partial y}$  and  $J = (0, \infty)$  that  $F''(y)$  exists for  $y > 0$  and

$$F''(y) = \int_0^\infty \frac{x^2 e^{-xy}}{1+x^3} dx.$$

Repeating this argument, it is possible to show that  $F$  is infinitely differentiable on  $(0, \infty)$  and to obtain integrals for all the derivatives.

## 7. DOUBLE INTEGRALS

In Section 6, we considered some properties concerning functions of two variables, but we confined integration to one of the variables. Now it is time to consider integration with respect to both variables. An example on Problem Sheet 1 shows that this is not just a matter of integrating first with respect to one variable, and then with respect to the other (repeated integration). What one has to do is to define the class  $\mathcal{L}^1(\mathbb{R}^2)$  of integrable functions on  $\mathbb{R}^2$ , and their (double) integrals, in a way which treats both variables simultaneously, then establish the theorem (Fubini's Theorem) which ensures that the double integrals coincide with the repeated integrals for functions in  $\mathcal{L}^1(\mathbb{R}^2)$ , and establish a practical method (Tonelli's Theorem) to determine whether a given function is integrable.

The first part of this is routine. The class  $\mathcal{L}^1(\mathbb{R}^2)$  of integrable functions on  $\mathbb{R}^2$  is defined in exactly the same way as  $\mathcal{L}^1(\mathbb{R})$ , except that intervals  $(a, b)$ , and their lengths  $b - a$ , are replaced by rectangles  $(a, b) \times (c, d)$  and their areas  $(b - a)(d - c)$ . Then one defines outer measure, null sets (line segments etc are null), measurable sets (all open sets etc are measurable), measurable functions, simple functions, integrable functions and (double) integrals just as in Sections 2–4. Moreover, the results of Sections 2–6, except Section 4 from Theorem 4.11 onwards, remain valid, with obvious changes of wording where necessary. More details may be found in Capinski & Kopp (Chap 6, but in greater generality) or Stein & Shakarchi (from beginning).

The (double) integral of an integrable function  $f$  over  $\mathbb{R}^2$  may be denoted by any of the following:

$$\int f, \quad \int_{\mathbb{R}^2} f, \quad \int f(x, y) d(x, y), \quad \int_{\mathbb{R}^2} f(x, y) d(x, y).$$

**Theorem 7.1. (Tonelli)** *Let  $f : \mathbb{R}^2 \rightarrow [0, \infty]$  be measurable. Then*

- (1)  $x \mapsto \int f(x, y) dy$  is measurable for almost all  $x$ ;

- (2)  $y \mapsto \int_{\mathbb{R}} f(x, y) dx$  (defined a.e.) is measurable;  
 (3)

$$\int_{\mathbb{R}^2} f(x, y) d(x, y) = \int_{\mathbb{R}} \left( \int_{\mathbb{R}} f(x, y) dx \right) dy.$$

Now we state two consequences of this in their traditional form.

**Theorem 7.2. [Fubini's Theorem]** *Let  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  be integrable. Then, for almost all  $y$ , the function  $x \mapsto f(x, y)$  is integrable. Moreover, if  $F(y)$  is defined (for almost all  $y$ ) by  $F(y) = \int f(x, y) dx$ , then  $F$  is integrable, and*

$$\int_{\mathbb{R}^2} f(x, y) d(x, y) = \int_{\mathbb{R}} \left( \int_{\mathbb{R}} f(x, y) dx \right) dy.$$

Similarly,

$$\int_{\mathbb{R}} \left( \int_{\mathbb{R}} f(x, y) dx \right) dy = \int_{\mathbb{R}^2} f(x, y) d(x, y) = \int_{\mathbb{R}} \left( \int_{\mathbb{R}} f(x, y) dy \right) dx,$$

where the first repeated integral exists in the sense described above.

*Proof.* Apply Theorem 7.1 to  $f^+$  and  $f^-$ , using Proposition 4.8(6) to get that  $\int_{\mathbb{R}} f^{\pm}(x, y) dx < \infty$  a.e.( $y$ ). □

**Theorem 7.3. [Tonelli's Theorem]** *Let  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  be a measurable function, and suppose that either of the following repeated integrals is finite:*

$$\int_{\mathbb{R}} \left( \int_{\mathbb{R}} |f(x, y)| dx \right) dy, \quad \int_{\mathbb{R}} \left( \int_{\mathbb{R}} |f(x, y)| dy \right) dx.$$

*Then  $f$  is integrable. Hence, Fubini's Theorem is applicable to both  $f$  and  $|f|$ .*

*Proof.* Apply Theorem 7.1 to get that  $\int_{\mathbb{R}^2} |f| < \infty$ . Then  $f \in \mathcal{L}^1(\mathbb{R}^2)$ , by Proposition 4.8(2). □

*Remark.* Note that, when applying Tonelli's Theorem, one must verify that a repeated integral of  $|f|$  is finite. It is not sufficient that the repeated integrals of  $f$  exist (see Example 7.4), nor is it sufficient that the repeated integrals of  $f$  both exist and are equal (see Example 7.7).

If  $E$  is a measurable subset of  $\mathbb{R}^2$  and  $f : E \rightarrow \mathbb{R}$  is any function, then  $f$  is said to be integrable over  $E$  if  $\tilde{f}$  is integrable over  $\mathbb{R}^2$ , where  $\tilde{f}(x, y) = f(x, y)$  if  $(x, y) \in E$ ,  $\tilde{f}(x, y) = 0$  otherwise. Then  $\int_E f$  is defined to be  $\int_{\mathbb{R}^2} \tilde{f}$ .

Fubini's Theorem and Tonelli's Theorem can be applied in this situation. However, when  $E$  is not a rectangle, great care must be taken to choose the correct limits of integration in the repeated integrals. If in any doubt draw a sketch of the region. See Example 7.5.

In repeated integrals, one often omits the brackets around the inner integral and writes  $\iint f(x, y) dy dx$ , etc., with appropriate limits of integration. This means that

one is integrating first with respect to  $y$  between the limits on the right-hand integral sign, which may be functions of  $x$ . Thus

$$\int_a^b \int_{\phi(x)}^{\psi(x)} f(x, y) dy dx$$

denotes the repeated integral over the region  $E$  bounded by curves  $y = \phi(x)$  and  $y = \psi(x)$  and by vertical lines  $x = a$ ,  $x = b$ .

**Example 7.4.** Let  $f(x, y) = \frac{x-y}{(x+y)^3}$  ( $0 < x < 1, 0 < y < 1$ ). It was an exercise in Problem Sheet 1 that the repeated integrals of  $f$  exist, but are not equal. It follows from the final part of Fubini's Theorem that  $f$  is not integrable over the square  $(0, 1) \times (0, 1)$ .

**Example 7.5.** Consider  $\int_0^1 \left( \int_0^x \left( \frac{1-y}{x-y} \right)^{1/2} dy \right) dx$ . As it stands, the inner integral is difficult. However, it turns out that when the order of integration is reversed, the other repeated integral is easily evaluated. To justify the equality of the repeated integrals, we apply Tonelli's Theorem; this is contained in the following discussion.

First, note that the integrand is continuous except on the line  $y = x$  which is null; it is non-negative throughout the range of integration, so that in applying Tonelli's Theorem, it is unnecessary to replace  $f$  by  $|f|$ . The next problem is to work out the limits of integration when we reverse the order. For this, we have to identify the region in  $\mathbb{R}^2$  over which the double integral is taken. For each  $x$ , between 0 and 1, we are integrating along the (vertical) line-segment from  $y = 0$  to  $y = x$ . As  $x$  runs from 0 to 1, this sweeps out the triangle shown. The integrand is continuous on the interior of the triangle (and we take it to be 0 outside the triangle), so it is measurable. If we fix a value of  $y$ , the values of  $x$  which give us points within the triangle are those between  $x = y$  and  $x = 1$ . This applies for  $y$  between 0 and 1; otherwise there are no points within the triangle. Thus the limits of the reversed repeated integral are  $x = y$  and  $x = 1$  in the inner integral, and  $y = 0$  and  $y = 1$  in the outer. This is confirmed by the following equalities of sets:

$$\begin{aligned} \{(x, y) \in \mathbb{R}^2 : 0 < y < x, 0 < x < 1\} &= \{(x, y) \in \mathbb{R}^2 : 0 < y < x < 1\} \\ &= \{(x, y) \in \mathbb{R}^2 : y < x < 1, 0 < y < 1\}, \end{aligned}$$

but the picture was more informative!

Now the reversed repeated integral is:

$$\begin{aligned} \int_0^1 \left( \int_y^1 \left( \frac{1-y}{x-y} \right)^{1/2} dx \right) dy &= \int_0^1 \left[ 2(1-y)^{1/2}(x-y)^{1/2} \right]_{x=y}^{x=1} dy \\ &= \int_0^1 2(1-y) dy = 1. \end{aligned}$$

Since the integrand is non-negative, and since this repeated integral is finite, it follows from Tonelli's Theorem that  $f$  is integrable over the triangle, and from Fubini's Theorem that

$$\int_0^1 \left( \int_0^x \left( \frac{1-y}{x-y} \right)^{1/2} dy \right) dx = 1.$$

The next example shows how it is both possible and useful to make changes of variable within the inner integral of a repeated integral. The same technique will be used in several subsequent examples.

**Example 7.6.** Let  $f(x, y) = ye^{-y^2(1+x^2)}$ . Since  $f$  is continuous, it is certainly measurable. We shall consider the integral of  $f$  over the positive quadrant  $(0, \infty) \times (0, \infty)$ . First we consider  $\int_0^\infty f(x, y) dy$  for a fixed  $x$ . Making the change of variable  $t = y(1+x^2)^{1/2}$  ( $x$  is a constant at this point),

$$\int_0^\infty f(x, y) dy = \int_0^\infty \frac{te^{-t^2}}{1+x^2} dt = \lim_{k \rightarrow \infty} \left[ -\frac{e^{-t^2}}{2(1+x^2)} \right]_{t=0}^{t=k} = \frac{1}{2(1+x^2)}.$$

This function is integrable with respect to  $x$ , and

$$\int_0^\infty \left( \int_0^\infty f(x, y) dy \right) dx = \frac{\pi}{4}.$$

Since  $f(x, y) \geq 0$  for  $y \geq 0$ , it follows from Tonelli's Theorem that  $f$  is integrable over  $(0, \infty) \times (0, \infty)$ , and by Fubini's Theorem,

$$\int_0^\infty \left( \int_0^\infty f(x, y) dx \right) dy = \frac{\pi}{4}.$$

In the inner integral, where  $y > 0$  is fixed, we can make the change of variable  $u = xy$ , and obtain

$$\begin{aligned} \frac{\pi}{4} &= \int_0^\infty \left( \int_0^\infty e^{-(y^2+u^2)} du \right) dy = \int_0^\infty e^{-y^2} \left( \int_0^\infty e^{-u^2} du \right) dy \\ &= \left( \int_0^\infty e^{-u^2} du \right) \left( \int_0^\infty e^{-y^2} dy \right) = \left( \int_0^\infty e^{-x^2} dx \right)^2. \end{aligned}$$

It follows that

$$\int_0^\infty e^{-x^2} dx = \frac{\sqrt{\pi}}{2}.$$

If  $f$  takes both positive and negative values, then to apply Tonelli's Theorem, it is necessary to consider  $|f|$ , or alternatively to consider separately the regions where  $f$  is positive and where it is negative.

**Example 7.7.** Let  $f(x, y) = \frac{xy}{x^4 + y^4}$ . Since  $f$  is odd both as a function of  $x$ , and also as a function of  $y$ ,  $\int_{-\infty}^\infty f(x, y) dy = 0$  for all  $x$ , and  $\int_{-\infty}^\infty f(x, y) dx = 0$  for all  $y$ . Hence

both repeated integrals exist and equal 0. However, if we consider  $f$  over the quadrant  $x > 0, y > 0$ , part of the region where  $f(x, y) > 0$ , then, putting  $y = xt$  ( $x > 0$  fixed),

$$\int_0^\infty f(x, y) dy = \int_0^\infty \frac{x^3 t}{x^4(1+t^4)} dt = \frac{c}{x},$$

where  $c$  is the constant  $\int_0^\infty \frac{t}{1+t^4} dt$ . Since  $cx^{-1}$  is not integrable with respect to  $x$  over  $(0, \infty)$ , it follows that  $f$  is not integrable over the quadrant, and therefore not integrable over the plane.

In practice, it often happens that one has no means of evaluating the repeated integrals of  $f$  or  $|f|$ , but can nevertheless decide whether  $f$  is integrable. One technique for this is to show that  $f$  is dominated by a simpler function which one can show to be integrable (or that  $f$  dominates a function which one can show not to be integrable).

**Example 7.8.** Let  $f(x, y) = \sin\left(\frac{1}{x^2 + y^4}\right) \cos(x^2 + y^3)$ . We wish to show that  $f$  is integrable over the positive quadrant  $(0, \infty) \times (0, \infty)$ . Since  $f$  is continuous in this region (although not continuous at  $(0, 0)$ ), it is measurable. Moreover,  $f$  is bounded, and hence integrable over any bounded region, in particular over the square  $(0, 1) \times (0, 1)$ . Thus it suffices to show that  $f$  is integrable over the regions  $[1, \infty) \times [0, \infty)$  and  $(0, 1) \times (1, \infty)$ .

Using the inequalities  $|\sin t| \leq |t|$  and  $|\cos t| \leq 1$ , it follows that  $|f(x, y)| \leq (x^2 + y^4)^{-1}$ , so it suffices to show that  $(x^2 + y^4)^{-1}$  is integrable over the regions. But

$$\int_1^\infty \left( \int_0^\infty \frac{dy}{x^2 + y^4} \right) dx = \int_1^\infty \left( \int_0^\infty \frac{dz}{x^{3/2}(1+z^4)} \right) dx < \infty,$$

where we made the substitution  $y = x^{1/2}z$  and used the integrability of  $x^{-3/2}$  over  $[1, \infty)$  and of  $(1+z^4)^{-1}$  over  $(0, \infty)$ . Also,

$$\int_1^\infty \left( \int_0^1 \frac{dx}{x^2 + y^4} \right) dy \leq \int_1^\infty \left( \int_0^1 \frac{dx}{y^4} \right) dy = \int_1^\infty \frac{dy}{y^4} = \frac{1}{3}.$$

It follows from Tonelli's Theorem that  $(x^2 + y^4)^{-1}$  is integrable over both strips, so  $f$  is integrable over the quadrant.

Another useful technique for testing functions for integrability, and for evaluating integrals, is to change variables. The reader will be familiar with this idea from courses in applied mathematics and in A3 Probability, and will know that one has to take account of the Jacobian of the transformation. The method is the extension to two variables of Theorem 4.15. We shall state the result and give examples for polar coordinates  $x = r \cos \theta$ ,  $y = r \sin \theta$ , when the Jacobian is  $r$ . This corresponds to the fact that a small rectangle with sides  $\delta r, \delta \theta$  (area  $\delta r \delta \theta$ ) in the  $(r, \theta)$ -space is transformed into an approximate rectangle of sides  $\delta r, r \delta \theta$  (area  $r \delta r \delta \theta$ ) in the  $(x, y)$ -space.



**Theorem 7.9.** Let  $E$  be a measurable subset of  $\mathbb{R}^2$ , and  $f : E \rightarrow \mathbb{R}$  be a function. Let  $E' = \{(r, \theta) : 0 \leq r, 0 \leq \theta < 2\pi, (r \cos \theta, r \sin \theta) \in E\}$  and  $g(r, \theta) = rf(r \cos \theta, r \sin \theta)$  ( $r, \theta \in E'$ ). Then  $f$  is integrable over  $E$  if and only if  $g$  is integrable over  $E'$ . In that case,

$$\int_E f(x, y) d(x, y) = \int_{E'} f(r \cos \theta, r \sin \theta) r d(r, \theta).$$

**Example 7.10.** In Example 7.6 we evaluated  $\int_0^\infty e^{-x^2} dx$ , using Fubini's Theorem. Here, we shall evaluate the same integral by the more common method of polar coordinates.

Let  $E = (0, \infty) \times (0, \infty)$  and  $f(x, y) = e^{-(x^2+y^2)}$ . Then

$$\int_0^\infty \int_0^\infty f(x, y) dy dx = \left( \int_0^\infty e^{-x^2} dx \right) \left( \int_0^\infty e^{-y^2} dy \right) = \left( \int_0^\infty e^{-x^2} dx \right)^2 < \infty.$$

It follows from Tonelli's Theorem that  $f$  is integrable over  $E$ . In the notation of Theorem 7.9,  $E' = \{(r, \theta) : 0 < r, 0 < \theta \leq \pi/2\}$ , so it follows from Theorem 7.9 and Fubini's Theorem that

$$\left( \int_0^\infty e^{-x^2} dx \right)^2 = \int_{E'} e^{-r^2} r d(r, \theta) = \int_0^{\pi/2} \int_0^\infty e^{-r^2} r dr d\theta = \frac{\pi}{4}.$$

This confirms that  $\int_0^\infty e^{-x^2} dx = \sqrt{\pi}/2$ .

**Example 7.11.** As in Example 7.7, let  $f(x, y) = \frac{xy}{x^4 + y^4}$ . In the notation of Theorem 7.9,  $g(r, \theta) = \frac{1}{r} \frac{\sin \theta \cos \theta}{\sin^4 \theta + \cos^4 \theta}$ . Since  $g$  is not integrable over  $[0, \infty) \times [0, 2\pi)$  (because  $r^{-1}$  is not integrable over  $[0, \infty)$ ),  $f$  is not integrable over  $\mathbb{R}^2$ .

**Example 7.12.** Let  $f(x, y) = \frac{x^2 - y^2}{(x^2 + y^2)^2}$ . The square  $(0, 1) \times (0, 1)$  is not very convenient for polar coordinates, but we can easily overcome this problem. Since  $f$  is bounded, hence integrable, over the bounded region  $\{(x, y) : 0 < x < 1, 0 < y < 1, 1 < x^2 + y^2\}$ ,  $f$  is integrable over the square if and only if it is integrable over the quadrant  $E = \{(x, y) : 0 < x < 1, 0 < y < 1, x^2 + y^2 \leq 1\}$ . In the notation of Theorem 7.9,  $E' = \{(r, \theta) : 0 < r \leq 1, 0 < \theta < \pi/2\}$  and

$$g(r, \theta) = r \frac{r^2(\cos^2 \theta - \sin^2 \theta)}{r^4} = \frac{\cos 2\theta}{r}.$$

Since  $r^{-1}$  is not integrable over  $(0, 1)$ ,  $g$  is not integrable over the rectangle  $E'$  (in  $(r, \theta)$ -space), so  $f$  is not integrable over  $E$ .

Now we state a version of Theorem 7.9 for general changes of coordinates. Let  $T : (u, v) \mapsto (x, y)$  be a change of variables, and suppose that  $x, y$  are differentiable functions of  $u, v$ . Let  $J_T$  be the Jacobian matrix:

$$J_T = \begin{pmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{pmatrix}.$$

Observe that  $J_{S \circ T} = J_S J_T$  (Chain Rule).

**Theorem 7.13.** *Let  $E'$  be an open subset of  $\mathbb{R}^2$ ,  $T : E' \rightarrow \mathbb{R}^2$  be a one-to-one differentiable function of  $E'$  onto a subset  $E$  of  $\mathbb{R}^2$ , and  $f : E \rightarrow \mathbb{R}$  be a function. Then  $f$  is integrable over  $E$  if and only if  $(f \circ T)|\det J_T|$  is integrable over  $E'$ . In that case,*

$$\int_E f = \int_{E'} (f \circ T)|\det J_T|.$$

Writing  $\frac{\partial(x,y)}{\partial(u,v)}$  for  $\det J_T$ , this formula becomes

$$\int_E f(x,y) d(x,y) = \int_{E'} f(u,v) \left| \frac{\partial(x,y)}{\partial(u,v)} \right| d(u,v).$$

To recover Theorem 7.9 from Theorem 7.13, take  $T(r, \theta) = (r \cos \theta, r \sin \theta)$ , so  $\frac{\partial(x,y)}{\partial(r,\theta)} = r$ .

In the situation of Theorem 7.13,  $E$  is always measurable (continuous image of a Borel set) although this is not obvious.

One can extend Section 7 to  $\mathbb{R}^n$  instead of  $\mathbb{R}^2$ . Moreover, for any ( $\sigma$ -finite) measure spaces  $(\Omega_1, \mathcal{F}_1, \mu_1)$  and  $(\Omega_2, \mathcal{F}_2, \mu_2)$ , one can define a product  $(\Omega_1 \times \Omega_2, \mathcal{F}_1 \otimes \mathcal{F}_2, \mu_1 \times \mu_2)$  such that Fubini's and Tonelli's theorems hold.

## 8. $L^p$ -SPACES

In this section we will write  $\mathcal{L}^1$  for the integrable functions on  $\mathbb{R}$ , or an interval, or  $\mathbb{R}^2$ , or any measure space). A useful measure of distance between two integrable functions  $f$  and  $g$  is:

$$d(f, g) = \int |f - g| =: \|f - g\|_1.$$

Then

- (i)  $\|f\|_1 = 0$  if and only if  $f = 0$  a.e. (Proposition 4.8(5),(7));
- (ii)  $\|\alpha f\|_1 = |\alpha| \|f\|_1$ ;
- (iii)  $\|f + g\|_1 \leq \|f\|_1 + \|g\|_1$ .

Consequently,

- (i)'  $d_1(f, g) = 0$  if and only if  $f = g$  a.e.
- (ii)'  $d_1(g, f) = d_1(f, g)$ ;
- (iii)'  $d_1(f, h) \leq d_1(f, g) + d_1(g, h)$ .

So  $\|\cdot\|_1$  is almost a norm and  $d_1$  is almost a *metric* (cf., Metric Spaces). The problems are that  $\mathcal{L}^1$ , as defined so far, is not a vector space, and  $\|f\|_1 = 0$  does not imply that  $f$  is the zero function.

We can get around these problems by identifying functions which are almost everywhere equal (actually, we have effectively been doing this for some time). Any function in  $\mathcal{L}^1$  is real-valued almost everywhere, so we will now take  $\mathcal{L}^1$  to be the space of all integrable functions with real values. This is a vector space, and we can define an equivalence relation on  $\mathcal{L}^1$  by

$$f \sim g \iff f = g \text{ a.e.}$$

Let  $[f]$  be the equivalence class of  $f$ , and  $\mathcal{N} = [0] = \{f : f = 0 \text{ so a.e.}\}$ . Then  $\mathcal{N}$  is a subspace of the vector space  $\mathcal{L}^1$ , and we can form the quotient space  $L^1 := \mathcal{L}^1/\mathcal{N}$  as a vector space whose elements are the equivalence classes  $[f]$  (cf., Linear Algebra). Let

$$\|[f]\|_1 = \int |f|.$$

Then  $\|\cdot\|_1$  is well-defined, and it is a norm on  $L^1$ . The distinction between  $[f]$  and  $f$  is usually a distracting nuisance, so we suppress it, and we just write  $\|f\|_1$  as the norm of  $f$ . However it is occasionally necessary to be aware of the difference.

Now we have a notion of convergence:

$$f_n \rightarrow f \text{ in } L^1\text{-norm} \iff \lim_{n \rightarrow \infty} \|f_n - f\|_1 = 0 \iff \int |f_n - f| \rightarrow 0.$$

In probability this may be called *convergence in mean*. Actually, convergence in mean square is more convenient in some respects. For that, one considers the space  $\mathcal{L}^2$  of all measurable functions  $f$  such that  $|f|^2$  is integrable. Suppose that  $f, g \in \mathcal{L}^2$ . Then simple inequalities for real/complex numbers give

$$|f + g|^2 \leq 2(|f|^2 + |g|^2), \quad |f\bar{g}| \leq \frac{1}{2}(|f|^2 + |g|^2)$$

So  $f + g \in \mathcal{L}^2$  and  $f\bar{g}$  is integrable. Thus  $\mathcal{L}^2$  is a vector space, and we can put

$$\langle f, g \rangle_2 = \int f\bar{g}.$$

Then  $\langle \cdot, \cdot \rangle_2$  is positive-semidefinite, linear in the first variable, and conjugate-symmetric, so it is almost an inner product. Again there is a small problem that  $\langle f, f \rangle_2 = 0$  implies only that  $f \in \mathcal{N}$ . So we form  $L^2 = \mathcal{L}^2/\mathcal{N}$ , and we obtain an inner product on  $L^2$ . Hence, we get a well-defined norm on  $L^2$  given by

$$\|[f]\|_2 = \|f\|_2 = \langle f, f \rangle_2^{1/2} = \left( \int |f|^2 \right)^{1/2}.$$

Now,  $\|f_n - f\|_2 \rightarrow 0$  (convergence in  $L^2$ -norm) corresponds exactly to convergence in mean square in the case of probability spaces.

Let's see what happens if the indices 1 and 2 are replaced by some other real  $p > 0$ . Let  $\mathcal{L}^p$  be the set of all measurable functions  $f$  such that  $|f|^p$  is integrable. Note that

$$(|f + g|)^p \leq (2 \max(|f|, |g|))^p = 2^p \max(|f|^p, |g|^p) \leq 2^p(|f|^p + |g|^p),$$

$\mathcal{L}^p$  is a vector space. Let  $L^p = \mathcal{L}^p/\mathcal{N}$ , and

$$\|f\|_p = \left( \int |f|^p \right)^{1/p}.$$

Now it is not obvious whether the triangle inequality holds.

**Proposition 8.1. [Minkowski's Inequality]** For  $p \geq 1$  and  $f, g \in \mathcal{L}^p$ ,  $\|f + g\|_p \leq \|f\|_p + \|g\|_p$ .

*Proof.* If  $f = 0$  a.e. or  $g = 0$  a.e., the inequality is trivial. So suppose that  $\alpha := \|f\|_p > 0$  and  $\beta := \|g\|_p > 0$ .

The function  $t \mapsto t^p$  is continuous on  $[0, \infty)$  and its second derivative  $p(p-1)t^{p-2}$  is positive on  $(0, \infty)$ . This implies that it is convex, i.e.

$$(\lambda s + (1-\lambda)t)^p \leq \lambda s^p + (1-\lambda)t^p$$

for  $0 \leq \lambda \leq 1$ ,  $s, t \geq 0$ . Apply this with

$$\lambda = \frac{\alpha}{\alpha + \beta}, \quad s = \frac{|f(x)|}{\alpha}, \quad t = \frac{|g(x)|}{\beta}.$$

This gives

$$\left( \frac{|f| + |g|}{\alpha + \beta} \right)^p \leq \frac{|f|^p}{\alpha^{p-1}} + \frac{|g|^p}{\beta^{p-1}}.$$

Using  $|f + g| \leq |f| + |g|$ , integrating, and taking  $p^{\text{th}}$  roots gives the required inequality.  $\square$

So  $L^p$  becomes a normed vector space, whenever  $p \geq 1$ .

A related result is:

**Proposition 8.2. [Hölder's Inequality]** *Let  $p, q \in (1, \infty)$  with  $1/p + 1/q = 1$ . Let  $f \in L^p$  and  $g \in L^q$ . Then  $fg \in L^1$  and  $\|fg\|_1 \leq \|f\|_p \|g\|_q$ .*

For  $p = q = 2$ , Hölder's Inequality is the Cauchy-Schwarz Inequality.

*Proof.* Note first that the function  $t \mapsto \log t$  is concave on  $[0, \infty)$ , because its second derivative  $-t^{-2}$  is negative. Hence

$$\frac{1}{p} \log s + \frac{1}{q} \log t \leq \log \left( \frac{s}{p} + \frac{t}{q} \right).$$

Exponentiate to obtain  $s^{1/p} t^{1/q} \leq \frac{s}{p} + \frac{t}{q}$ . Let  $s = (|f(x)|/\|f\|_p)^p$  and  $t = (|g(x)|/\|g\|_q)^q$ . This gives

$$\frac{|fg|}{\|f\|_p \|g\|_q} \leq \frac{|f|^p}{p \|f\|_p^p} + \frac{|g|^q}{q \|g\|_q^q}.$$

Integrate.  $\square$

**Corollary 8.3.** *If  $1 \leq p_1 < p_2 < \infty$  and  $f \in L^{p_2}(a, b)$ , then  $f \in L^{p_1}(a, b)$  and*

$$\|f\|_{p_1} \leq (b-a)^{\frac{1}{p_1} - \frac{1}{p_2}} \|f\|_{p_2}.$$

*Hence if  $f_n \in L^{p_2}(a, b)$  and  $\|f_n\|_{p_2} \rightarrow 0$ , then  $\|f_n\|_{p_1} \rightarrow 0$ .*

*Proof.* Apply Proposition 8.2 to the functions  $|f|^{p_1}$  and  $\chi_{(a,b)}$ , with  $p = p_2/p_1$ .  $\square$

The inclusion  $L^{p_2}(a, b) \subset L^{p_1}(a, b)$  in Corollary 8.3 is strict: consider  $x^\alpha$  on  $(0, 1)$ .

Corollary 8.3 holds if  $(a, b)$  is replaced by any *finite* measure space. However,  $L^{p_1}(1, \infty)$  is not contained in  $L^{p_2}(1, \infty)$  (exercise).

For  $p \geq 1$ ,  $L^p$  is a normed space and hence a metric space for  $d_p(f, g) = \|f - g\|_p$ . How does convergence in  $L^p$ -norm compare with pointwise a.e. convergence?

**Examples 8.4.** 1. Convergence a.e. does not imply convergence in  $L^p$ -norm: If  $f_n(x) = n^2 x^n (1-x)$  ( $0 \leq x \leq 1$ ), then  $f_n(x) \rightarrow 0$  a.e., but  $\|f_n\|_1 \rightarrow 1$ .

2. Convergence in  $L^p$ -norm does not imply convergence a.e.: For  $n = 2^r + k$ , where  $0 \leq k < 2^r$ , let  $f_n$  be the characteristic function of  $[k2^{-r}, (k+1)2^{-r}]$ . Then  $\|f_n\|_1 = 2^{-r} \leq 2/n \rightarrow 0$ , but for each  $x \in [0, 1]$ ,  $f_n(x)$  takes the values 0 and 1 infinitely often.

**Theorem 8.5.** *Let  $p \in [1, \infty)$ , and let  $(f_n)$  be a sequence in  $\mathcal{L}^p$  which is Cauchy, i.e., for each  $\varepsilon > 0$ , there exists  $N$  such that  $\|f_n - f_m\|_p < \varepsilon$  whenever  $m, n \geq N$ . Then there exists  $f \in \mathcal{L}^p$  such that*

1. *There is a subsequence  $(f_{n_k})$  such that  $\lim_{k \rightarrow \infty} f_{n_k}(x) = f(x)$  a.e.*
2.  $\lim_{n \rightarrow \infty} \|f_n - f\|_p = 0$ .

*Thus  $L^p$  is a complete metric space.*

*Proof.* [For  $p = 1$ .] By assumption, there exist  $N_1 < N_2 < N_3 < \dots$  such that  $\int |f_n - f_m| < 2^{-r}$  whenever  $n, m \geq N_r$ . In particular,  $\int |f_{N_{r+1}} - f_{N_r}| < 2^{-r}$ . Let  $g_1 = f_{N_1}$  and  $g_r = f_{N_r} - f_{N_{r-1}}$  for  $r = 2, 3, \dots$ . By Lebesgue's Series Theorem 5.9,  $\sum_{r=1}^{\infty} g_r$  converges a.e. to  $f \in \mathcal{L}^1$ . Now

$$f_{N_k} = \sum_{r=1}^k g_r \rightarrow f \quad \text{a.e.,}$$

$$\|f_{N_k} - f\|_1 \leq \int \sum_{r=k+1}^{\infty} |g_r| = \sum_{r=k+1}^{\infty} \int |g_r| < 2^{-k} \rightarrow 0.$$

If a Cauchy sequence has a convergent subsequence, then the whole sequence is convergent. See Prelims proof that every Cauchy sequence in  $\mathbb{R}$  is convergent.

For general  $p$ , the use of LST has to be replaced by Minkowski's inequality plus Fatou's Lemma. □

- Corollary 8.6.** 1. *If  $\|f_n - f\|_p \rightarrow 0$ , then there is a subsequence  $(f_{n_r})$  which converges to  $f$  a.e.*  
 2. *If  $\|f_n - f\|_p \rightarrow 0$  and  $f_n \rightarrow g$  a.e., then  $f = g$  a.e.*

The Convergence Theorems provide situations when a.e. convergence implies convergence in  $L^p$ -norm. Here is a general result in that direction with a weaker conclusion.

**Theorem 8.7. [Egorov's Theorem]** *Suppose that  $f_n \rightarrow f$  a.e. Let  $E$  be a measurable set with  $m(E) < \infty$  and let  $\varepsilon > 0$ . Then there is a measurable subset  $F$  of  $E$  with  $m(E \setminus F) < \varepsilon$  such that  $f_n \rightarrow f$  uniformly on  $F$ . In particular,  $\|f_n - f\|_{L^p(F)} \rightarrow 0$  for all  $p \geq 1$ .*

Another very useful theorem is the following.

**Theorem 8.8.** *If  $f \in L^p(\mathbb{R})$  where  $1 \leq p < \infty$ , there is a sequence of step functions  $\psi_n$  such that  $\lim_{n \rightarrow \infty} \|f - \psi_n\|_p = 0$ .*

This result is closely related to Theorem 3.10, that measurable functions are pointwise (a.e.) limits of sequences of step functions. For a proof when  $p = 1$ , see Stein & Shakarchi, Theorem 2.4, p.71.

Let  $f \in \mathcal{L}^1(\mathbb{R})$ . The *Fourier transform* of  $f$  is the function  $\widehat{f}: \mathbb{R} \rightarrow \mathbb{C}$  defined by

$$\widehat{f}(s) = \int_{\mathbb{R}} f(x)e^{-isx} dx.$$

This definition appeared in the short option Integral Transforms, and now we can give rigorous proofs of some of the properties shown in that course without full rigour.

**Theorem 8.9.** *Let  $f \in \mathcal{L}^1(\mathbb{R})$ .*

1.  $|\widehat{f}(s)| \leq \|f\|_1$  for all  $s$ ,
2.  $\widehat{f}$  is continuous,
3.  $\widehat{f}(s) \rightarrow 0$  as  $s \rightarrow \pm\infty$ . [**Riemann–Lebesgue Lemma**]
4. Let  $g(x) = xf(x)$ . If  $g \in \mathcal{L}^1(\mathbb{R})$  then  $\widehat{f}$  is differentiable everywhere and  $(\widehat{f})'(s) = -i\widehat{g}(s)$ .
5. If  $f$  has a continuous derivative  $f' \in \mathcal{L}^1(\mathbb{R})$ , then the Fourier transform of  $f'$  is  $is\widehat{f}(s)$ .

*Proof.* (1) follows from  $|f(x)e^{-isx}| = |f(x)|$ . (2) follows from the continuous-parameter DCT (Theorem 6.2) with  $g(x) = |f(x)|$ .

For (3), For  $f = \chi_{(a,b)}$ ,  $\widehat{f}(s) = \frac{i(e^{-isb} - e^{-isa})}{s} \rightarrow 0$  as  $|s| \rightarrow \infty$ . This extends to step functions, by linearity. For general  $f \in \mathcal{L}^1(\mathbb{R})$  and  $\varepsilon > 0$ , there is a step function  $\varphi$  such that  $\|f - \varphi\|_1 < \varepsilon$  by Theorem 8.8, and there exists  $K$  such that  $|\widehat{\varphi}(s)| < \varepsilon$  whenever  $|s| > K$ . Then

$$|\widehat{f}(s)| \leq |\widehat{f}(s) - \widehat{\varphi}(s)| + |\widehat{\varphi}(s)| \leq \|f - \varphi\|_1 + |\widehat{\varphi}(s)| < 2\varepsilon.$$

[Note that (2) can be proved by observing that  $\widehat{f}$  is a uniform limit of continuous functions  $\widehat{\varphi}_n$  where  $\varphi_n$  are step functions converging to  $f$  in  $L^1$ -norm.]

(4) can be proved by applying Theorem 6.5 with  $|g|$  as dominating function. (5) can be proved by using integration by parts over intervals  $[a_n, b_n]$  where  $a_n \rightarrow -\infty$ ,  $f(a_n) \rightarrow 0$ ,  $b_n \rightarrow \infty$  and  $f(b_n) \rightarrow 0$ .  $\square$

The theorem about the Fourier transform of the convolution of two integrable functions is an application of Fubini/Tonelli.

## 9. ABSOLUTELY CONTINUOUS FUNCTIONS

Recall from Section 4 that the Fundamental Theorem of Calculus is true for functions with a continuous derivative on  $[a, b]$  (Theorem 4.11, but proved in Prelims), but it is false for the Cantor-Lebesgue function  $\Phi$  whose derivative exists and equals 0 a.e. on  $[0, 1]$  (Example 4.12).

The ideal Fundamental Theorem of Calculus would identify a class  $\mathcal{A}$  of functions  $F$  on  $[a, b]$  with both the following properties:

- (i) If  $F \in \mathcal{A}$ , then  $F$  is differentiable a.e.,  $F' \in L^1(a, b)$ , and  $\int_a^x F'(y) dy = F(x) - F(a)$  for all  $x \in [a, b]$ .
- (ii) If  $f \in L^1(a, b)$  and  $F(x) = \int_a^x f(y) dy$  for  $x \in [a, b]$ , then  $F \in \mathcal{A}$  and  $F' = f$  a.e.

It is not obvious that such a class exists—its existence implies that the indefinite integral  $F$  of an integrable function  $f$  is differentiable a.e. and  $F' = f$  a.e.

In fact, this is true. Then  $\mathcal{A}$  is the class of all functions of the form  $F(x) := c + \int_a^x f$  for some  $c \in \mathbb{R}$  and some  $f \in L^1(a, b)$ . Remarkably there is an intrinsic characterisation of such functions.

Let  $I$  be an interval. A function  $F : I \rightarrow \mathbb{R}$  is said to be *absolutely continuous* on  $I$  if, for each  $\varepsilon > 0$ , there exists  $\delta > 0$  such that

$$\sum_{r=1}^n |F(b_r) - F(a_r)| < \varepsilon$$

whenever  $n \in \mathbb{N}$ ,  $(a_r, b_r)$  ( $r = 1, \dots, n$ ) are disjoint subintervals of  $I$  and  $\sum_{r=1}^n (b_r - a_r) < \delta$ .

If we only allowed  $n = 1$  in this definition, we would have the definition of uniform continuity on  $I$ . Recall from Prelims that any continuous function on  $[a, b]$  is uniformly continuous.

**Examples 9.1.** 1. Recall that  $F$  is Lipschitz if there exists  $c$  such that  $|F(y) - F(x)| \leq c|y - x|$  for all  $x, y$ . Any Lipschitz function is absolutely continuous (take  $\delta = \varepsilon/c$ ).  
 2. If  $f$  is a bounded measurable function and  $F(x) = \int_a^x f(y) dy$ , then  $F$  is Lipschitz.  
 3. The Cantor-Lebesgue function is not absolutely continuous on  $[0, 1]$ .

**Theorem 9.2.** *Let  $f \in L^1(I)$  and  $F(x) = \int_a^x f(y) dy$ . Then  $F$  is absolutely continuous on  $I$ .*

**Theorem 9.3.** *Let  $F$  be an absolutely continuous function on  $[a, b]$ . Then  $F$  is differentiable a.e.,  $F' \in L^1(a, b)$  and  $F(x) - F(a) = \int_a^x F'(y) dy$  for all  $x \in [a, b]$ .*

One way to a proof of Theorem 9.2 is outlined in an optional exercise on the Supplementary Problem Sheet. There are various other proofs.

Theorem 9.3 is rather hard to prove. There is a proof in Capinski & Kopp, Section 7.3. It is a remarkable theorem as differentiability (a.e.) is inferred from an assumption that seems to be only a type of continuity.

A corollary of Theorem 9.3 is that every Lipschitz function is differentiable a.e. Thus the Lipschitz functions are precisely the indefinite integrals of bounded measurable functions.