

# Projective Geometry

## Lecture 7

Balázs Szendrői, University of Oxford, Trinity term 2020

## Bilinear forms over arbitrary fields

---

A **symmetric bilinear form** on a vector space  $V$  over  $\mathbb{F}$  is a map

$$B : V \times V \rightarrow \mathbb{F}$$

such that  $B(v, w) = B(w, v)$  and  $B$  is linear in  $v$  and  $w$ .

The form is **nondegenerate** or **nonsingular** if in addition we have that if  $B(v, w) = 0$  for all  $w \in V$ , then  $v = 0$ .

We have the associated **quadratic form**  $Q(v) = B(v, v)$ .

A bilinear form is given by  $B(v, w) = v^t X w$  for a symmetric matrix  $X$  given by  $X_{ij} = B(e_i, e_j)$ .

$B$  is nondegenerate if and only if  $X$  is invertible.

## Projective quadrics and conics

---

A **projective quadric** is

$$C = \{[v]: Q(v) = 0\} \subset \mathbb{P}(V)$$

where  $v \mapsto Q(v) = B(v, v)$  is a (not identically zero) quadratic form on  $V$ .

In matrix form, we get

$$C = \{[v]: v^t X v = 0\} \subset \mathbb{P}(V)$$

where  $X$  is a nonzero symmetric matrix over  $F$ .

A **projective conic** is a projective quadric in a projective plane  $\mathbb{P}(V) = \mathbb{FP}^2$ .

$$C = \{[v]: Q(v) = 0\} \subset \mathbb{FP}^2$$

where  $v \mapsto Q(v) = B(v, v)$  is a (not identically zero) quadratic form on a three-dimensional vector space  $V$ .

## Projective quadrics: some examples

---

**Example 1** The simplest example is the identity matrix  $X = I$ , for which we get the quadric

$$C_1 = \left\{ [x_0 : x_1 : \dots : x_n] \left| \sum_{i=0}^n x_i^2 = 0 \right. \right\} \subset \mathbb{F}\mathbb{P}^n.$$

Note that in the most familiar example  $\mathbb{F} = \mathbb{R}$ , this quadric is **empty**.

However, it has plenty of points over other fields such as  $\mathbb{F} = \mathbb{C}$ .

**Extended exercise** What happens over  $\mathbb{F} = \mathbb{F}_p$ ?

## Projective quadrics: some examples

---

**Example 2** Consider the matrix

$$X = \begin{pmatrix} -1 & 0 \\ 0 & I_n \end{pmatrix},$$

with  $I_n$  the  $n \times n$  identity matrix. We get the quadric

$$C_2 = \left\{ [x_0 : x_1 : \dots : x_n] \left| -x_0^2 + \sum_{i=1}^n x_i^2 = 0 \right. \right\} \subset \mathbb{FP}^n.$$

For  $\mathbb{F} = \mathbb{R}$ , in affine coordinates  $x_i/x_0$ , this would give back the equation of the sphere in  $\mathbb{R}^n$ . So this has plenty of points over  $\mathbb{F} = \mathbb{R}$ .

Note that for  $n = 2$ , all the familiar plane conics were projectively equivalent to this conic.

## Projective quadrics: some examples

---

**Example 3** Let  $n = 2$  and consider the matrix

$$X = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}.$$

We get the plane conic

$$C_3 = \{[x_0 : x_1 : x_2] \mid x_1^2 - x_2^2 = 0\} \subset \mathbb{P}^2.$$

Note that the equation now **factorizes**:

$$C_3 = \{[x_0 : x_1 : x_2] \mid (x_1 - x_2)(x_1 + x_2) = 0\} \subset \mathbb{P}^2.$$

So we can write


$$C_3 = L_- \cup L_+ \subset \mathbb{P}^2,$$

a union of the lines  $L_{\pm} = \{x_1 \pm x_2 = 0\} \subset \mathbb{P}^2$ . These lines meet at the point  $[1 : 0 : 0] \in \mathbb{P}^2$ .

## Projective quadrics: some examples

---

$$C_3 = \{[x_0 : x_1 : x_2] \mid x_1^2 - x_2^2 = 0\} = L_- \cup L_+ \subset \mathbb{P}^2.$$



## Nonsingular quadrics

---

We say that a quadric is **nonsingular**, if the associated bilinear form  $B$  is nondegenerate, equivalently the matrix  $X$  is invertible.

A **singular point** of a quadric

$$C = \{[v] : Q(v) = 0\} \subset \mathbb{P}(V)$$

is  $[v] \in \mathbb{P}(V)$  for nonzero  $v$  such that  $B(v, w) = 0$  for all  $w \in V$ , equivalently nonzero  $v \in \ker X$ .

The quadrics  $C_1, C_2$  in Examples 1-2 above are nonsingular.

The plane conic  $C_3 = L_- \cup L_+$  is singular at  $[1 : 0 : 0] = L_- \cap L_+$ .

**Remark** The terminology comes from thinking about conics as **submanifolds** (locally) of Euclidean space (see parallel Part A course). At a nonsingular point, a quadric is a submanifold. At singular points, the Jacobian condition fails; hence the name.



## Singular and nonsingular projective conics


---

$$C_2 = \{-x_0^2 + x_1^2 + x_2^2 = 0\}$$



$C_2$

$$C_3 = \{x_1^2 - x_2^2 = 0\} = L_- \cup L_+$$



$C_3 = L_- \cup L_+$

## Diagonalizing quadratic forms

---

Over the fields  $\mathbb{R}$  and  $\mathbb{C}$ , we can diagonalise quadratic forms.

**Theorem** Let  $v \mapsto Q(v) = B(v, v)$  be a quadratic form defined on an  $(n + 1)$ -dimensional vector space  $V$ .

- (i) Over the base field  $\mathbb{F} = \mathbb{C}$ , there is a basis  $\{e_0, \dots, e_n\}$  of  $V$ , with respect to which

$$Q(v) = \lambda_0^2 + \dots + \lambda_r^2,$$

where  $v = \sum_{i=0}^n \lambda_i e_i$ .

- (ii) Over the base field  $\mathbb{F} = \mathbb{R}$ , there is a basis  $\{e_0, \dots, e_n\}$  of  $V$ , with respect to which

$$Q(v) = \lambda_0^2 + \dots + \lambda_r^2 - \lambda_{r+1}^2 - \dots - \lambda_{r+s}^2,$$

where  $v = \sum_{i=0}^n \lambda_i e_i$ .

**Remark** Note that  $Q$  is non-degenerate if and only if  $r = n$ , respectively  $r + s = n$ . More generally,  $\text{rk } X = r + 1$ ,  $\text{rk } X = r + s + 1$  in the two cases.

## Proof of the diagonalization theorem: Step 1

---

Write

$$Q(v) = v^t X v = \sum_{i,j} X_{ij} v_i v_j$$

in some basis, where  $X$  is a nonzero, symmetric matrix.

**Step 1** We can assume that some  $X_{ii}$  is nonzero. For if all  $X_{ii} = 0$ , find a nonzero  $X_{ij}$ . Introduce new variables

$$y_i = \frac{1}{2}(v_i + v_j), \quad y_j = \frac{1}{2}(v_i - v_j).$$

Now  $Q$  has the term

$$X_{ij} v_i v_j = X_{ij} y_i^2 - X_{ij} y_j^2$$

with nonzero diagonal terms in the new basis.

## Proof of the diagonalization theorem: Steps 2-3

---

### Step 2

Now we complete the square.

$$\frac{1}{X_{ii}} \left( \sum_j X_{ij} v_j \right)^2 = X_{ii} v_i^2 + 2 \sum_{j \neq i} X_{ij} v_j v_i + \text{terms in } v_j \ (j \neq i)$$

so by introducing the new variable  $y_i = \sum X_{ij} v_j$ , we can put  $Q$  into the form

$$Q(v) = \frac{1}{X_{ii}} y_i^2 + Q'(v_0, \dots, v_{i-1}, v_{i+1}, \dots, v_n)$$

for some quadratic form  $Q'$  with one less variable.

### Step 3

Now we repeat the process until we have diagonalised  $Q$ . Finally rescaling the variables appropriately, we can assume that we have the stated forms; note that over  $\mathbb{R}$ , we cannot change the sign of  $y_i^2$  by rescaling.

## Classification of complex projective quadrics

---

Turning to quadrics, for the field  $\mathbb{F} = \mathbb{C}$ , our diagonalization theorem says that every complex projective quadric in  $\mathbb{CP}^n$  is projectively equivalent to

$$D_r = \left\{ [x_0 : x_1 : \dots : x_n] \left| \sum_{i=0}^r x_i^2 = 0 \right. \right\} \subset \mathbb{CP}^n.$$

for some  $0 \leq r \leq n$ .

The quadric  $D_r$  is nonsingular if and only if  $r = n$ .

## Classification of complex projective conics

---

Specializing to  $n = 2$ , we get that every complex projective conic is projectively equivalent to one of the following:

- (i) The **nonsingular conic**  $D_2 = \{[x_0 : x_1 : x_2] \mid x_0^2 + x_1^2 + x_2^2 = 0\} \subset \mathbb{CP}^2$ .
- (ii) The **line pair**  $D_1 = \{[x_0 : x_1 : x_2] \mid x_0^2 + x_1^2 = 0\} \subset \mathbb{CP}^2$ .

Indeed, as before, we can write

$$D_1 = M_+ \cup M_- \subset \mathbb{CP}^2$$

for

$$M_{\pm} = \{x_0 \pm ix_1 = 0\} \subset \mathbb{CP}^2.$$

- (iii) The **double line**  $D_0 = \{[x_0 : x_1 : x_2] \mid x_0^2 = 0\} \subset \mathbb{CP}^2$ .

Indeed, this is “twice the line”  $M$ , where

$$M = \{x_0 = 0\} \subset \mathbb{CP}^2.$$

## Classification of nonsingular real projective conics

---

Let us now see what we get in the case of conics with  $\mathbb{F} = \mathbb{R}$ , restricting to the nonsingular case. (Exercise: think about the singular cases!)

Writing  $v = \sum x_i e_i$ , there are four nonsingular quadratic forms to consider.

- (i)  $Q_1(v) = x_0^2 + x_1^2 + x_2^2$ .
- (ii)  $Q_2(v) = x_0^2 + x_1^2 - x_2^2$ .
- (iii)  $Q_3(v) = x_0^2 - x_1^2 - x_2^2$ .
- (iv)  $Q_4(v) = -x_0^2 - x_1^2 - x_2^2$ .

The conics corresponding to  $Q_1$  and  $Q_4$  are **empty**. The conics corresponding to  $Q_2, Q_3$  are **the same**, as the forms are constant multiples of each other (up to change of variables).

Hence indeed, up to projective equivalence **there is a unique nonempty nonsingular real projective conic**

$$C = \{-x_0^2 + x_1^2 + x_2^2 = 0\} \subset \mathbb{RP}^2.$$


## Parametrising conics

---

Work over an arbitrary field  $\mathbb{F}$  again, only assuming  $\text{char } \mathbb{F} \neq 2$ .

Consider a nonsingular conic  $C \subset \mathbb{P}^2$ , a point  $X \in C$ , and a projective line  $L \subset \mathbb{P}^2$  not containing  $X$ .

We will give a description of  $C$  using the following geometric idea: projection from  $X$  onto the line  $L$  sets up a bijection between the conic  $C$  and the line  $L$ .






## Parametrising conics: the theorem

---

**Theorem** Let  $C$  be a nonsingular conic in the projective plane  $\mathbb{P}(V) = \mathbb{F}\mathbb{P}^2$ , over a field  $\mathbb{F}$  with  $\text{char } \mathbb{F} \neq 2$ . Let  $X$  be a point of  $C$ . Let  $L = \mathbb{P}(U)$  be a projective line in the plane not containing  $X$ . Then there is a bijection

$$\alpha : L \rightarrow C$$

such that  $X, Y, \alpha(Y)$  are collinear for each  $Y \in L$ .



## Parametrising conics: the proof

---

**Proof** Let  $B$  denote the nondegenerate bilinear form whose quadratic form  $Q$  defines the conic  $C$ . Let  $X = [x]$  be a point on  $C$ , so that  $B(x, x) = 0$ .

For each  $Y \in \mathbb{P}(U)$ , we want to see where (other than at  $X$ ) the projective line  $XY$  meets the conic. We will find that there is a unique such point and this will be  $\alpha(Y)$ .

Let  $Y \in \mathbb{P}(U)$  have representative vector  $y \in U$ , so that  $x, y$  are linearly independent, as we are assuming  $X \notin L = \mathbb{P}(U)$ .

Consider the 2-dimensional subspace  $W_y = \langle x, y \rangle \subset V$ , so the projective line we are considering is  $XY = \mathbb{P}(W_y)$ .

**Key Claim:** The bilinear form  $B$  cannot be identically zero on the space  $W_y$ . (See Lecture 8.)

## Parametrising conics: the proof

---

With respect to the basis  $\{x, y\}$ , the form  $Q$  restricted to  $W_y$  is

$$Q(\lambda_0 x + \lambda_1 y) = 2\lambda_0 \lambda_1 B(x, y) + \lambda_1^2 B(y, y).$$

$B(x, y), B(y, y)$  are not both zero by the Key Claim. So the projective line  $\mathbb{P}(W_y)$  meets the conic  $C$  at two points, corresponding to the solutions  $[\lambda_0 : \lambda_1]$ .

One intersection point is the basepoint  $X = [x]$ , corresponding to

$$[\lambda_0 : \lambda_1] = [1 : 0].$$

Defined  $\alpha(Y)$  to be the other intersection point, corresponding to

$$[\lambda_0 : \lambda_1] = [B(y, y) : -2B(x, y)].$$


So  $X, Y, \alpha(Y)$  are collinear by construction.

$\alpha$  is bijective: given any point  $Z \neq X$  on the conic, the projective line  $XZ$  meets the line  $L$  in a unique point  $Y$ , and then  $\alpha(Y) = Z$ .

## Parametrising conics: the proof

---

For  $Z = X$  itself, the image  $\alpha(X) = X$ , coming from the intersection point  $Y'$  of the **tangent line** at  $X$  with the line  $L$ .



This corresponds to the case then the quadratic above has a double root. (Think about this!)