

BO1 History of Mathematics
Lecture II
Dissemination and development
(AD 500 – AD 1600)

MT 2019 Week 1

Summary

- ▶ Influence of the ancient world
- ▶ The Renaissance (15th and 16th centuries)
- ▶ The 16th century
- ▶ A case study: Napier's invention of logarithms 1614

Remnants of the collapse of the ancient world

in Greek: manuscripts preserved at Constantinople and in libraries or collections around the Mediterranean

in Latin: writings by Boethius (c. 480–524) on philosophy, arithmetic, geometry, music

The spread of Islam and Islamic learning

- 632–732: Islam spreads throughout Middle East, north Africa, and into Spain and Portugal
- c. 820: *Bayt al-Ḥikma*, the House of Wisdom, founded in Baghdad under Caliph al-Ma'mūn; it became a centre for translation into Arabic from Greek, Persian, Sanskrit
- c. 825: al-Khwārizmī active in Baghdad
- 9th century: texts on arithmetic, algebra, astronomy reach Spain
- 12th century: translations from Arabic to Latin

Oxford in the 14th century

The Merton School, a.k.a. the Merton Calculators (principally, Thomas Bradwardine, William Heytesbury, Richard Swineshead, John Dumbleton):

- ▶ arithmetic using Hindu-Arabic numerals
- ▶ translations of Euclid (some partial)
- ▶ possibly a little algebra
- ▶ computus texts (calculation of time)
- ▶ astronomy and astrology

The mid-Renaissance (15th and 16th centuries)


Classical mathematical texts more widely available due to:

- ▶ rediscovery of manuscripts
- ▶ revival of knowledge of Greek
- ▶ (Western) invention of printing (Gutenberg, c. 1436)

Euclid's *Elements*: transmission history


- ▶ commentaries written by Pappus (c. AD 320), Theon (c. AD 380), Proclus (c. AD 450)
- ▶ a few propositions in Boethius (c. AD 500)
- ▶ copies in Greek (earliest from Constantinople, AD 888)
- ▶ many translations or commentaries in Arabic (AD 750–1250)
- ▶ mediaeval translations from Arabic to Latin: Adelard of Bath (1130), Robert of Chester (1145), Gerard of Cremona (mid-12th century)
- ▶ printed editions in Latin or Greek from 1482 onwards

Euclid in Arabic



Translated from the Greek by Ishaq ibn Hunayn, AD 1466


Euclid I.47 from Bodleian ms. dated 888



Whole manuscript is digitised:

<http://www.claymath.org/library/historical/euclid/>

Euclid I.47 from Bodleian ms. dated 888




<http://www.claymath.org/library/historical/euclid/files/elem.1.47.html>

Treatises by Archimedes: transmission history



- ▶ quoted or explained by Pappus (c. 320 AD), Theon (c. 380 AD), Eutocius (c. 520 AD)
- ▶ 6th-century Byzantine 'collected works' (Isidore of Miletus)
- ▶ several translations of individual treatises into Arabic
- ▶ translations from Arabic into Latin
- ▶ a new find in the twentieth century:
www.archimedespalimpsest.org/

Netz & Noel: *The Archimedes Codex*



(Weidenfeld & Nicolson, 2007)

The Archimedes palimpsest



Apollonius' *Conics* (c. 180 BC): transmission history

- ▶ Books I–IV survived in Greek
- ▶ Books V–VII survived only in Arabic
- ▶ Book VIII is lost, known only from commentaries
- ▶ early (Latin) printed edition, 1566

(See: *Mathematics emerging*, §1.2.4.)

Apollonius, Oxford, 1710

APOLLONII PERGÆI
C O N I C O R U M
LIBRI OCTO,
ET
SERENI ANTISSENSIS
DE SECTIONE
CYLINDRI & CONI
LIBRI DUO.



O X O N I Æ,
E THEATRO SHELDONIANO, An. Dom. MDCCX.

16th century change

New forces at work in the 16th century:

- ▶ global exploration
- ▶ growth of international commerce
- ▶ new technology (in printing, shipping, military engineering, instrumentation, etc.)

Simon Stevin (1548–1620), Leiden

Under the patronage of Maurice of Nassau, Prince of Orange, Stevin wrote on:

- ▶ accounting (1581)
- ▶ tables of interest (1582)
- ▶ geometry (1583)
- ▶ decimal fractions (1585)
- ▶ arithmetic (1585)
- ▶ weight and hydrostatics (1586)
- ▶ algebra (1594)
- ▶ fortification (1594)
- ▶ navigation (1599)
- ▶ mathematics (1608), including cosmography, geography, tides, heavenly motions, optics, perspective, refraction (Snell's law), pulleys, floating bodies, bookkeeping
- ▶ locks and sluices (1617)



Thomas Harriot (1560–1621), London

Under the patronage of the Earl of Northumberland, Harriot worked on:

- ▶ navigation
- ▶ optics, refraction (Snell's law)
- ▶ rates of fall
- ▶ calculations of density
- ▶ alchemy
- ▶ geometry
- ▶ algebra
- ▶ astronomy

none of it published

Harriot papers online:

http://echo.mpiwg-berlin.mpg.de/content/scientific_revolution/harriot



A case study of a text from 1614

Napier's invention of logarithms:

- ▶ what did 17th-century mathematics look like?
- ▶ how can we begin to read historical texts?

Napier's definition of a logarithm (of a sine)

*The **Logarithme** therefore of any sine is a number very neerely expressing the line, which increased equally in the meane time, whiles the line of the whole sine decreased proportionally into that sine, both motions being equal-timed, and the beginning equally swift.*

?

Context, content, significance

Context: who? when? where? why?

Content: what is it about? how is it written?

Significance: why did/does it matter?

Context — who?

John Napier (1550–1617), Merchiston,
Scotland

Scottish landowner with interests in:

- ▶ mining
- ▶ calculating aids
- ▶ astrology/astronomy
- ▶ The Revelation of St John



See *Oxford Dictionary of National Biography*:
<http://www.oxforddnb.com/view/article/19758>

Context — why?

From Napier's preface to the translation of 1616:

Seeing there is nothing (right well-beloved Students of the Mathematics) that is so troublesome to mathematical practice, nor that doth more molest and hinder calculators, than the multiplications, divisions, square and cubical extractions of great numbers, which besides the tedious expense of time are for the most part subject to many slippery errors, I began therefore to consider in my mind by what certain and ready art I might remove those hindrances.

Context — why?

Inspired by the 16th-century technique of **prosthaphaeresis**:

the use of trigonometric identities such as

$$\cos x \cos y = \frac{1}{2} [\cos(x + y) + \cos(x - y)]$$

$$\sin x \sin y = \frac{1}{2} [\cos(x - y) - \cos(x + y)]$$

to convert multiplication into addition.

Context — in what form, and in which language?

Original Latin text of 1614:

Mirifici logarithmorum canonis descriptio


translated into English by Edward Wright in 1616 as

A description of the admirable table of logarithms

Transcribed text available at:

http://www.johndnapier.com/table_of_logarithms_001.htm

Napier's 1616 title-page decoded



Inventor:

John Napier (1550–1617)

Translator:

Edward Wright (?1558–1615)
(interests: navigation, charts
and tables)

Additional material:

Henry Briggs (1561–1630)
Gresham Professor of Geometry,
later Savilian Professor of
Geometry at Oxford
(interests: navigation)

Printer:

Nicholas Okes

Readers:

Thomas Hulcher,
Thomas Panner

Napier's logarithms: content

Recall:

*The **Logarithme** therefore of any sine is a number very neerely expressing the line, which increased equally in the meane time, whiles the line of the whole sine decreased proportionally into that sine, both motions being equal-timed, and the beginning equally swift.*

Napier's logarithms

4 The first Booke. CHAP. I

peare by the 19 Prop. 5. and II. Prop. 7, Euclid.

3 Def. *Surd quantities, or unexplicable by numbers, are said to be defined, or expressed by numbers very neere, when they are defined or expressed by great numbers which differ not so much as one vnite, from the true value of the Surd quantity.*

As for example. Let the semidiameter, or whole sine be the rational number; 1000000 the sine of 45 degrees shall be the square root of 50,000,000,000,000, which is surd, or irrational and inexplicable by any number, & is included between the limits of 7071067 the lesse, and 7071068 the greater: therefore, it differeth not an vnite from either of these. Therefore that surd sine of 45 degrees, is said to be defined and expressed very neere, when it is expressed by the whole numbers, 7071067, or 7071068, not regarding the fractions. For in great numbers there ariseth no sensible error, by neglecting the fragments, or parts of an vnite.

4 Def. *Equal-timed motions are those which are made together, and in the same time.*



As in the figures following, admit that B be moued from A to C, in the same time, wherein b is moued from a to c the right lines AC & ac, shall be sayd to be described with an equal-timed motion.

5 Def. *Seeing, that there may bee a slower and a swifter motion giuen then any motion, it shall necessarily follow, that there may be a motion giuen of equal swiftnesse to any motion (which wee define to be neither swifter nor slower.)*

6 Def. *The Logarithme therefore of any sine is a number very neerely expressing the line, which increased*

CHAP. 2. The first Booke. 5


sed equally in the meane time, whiles the line of the whole sine de creased proportionally into that sine, both motions being equal-timed, and the beginning equally swift.




As for example. Let the 2 figures going afore bec here repeated, and let B bec moued alwayes, and euery where with equal, or the same swiftnesse wherewith b beganne to be moued in the beginning, when it was in a. Then in the first moment let B proceed from A to C, and in the same time let b moue proportionally from a to c, the number defining or expressing AC shal be the *Logarithme* of the line, or sine c Z. Then in the second moment let B bec moued forward from C to D. And in the same moment or time let b be moued proportionally from c to d, the number defining A D, shall be the *Logarithme* of the sine d Z. So in the third moment let B go forward equally from D to E, and in the same moment let b be moued forward proportionally from d to e, the number expressing A E the *Logarithme* of the sine e Z. Also in the fourth moment, let B proceed

B 3 cced

Napier's logarithms




Napier's logarithms



Napier's logarithms

Logarithms



Numbers



Naper's logarithms (1614)

In modern terms (i.e., **not Napier's**):

$$\text{if } y = 10^7 (1 - 10^{-7})^x, \text{ then Nap log } y = x$$

Nap log $10^7 = 0$, Nap log 0 is infinite, Nap log 1 = 161,180,956

$$\text{Nap log } \left(\frac{p \times q}{10^7} \right) = \text{Nap log } p + \text{Nap log } q$$

$$\text{Nap log } (p \times q) = \text{Nap log } p + \text{Nap log } q - \text{Nap log } 1$$

$$\text{Note that Nap log } x = 10^7 \ln \left(\frac{10^7}{x} \right)$$

No notion of base, although Nap log 'nearly' has base $\frac{1}{e}$ — see: Robin Wilson, *Euler's Pioneering Equation*, OUP, 2019, p. 101

Modifications by Napier and Briggs (1617)

Definition revised to remove the need to subtract Nap $\log 1$

'Briggsian' logarithms have base 10 and $\text{Log } 1 = 0$, so that

$$\text{Log}(p \times q) = \text{Log } p + \text{Log } q$$

Briggs produced *Logarithmorum chilias prima* (*The first thousand logarithms*) in 1617, followed by his *Arithmetica logarithmica* in 1624, which contained logarithms of 1 to 20,000 and 90,000 to 100,000, all to 14 decimal places (calculated by hand); the gap in the table was filled by Adriaan Vlacq in 1628

Napier's logarithms

One last time:

*The **Logarithme** therefore of any sine is a number very neerely expressing the line, which increased equally in the meane time, whiles the line of the whole sine decreased proportionally into that sine, both motions being equal-timed, and the beginning equally swift.*

