

BO1 History of Mathematics
Lecture VII
Infinite series

MT 2019 Week 4

Summary

- ▶ A non-Western prelude
- ▶ Newton and the Binomial Theorem
- ▶ Other 17th century discoveries
- ▶ Ideas of convergence
- ▶ Much 18th century progress: power series
- ▶ Doubts — and more on convergence

The Kerala School

Flourished in Southern India from the 14th to the 16th centuries, working on mathematical and astronomical problems

Names associated with the school: Narayana Pandita, Madhava of Sangamagrama, Vatasseri Parameshvara Nambudiri, Kelallur Nilakantha Somayaji, Jyeṣṭhadeva, Achyuta Pisharati, Melpathur Narayana Bhattathiri, Achyutha Pisharodi, Narayana Bhattathiri

Treatises on arithmetic, algebra, geometry, inc. methods for approximation of roots of equations, discussion of magic squares, infinite series, . . .

The Kerala School

Flourished in Southern India from the 14th to the 16th centuries, working on mathematical and astronomical problems

Names associated with the school: Narayana Pandita, Madhava of Sangamagrama, Vatasseri Parameshvara Nambudiri, **Kelallur Nilakantha Somayaji**, **Jyeṣṭhadeva**, Achyuta Pisharati, Melpathur Narayana Bhattathiri, Achyutha Pisharodi, Narayana Bhattathiri

Treatises on arithmetic, algebra, geometry, inc. methods for approximation of roots of equations, discussion of magic squares, **infinite series**, . . .

Tantrasamgraha (1501)

Completed by Kelallur Nilakantha Somayaji (1444–1544) in 1501;
concerns astronomical computations



Keralan series

Infinite series for trigonometric functions appear in Sanskrit verse in an anonymous commentary on the *Tantrasamgraha*, entitled the *Tantrasamgraha-vyakhya*, of c. 1530:

दृष्टज्यात्रिज्ययोर्घातात् कोटचाप्तं प्रथमं फलम् ।
ज्यावर्गं गुणकं कृत्वा कोटिवर्गं च हारकम् ॥
प्रथमादिफलेभ्योऽथ नेया फलततिर्मुहुः ।
एकत्रचाद्योज संख्याभिर्भक्ते ष्वेतेष्वनुक्रमात् ॥
ओजानां संयुतेस्त्यक्त्वा युग्मयोगं धनुर्भवेत् ।
दोःकोटघोरल्पमेवेह कल्पनीयमिह स्मृतम् ।
लब्धीनामवसानं स्यान्नान्यथापि मुहुः कृते ॥

Proof supplied by Jyeṣṭhadeva in his *Yuktibhāṣā* (1530)

Keralan series

From the *Tantrasamgraha-vyakhya*:

The product of the given Sine and the radius divided by the Cosine is the first result. From the first, [and then, second, third] etc., results obtain [successively] a sequence of results by taking repeatedly the square of the Sine as the multiplier and the square of the Cosine as the divisor. Divide [the above results] in order by the odd numbers one, three, etc. [to get the full sequence of terms]. From the sum of the odd terms, subtract the sum of the even terms. [The results] become the arc. In this connection, it is laid down that the [Sine of the] arc or [that of] its complement, whichever is smaller, should be taken here [as the given Sine]; otherwise, the terms obtained by the [above] repeated process will not tend to the vanishing magnitude.

Modern interpretation:

$$R\theta = \frac{R(R \sin \theta)^1}{1(R \cos \theta)^1} - \frac{R(R \sin \theta)^3}{3(R \cos \theta)^3} + \frac{R(R \sin \theta)^5}{5(R \cos \theta)^5} - \dots \quad (R \sin \theta < R \cos \theta)$$

But these results were unknown in the West until the 1830s

As we will see, the series for arctan was reproduced independently in Scotland in the 1670s

(509)

XXXIII. *On the Hindú Quadrature of the Circle, and the infinite Series of the proportion of the circumference to the diameter exhibited in the four Sástras, the Tantra Sangraham, Yucti Bháshá, Carana Padhati, and Sadratnamála. By CHARLES M. WILSE, Esq., of the Hon. East-India Company's Civil Service on the Madras Establishment.*

(Communicated by the MADRAS LITERARY SOCIETY and AUXILIARY ROYAL ASIATIC SOCIETY.)

Read the 13th of December 1832.

A'RYAB'HATTA, who flourished in the beginning of the thirty-seventh century of the *Cálí Yuga*,* of which four thousand nine hundred and twenty years have passed, has in his work, the *Aryab'hatiyam*, in which he mentions the period of his birth, exhibited the proportion of the diameter to the circumference of the circle as 20000 to 62832, in the following verse :

*Chaturáddicam satamashágunandwásháñtathá sahasráñám
Ayutadwaya visahcambházyasannó vritta pariñdhak.†*

Which is thus translated :

“ The product of one hundred increased by four and multiplied by eight, added to sixty and two thousands, is the circumference of a circle whose diameter is twice ten thousand.”

The author of the *Lilávatí*, who lived six centuries after A'RYAB'HATTA, states the proportion as 7 to 22, which, he adds, is sufficiently exact for common purposes. As a more correct or precise circumference, he proposes that the diameter be multiplied by 3927, and the product divided by 1250; the quotient will be a very precise circumference. This proportion is the same with that of A'RYAB'HATTA, which is less correct than that of

* Or the sixth century of the Christian era.

† This verse is in the variety of the *Arjaváritam* measure, called *Típála*.

Infinite series 1600–1900: an overview


Lecture VII:

- ▶ mid–late 17th century: many discoveries
- ▶ early 18th century: much progress
- ▶ later 18th century: doubts and questions

Lecture VIII:

- ▶ early 19th century: Fourier series
- ▶ early 19th century: convergence better understood

Newton and the general binomial theorem



CUL Add. MS 3958.3, f. 72

(See lecture IV)

Recall: Newton's integration of $(1+x)^{-1}$

	$(1+x)^{-1}$	$(1+x)^0$	$(1+x)^1$	$(1+x)^2$	$(1+x)^3$	$(1+x)^4$...
x	1	1	1	1	1	1	...
$\frac{x^2}{2}$	-1	0	1	2	3	4	...
$\frac{x^3}{3}$	1	0	0	1	3	6	...
$\frac{x^4}{4}$	-1	0	0	0	1	4	...
$\frac{x^5}{5}$	1	0	0	0	0	1	...
\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\ddots

The entry in the row labelled $\frac{x^m}{m}$ and the column labelled $(1+x)^n$ is the coefficient of $\frac{x^m}{m}$ in $f(1+x)^n dx$. (NB.

Newton did **not** use the notation $f(1+x)^n dx$.)

Newton's method of extrapolation

In fact, this method extends easily to any integer n


Newton's explanation:

The property of which table is y^t y^e sum of any figure and y^e figure above it is equal to y^e figure next after it save one. Also y^e numerall progressions are of these forms.

a	a	a	a	
b	$a + b$	$2a + b$	$3a + b$	
c	$b + c$	$a + 2b + c$	$3a + 3b + c$	&c.
d	$c + d$	$b + 2c + d$	$a + 3b + 3c + d$	
e	$d + e$	$c + 2d + e$	$b + 3c + 3d + e$	

(See: *Mathematics emerging*, §8.1.1.)

Newton and the general binomial theorem



CUL Add. MS 3958.3, f. 72

Newton's method of interpolation

	$=f_0$	$=f_1$	$=f_2$	$=f_3$	$=f_4$	$=f_5$	$=f_6$	$=f_7$	$=f_8$	$=f_9$	$=f_{10}$	$=f_{11}$	$=f_{12}$	$=f_{13}$	$=f_{14}$	$=f_{15}$
$x^0 X$	1.	1.	1.	1.	1.	1.	1.	1.	1.	1.	1.	1.	1.	1.	1.	1.
$-x^1 X$	-1.	$-\frac{1}{2}$	0.	$\frac{1}{2}$	1.	$\frac{3}{2}$	2.	$\frac{5}{2}$	3.	$\frac{7}{2}$	4.	$\frac{9}{2}$	5.	$\frac{11}{2}$	6.	
$x^2 X$	1.	$\frac{3}{8}$	0.	$-\frac{1}{8}$	0.	$\frac{3}{8}$	1.	$\frac{15}{8}$	3.	$\frac{35}{8}$	6.	$\frac{63}{8}$	10.	$\frac{99}{8}$	15.	
$-x^3 X$	-1.	$-\frac{5}{16}$	0.	$\frac{3}{48}$	0.	$-\frac{1}{16}$	0.	$\frac{5}{16}$	1.	$\frac{35}{16}$	4.	$\frac{105}{16}$	10.	$\frac{231}{16}$	20.	
$x^4 X$	1.	$+\frac{75}{128}$	0.	$-\frac{15}{384}$	0.	$\frac{3}{128}$	0.	$-\frac{5}{128}$	0.	$\frac{35}{128}$	1.	$\frac{315}{128}$	5.	$\frac{1155}{128}$	15.	
$-x^5 X$	-1.	$-\frac{63}{256}$	0.	$\frac{105}{3840}$	0.	$-\frac{3}{256}$	0.	$\frac{3}{256}$	0.	$-\frac{7}{256}$	0.	$\frac{63}{256}$	1.	$\frac{693}{256}$	6.	
$x^6 X$	1.	$\frac{231}{1024}$	0.	$-\frac{245}{46080}$	0.	$\frac{7}{1024}$	0.	$-\frac{5}{1024}$	0.	$\frac{7}{1024}$	0.	$-\frac{31}{1024}$	0.	$\frac{231}{1024}$	1.	

... of intermediate terms may be ...

Newton's method of interpolation

	$(1-x^2)^{-1}$	$(1-x^2)^{-\frac{1}{2}}$	$(1-x^2)^0$	$(1-x^2)^{\frac{1}{2}}$	$(1-x^2)^1$	$(1-x^2)^{\frac{3}{2}}$	$(1-x^2)^2$...
x	1	1	1	1	1	1	1	...
$-\frac{x^3}{3}$	-1	$-\frac{1}{2}$	0	$\frac{1}{2}$	1	$\frac{3}{2}$	2	...
$\frac{x^5}{5}$	1	$\frac{3}{8}$	0	$-\frac{1}{8}$	0	$\frac{3}{8}$	1	...
$-\frac{x^7}{7}$	-1	$-\frac{5}{16}$	0	$\frac{3}{48}$	0	$-\frac{1}{16}$	0	...
$\frac{x^9}{9}$	1	$\frac{35}{128}$	0	$-\frac{15}{384}$	0	$\frac{3}{128}$	0	...
\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\ddots

The entry in the row labelled $\pm \frac{x^m}{m}$ and the column labelled $(1-x^2)^n$ is the coefficient of $\pm \frac{x^m}{m}$ in $\int (1-x^2)^n dx$.

(NB: possible slips in the last two rows of the original table)

Newton's method of interpolation

Can fill in some initial values by other methods

Newton applied the formula

$$\binom{n}{k} = \frac{n(n-1)(n-2)\cdots(n-k+1)}{k!}$$

to fractional n , so that

$$\binom{1/2}{1} = \frac{1}{2}, \quad \binom{1/2}{2} = \frac{1/2(1/2-1)}{2!} = -\frac{1}{8}$$

and so on


Newton's and the general binomial theorem

Newton went on to extend this method to other fractional powers, and also to $(a + bx)^n$, thereby convincing himself of the truth of the **general binomial theorem** — but this was not **proved** until the 19th century

On Newton and the binomial theorem, see

https://www.youtube.com/watch?v=xv_PWwdDWDk

One more table



The table at the bottom of the page gives the interpolations for $(1+x)^n$ for half-integer n

Further discoveries by Newton

By further interpolations and integrations (based on strong geometric intuition) Newton found further series for:

- ▶ $(1 + x)^{p/q}$
- ▶ log, antilog
- ▶ sin, tan, ... (NB: cosine was not yet much in use)
- ▶ arcsin, arctan, ...

(See: *Mathematics emerging*, §§8.1.2–8.1.3.)

Newton on the move from finite to infinite series


And whatever common analysis performs by equations made up of a finite number of terms (whenever it may be possible), this method may always perform by infinite equations: in consequence, I have never hesitated to bestow on it also the name of analysis.

(*De analysi*, 1669; Derek T. Whiteside, *The mathematical papers of Isaac Newton*, CUP, 1967–1981, vol. II, p. 241)

Other 17th-century discoveries (1a)

Brouncker, c. 1655, published 1668: area under the hyperbola

given by $\frac{1}{1 \times 2} + \frac{1}{3 \times 4} + \frac{1}{5 \times 6} + \dots$



Other 17th-century discoveries (1b)

$$\begin{aligned}
 \text{I say } ABCdEA &= \frac{1}{1 \times 2} + \frac{1}{3 \times 4} + \frac{1}{5 \times 6} + \frac{1}{7 \times 8} + \frac{1}{9 \times 10} \&c. \\
 EdCDE &= \frac{1}{2 \times 3} + \frac{1}{4 \times 5} + \frac{1}{6 \times 7} + \frac{1}{8 \times 9} + \frac{1}{10 \times 11} \&c. \\
 EdCyE &= \frac{1}{2 \times 3 \times 4} + \frac{1}{4 \times 5 \times 6} + \frac{1}{6 \times 7 \times 8} + \frac{1}{8 \times 9 \times 10} \&c.
 \end{aligned}
 \left. \vphantom{\begin{aligned} \text{I say } ABCdEA \\ EdCDE \\ EdCyE \end{aligned}} \right\} \text{in infinitum.}$$

(647)


And that therefore in the first series half the first term is greater than the sum of the two next, and half this sum of the second and third greater than the sum of the four next, and half the sum of those four greater than the sum of the next eight, &c. in infinitum. For $\frac{1}{2} dD = br + bn$; but $bn > fg$, therefore $\frac{1}{2} dD > br + fg$, &c. And in the second series half the first term is less than the sum of the two next, and half this sum less than the sum of the four next, &c. in infinitum.

That the first series are the even terms, viz. the 2^d, 4th, 6th, 8th, 10th, &c. and the second, the odd, viz. the 1st, 3^d, 5th, 7th, 9th, &c. of the following series, viz. $\frac{1}{1 \times 2}, \frac{1}{2 \times 3}, \frac{1}{3 \times 4}, \frac{1}{4 \times 5}, \frac{1}{5 \times 6}, \frac{1}{6 \times 7}$, &c. in infinitum = 1. Whereof a being put for the number of terms taken at pleasure, $\frac{1}{a-1}$ is the last, $\frac{a}{a-1}$ is the sum of all those terms from the begin-

ning, and $\frac{1}{a-1}$ the sum of the rest to the end.

That $\frac{1}{4}$ of the first term in the third series is less than the sum of the two next, and a quarter of this sum, less than the sum of the four next, and one fourth of this last sum less than the next eight, I thus demonstrate.

Let a = the 3^d or last number of any term of the first Column, viz. of Divisors,




Other 17th-century discoveries (2)

Mercator's series (1668), found by long division:

$$\frac{1}{1+a} = 1 - a + aa - a^3 + a^4 \text{ (&c.)}$$

Gives rise to series for log



Other 17th-century discoveries (3)



James Gregory (1671):

- ▶ general binomial expansion
- ▶ series for tan, sec, and others, including

$$\theta = \tan \theta - \frac{1}{2} \tan^3 \theta + \frac{1}{5} \tan^5 \theta - \dots$$

$$\text{for } -\frac{\pi}{4} \leq \theta \leq \frac{\pi}{4}$$

Gregory to Collins, 23rd November 1670:

I suppose these series I send here enclosed, may have some affinity with those inventions you advertise me that Mr. Newton had discovered.

(On Gregory's work, see: *Mathematics emerging*, §8.1.4.)

Other 17th-century discoveries (4)

Gottfried Wilhelm Leibniz (1675):

The area of a circle with unit diameter is given by

$$\frac{\pi}{4} = 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \frac{1}{9} - \frac{1}{11} + \&c.$$

The error in the sum is successively less than $\frac{1}{3}$, $\frac{1}{5}$, $\frac{1}{7}$, etc.

Therefore the series as a whole contains all approximations at once, or values greater than correct and less than correct: for according to how far it is understood to be continued, the error will be smaller than a given fraction, and therefore also less than any given quantity. Therefore the series as a whole expresses the exact value.

(See: *Mathematics emerging*, §8.3.)

Series in the 17th century: 'convergence'

John Wallis (1656), *Arithmetica infinitorum*:

$$\square = \frac{4}{\pi} = \frac{3 \times 3 \times 5 \times 5 \times 7 \times 7 \times \dots}{2 \times 4 \times 4 \times 6 \times 6 \times 8 \times \dots}$$

(Determined that

$$\square > \sqrt{\frac{3}{2}}, \quad \square < \frac{3}{2} \sqrt{\frac{3}{4}}, \quad \square > \left(\frac{3 \times 3}{2 \times 4} \right) \sqrt{\frac{5}{4}},$$

and so on)

Brouncker (1668): grouping of terms

Leibniz (1675): 'alternating' series

Power series in the 17th century


Power series (infinite polynomials):

- ▶ enabled term-by-term integration for difficult quadratures;
- ▶ helped establish sine, log, ... as 'functions' (transcendental);
- ▶ encouraged a move from geometric to algebraic descriptions;
- ▶ for Newton (and others) inextricably linked with calculus.

Power series rank with calculus as a major advance of the 17th century

Calculus and series combined

Newton's treatise of 1671,
published 1736




Move on to the 18th century

Eighteenth century:

- ▶ as in 17th century, much progress;
- ▶ also many questions and doubts

Taylor series



Brook Taylor,
*The method of direct and
inverse increments* (1715)

Taylor series

Taylor denoted a small change in x by \dot{x} (our δx), a small change in \dot{x} by \ddot{x} (our $\delta(\delta x)$), and so on

Dependent variable x ; independent variable z increases uniformly with time

x increases to $x + \delta x$ in time δt ; after a further interval of δt , x has become $x + \delta x + \delta(x + \delta x) = x + 2\delta x + \delta(\delta x)$; continuing:

$$\begin{aligned} & x + \frac{n}{1}\delta x + \frac{n(n-1)}{1 \cdot 2}\delta(\delta x) + \frac{n(n-1)(n-2)}{1 \cdot 2 \cdot 3}\delta(\delta(\delta x)) + \dots \\ &= x + \delta x \frac{n\delta z}{1\delta z} + \delta(\delta x) \frac{n\delta z(n-1)\delta z}{1 \cdot 2 \cdot (\delta z)^2} + \delta(\delta(\delta x)) \frac{n\delta z(n-1)\delta z(n-2)\delta z}{1 \cdot 2 \cdot 3(\delta z)^3} + \dots \end{aligned}$$

Taylor series

$$x + \delta x \frac{n\delta z}{1\delta z} + \delta(\delta x) \frac{n\delta z(n-1)\delta z}{1 \cdot 2 \cdot (\delta z)^2} + \delta(\delta(\delta x)) \frac{n\delta z(n-1)\delta z(n-2)\delta z}{1 \cdot 2 \cdot 3(\delta z)^3} + \dots$$

Assumptions:

- ▶ $(n-k)\delta z \approx n\delta z$, since δz is small, so replace each $(n-k)\delta z$ by v , a constant
- ▶ $\delta x \propto \dot{x}$ and $\delta z \propto \dot{z}$, so in each case the former can be replaced by the latter

In essence (in modern terms): $\frac{\delta x}{\delta z} \rightarrow \frac{dx}{dz}$, $\frac{\delta(\delta x)}{(\delta z)^2} \rightarrow \frac{d^2x}{dz^2}$, and so on

Again in modern terms, we arrive at:

$$x + \frac{dx}{dz} v + \frac{d^2x}{dz^2} \frac{v^2}{1 \cdot 2} + \frac{d^3x}{dz^3} \frac{v^3}{1 \cdot 2 \cdot 3} + \dots$$

Cf. Taylor's notation in *Mathematics Emerging*, §8.1.2

Maclaurin's *Treatise of fluxions*, vol. II, p. 610

610 *Of the inverse method of Fluxions.* Book II.

ties multiplied by $k + 1x^n + mx^{2n}$ &c. raised to a power of any exponent k . *De quadrat. curvar.* prop. 5. & 6.

751. The following theorem is likewise of great use in this doctrine. Suppose that y is any quantity that can be expressed by a series of this form $A + Bz + Cz^2 + Dz^3 + \&c.$ where $A, B, C, \&c.$ represent invariable coefficients as usual, any of which may be supposed to vanish. When z vanishes, let E be the value of y , and let $\dot{E}, \ddot{E}, \ddot{\dot{E}}, \&c.$ be then the respective values of $\dot{y}, \ddot{y}, \ddot{\dot{y}}, \&c.$ z being supposed to flow uniformly.

Then $y = E + \frac{\dot{E}z}{1} + \frac{\ddot{E}z^2}{1 \times 2z^2} + \frac{\ddot{\dot{E}}z^3}{1 \times 2 \times 3z^3} + \frac{\ddot{\ddot{E}}z^4}{1 \times 2 \times 3 \times 4z^4} + \&c.$

the law of the continuation of which series is manifest. For since $y = A + Bz + Cz^2 + Dz^3 + \&c.$ it follows that when $z = 0$, A is equal to y ; but (by the supposition) E is then equal to y ; consequently $A = E$. By taking the fluxions, and dividing by \dot{z} , $\frac{\dot{y}}{z} = B + 2Cz + 3Dz^2 + \&c.$ and when

$z = 0$, B is equal to $\frac{\dot{y}}{z}$, that is to $\frac{\dot{E}}{z}$. By taking the fluxions again, and dividing by \dot{z} , (which is supposed invariable) $\frac{\ddot{y}}{z^2} =$

$2C + 6Dz + \&c.$ let $z = 0$, and substituting \ddot{E} for \ddot{y} , $\frac{\ddot{E}}{z^2} =$

$2C$, or $C = \frac{\ddot{E}}{2z^2}$. By taking the fluxions again, and dividing by \dot{z} , $\frac{\ddot{\dot{y}}}{z^3} = 6D + \&c.$ and by supposing $z = 0$, we have $D = \frac{\ddot{\dot{E}}}{6z^3}$.

Thus it appears that $y = A + Bz + Cz^2 + Dz^3 + \&c. =$

$E + \frac{\dot{E}z}{1} + \frac{\ddot{E}z^2}{1 \times 2z^2} + \frac{\ddot{\dot{E}}z^3}{1 \times 2 \times 3z^3} + \frac{\ddot{\ddot{E}}z^4}{1 \times 2 \times 3 \times 4z^4} + \&c.$ This proposition may be likewise deduced from the binomial theorem. Let

Suppose that y can be expressed as $A + Bz + Cz^2 + Dz^3 + \dots$

When z vanishes, $y = E$, $\dot{y} = \dot{E}$,

$\ddot{y} = \ddot{E}$, $\ddot{\dot{y}} = \ddot{\dot{E}}$, and so on

z is assumed to flow uniformly, so that $\dot{z} = \text{const}$

By repeatedly taking fluxions, we may calculate in turn: $A = E$,

$B = \dot{E}\dot{z}$, $C = \frac{\ddot{E}}{2\dot{z}^2}$, $D = \frac{\ddot{\dot{E}}}{6\dot{z}^3}$, etc.

“the law of the continuation of [the] series is manifest”

(*Mathematics emerging*, §8.2.2.)

Euler's *Introductio*

Leonhard Euler, *Introduction to analysis of the infinite* (1748)

INTRODUCTIO
IN ANALYSIN
INFINITORUM.

AUCTORE

LEONHARDO EULERO,

Professore Regio BEROLINENSI, & Academiæ Imperialis Scientiarum PETROPOLITANÆ
Socio.

TOMUS PRIMUS.



LAUSANNÆ,

Apud MARCUM-MICHAELEM BOUSQUET & Socios.

MDCCLVIII

Euler's *Introductio*

Incorporated power series into the definition of a **function**:


*A **function** of a variable quantity is an analytic expression composed in any way whatsoever of the variable quantity and numbers or constant quantities.*

Euler derived series for sine, cosine, exp, log, etc.;

he also discovered relationships between them, for example:

$$\cos v = \frac{1}{2}(e^{iv} + e^{-iv})$$

An application of series



Abraham de Moivre posed this problem about confidence intervals:

What are the Odds that after a certain number of Experiments have been made concerning the happening or failing of Events, the Accidents of Contingency will not afterwards vary from those of Observation beyond certain Limits?

His answer involved clever (but non-rigorous) summation and manipulation of infinite series.

(Mathematics emerging, §7.1.3.)



XXXV^{ME} MÉMOIRE.

Reflexions sur les Suites & sur les Racines imaginaires.

S. I.

Reflexions sur les suites divergentes ou convergentes.

1. SI on éleve $1 + \mu$ à la puissance m , le terme n^e de la serie sera $\mu^{n-1} \times \frac{m(m-1)\dots(m-n+2)}{2 \cdot 3 \cdot 4 \dots n-1}$, & le suivant, c'est-à-dire le $(n+1)^e$, sera $\mu^n \times \frac{m(m-1)\dots(m-n+2)(m-n+1)}{2 \cdot 3 \cdot 4 \dots n-1 \cdot n}$; donc le rapport du $(n+1)^e$ terme au n^e sera $\frac{\mu(m-n+1)}{n}$; or pour que la serie soit convergente, il faut que ce rapport (abstraction faite du signe qu'il doit avoir) soit < que l'unité.

2. Remarquons d'abord que la formule précédente donnera le moyen de former très-prompement les termes d'une suite: par exemple, si $m = \frac{1}{2}$, il faudra multiplier le premier terme par $\mu \times \frac{1}{2}$ pour avoir le second;

Y ij

D'Alembert, 1761:

... all reasoning and calculation based on series that do not converge, or that one may suppose not to, always seems to me extremely suspect, even when the results of this reasoning agree with truths known in other ways.


Introduced, without proof, what came to be known (in a more general setting) as **d'Alembert's ratio test**.

(See: *Mathematics emerging*, §8.3.1.)

Lagrange's use of series

J.-L. Lagrange, *Théorie des fonctions analytiques* (1797)

Lagrange's use of series: an attempt to liberate calculus from infinitely small quantities (essentially by treating only those functions that may be described by power series)



Lagrange and convergence

... [one needs] a way of stopping the expansion of the series at any term one wants and of estimating the value of the remainder of the series.

This problem, one of the most important in the theory of series, has not yet been resolved in a general way

Lagrange found bounds for the 'remainder' ...

and applied his findings to the binomial series ...

thus proving what Newton had taken for granted

(See: *Mathematics emerging*, §8.3.2.)