

## C3.1 Algebraic Topology

Please be aware there are likely typos in these notes: comments/corrections are welcome!

### Course Book

- **Hatcher, Algebraic Topology** — Chp. 2 & 3  
This is also freely available from the author's website.  
Expectations
- You are expected to read chapters 2 & 3 of Hatcher
- You should read the technical remarks about orientation signs in these notes: we will likely not have time for those in lectures.
- This course will not discuss intersection numbers rigorously. The notes often mention these in order to develop your intuition.  
The books by Bott & Tu and Guillemin & Pollack discuss these ideas rigorously

### Other references

- Ulrike Tillmann's C3.1 notes — see course page
- The discussion of orientations in my B3.2 Geometry of Surfaces notes may be helpful.

### Other books

**Massey**, A basic course in Algebraic Topology

(Not "Algebraic Topology: An Introduction" ← does not treat homology)

MORE BASIC but full of ideas:

**Fulton**, Algebraic Topology: a first course.

### MORE ADVANCED:

**May**, A concise course in Algebraic Topology

**Davis & Kirk**, Lecture Notes in Algebraic Topology

**Bredon**, Topology and Geometry

Classics by **Spanier**, **Dold**, also see references in May's book

**Bott & Tu**, **Differential forms in Algebraic Topology**

**Guillemin & Pollack**, **Differential Topology**

## CONTENTS

### 0. OVERVIEW OF THE COURSE

Motivation, category theory, functors  $H_*$  and  $H^*$ : some computations why functors are useful: Invariance of dimension, Brouwer fixed pt thm

### 1. ALGEBRAIC PRELIMINARIES: CHAIN COMPLEXES

Graded abelian groups

chain complexes, chain maps, subcomplex, quotient complex

chain map induces map on homology

exact sequence, snake lemma: SES induces LES on  $H_*$ , naturality of LES

5-Lemma, SES splits  $\Leftrightarrow$  direct sum

### 2. $\Delta$ -COMPLEXES AND SIMPLICIAL HOMOLOGY

$\Delta^n$ ,  $n$ -simplices,  $\Delta$ -complex (structure), simplicial cx, triangulation

simplicial chain complex,  $H_*(S^n)$ ,  $H_*(T^2)$ , remark about orientations

$H_*^{\Delta}(\sqcup \text{conn.comp.}) \cong \bigoplus H_*^{\Delta}(\text{conn.comp.})$ ,  $H_0^{\Delta}(X) \cong \mathbb{Z}^{\# \text{conn.comp}}$

### 3. SINGULAR HOMOLOGY

Motivation, singular chain cx

naturality / functoriality,  $H_*(\text{point})$

### 4. CHAIN HOMOTOPIES AND HOMOTOPY INVARIANCE

chain homotopy, prism operator

homotopic maps  $f \simeq g$  (relative  $A$ ), homotopy equivalent spaces  $X \simeq Y$

contractible space, homotopic maps induce equal maps on homology

homotopy equivalence induces iso on  $H_*$ ,  $H_*(\mathbb{R}^n) = H_*(D^n) = H_*(pt)$

pairs of spaces, relative homology  $H_*(X, A)$ , LES in  $H_*$  for pair

reduced homology  $\tilde{H}_*(X)$ , LES for  $\tilde{H}_*$ ,  $H_*(D^n, S^{n-1}) \cong \tilde{H}_{*-1}(S^{n-1})$

naturality of LES for pairs

### 5. EXCISION THEOREM AND QUOTIENTS

retractions, deformation retractions, excision thm, quotients,

good pairs  $\Rightarrow H^*(X/A) \cong \tilde{H}^*(X/A)$ , generator of  $H_n(S^n) \cong H_n(D^n, S^{n-1}) \cong \mathbb{Z}$

another remark about orientations

Oxford 2019

Prof. Alexander Ritter

ritter@maths.ox.ac.uk

## 6. MAYER-VIETORIS SEQUENCE

MV LES,  $H_*(S^n)$

wedge sum  $X \vee Y$ , cone  $CX$ , suspension  $\Sigma X$ , connected sum  $\#X$

## 7. DEGREE OF MAPS OF SPHERES

degree, application to tangent vector-fields on sphere, hairy ball theorem  
local degree, proof of fundamental thm of algebra

## 8. CELLULAR HOMOLOGY

CW complexes, cellular complex, rank  $H_n^{CW} \leq \#n\text{-cells}$   
 $H_*^{CW}(D^1 \times D^1)$ ,  $H_*^{CW}(\mathbb{R}P^n)$ ,  $H_*^{CW}(S^n)$ ,  $H_*^{CW}(\Sigma g)$

$\Delta\text{-cx} \Rightarrow CW\text{ cx}$ ,  $H_*^{CW}(X) \cong H_*^{\Delta}(X) \cong H_*^{\Delta}(X)$ , Axioms for homology

## 9. COHOMOLOGY

cochains, cohomology,  $H^*(X)$ ,  $H_{CW}^*(X)$ ,  $H_{\Delta}^*(X)$ ,  $H^*(\mathbb{R}P^3)$   
functoriality, homotopy invariance, cochain homotopy, dual of a SES  
excision, LES, Mayer-Vietoris for  $H^*$ , axioms for cohomology

## 10. CUP PRODUCT

Cup product,  $H^*(T^2)$ ,  $H^*(\Sigma_2)$ , remarks about intersection theory  
examples:  $H^*(T^2)$ ,  $H^*(\Sigma_2)$ , remarks about intersection theory

## 11. KÜNNETH FORMULA AND PRODUCT SPACES

Tensor products of  $R$ -mods, tensor product of chain cxes,  
algebraic Künneth thm, product spaces  $X \times Y$ , Euler characteristic  $\chi$   
CW-cx for product space, Künneth thm,  $H^*(S^n \times S^m)$ ,  $H^*(T^n)$

## 12. UNIVERSAL COEFFICIENTS THEOREM

Universal coeff. thm, Background on Ext groups and free resolutions  
(Co)homology with coefficients in a ring/field/module,  $H^*(\mathbb{R}P^2; \mathbb{Z}/2)$

Univ. coeff. thm for PID  $R$ , Duality  $H^*(X; \mathbb{F}) \cong H_*(X; \mathbb{F})$  over fields

Structure thm for f.g. mods  $M$  over PID  $R$ ,  $\text{Ext}_R^1(M; R)$ , torsion shift  $H_*$  to  $H^{*+1}$

## 13. MANIFOLDS: POINCARÉ-LEFSCHETZ DUALITY

Local orientation, orientation, fundamental class, degree, P. duality, L. duality,  
Locally finite homology  $H_*^{\text{lf}}$ , cohomology with compact supports  $H_c^*$ , Cap product and P.D.,  
Alexander duality, knot complements, Jordan curve thm

## 0. OVERVIEW OF THE COURSE

### Motivation

Space  $X$  associate  $\implies$  Algebraic object  $A(X)$   
like numbers, groups, rings, ...

Isomorphism of spaces  $X \cong Y \implies$  Isomorphism  $A(X) \cong A(Y)$

PAY-OFF Can tell spaces apart:

compute  $A(X), A(Y) \rightsquigarrow$  if  $A(X) \neq A(Y)$  then  $X \neq Y$

### Examples

1) Set  $X \longrightarrow A(X) = \#X \in \mathbb{N}$   
same size

(bijection  $X \rightarrow Y$ )  $\implies$

2) Vector space  $X \longrightarrow A(X) = \dim X \in \mathbb{N}$   
(linear iso  $X \rightarrow Y$ )  $\implies$  same dim

3) Topological Space  $X \longrightarrow \# \pi_0(X) = \# \text{path components} \in \mathbb{N}$   
 $\longrightarrow \# \text{Connected components} \in \mathbb{N}$

$\chi(X) = \text{Euler characteristic} \in \mathbb{Z}$   
 $\longleftarrow \text{loops} = C^0(S^1, X)$

Function  $X \times \mathbb{Z}X \longrightarrow \mathbb{Z}$

$(P, \gamma) \longmapsto W(\gamma; P)$

Winding number of  $\gamma$  around  $P$ .

(Homeomorphism  $X \rightarrow Y$ )  $\longrightarrow A(X) = A(Y)$



CONVENTION for this course:

Space = topological space

Map of spaces = continuous map = cts map

For spaces, " $\cong$ " means homeomorphism

"id" = identity map


All diagrams commute unless we say otherwise, e.g.

$A \xrightarrow{\alpha} B$  means  
 $\delta \downarrow \delta \downarrow \beta \circ \alpha = \delta \circ \beta$

# Category theory is the best language to phrase all this

Algebraic topology builds functors

such as



We will not use much category theory, just basic terminology:

Def A category C consists of the data:

Ob(C) = a collection of objects

Hom(A, B) = a set of morphisms between any  $A, B \in \text{Ob } C$  ("arrows")

• with composition rule  $\text{Hom}(B, C) \times \text{Hom}(A, B) \rightarrow \text{Hom}(A, C)$   
 $A \xrightarrow{f} B \xrightarrow{g} C$   
 $\quad \quad \quad \text{g} \circ f$

• with identity morphs  $\text{id}_A \in \text{Hom}(A, A)$  s.t.  $f \circ \text{id}_A = \text{id}_B \circ f = f$

$\forall (f: A \rightarrow B) \in \text{Hom}(A, B)$

Example Sets = {sets with all maps between sets}  
 Top = {topological spaces with continuous maps}  
 Gps = {groups with group homs}

Def A (covariant) functor  $F: C_1 \rightarrow C_2$  is the data:

- an assignment  $(A \in \text{Ob } C_1) \mapsto (F(A) \in \text{Ob } C_2)$
- an assignment  $(A \xrightarrow{f} B) \mapsto (F(A) \xrightarrow{F(f)} F(B))$

$\text{Hom}_{C_1}(A, B) \quad \text{Hom}_{C_2}(F(A), F(B))$

Compatible with identities and compositions.

$F(\text{id}_A) = \text{id}_{F(A)} \quad F(g \circ f) = F(g) \circ F(f)$


A contravariant functor is defined similarly except it reverses the direction of arrows:  $(F(A) \xleftarrow{F(f)} F(B)) \in \text{Hom}(F(B), F(A))$   
 (so  $F(g \circ f) = F(f) \circ F(g)$  reverses order of compositions)

## Examples

- 1)  $F: \text{Top} \rightarrow \text{Sets}, A \mapsto A, f \mapsto f$  "forget the topology and continuity"
- 2)  $F: \text{Sets} \rightarrow \text{Gps}, A \mapsto \text{free abelian group generated by } A$

$Z\langle A \rangle = \left\{ \sum_{\text{finite}} n_i \cdot a_i : a_i \in A, n_i \in \mathbb{Z} \right\}$   
 $(A \xrightarrow{f} B) \mapsto (F(A) : Z\langle A \rangle \rightarrow Z\langle B \rangle \xrightarrow{\sum n_i \cdot a_i} \sum n_i \cdot f(a_i))$

When we say a construction is natural we mean functorial:



A: (a category of spaces)  $\rightarrow$  (a cat. of algebraic objects)  
 The algebraic objects we assigned are assigned compatibly with maps of spaces, and the compatibility maps  $A(f)$  are also compatible w.r.t. composition.  
 So we made compatible choices in constructing A.

Not to be confused with natural transformations of functors (later) which is about relating two such constructions  $A_1, A_2$  in a compatible way


## Example of a functor in algebraic topology (see B.3.5 Topology and Groups course)

$\Pi_1(X, p) = \text{Fundamental group} = \left\{ \gamma \in \mathcal{L}X \mid \gamma(1) = p \right\} / \sim$   
 topological space

Group multiplication: concatenate loops  $\delta_1 * \delta_2$  (each travelling twice as fast)  
 deform:  $h: S^1 \times [0, 1] \rightarrow \mathbb{R}^n, h(t, s) = (1-s)\gamma(t)$  (total # times wind around circle)



Examples:  $\Pi_1(\mathbb{R}^n) = 0$ ,  $\Pi_1(S^1) \cong \mathbb{Z}$ ,  $\Pi_1(S^n) \cong 0$  ( $n \geq 2$ ),  $\Pi_1(\text{torus}) \cong \mathbb{Z}^2$  (not obvious)  
 Those loops generate  $\Pi_1$



Based Top = {Topological spaces with choice of base point, and continuous basepoint-preserving maps}  $\xrightarrow{\Pi_1} \text{Gps}$   
 $(X, p) \mapsto \Pi_1(X, p)$   
 $((X, x) \xrightarrow{f} (Y, y)) \mapsto (\Pi_1(X, x) \xrightarrow{\text{gp.hom.}} \Pi_1(Y, y))$

Lemma Functors map isomorphisms to isomorphisms (iso. means  $\exists$  inverse w.r.t. composition)  
 Pf  $A \xrightarrow{f} B \xrightarrow{g} A \Rightarrow FA \xrightarrow{Ff} FB \xrightarrow{Fg} FA$ , similarly for  $B \xrightarrow{g} A \xrightarrow{f} B$ .  $\square$

Def **Natural transformation**  $\alpha: F \rightarrow G$  between functors  $C \xrightarrow{F} D \xrightarrow{G} E$   
 is an association  $(A \in \text{Ob } C) \mapsto (\alpha_A: F(A) \rightarrow G(A)) \in \text{Hom}_E(F(A), G(A))$

such that  $(A \xrightarrow{f} B) \Rightarrow \begin{matrix} F(A) \xrightarrow{\alpha_A} G(A) \\ \downarrow F(f) \quad \downarrow G(f) \\ F(B) \xrightarrow{\alpha_B} G(B) \end{matrix}$  (commutes)

It is called a **natural isomorphism** if each  $\alpha_A$  is an isomorphism in  $C_2$

Example of a natural transformation in algebraic topology

Let  $H_1(X, P) = \text{abelianisation of } \pi_1(X, P)$  (want to identify  $ab=ba$  so quotient by  $\langle aba^{-1}b^{-1} \rangle$ )  
 $\Rightarrow$  natural trans.  $(\text{Based Top } \xrightarrow{\pi_1} \text{Gps}) \xrightarrow{\alpha} (\text{Based Top } \xrightarrow{H_1} \text{Gps})$  **Commutators**  
 which associates  $(X, P) \mapsto (\alpha_{(X,P)}: \pi_1(X, P) \xrightarrow{\text{quotient}} H_1(X, P))$

Cultural link higher homotopy groups  $\pi_n(X, P) = \left\{ S^n \xrightarrow{\text{cts}} X \right\} / \left\{ \text{basept } \uparrow P \right\} / \text{deform}$   
 FACT abelian for  $n \geq 2$ , but hard: e.g.  $\pi_k(S^n)$  not all known.

We will not study these in this course.

We will study simpler invariants called HOMOLOGY groups  $H_n(X)$

FACT (Hurewicz)  $\exists$  natural transformation  $\pi_n \rightarrow H_n$  which will make sense at the end of course:  
 $f: S^n \xrightarrow{\text{cts}} X$  gives rise to a class  $f_*[S^n] \in H_n(X)$ .

Exercise to practice these notions from category theory:

- Summarise your undergraduate linear algebra as follows:
- $\exists$  functor  $F: \left\{ \begin{matrix} \mathbb{R}^n \text{ for } n \in \mathbb{N} \\ \text{Hom}(\mathbb{R}^m, \mathbb{R}^n) = \text{matrices} \end{matrix} \right\} \rightarrow \left\{ \begin{matrix} \text{finite dimensional real vector spaces} \\ \text{and linear maps} \end{matrix} \right\}$   
 Mat  $\xrightarrow{F} \text{Vect}$
  - A choice of basis for each vector space  $V$  determines a functor  $G: \text{Vect} \rightarrow \text{Mat}$
  - Construct natural isomorphisms  $G \circ F \xrightarrow{\alpha} \text{Id}_{\text{Mat}}$ ,  $F \circ G \xrightarrow{\beta} \text{Id}_{\text{Vect}}$
- When functors satisfying such natural isos exist, the categories are called **equivalent** (not isomorphic). So  $\text{Mat}, \text{Vect}$  are equivalent categories.

Aim of the course: build a functor

HOMOLOGY  $H_*: \text{Top} \rightarrow \text{Graded abelian groups}$   
 $(X \rightarrow Y) \mapsto (H_*(X) \rightarrow H_*(Y))$   
 (grading preserving hom)

and a contravariant functor

COHOMOLOGY  $H^*: \text{Top} \rightarrow \text{Graded rings}$   
 $(X \rightarrow Y) \mapsto (H^*(X) \leftarrow H^*(Y))$

Rough idea:

$H_*X$  is generated by "nice" subspaces  $C \subseteq X$  which have no boundary:  $\partial C = \emptyset$ , modulo identify  $C_1, C_2$  if  $C_1 \cup C_2$  arises as a boundary  $\partial B$ .  
 Call such  $C_1, C_2$  **homologous**.

FACTS


- $H_0(X) \cong \bigoplus_{\text{pts } X} \mathbb{Z} \leftarrow \pi_0 X = \{\text{path-connected components}\} \leftarrow$  generated by a point in each path-comp.
- $X = \sqcup X_i$ : path-components  $\Rightarrow H_*(X) \cong \bigoplus H_*(X_i)$
- $\chi(X) = \sum_{d \geq 0} (-1)^d \text{rank } H_d(X)$  max #  $\mathbb{Z}$ -linearly independent elements

Euler characteristic


Example: compact surfaces

$H_*(\Sigma_g) \cong \begin{cases} \mathbb{Z} & * = 0 \\ \mathbb{Z}^{2g} & * = 1 \\ \mathbb{Z} & * = 2 \\ 0 & \text{else} \end{cases}$   
 orientable surface genus  $g$   
 $\chi = 2 - 2g$

$H_*(N_k) \cong \begin{cases} \mathbb{Z} & * = 0 \\ \mathbb{Z}_2 \oplus \mathbb{Z}^{k-1} & * = 1 \\ 0 & \text{else} \end{cases}$   
 non-orientable surface  $S^2$  with  $k$  Möbius bands attached  
 $\chi = 2 - k$



We will show that those 4 loops generate  $H_1(\Sigma_2)$



Notice  $\gamma$  is a loop. It generates  $H_1(N_1)$



## Example of why such functors are useful

Suppose  $\exists F_*: \text{Top} \rightarrow \text{Gps}$  functors s.t.

①  $F_*(S^n) \neq 0 \iff * = n$  and ②  $F_*(D^n) = 0$  all  $*$

Rmk we'll build such an  $F_*$ :  $\text{reduced homology } \tilde{H}_*$   
 s.t.  $\tilde{H}_* = H_*$  for  $* \neq 0$ , and  $\tilde{H}_0 \cong \mathbb{Z}^{\#(\text{path-components})-1}$

## Theorem Invariance of dimension

$$\begin{matrix} S^n \cong S^m & \iff & n=m \\ \mathbb{R}^n \cong \mathbb{R}^m & \iff & n=m \end{matrix}$$

by ①

Pf Lemma  $\Rightarrow F_n(S^n \cong S^m)$  is iso  $F_n(S^n) \cong F_n(S^m)$  of gps.

If  $\mathbb{R}^n \xrightarrow{\cong} \mathbb{R}^m$ , then can extend  $\times 0$  if  $n \neq m$  ✓

$\varphi$  to the one-point compactifications:  $S^n \cong \mathbb{R}^n \cup \{\infty\} \xrightarrow{\cong} \mathbb{R}^m \cup \{\infty\} \cong S^m$   
 ("Alexandroff extension")  $\xrightarrow{\text{stereographic projection } (x_0, \dots, x_n) \mapsto \frac{(x_1, \dots, x_n)}{1-x_0}}$

Rmk new open neighbourhoods at  $\infty$  are  $\{\infty\} \cup (\mathbb{R}^n \setminus C)$  where  $C$  is (closed B) compact.  
 The extended map is cts since  $\varphi^{-1}(C)$  is (closed B) compact since  $\varphi^{-1}$  is homeo.

## Theorem Brouwer fixed point thm by ① & ②

$f: D^n \rightarrow D^n$  continuous  $\Rightarrow f$  has a fixed point ( $f(p) = p$  some  $p$ )

Proof Suppose not. Let  $r(x) = (\text{ray from } f(x) \text{ to } x) \cap \partial D^n$


notice:  $r: D^n \rightarrow \partial D^n = S^{n-1}$  continuous

$$r|_{\partial D^n} = \text{id}_{S^{n-1}} \quad \text{continuous}$$

$$S^{n-1} = \partial D^n \xrightarrow{\text{inclusion } i} D^n \xrightarrow{r} S^{n-1}$$

apply  $F_{n-1}$   $F_{n-1}(r) \circ F_{n-1}(i) = \text{id} \Rightarrow F_{n-1}(i)$  injective  $F_{n-1}(S^{n-1}) \rightarrow F_{n-1}(D^n) \xrightarrow{\cong} 0$

Example  $A = n \times n$  matrix,  $A_{ij} > 0$  real  $\Rightarrow \exists$  eval  $\lambda > 0$  with real evector  $(v_1, \dots, v_n)$  with  $v_i > 0$



Hint  $\mathbb{R}^3 \cong \Delta^3 = \{x \in \text{octant} : \sum x_i = 1\} \cong D^3$   
 ray  $\mapsto$  ray  $\cap \Delta^n$

## 1. ALGEBRAIC PRELIMINARIES: CHAIN COMPLEXES

### Graded abelian groups

Def A  $\mathbb{Z}$ -graded abelian group  $C$  is an abelian group together

with a direct sum decomposition

$$C = \bigoplus_{n \in \mathbb{Z}} C_n$$

abelian group

Convention: always grade by  $\mathbb{Z}$  unless say otherwise.

Example  $C = \mathbb{Z}[x]$  = integer polynomials in  $x$ ,  $C_n = \mathbb{Z} \cdot x^n \leftarrow$  so grading by degree

A graded ab. gp.  $A$  is a graded subgp of  $C$  if .subgp  
 $\cdot A_n \subseteq C_n$ .

A homomorphism  $h: C \rightarrow D$  of gr.ab.gps is hom of gps s.t.

$$h(C_n) \subseteq D_n$$

A hom of degree  $k$  is hom with

$$h(C_n) \subseteq D_{n+k}$$

Shift by  $k$ :  $\mathbb{Z}$ -gr.ab.gp.  $C[k]$  with

$$C[k]_n = C_{k+n}$$

Notice:

$$C[k]_0 = C_k$$

is now in degree zero, so shifted down by  $k$

$\Rightarrow$  Can view gr. hom of deg  $k$  as a gr. hom  $h: C \rightarrow D[k]$

Abelian groups which are finitely generated

FACT Finitely generated abelian groups are classified:

$$G \cong \mathbb{Z}^r \oplus \mathbb{Z}/p_1^{n_1} \oplus \dots \oplus \mathbb{Z}/p_k^{n_k}$$

free part  $\rightarrow$  called rank  $G$

torsion part

$n_i \in \mathbb{Z}$

$p_i$  primes (possibly not distinct)

Compare finite dimensional vector spaces/field  $\mathbb{F}$ :  $V \cong \mathbb{F}^r$   $r = \dim V$

"homeomorphisms preserve dimension"

Non-trivial result because there are space-filling curves.

e.g. Peano (1890)

$\exists$  cts surjection  $[0,1] \rightarrow [0,1]^2$

interval square


The theorem implies this is not injective.

(cts, bij, compact  $\rightarrow$  Hausdorff)  $\Rightarrow$  homeo

# Chain complexes

differential or boundary homomorph

Def A chain complex  $(C_*, \partial_*)$  is a gr. ab. gp.  $C$  together with a hom  $\partial$  of degree  $-1$  such that  $\partial \circ \partial = 0$ .



Now consider "cycles modulo boundaries":

Def The homology of  $(C_*, \partial_*)$  is the gr. ab. gp.

$$H_n(C_*, \partial_*) = \frac{\text{Ker } \partial_n}{\text{Im } \partial_{n+1}}$$

often abbreviate by  $H_n(C)$

Two cycles are called homologous if they differ by a boundary

Def A chain map  $h : (C_*, \partial_*) \rightarrow (\tilde{C}_*, \tilde{\partial}_*)$

is a graded hom such that

$$h \circ \partial_* = \tilde{\partial}_* \circ h$$

Example A chain subcomplex  $C_* \subseteq \tilde{C}_*$  is a

graded subgp with  $\partial_* = \text{restriction of } \tilde{\partial}_*$  to  $C_*$ .

So the inclusion  $\text{incl} : (C_*, \partial_*) \rightarrow (\tilde{C}_*, \tilde{\partial}_*)$  is a chain map.

Also get quotient complex  $\tilde{C}_*/C_*$

with  $\tilde{\partial}_*[\tilde{c}] = [\tilde{\partial}_*\tilde{c}]$

(well-defined:  $\tilde{\partial}_*C_* = \partial_*C_* \subseteq C_*$ )

Lemma A chain map induces a hom on homology

$$H_*(h) = h_* : H_*(C_*, \partial_*) \rightarrow H_*(\tilde{C}_*, \tilde{\partial}_*)$$

$$[x] \mapsto [h(x)]$$

Proof  $\text{Ker } \partial_n \xrightarrow{\text{hom}} \text{Ker } \tilde{\partial}_n$  since  $\tilde{\partial}(h(x)) = h(\partial x) = 0$

Need  $\text{Im } \partial_n \rightarrow \text{Im } \tilde{\partial}_n$  to get well-defined hom

$$H_n(C) = \text{Ker } \partial_n / \text{Im } \partial_{n+1} \rightarrow \text{Ker } \tilde{\partial}_n / \text{Im } \tilde{\partial}_{n+1} = H_n(\tilde{C})$$

Proof:  $h(b) = \tilde{h}(\partial c) = \tilde{\partial} h(c) \in \text{Im } (\tilde{\partial})$ .  $\square$

The last step was a very simple example of a proof by "diagram chasing"

$$\begin{array}{ccccccc} \dots & \rightarrow & C_{n+1} & \xrightarrow{\partial_{n+1}} & C_n & \xrightarrow{\partial_n} & C_{n-1} \rightarrow \dots \\ & & \downarrow h_{n+1} & & \downarrow h_n & & \downarrow h_{n-1} \\ \dots & \rightarrow & \tilde{C}_{n+1} & \xrightarrow{\tilde{\partial}_{n+1}} & \tilde{C}_n & \xrightarrow{\tilde{\partial}_n} & \tilde{C}_{n-1} \rightarrow \dots \end{array}$$

$$c \xrightarrow{\partial} \partial c = b$$

$$h \downarrow \quad \downarrow h$$

$$h c \xrightarrow{\tilde{\partial}} \tilde{\partial}(h c) = h \partial c = h(b) \quad \square$$

Def  $(C_*, \partial_*)$  is exact (or acyclic) if  $H_*(C) = 0$

so  $\text{Im } \partial_{n+1} = \text{Ker } \partial_n$

In general exact sequence means " $\text{Im}(\text{previous map}) = \text{Ker}(\text{next map})$ "

A short exact sequence (SES) is an exact sequence

$$0 \rightarrow A \xrightarrow{i} B \xrightarrow{\pi} C \rightarrow 0$$





### 5-Lemma

$$\begin{array}{c}
 A \rightarrow B \rightarrow C \rightarrow D \rightarrow E \\
 \cong \downarrow \alpha \cong \downarrow \beta \quad \downarrow \gamma \quad \cong \downarrow \delta \cong \downarrow \epsilon \\
 A' \rightarrow B' \rightarrow C' \rightarrow D' \rightarrow E'
 \end{array}$$

exact rows  $\implies \gamma$  also iso.

Pf exercise (diagram chase)  $\square$

### Splitting Lemma

Cor  $0 \rightarrow A \xrightarrow{\alpha} B \xrightarrow{\beta} C \rightarrow 0$  SES of abelian gps

If  $B \xrightarrow{\beta} C$  s.t.  $\beta \circ \gamma = \text{id}_C$  then the SES splits:  $B \cong A \oplus C$   
(converse is obvious)

Pf  $0 \rightarrow A \rightarrow A \oplus C \rightarrow C \rightarrow 0$   
 $\parallel \quad \downarrow \alpha + \gamma \quad \parallel \quad \parallel$   
 $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0 \quad \square$

Exercise If  $A \xrightarrow{\alpha} B$  s.t.  $\mu \circ \alpha = \text{id}_A$  then it splits:  $B \cong A \oplus C$   
 $\mu \oplus \beta$

Exercise If  $C$  is a free abelian group ( $C \cong \bigoplus_{i \in I} \mathbb{Z}$ ) then the SES splits.

Remark A free  $\neq$  splits, e.g.  $0 \rightarrow \mathbb{Z} \xrightarrow{2} \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$


Cultural Remark Splitting Lemma generalises the rank-nullity theorem from linear algebra:  $V \xrightarrow{\alpha} W$  linear map of vector spaces  $\implies \text{Im } \alpha \oplus \text{Ker } \alpha \cong V$   
 Pf  $0 \rightarrow \text{Ker } \alpha \xrightarrow{\text{incl}} V \xrightarrow{\alpha} \text{Im } \alpha \rightarrow 0$  is SES, and splits since  $\text{Im } \alpha$  free.

## 2. $\Delta$ -COMPLEXES AND SIMPLICIAL HOMOLOGY

standard n-simplex  $\Delta^n = \left\{ (t_0, \dots, t_n) \in \mathbb{R}^{n+1} : t_i \geq 0, \sum t_i = 1 \right\}$   
 $\parallel \sum t_i e_i$

standard basis of  $\mathbb{R}^{n+1}$   
 $(e_0 = (1, 0, \dots, 0), \dots, e_n)$

Examples



Def For  $\{v_0, \dots, v_n\} \subseteq \mathbb{R}^{n+k}$  s.t. any  $k \geq 0$

$v_1, \dots, v_n$   $\mathbb{R}$ -linearly independent


$[v_0, \dots, v_n] = n$ -Simplex spanned by  $v_0, \dots, v_n$

= convex hull of  $v_0, \dots, v_n$

=  $\{ \sum t_i v_i : t_i \geq 0 \text{ and } \sum t_i = 1 \}$

= image of linear homeo  $\sigma: \Delta^n \rightarrow \mathbb{R}^{n+k}$

canonical homeomorphism  $\sigma(e_i) = v_i$



(Solid prism: includes inside)

Will often blur the distinction between map  $\sigma$  and its image,  
 $\sigma \equiv [\sigma e_0, \dots, \sigma e_n]$

but the ordering of the  $v_j$  will be important (so the map  $\sigma$  is more precise)

We encode this extra data by orienting the edges  $v_i \rightarrow v_j$  if  $i < j$

Def d-dimensional faces  $[v_{i_0}, \dots, v_{i_d}]$  for  $i_0 < \dots < i_d$


Example 0-dim faces are the vertices  $v_0, \dots, v_n$

facets =  $(n-1)$ -dimensional faces


=  $[v_0, \dots, \hat{v}_k, \dots, v_n]$  where we omit  $v_k$

=  $\{ \sum t_i v_i \in [v_0, \dots, v_n] : t_k = 0 \}$

=  $\text{Image } \sigma|_{\Delta_k^{n-1}} : \Delta_k^{n-1} \rightarrow \mathbb{R}^{n+k}$   
 $\parallel \{ t \in \Delta^n : t_k = 0 \}$



Example Can build a torus out of simplices:



Def  $\Delta$ -complex is determined by data

- indexing set  $I_n$ , for each  $n \in \mathbb{N}$
- choice of  $n$ -simplex  $\sigma_\alpha^n$  (not necessarily standard) for each  $\alpha \in I_n$
- gluing data: for each  $\alpha \in I_n$ ,  $0 \leq i \leq n$ , associate some  $\beta(\alpha, i) \in I_{n-1}$
- consistency condition (see later)

The  $\Delta$ -complex is the quotient space

$$X = \bigsqcup_{\substack{\alpha \in I_n \\ n \in \mathbb{N}}} \sigma_\alpha^n / \sim$$

$\sim$   $i$ -th facet of  $\sigma_\alpha^n$  is identified with  $\sigma_{\beta(\alpha, i)}^{n-1}$   
 via the order-preserving canonical linear homeo

(quotient topology:  $U \subseteq X$  is open  $\Leftrightarrow U$  intersects  $\sigma_\alpha^n$  in an open set,  $\forall \alpha, n$ )

A  $\Delta$ -complex structure on a top. space  $Y$  is a homeo from a  $\Delta$ -cx  $X \cong Y$ .

Explicit description of the facet identification

$$\left\{ \sum s_i w_i \right\} = [w_0, \dots, w_{n-1}] \longrightarrow [\sigma_0, \dots, \sigma_n] = \left\{ \sum t_i v_i \right\}$$

$$\begin{matrix} \sigma_{\beta(\alpha, i)}^{n-1} \uparrow & \sigma_\alpha^n \uparrow & \sigma_\alpha^{n-1} \Delta_i^{n-1} \\ \sigma_{\beta(\alpha, i)}^{n-1} \uparrow & \sigma_\alpha^n \uparrow & \sigma_\alpha^{n-1} \Delta_i^{n-1} \\ \sigma_{\beta(\alpha, i)}^{n-1} \uparrow & \sigma_\alpha^n \uparrow & \sigma_\alpha^{n-1} \Delta_i^{n-1} \\ \sigma_{\beta(\alpha, i)}^{n-1} \uparrow & \sigma_\alpha^n \uparrow & \sigma_\alpha^{n-1} \Delta_i^{n-1} \\ \sigma_{\beta(\alpha, i)}^{n-1} \uparrow & \sigma_\alpha^n \uparrow & \sigma_\alpha^{n-1} \Delta_i^{n-1} \\ \sigma_{\beta(\alpha, i)}^{n-1} \uparrow & \sigma_\alpha^n \uparrow & \sigma_\alpha^{n-1} \Delta_i^{n-1} \\ \sigma_{\beta(\alpha, i)}^{n-1} \uparrow & \sigma_\alpha^n \uparrow & \sigma_\alpha^{n-1} \Delta_i^{n-1} \\ \sigma_{\beta(\alpha, i)}^{n-1} \uparrow & \sigma_\alpha^n \uparrow & \sigma_\alpha^{n-1} \Delta_i^{n-1} \\ \sigma_{\beta(\alpha, i)}^{n-1} \uparrow & \sigma_\alpha^n \uparrow & \sigma_\alpha^{n-1} \Delta_i^{n-1} \\ \sigma_{\beta(\alpha, i)}^{n-1} \uparrow & \sigma_\alpha^n \uparrow & \sigma_\alpha^{n-1} \Delta_i^{n-1} \end{matrix}$$

$$\sigma_\alpha^n = \{ s_0 v_0 + \dots + s_{i-1} v_{i-1} + s_i v_i + \dots + s_{n-1} v_{n-1} \}$$


$$\sigma_\alpha^{n-1} \Delta_i^{n-1} = \{ v_0, \dots, v_{i-1}, v_{i+1}, \dots, v_n \}$$

$$\Delta^{n-1} \longrightarrow \Delta_i^{n-1} \subseteq \Delta^n$$

$$(s_0, \dots, s_{n-1}) \mapsto (s_0, \dots, s_{i-1}, 0, s_i, \dots, s_{n-1})$$

Non-example

This decomposition for  $T^2$  is not a  $\Delta$ -complex.



Consistency condition

We want to additionally ensure that each point of  $X$  lies in the interior of exactly one  $\sigma_\alpha^n$ , because we want to avoid unexpected identifications.



then glue  $\sigma_1^2 = \sigma_2^2$  via  $\sigma_1^2 \times \sigma_2^2$

notice how  $\sigma_3, \sigma_4$  get identified in the quotient, but we only notice this after gluing  $\sigma_1^2$  (if you try to run the definition of simplicial homology-defined later-you notice that the differential cannot satisfy  $\partial_0 \partial_2 = 0$ )

Equivalently: the facet gluing maps are compatible under double restriction:  $\forall i, j$

$$\begin{matrix} \text{facet} & \text{facet} & \text{facet} & \text{facet} \\ \text{[} v_0, \dots, v_n \text{]} & \text{[} v_0, \dots, v_{n-1} \text{]} & \text{[} w_0, \dots, w_{n-1} \text{]} & \text{[} x_0, \dots, x_{n-2} \text{]} \\ \text{[} v_0, \dots, v_n \text{]} & \xrightarrow{\text{identity}} & \text{[} w_0, \dots, w_{n-1} \text{]} & \xrightarrow{\text{identity}} & \text{[} x_0, \dots, x_{n-2} \text{]} \\ \text{[} v_0, \dots, v_n \text{]} & \xrightarrow{\text{identity}} & \text{[} z_0, \dots, z_{n-1} \text{]} & \xrightarrow{\text{identity}} & \text{[} x_0, \dots, x_{n-2} \text{]} \end{matrix}$$

this ensures that  $[v_0, \dots, v_i, \dots, v_j, \dots, v_n]$  is identified with the same  $[x_0, \dots, x_{n-2}]$  whether we first restrict to  $t_i=0$  (omit  $v_i$ ) or first restrict to  $t_j=0$  (omit  $v_j$ ).

Another equivalent condition: can define the  $k$ -th skeleton of  $\Delta$ -cx  $X$ ,

$X^k =$  quotient space you get by gluing all simplices of dimensions  $\leq k$ . Consistency is the condition that the boundary of each  $\sigma_\alpha^n$  should map continuously into  $X^{n-1}$


(in the above Example consider the vertex  $\Delta = \partial \sigma_1^2$ )

Rmk (see part A) A simplicial complex is a  $\Delta$ -complex in which (more precisely, the topological realisation" of a simplicial complex)

each  $d$ -dim face is uniquely determined by  $d$  distinct vertices.

A homeo from such a complex to  $X$  is a triangulation of  $X$ .

Non-example



both 2-simplices have vertices  $v, v, v$

$$T^2 =$$



whereas  $T^2 =$  is a triangulation.

Simplicial chain complex

Def For a  $\Delta$ -complex  $X$ , let  $X_n =$  set of  $n$ -simplices of  $X$

$C_n^{\Delta}(X) =$  free abelian group generated by the set  $X_n$

$$= \left\{ \sum_{\alpha \in I_n} c_\alpha \sigma_\alpha^n : c_\alpha \in \mathbb{Z} \text{ and only finitely many } c_\alpha \neq 0 \right\}$$

differential:  $\partial_n \sigma_\alpha = \sum (-1)^i \cdot \sigma_{\beta(\alpha, i)}$

so:  $\partial_n [v_0, \dots, v_n] = \sum (-1)^i \cdot [v_0, \dots, \hat{v}_i, \dots, v_n]$  and extend linearly

will show  $\partial \circ \partial = 0$ , so get simplicial homology:  $H_*^\Delta(X) = H_*(C_*^\Delta, \partial_*)$

Examples

$\partial_1 \begin{pmatrix} v_0 \\ \rightarrow \\ v_1 \end{pmatrix} = \begin{pmatrix} \bullet \\ -v_0 \\ +v_1 \end{pmatrix}$

$\partial_2 \begin{pmatrix} \triangle \\ \nearrow v_0 \\ \searrow v_1 \\ \rightarrow v_2 \end{pmatrix} = \begin{pmatrix} \bullet \\ +v_0 \\ -v_1 \\ +v_2 \end{pmatrix}$

$\partial_2 \circ \partial_1$  (this)  $= + (v_2 - v_1) - (v_2 - v_0) + (v_1 - v_0) = 0$

$\partial \circ \partial = 0$  fails for  $\triangle$  (not  $\Delta$ -complex), try!

Lemma  $\partial \circ \partial = 0$

Pf  $\partial_{n-1} (\partial_n [v_0, \dots, v_n]) = \sum (-1)^i \partial_{n-1} [v_0, \dots, \hat{v}_i, \dots, v_n]$

$= \sum (-1)^i (-1)^{j-1} [v_0, \dots, \hat{v}_i, \dots, \hat{v}_j, \dots, v_n]$  antisymmetric if swap  $v_i, v_j$

$= 0$

Example  $S^1 = \text{circle}$

$\Delta$ -cx:  $X_0 = 1$  0-simplex  $\bullet$   $e_i^0 = e_{\beta(i,0)} = e_{\beta(i,1)}$

$X_1 = 1$  1-simplex  $\rightarrow e_i^1$

$0 \rightarrow C_1^\Delta \rightarrow C_0^\Delta \rightarrow 0$

$\cong \mathbb{Z} \xrightarrow{e} \mathbb{Z} \rightarrow 0$

$e \mapsto v - v = 0$

$\Rightarrow H_*^\Delta(S^1) = \begin{cases} \mathbb{Z} & * = 0, 1 \\ 0 & \text{else} \end{cases}$

Example  $\Delta$ -cx structure on  $S^n$ :


$S^n = \Delta^n \cup \Delta^n / \text{glue along } \partial \Delta^n$

call this  $\Delta_1$  this  $\Delta_0$

One can deduce:  $H_* S^n = \begin{cases} \mathbb{Z} \langle \text{point} \rangle \cong \mathbb{Z} & * = 0 \\ \mathbb{Z} \langle \Delta_1 - \Delta_0 \rangle \cong \mathbb{Z} & * = n \\ 0 & \text{else} \end{cases}$

pick any vertex

Example



$0 \rightarrow C_2^\Delta \xrightarrow{\cong} C_1^\Delta \xrightarrow{\cong} C_0^\Delta \rightarrow 0$

$\mathbb{Z}f_1 + \mathbb{Z}f_2 \xrightarrow{\cong} \mathbb{Z}e_1 + \mathbb{Z}e_2 + \mathbb{Z}e_3 \xrightarrow{\cong} \mathbb{Z}v$

$f_1 \mapsto e_1 - e_3 + e_2$

$f_2 \mapsto e_2 - e_3 + e_1$

$e_1, e_2, e_3 \mapsto v - v = 0$

$H_*^\Delta(T^2) = \begin{cases} \mathbb{Z}v & * = 0 \\ (\mathbb{Z}e_1 + \mathbb{Z}e_2 + \mathbb{Z}e_3) / \mathbb{Z}(e_1 - e_3 + e_2) & * = 1 \\ \mathbb{Z}(f_1, -f_2) & * = 2 \\ 0 & \text{else} \end{cases}$


$* = 1 \leftarrow$  freely generated by  $e_1, e_2$

Smith normal form of  $\partial_2$ :  $\begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ -1 & -1 & 0 \end{pmatrix} \xrightarrow{\text{row ops}} \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{pmatrix} \xrightarrow{\text{col ops}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}$

so after  $\mathbb{Z}$ -isos of  $C_2, C_1$ , we get  $\mathbb{Z}^2 \rightarrow \mathbb{Z}^3, (a, b) \mapsto (a, 0, 0)$

Remark about orientations (see also my B3.2 Geometry of Surfaces notes)

For vector space an orientation is a choice of basis modulo linear endomorphisms of  $\det > 0$




Fact  $GL(n, \mathbb{R})$  has 2 path-components  $\langle A : \det A > 0 \rangle$  so can always continuously deform a basis to another within same orientation

Canonical orientation on  $\mathbb{R}^n$ :  $e_1, \dots, e_n$  standard basis  $\leftarrow$  "positive orientation"

Example  $[v_0, \dots, v_n]$  simplex  $\Rightarrow v_1 - v_0, \dots, v_n - v_0$  is a basis of vector subspace  $V = \{ \sum a_i v_i : \sum a_i = 0 \} \subseteq \mathbb{R}^{n+1}$

hence a choice of orientation of  $V$ , and each transposition of vertices  $v_0, \dots, v_n$  switches the orientation class.

If  $v_0, v_1 \in \mathbb{R}^n$  then  $V = \mathbb{R}^n$  so simplex's orientation can be compared with  $\mathbb{R}^n$ -orient.




- No canonical choice of orientation for abstract vector space.
- Need choose basis  $v_1, \dots, v_n$  then declare another basis positively oriented if the change of basis matrix has  $\det > 0$ .

For hyperplane  $H \subseteq \mathbb{R}^n$  with choice of normal can declare orientation of basis  $w_1, \dots, w_{n-1}$  of  $H$  positive if normal,  $w_1, \dots, w_{n-1}$  is positive  $\mathbb{R}^n$ -basis convention "outward normal first"




Example  $\xrightarrow{e_1}$   $H \subseteq \mathbb{R}^2 \Rightarrow e_1$  positive basis for  $H$   
 $(\text{normal}, e_1) = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \det = +1 > 0$

Example  $\Delta^n \subseteq \mathbb{R}^{n+1}$  with normal  $(1, 1, \dots, 1)$  is positively oriented.  
UPSHOT For an  $n$ -simplex  $[v_0, \dots, v_n]$  in  $\mathbb{R}^n$ , each facet lies in a hyperplane and have canonical choice of normal: outward normal. Hence facets are canonically oriented.



Any reflection of  $\mathbb{R}^n$  will swap orientation: after  $\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$  get clockwise



UPSHOT  $(-1)^i$  in  $(-1)^i [v_0, \dots, \hat{v}_i, \dots, v_k]$  in definition of simplicial  $\partial$  is there to ensure that orientations are consistent (crucial for  $\partial \partial = 0$ )


Lemma  $H_*^\Delta(X) \cong \bigoplus H_*^\Delta(X_i)$  where  $X_i$  are the path-components of  $X$ .

Pf  $\bigoplus C_*^\Delta(X_i) \rightarrow C_*^\Delta(X), \oplus c_i \mapsto \Sigma c_i$  since  $\Delta^k$  path-conn.  
 is chain isomorphism since any simplex  $\sigma: \Delta^k \rightarrow X$  has path-connected image, so  $\subseteq X_i$  some  $i$ .  $\square$

Theorem  $X$  has  $\Delta$ -cx structure  $\Rightarrow H_0^\Delta(X) \cong \bigoplus \mathbb{Z}$  Path-Conn. components

Pf By Lemma, wlog  $X$  path-connected

- vertex  $v \Rightarrow v \in C_0(X) \xrightarrow{\partial} C_{-1}(X) \cong 0 \Rightarrow [v] \in H_0(X)$
- vertices  $v_0, v_1 \in X \Rightarrow \exists$  path  $\gamma$  from  $v_0$  to  $v_1$ , can homotope path so that going edges (continuously deform)  $\Rightarrow \gamma$  is sum of 1-chains s.t.  $\partial \gamma = v_1 - v_0$
- $\Rightarrow [\gamma] \in H_0(X)$  independent of choice of  $\gamma$
- $\Rightarrow H_0(X) = \langle [v] \rangle$



$H_0(X) = \langle [v] \rangle \leftarrow \mathbb{Z}$  is injective?

$n \nu \leftarrow n$  suppose  $n \nu = \partial c$  some  $c \in C_1(X)$   
 consider the augmentation hom

$$C_1(X) \xrightarrow{\partial} C_0(X) \xrightarrow{\epsilon} \mathbb{Z}$$

$\xrightarrow{\sum n_i \sigma_i} \xrightarrow{\sum n_i}$   
 0-simplices


notice composite is 0 since  $\partial \begin{pmatrix} 1\text{-simplex} \\ \sigma_0 \rightarrow \sigma_1 \end{pmatrix} = \sigma_1 - \sigma_0 \xrightarrow{\epsilon} 1 - 1 = 0$   
 $\Rightarrow n = \epsilon(n \nu) = \epsilon \partial c = 0$ .

Rmk  $X$  top space  $\Rightarrow$  path conn. component  $\subseteq$  connected component since path-conn.  $\Rightarrow$  connected. For  $\Delta$ -cx, these are same (since connected + locally path-conn.  $\Rightarrow$  path-connected).

However for top. space they need not be the same, e.g. Topologist's sine curve



### 3. SINGULAR HOMOLOGY

Motivation Not obvious that  $H_*^\Delta$  is functorial:  $\Delta^k \xrightarrow{\sigma} X \xrightarrow{f} Y$   
 then  $f \circ \sigma$  typically not a simplex:  $\Delta \xrightarrow{\sigma} X \xrightarrow{f} Y$    $\xrightarrow{\text{continuous map}}$

Solution 1: only allow simplical maps  $f: X \rightarrow Y$  (so for simplex  $\forall \sigma$ )

Solution 2: show that any cts map  $f: X \rightarrow Y$  can be approximated arbitrarily well by a simplicial map after having performed barycentric subdivision on  $X, Y$  enough times. Also any two such approximations induce the same map  $H_*^\Delta(X) \rightarrow H_*^\Delta(Y)$

Solution 3: develop new homology  $H_*(X)$  which allows any cts map  $\Delta^n \rightarrow X$  will do THIS. and prove  $H_*^\Delta(X) \cong H_*(X)$  for  $\Delta$ -complexes  $X$ .

Def Singular n-simplex is any  $\boxed{\text{continuous map } \sigma: \Delta^n \rightarrow X}$   
 $X$  is any top. space

Singular n-chains  $C_n(X) =$  free abelian group generated by  $\sum_{\text{singular } n\text{-simplices } \sigma} c_\sigma \cdot \sigma$  only finitely many  $c_\sigma \neq 0$

$\partial_n \sigma = \sum_{i=0}^n (-1)^i \cdot \sigma|_{\Delta_i^{n-1}}$  (and extend linearly)  
 Rmk Here  $\Delta_i^{n-1} = [e_0, \dots, \hat{e}_i, \dots, e_n]$  is identified canonically with  $\Delta^{n-1}$  (send  $e_k \rightarrow e_{k-1}$  for  $k < i$ )

Will show  $\partial \circ \partial = 0$ , so get singular homology:  $\boxed{H_*(X) = H_*(C_*, \partial_*)}$

For  $\Delta$ -complex  $X$  have inclusion of subcomplex  $C_*^\Delta \rightarrow C_*$   
 $\Rightarrow$  induces  $H_*^\Delta(X) \rightarrow H_*(X)$  Fact: isomorphism (proof later, see cellular  $H_*^{CW} \cong H_*$ )

Corollary  $H_*^\Delta(X)$  is independent of choice of  $\Delta$ -cx structure on  $X$

Lemma  $\partial \circ \partial = 0$

Proof  $\partial_{n+1}(\partial_n \sigma) = \partial_{n+1}(\sum_{i=0}^n (-1)^i \sigma|_{\Delta_i^{n-1}})$   
 $= \sum_{j < i} (-1)^j (-1)^i \sigma|_{[e_0, \dots, \hat{e}_j, \dots, \hat{e}_i, \dots, e_n]}$   
 $+ \sum_{j > i} (-1)^j (-1)^i \sigma|_{[e_0, \dots, \hat{e}_i, \dots, \hat{e}_j, \dots, e_n]} = 0$   $\square$

Example  $X = \text{point} \Rightarrow C_n(X) = \mathbb{Z} \cdot (\sigma_n: \Delta^n \rightarrow X \text{ the constant map})$   
 $\partial \sigma_n = \sum_{i=0}^n (-1)^i \sigma_n|_{\Delta_i^{n-1}} = \sum_{i=0}^n (-1)^i \sigma_{n-1} \Rightarrow \dots \Rightarrow \mathbb{Z} \xrightarrow{\partial} \mathbb{Z} \xrightarrow{\partial} \mathbb{Z} \xrightarrow{\partial} \dots \xrightarrow{\partial} \mathbb{Z} \xrightarrow{\partial} \mathbb{Z} \xrightarrow{\partial} \dots \xrightarrow{\partial} \mathbb{Z} \xrightarrow{\partial} \mathbb{Z} \xrightarrow{\partial} \dots$   
 $\begin{cases} 1 & \text{if } n \text{ even} \\ 0 & \text{if } n \text{ odd} \end{cases} \Rightarrow H_*(\text{pt}) \cong \begin{cases} \mathbb{Z} & * = 0 \\ 0 & \text{else} \end{cases}$

Lemma  $\boxed{H_*(X) \cong \bigoplus_i H_*(X_i)}$  where  $X_i$  are path-components of  $X$   
Pf Image of cts map  $\Delta^n \rightarrow X$  is path conn. so lies in some  $X_i$ .  $\square$

Cor  $\boxed{H_0(X) = \bigoplus_i \mathbb{Z}}$   $\leftarrow$  generators of  $C_0(X)$

Pf By Lemma, wlog  $X$  path-connected.  $\Delta^0 = \text{pt} \rightarrow X$  is cycle since  $C_{-1}(X) = \emptyset$   
 Given 2 points  $x, y \in X$ , a path  $\Delta^1 = [0, 1] \xrightarrow{\gamma} X$ ,  $\gamma(0) = x, \gamma(1) = y$  is also a 1-chain!  
 So  $x - y = \partial \gamma$ , so  $x, y$  are homologous. Finally if  $n \cdot [x] = 0 \in H_0(X)$  then  $n \cdot x = \partial c$  some  $c \in C_1(X)$  generated by paths. Now run the augmentation hom-trick like we did for  $H_0^{\Delta}$ :  $n = \varepsilon(n \cdot x) = \varepsilon \partial c = 0$  as  $\varepsilon \partial = 0$ .  $\square$

### Naturality (i.e. functoriality)

Lemma  $f: X \rightarrow Y$  continuous

$\Rightarrow H_*(f) = f_*: H_*(X) \rightarrow H_*(Y)$  induced by chain map  
 $f_*: C_*(X) \rightarrow C_*(Y)$  and extend linearly  
 $\boxed{f_*(\sigma) = f \circ \sigma}$

Pf  $\partial_n(f_* \sigma) = \sum_{i=0}^n (-1)^i f \circ \sigma|_{\Delta_i^{n-1}} = f_* (\sum_{i=0}^n (-1)^i \sigma|_{\Delta_i^{n-1}}) = f_* (\partial_n \sigma) = \partial_n(f_* \sigma)$   $\square$

Properties 1)  $X \xrightarrow{f} Y \xrightarrow{g} Z \Rightarrow (g \circ f)_* = g_* \circ f_*$   
 2)  $\text{id}_X = \text{id} \Rightarrow \text{id}_*(\sigma) = \sigma$

Pf 1)  $(g \circ f)_* \sigma = g_* \circ f_* \sigma = g_* (f_* \sigma) = g_* (\sigma)$   $\checkmark$   
 2)  $\text{id}_*(\sigma) = \text{id} \circ \sigma = \sigma$   $\checkmark$   $\square$

Cor  $H_*: \left\{ \begin{array}{l} \text{topological spaces} \\ \& \text{cts maps} \end{array} \right\} \rightarrow \left\{ \begin{array}{l} \text{graded abelian groups} \\ \& \text{graded homs} \end{array} \right\}$  is a functor  
Cor  $X \cong Y \Rightarrow H_*(X) \cong H_*(Y)$

# 4. CHAIN HOMOPIES AND HOMOLOGY INVARIANCE

Algebra: chain homotopies

$f_*, g_* : (C_*, \partial_*) \rightarrow (\tilde{C}_*, \tilde{\partial}_*)$  chain maps


Def  $f_*, g_*$  are chain homotopic if  $\exists$  (degree +1)

hom  $h : C_* \rightarrow \tilde{C}_*[1]$  s.t.

$$\tilde{\partial} \circ h + h \circ \partial = f - g$$

$h$  is called a chain homotopy

Consequence  $f_* = g_* : H_+(C_*, \partial_*) \rightarrow H_+(\tilde{C}_*, \tilde{\partial}_*)$  on homology



$c$  cycle  $\in C_n$

$$\Rightarrow f_n(c) - g_n(c) = \underbrace{\tilde{\partial}_{n+1} h_n(c)}_{\text{boundary}} + h_{n-1} \circ \underbrace{\partial_n(c)}_0 = 0$$

$$\Rightarrow f_n(c) = g_n(c) \text{ in } H_*(\tilde{C}) \quad \square$$

Theorem  $i_0 : X \rightarrow X \times I, i_0(x) = (x, 0)$  where  $I = [0, 1]$

$i_1 : X \rightarrow X \times I, i_1(x) = (x, 1)$

$\Rightarrow i_{0*}, i_{1*} : C_*(X) \rightarrow C_*(X \times I)$  are chain hpic.

Key idea Need "prism operator" which cuts  $\Delta^n \times I$  into a sum  $\Gamma_n$

of  $(n+1)$ -simplices in  $\Delta^n \times I$ :

$$(\sigma : \Delta^n \rightarrow X) \mapsto \sigma \text{ xid} : \Delta^n \times I \rightarrow X \times I$$

$\Gamma_n = \text{combo of maps } \Delta^{n+1}$

$$\text{Prism operator } P_n \rightarrow (\sigma \text{ xid}) \circ \Gamma_n : \Delta^{n+1} \rightarrow X \times I$$

What is  $\partial$  of  $P_n \circ \sigma$ ?




hence  $P$  is chain hpic

Pf  $\leftarrow$  Non-examinable

$$\text{bottom facet } \Delta^n \times 0 = [v_0, \dots, v_n] \leftarrow v_i = e_i \times 0 \subseteq \Delta^n \times [0, 1] \subseteq \mathbb{R}^{n+1}$$

$$\text{top facet } \Delta^n \times 1 = [w_0, \dots, w_n] \leftarrow w_i = e_i \times 1$$

Examples




$n=1$ :

two 2-simplices

$$[v_0, v_1, w_1]$$

$$[v_0, w_0, w_1]$$

$n=2$ :



three 3-simplices:

$$[v_0, v_1, v_2, w_2]$$

$$[v_0, v_1, w_1, w_2]$$

$$[v_0, w_0, w_1, w_2]$$

Let  $s_i = [v_0, \dots, v_i, w_i, \dots, w_n]$

Claim The  $s_i$  cover  $\Delta \times [0, 1]$  and give  $\Delta$ -cx structure on  $\Delta^n \times I$

$$\text{Pf } \sum_{k \leq i} t_k v_k + \sum_{k > i} s_k w_k = (t_0, \dots, t_{i-1}, t_i + s_i, s_{i+1}, \dots, s_n, s_i + \dots + s_n)$$

So given  $(x_0, \dots, x_n, a) \in \Delta^n \times I$ , equate and solve:

$$t_0 = x_0, \dots, t_{i-1} = x_{i-1}, s_{i+1} = x_{i+1}, \dots, s_n = x_n, \text{ and } \begin{cases} s_i = a - x_{i+1} - \dots - x_n \\ t_i = x_i - s_i \end{cases}$$

Note  $x_k \geq 0, \sum x_k = 1, a \in [0, 1]$  hence  $\sum t_k + \sum s_k = 1 \checkmark$

but  $s_i \geq 0 \Leftrightarrow \begin{cases} a \geq x_{i+1} + \dots + x_n \\ t_i \geq 0 \end{cases}$ . Thus a solution exists if we pick  $i = \min\{k : a \geq x_{k+1} + \dots + x_n\}$

There are multiple solutions if  $x_{i+1} = x_{i+2} = \dots = x_j = 0$ , but that is as expected: those points of  $\Delta^n \times I$  belong to the faces of  $s_i, s_{i+1}, \dots, s_j$ .  $\square$


Def

$$\Gamma_n = \sum (-1)^i s_i \in C_{n+1}(\Delta^n \times [0, 1]) \leftarrow \text{geometrically this "represents" } \partial(\Delta^n \times I) \text{ as a simplicial chain}$$

$$\Rightarrow \partial \Gamma_n = \sum_{j \leq i} (-1)^i (-1)^j [v_0, \dots, v_j, \dots, v_i, w_i, \dots, w_n] \leftarrow \text{geometrically, this "represents" } \partial(\Delta^n \times I)$$

$$+ \sum_{i > j} (-1)^i (-1)^j [v_0, \dots, v_i, w_i, \dots, w_j, \dots, w_n] = \partial(\Delta^n \times I) \cup (\Delta^n \times \partial I)$$

Example




$$\Gamma_1 = [v_0, w_0, w_1] - [v_0, v_1, w_1] \text{ "is the square"}$$

$$\partial \Gamma_1 = [w_0, w_1] - [v_0, w_1] + [v_0, w_0] - [v_1, w_1] + [v_0, v_1] - [v_0, v_1] \text{ "inside facets" cancel}$$

"is  $\partial$  of square"

Def  $X \simeq Y$  homotopy equivalent spaces if  $\exists$  maps



Rmk homeo  $\Rightarrow$  hpy equivalent


Def  $X$  contractible if  $X \simeq \text{pt}$

equivalently  $(X \xrightarrow{\text{id}} X) \simeq (X \xrightarrow{\text{const}} \text{point} \in X)$

Example.  $\mathbb{R}^n \simeq \text{pt}$

$F(x, t) = tx$  then  $f_0 \equiv 0, f_1 = \text{id}$ .

(star-shaped subsets of  $\mathbb{R}^n \simeq \text{pt}$ )



Theorem  $f_0 \simeq f_1 \Rightarrow f_{0*} = f_{1*}$

Pf  $f_{1*} - f_{0*} = F_* i_{1*} - F_* i_{0*} = F_* (i_{1*} - i_{0*}) = F_* (\partial P + P \partial)$

previous  $\xrightarrow{\text{Thm}} = \partial_0(F_* P) + (F_* P) \partial$

$F_*$  chain map  $\Rightarrow F_* \partial P$  is chain hpy from  $f_{0*}$  to  $f_{1*}$   $\square$

Cor  $X \simeq Y \Rightarrow H_* X \cong H_* Y$

Pf  $f_* g_* = \text{id}_*$ ,  $g_* f_* = \text{id}_*$   $\square$

Example  $X$  contractible  $\Rightarrow H_* X \cong H_*(\text{pt}) \cong \begin{cases} \mathbb{Z} & * = 0 \\ 0 & \text{else} \end{cases}$

Cultural Rmk (Whitehead's theorem) For nice topological spaces  $\leftarrow$  (CW complexes - see later in course) if  $X, Y$  are simply connected and  $\exists f: X \rightarrow Y$  inducing isomorphisms on  $H_*$  then  $X \simeq Y$  are homotopy equivalent.

Prism operator  $P: C_n(X) \rightarrow C_{n+1}(X \times [0, 1])$

$$P(\sigma) = (\sigma \times \text{id})_* (\bar{\Gamma}_n)$$

$\sigma: \Delta^n \rightarrow X$   
 $\sigma \times \text{id}: \Delta^n \times [0, 1] \rightarrow X \times [0, 1]$   
 $(\sigma \times \text{id})(x, t) = (\sigma(x), t)$

this abbreviated notation means the map

$$\partial P(\sigma) = \partial(\sigma \times \text{id})_* (\bar{\Gamma}_n) = (\sigma \times \text{id})_* (\partial \bar{\Gamma}_n)$$

$$= \sum_{i \leq n} (-1)^i (-1)^i [i_0 \sigma e_0, \dots, i_0 \sigma e_i, \dots, i_0 \sigma e_n]$$

$$+ \sum_{j \geq 1} (-1)^j (-1)^{j+1} [i_0 \sigma e_0, \dots, i_0 \sigma e_j, \dots, i_0 \sigma e_n]$$

$$= i_0 \sigma - i_n \sigma - P \partial \sigma$$

$$\uparrow \quad \uparrow \quad \uparrow$$

$$i=j=0 \quad i=j=n$$

$$1^{\text{st}} \text{ sum} \quad 2^{\text{nd}} \text{ sum}$$

$$((\partial \sigma) \times \text{id})_* \bar{\Gamma}_{n-1}$$

$$\sum_i (-1)^i [v_0, \dots, v_i, w_i, \dots, w_{n-1}]$$

now use  $\otimes$  and

$$\partial \sigma = \sum (-1)^k [\sigma e_0, \dots, \sigma e_k, \dots, \sigma e_n]$$

Def  $f_0 \simeq f_1$  homotopic if  $\exists$  continuous map  $F: X \times [0, 1] \rightarrow Y$

s.t.  $f_0 = F \circ i_0$   
 $f_1 = F \circ i_1$   
 called homotopy

Idea Think of this as a continuous family of maps from  $f_0$  to  $f_1$ .

Exercise  $\simeq$  is an equivalence relation.

Homotopic relative to A  $\subseteq X$  if  $F(a, t) = f_0(a) = f_1(a)$  all  $a \in A$  all  $t$ .  
 write "f  $\simeq_g$  rel A"


### Relative homology

Def  $(X, A)$  pair of spaces if  $A \subseteq X$  topological subspace

$\Rightarrow \hat{i} = \text{incl}: A \hookrightarrow X$  induces a subcx  $\hat{i}_*: C_*A \rightarrow C_*X$

$\Rightarrow C_*X/C_*A$  quotient chain cx (recall  $\partial[x] = [\partial x]$ )

$$H_*(X, A) = H_*(C_*X/C_*A)$$




Idea: relative cycles:  $c \in C_*X$  s.t.  $\exists x \in C_{*+1}X$   $\partial x = b + a \in C_*A$

relative boundaries:  $b \in C_*X$  s.t.  $\exists x \in C_{*+1}X$   $\partial x = b + a \in C_*A$

$$\Rightarrow 0 \rightarrow C_*A \rightarrow C_*X \rightarrow C_*X/C_*A \rightarrow 0 \text{ SES}$$

$$\text{Cor } \dots \rightarrow H_n(A) \xrightarrow{\hat{i}_*} H_n(X) \rightarrow H_n(X, A) \xrightarrow{\delta} H_{n-1}(A) \xrightarrow{\hat{i}_*} \dots$$

LES for the pair



### Reduced homology

$\tilde{H}_*X = H_*$  of augmented chain complex

$$\dots \rightarrow C_n(X) \rightarrow \dots \rightarrow C_0(X) \xrightarrow{\epsilon} \mathbb{Z} \rightarrow 0$$

augmentation  $\epsilon(\sum n_i \cdot p_i) = \sum n_i$

can view  $C_{-1}(X) = \mathbb{Z}$  (map  $\phi \rightarrow X$ ) where allow the empty simplex  $\emptyset$

For  $X \neq \emptyset$ ,  $\tilde{H}_*X = \text{Ker } H_*X \rightarrow H_*(pt)$  induced by  $X \rightarrow pt$

Example  $\tilde{H}_*(\mathbb{R}^n) \cong \tilde{H}_*(\mathbb{D}^n) \cong \tilde{H}_*(pt) = 0$

Check:  $H_*X = \tilde{H}_*X$   $*$   $\neq 0$ , and  $H_0X \cong \tilde{H}_0X \oplus \mathbb{Z}$  for  $X \neq \emptyset$

$f: X \rightarrow Y \Rightarrow f_*: \tilde{H}_*X \rightarrow \tilde{H}_*Y$

Lemma  $(X, A)$  pair  $\Rightarrow \exists$  LES


$$\dots \rightarrow \tilde{H}_n(A) \xrightarrow{\hat{i}_*} \tilde{H}_n(X) \rightarrow H_n(X, A) \rightarrow \tilde{H}_{n-1}(A) \xrightarrow{\hat{i}_*} \dots \rightarrow H_0(X, A) \rightarrow 0$$

Pf we augmented ch. cx. and  $\mathbb{Z}/\mathbb{Z} = 0 \square$

Cor  $H_*(X, pt) \cong \tilde{H}_*(X)$

Pf  $\tilde{H}_*(pt) = 0. \square$

Example LES:  $\tilde{H}_*(S^{n-1}) \rightarrow \tilde{H}_*(\mathbb{D}^n) = 0$



### Naturality of the LES for pairs

Def A map of pairs of spaces  $(X, A) \xrightarrow{f} (Y, B)$

means  $f: X \rightarrow Y$  and  $f(A) \subseteq B$ .

$$\text{Lemma } \dots \rightarrow H_*A \rightarrow H_*X \rightarrow H_*(X, A) \rightarrow H_{*-1}A \rightarrow \dots$$

$$\begin{matrix} f_* \downarrow & & f_* \downarrow & & f_* \downarrow \\ \dots \rightarrow & H_*B & \rightarrow & H_*Y & \rightarrow & H_*(Y, B) & \rightarrow & H_{*-1}B & \rightarrow & \dots \end{matrix}$$

and similarly for  $\tilde{H}_*$ .

Pf  $0 \rightarrow C_*A \rightarrow C_*X \rightarrow C_*X/C_*A \rightarrow 0 \Rightarrow$  claim follows by naturality of LES induced by SES of chain cxs.  $\square$

### 5. EXCISION THEOREM AND QUOTIENTS

$(X, A)$  pair

Def  $r: X \rightarrow X$  retraction onto  $A$  if  $\begin{cases} r(X) = A \\ r|_A = \text{id}_A \end{cases}$



Example  $X = S^2 \vee S^2 =$  two spheres glued at one point  $v$   
 $r: X \rightarrow A$  map second sphere to  $v$  (wedge sum)

Example In Pf of Brouwer fixed pt thm we built a retraction  $r$  by contradiction


Cor  $r$  retraction  $\Rightarrow r_*: H_*X \rightarrow H_*A$  surjective

$\text{incl}_*: H_*A \rightarrow H_*X$  injective

Pf  $A \xrightarrow{\text{incl}} X \xrightarrow{r} A$  now use  $H_*$  functorial  $\square$



Def  $r: X \rightarrow X$  deformation retraction onto  $A$  if  $\begin{cases} r \text{ retraction} \\ r \simeq \text{id}_X \text{ rel } A \end{cases}$



Example  $X = \text{Möbius strip}$   
 $A = \text{equator}$


Lemma def. retr.  $\Rightarrow A \xrightarrow{\text{incl}} X$  is a homotopy equivalence.

Pf  $A \xrightarrow{\text{incl}} X$   $\text{incl} \circ r = r \simeq \text{id}_X$ ,  $r \circ \text{incl} = r|_A = \text{id}_A$   $\square$

Example  $S^n \setminus \text{pt}$  def. retracts to  $D^n \cong \text{lower hemisphere}$ :  
 $\Rightarrow S^n \setminus \text{pt} \simeq D^n$  (move pt along great circles)  
 $\Rightarrow S^n \setminus \{2 \text{ points}\} \simeq D^n \setminus \text{pt} \simeq D^n \setminus 0 \simeq S^{n-1}$   
 $\Rightarrow S^n \setminus \{3 \text{ points}\} \simeq_{\text{def. retr.}} S^{n-1} \vee S^{n-1}$  (move out radially)


Good pairs and quotients

$(X, A)$  pair  
 • Quotient  $X/A = X/\sim \leftarrow \text{equiv. relation } x \sim y \Leftrightarrow \begin{cases} x=y \\ \text{or} \\ x, y \in A \end{cases}$   
 •  $(X, A)$  good pair if  $\begin{cases} A \neq \emptyset \\ A \text{ closed} \\ A \text{ deformation retract of nbhd } V \text{ of } A \end{cases}$



Example  $X = S^1 \vee S^1 = \bigcirc \geq V = \bigcirc \geq A = \bigcirc \cong S^1$   
 $X/A \cong \bigcirc \leftarrow \text{all points of } A \text{ are identified with the node}$

Non-example Topologist's sine curve  
 $\{(x, \sin \frac{1}{x}) : x \in (0, 1]\} \cup \{0\} \times [0, 1] \subseteq \mathbb{R}^2$   
Cultural Rmk connected, not path-connected, not locally connected, not locally path-connected



Smooth submanifold  $\subseteq$  Smooth manifold is a good pair (tubular neighbourhood theorem)


Cor  $(X, A)$  good  $\Rightarrow (X, A) \rightarrow (X/A, \text{pt})$  induces iso

$$H_*(X, A) \rightarrow H_*(X/A, \text{pt}) = \tilde{H}_*(X/A)$$

Pf good  $\Rightarrow \exists$  nbhd  $V$  of  $A$ , and  $A \xrightarrow{\cong} V$ .

LES for pair  
 $H_n(X, A) \xrightarrow{\text{quot.}} H_n(X, V) \xrightarrow{\text{quot.}} H_n(X/A, V/A) \xrightarrow{\text{quot.}} H_n(X/A, \text{pt})$   
 $H_n(X/A, A/A) \xrightarrow{\cong} H_n(X/A, V/A) \xrightarrow{\cong} H_n(X/A, \text{pt})$   
 since  $A=V, A \simeq V/A$   
 $\Rightarrow \text{identity}$

Hence all arrows are isos.  $\square$



Example  $D^n \geq S^{n-1}$  good:  $D^n / S^{n-1}$  quotient points of  $A=S^{n-1}$  identified

$\Rightarrow H_*(D^n, S^{n-1}) \cong \tilde{H}_*(D^n / S^{n-1}) \cong \tilde{H}_*(S^n)$

Excision theorem

$E \subseteq A \subseteq X$  subspaces  $\Rightarrow (X \setminus E, A \setminus E) \rightarrow (X, A)$  induces iso  
 with  $E \subseteq A^\circ$

$$H_*(X \setminus E, A \setminus E) \xrightarrow{\cong} H_*(X, A)$$

Proof Later.

Example  $X = S^1 \vee S^1 = \bigcirc \geq A = \bigcirc \geq E = \bigcirc \cong S^1$   
 $\Rightarrow H_*(X, A) \cong H_*(C, \cdot) \cong H_*(D^1, \partial D^1) \cong \tilde{H}_0(S^0) \cong \mathbb{Z}$   
 hpy invce  $\xrightarrow{\text{iso}}$  2 points

Rephrasing of Excision Thm

$X = A^\circ \cup B^\circ \Rightarrow H_*(X, A) \cong H_*(B, A \cap B)$   
 induced by inclusion  $(X, A) \leftarrow (B, A \cap B)$



Pf Take  $E = X \setminus B$  so  $X \setminus E = B$  and  $A \cap B = A \cap E$ .  $\square$

Idea why excision holds:  $C_*(A) + C_*(B) \rightarrow C_*(X)$  is a homotopy equivalence and  $C_*(A) \cap C_*(B) = C_*(A \cap B)$ . Idea: can subdivide chains in  $X$  many times, and small enough chains lie either in  $A$  or in  $B$  (or in both).

Recall we proved  $\widetilde{H}_*(\mathbb{D}^n, S^{n-1}) \cong \widetilde{H}_{*-1}(S^{n-1})$  (from LES &  $\widetilde{H}_*(\mathbb{D}^n) = 0$ )  
 $\Rightarrow \widetilde{H}_k(S^n) \cong \widetilde{H}_{k-1}(S^{n-1}) \cong \dots \cong \widetilde{H}_{k-n}(S^0) \cong \begin{cases} \mathbb{Z} & n=k \\ 0 & \text{else} \end{cases}$   
 inductively, using Example

Generator of  $H_n(S^n) \cong \widetilde{H}_n(\mathbb{D}^n/S^{n-1}) \cong H_n(\mathbb{D}^n, S^{n-1}) \cong \mathbb{Z}$

Observe  $\exists$  homeo  $e^n: \Delta^n \cong \mathbb{D}^n$  (homework) inducing  $\Delta$ -cx structure on  $S^{n-1}$ :  
 $\partial \Delta^n \cong \partial \mathbb{D}^n = S^{n-1}$   
 Example  $\mathbb{D}^2 \cong \begin{matrix} \nearrow v_2 \\ \Delta \\ \searrow v_1 \end{matrix} \xrightarrow{\partial} \begin{matrix} \triangleleft \\ \Delta^+ \\ \triangleright \end{matrix} \cong \mathbb{S}^1$   
 stretch cktly outwards from barycentre ( $\Delta^n$ )


Upshot ( $n \geq 2$ )  
 $H_n(\mathbb{D}^n, S^{n-1}) = \mathbb{Z} \cdot e^n$   
 $H_{n-1}(S^{n-1}) = \mathbb{Z} \cdot \partial e^n$   
 $\widetilde{H}_n(\mathbb{D}^n/S^{n-1}) = \mathbb{Z} \cdot [e^n]$   
 LES for  $n \geq 2$   
 by Cor  $[e^n]$  really lives in  $H_n(\mathbb{D}^n, S^{n-1}) \cong H_n(\mathbb{D}^n/S^{n-1}, S^{n-1})$   
 $H_n(\mathbb{D}^n, \partial \mathbb{D}^n) \cong H_n(\mathbb{D}^n/S^{n-1}, \partial \mathbb{D}^n)$   
 $\Delta_1 - \Delta_0 \rightarrow \Delta_1$

Exercise Recall another  $\Delta$ -cx structure on  $S^n$ :  
 $S^n = \Delta^n \cup \Delta^n / \text{glue along } \partial \Delta^n$   
 call this  $\Delta_1$  this  $\Delta_0$   
 then  $H_n S^n = \mathbb{Z} \cdot (\Delta_1 - \Delta_0)$  and  $H_n(S^n, \Delta_0) \cong H_n(\Delta_1, \partial \Delta_1)$   
 $\Delta_1 - \Delta_0 \rightarrow \Delta_1$


Another remark about orientations

Fact {homeos  $\Delta^n \rightarrow \mathbb{D}^n$ } has 2 path-components  
 Above we chose a path-component by constructing  $e^n$ .  
 If  $\tau$  is any reflection in  $\mathbb{R}^{n+1}$  then  $e^n \circ \tau$  is in the other path-component  
 $H_n(S^n) \cong H_n(\mathbb{D}^n, S^{n-1}) \xrightarrow{\cong} \mathbb{Z}$   
 e.g. swap 2 coordinates in  $\Delta^n$   
 $e^n \circ \tau \mapsto +1$   
 $e^n \circ \tau \mapsto -1$

We will see later in the course that this corresponds to a choice of orientation of  $\mathbb{D}^n$  and  $S^n$ .  
 Our choice is consistent with the inclusion  $\mathbb{D}^n \subseteq \mathbb{R}^n$  (with the positive (canonical) orientation of  $\mathbb{R}^n$ ) and the inclusion  $(\Delta^n \subseteq \mathbb{R}^{n+1}) \cong \{(t_1, \dots, t_n) \in \mathbb{R}^n : t_i \geq 0, \sum_{i=1}^n t_i \leq 1\}$   
 $(t_0, \dots, t_n) \mapsto (t_1, \dots, t_n)$   
 $t_i \geq 0, \sum t_i = 1$



Our choice is also consistent with the "normal first" Convention for orienting hyperplanes with a given choice of normal:  
 $\Delta^n \subseteq$  hyperplane  $\{(t_0, \dots, t_n) : \sum t_j = 1\} \subseteq \mathbb{R}^{n+1}$  normal  $(1, 1, \dots, 1)$  (so pointing to  $\infty$  in positive quadrant)



Compare  $\partial \Delta = +[e_0, e_1, e_2] - [e_0, \hat{e}_1, e_2] + [e_0, e_1, \hat{e}_2]$   
 This  $-[e_0, e_2]$  is not equal to singular chain  $[e_2, e_0]$  since they are different maps and we take free abelian group generated by maps.  
 But  $[e_0, e_2] + [e_2, e_0]$  is homologous to 0 (Homework).

Locality (or "small simplices theorem")

$\mathcal{U} = \{ \text{subspaces } U_i \subseteq X \}$  whose interior cover  $X$ :  
 $X = \bigcup U_i$

Def  $C_*^U(X) \subseteq C_*(X)$  subcomplex generated by  $n$ -simplices  $\sigma$  with  $\sigma(\Delta^n) \subseteq U_i$  some  $i$


Theorem  

$$H_* (C_*^U(X)) \cong H_* (C_*(X)) = H_* X$$

barycentre of  $[v_0, \dots, v_n]$  is  $\frac{1}{n+1}(v_0 + \dots + v_n)$

Sketch Pf ① Barycentric subdivision


Non-examinable



$\Rightarrow$  chain map  $S: C_*(X) \rightarrow C_*(X)$  and  $S(C_*^U) \subseteq C_*^U$   
 Construction of " $\sigma \circ S$ " is inductive:  
 On linear simplices (them for maps  $\sigma$  you restrict to...)  
 $S[e_0] = [e_0]$   
 $S[e_0, e_1] = [b_1, e_1] - [b_1, e_0]$  (geometrically  $\vec{e}_0 + \vec{e}_1 = \vec{b}_1$ )  
 $S[e_0, e_1, e_2] = [b_2, S\partial[e_0, e_1, e_2]]$   
 $= [b_2, S[e_1, e_2]] - [b_2, S[e_0, e_2]] + [b_2, S[e_0, e_1]]$   
 $= ([b_2, b_{12}, e_2] - [b_2, b_{02}, e_2]) - ([b_2, b_{02}, e_1] - [b_2, b_{02}, e_0]) + ([b_2, b_{01}, e_1] - [b_2, b_{01}, e_0])$   
 (geometrically:  $b_{02} + b_{01} = b_2$ )

so for  $\sigma: \Delta^2 \rightarrow X$  you take  $S(\sigma) = \sigma \circ S$   
 ② S chain hpic to id:  
 $T: C_n(X) \rightarrow C_{n+1}(X)$   
 $T(\sigma) = \Delta^n \times I \xrightarrow{\text{project}} \Delta^n \xrightarrow{\sigma} X$   
 exercise:  $\partial T + T\partial = S - \text{id}$

$$\left. \begin{aligned} T: C_n(X) &\rightarrow C_{n+1}(X) \\ T(\sigma) &= \Delta^n \times I \xrightarrow{\text{project}} \Delta^n \xrightarrow{\sigma} X \end{aligned} \right\} \Rightarrow S_*: H_*(X) \xrightarrow{\text{id}} H_*(X)$$



③  $\forall n$ -simplex  $\sigma: \Delta^n \rightarrow X$ , apply  $S(\cdot)$  enough times until  $\sigma$  (each  $n$ -simplex of subdivision)  $\subseteq U_i$  some  $i$   
 $\forall$  cycle  $c, \exists n$  s.t.  $S^n(c) \in C_*^U(X)$  cycle  
 $\Rightarrow H_*^U(c) \rightarrow H_*(X)$  surjective  
 $[S^n(c)] \rightarrow S_*^U[c] = [c]$  by ②

③  $\forall$  bdry  $c = \partial b, \exists n$  s.t.  $S^n(b) \in C_*^U(X)$   
 claim:  $H_*^U(c) \rightarrow H_*(X)$  injective  
 suppose  $[c] \mapsto 0$  then  $c = \partial b$  for  $b \in C_*^U(X)$   
 now  $S^n c, S^n b \in C_*^U(X)$  for large  $n$   
 $\Rightarrow \partial S^n b = S^n \partial b = S^n c$  in  $C_*^U(X)$   
 $\Rightarrow [c] \in S_*^U[c] = [S^n c] = [\partial S^n b] = 0$  in  $H_*^U(X) \checkmark \square$

Proof of excision theorem

Let  $B = X \setminus E$  use  $\mathcal{U} = \{A, B\}$  so  $C_*^U(X) = C_*(A) + C_*(B) \subseteq C_*(X)$

$$\Rightarrow C_*(X \setminus E) = C_*(B) + C_*(A \setminus E) \cong C_*(B) + C_*(A) \cap C_*(B) \cong C_*^U(X) / C_*(A)$$

$$\Rightarrow \text{Compare LES's: } H_*(X \setminus E, A \setminus E) \xrightarrow{\text{locality}} H_*(C_*^U X) \rightarrow H_*(C_*^U X / C_* A) \rightarrow H_{*-1}(A) \rightarrow H_{*-1}(C_*^U X)$$

$$\parallel \text{locality} \cong H_*(A) \rightarrow H_*(X) \rightarrow H_*(C_* X / C_* A) \rightarrow H_{*-1}(A) \rightarrow H_{*-1}(X)$$

(we are using naturality of LES's induced by SES's)


$$\parallel \text{iso by 5-lemma} \parallel H_*(X, A) \rightarrow H_{*-1}(A) \rightarrow H_{*-1}(X) \rightarrow H_{*-1}(X)$$

$$\parallel \text{locality} \parallel H_*(X, A)$$

2<sup>nd</sup> isomorphism theorem for groups

6. MAYER-VIETORIS SEQUENCE ← Key computational tool

$X = A \cup B$  s.t.  $X = A \cup B^o$   
 any subspaces



MV Theorem  $\exists$  LES:

$$\dots \rightarrow H_*(A \cap B) \xrightarrow{i_*} H_*(A) \oplus H_*(B) \rightarrow H_*(X) \rightarrow H_{*-1}(A \cap B) \xrightarrow{i_*} \dots$$

& same holds for  $\tilde{H}_*$  provided  $A \cap B \neq \emptyset$ .

Pf SES  $0 \rightarrow C_*(A \cap B) \rightarrow C_*(A) \oplus C_*(B) \rightarrow C_*(X) \rightarrow 0$   
 $\sigma \mapsto (\sigma, -\sigma)$   
 $(\alpha, \beta) \mapsto \alpha + \beta$

$\Rightarrow$  induces the LES (using locality  $H_*^u X \cong H_* X$ ). D

Exercise connecting map is  $\delta: H_*(X) \rightarrow H_{*-1}(A \cap B)$

$[\alpha + \beta] \mapsto [\partial\alpha] = -[\partial\beta]$



$\dots \rightarrow H_2(\text{pt}) \oplus H_2(\text{pt}) \rightarrow H_2(S^2) \rightarrow H_1(S^1) \rightarrow H_1(\text{pt}) \oplus H_1(\text{pt}) \rightarrow \dots$   
 $\parallel \quad \parallel \quad \parallel \quad \parallel \quad \parallel$   
 $0 \quad \text{hence } \mathbb{Z} \quad \mathbb{Z} \quad 0 \quad 0$

Exercise Compute  $H_* S^n = \begin{cases} \mathbb{Z} & * = 0 \\ \mathbb{Z} & * = n \\ 0 & \text{else} \end{cases}$  using MV

Example wedge sum of  $X, Y$  with basepoints  $x, y$   
 $X \vee Y = \frac{X \times Y}{x \sim y}$



$0 \rightarrow \mathbb{Z} \oplus \mathbb{Z} \xrightarrow{\cong} H_n(X) \rightarrow 0 \rightarrow \dots \rightarrow 0 \rightarrow \mathbb{Z} \rightarrow \mathbb{Z} \oplus \mathbb{Z} \rightarrow H_0(X) \rightarrow 0$   
 $\parallel \quad \parallel \quad \parallel$   
 $1 \mapsto (1, -1) \quad \parallel \quad \parallel$   
 $\mathbb{Z} \quad \mathbb{Z} \quad \mathbb{Z}$

Similarly  $[H_*(X \vee Y) \cong H_*(X) \oplus H_*(Y)]$  for  $* \neq 0$  if  $\exists$  contractible nbhds of  $x \in X, y \in Y$ .


Cones and suspensions

$CX = (X \times [0, 1]) / (x, s) \sim (x, t) \text{ iff equal or } s=t=1$   
 $\simeq \text{pt}$

$\Sigma X = (X \times [0, 1]) / (x, s) \sim (y, t) \text{ iff equal}$

Example  $CS^n \cong \mathbb{D}^{n+1}, \Sigma S^n \cong S^{n+1}$   
 or  $s=t=0$   
 or  $s=t=1$

Lemma  $\tilde{H}_*(\Sigma X) \cong \tilde{H}_{*-1}(X)$



now apply MV.  $\square$

Rmk  $\phi \neq A \subseteq X \Rightarrow \tilde{H}_*(X \setminus A) \stackrel{\text{LES}}{\cong} H_*(X \setminus A, CA) \stackrel{\text{exc.}}{\cong} H_*(X, A)$

Connected sum

$M, N$  connected  $\Rightarrow M \# N = (M \setminus \text{open } n\text{-ball}) \cup (N \setminus \text{open } n\text{-ball})$   
 identify  $\partial$  balls via a homeo



Fact compact connected orientable surfaces are homeo to  $S^2$  or  $T^2 \# \dots \# T^2$   
 $g = \# \text{ copies}$   
 and " " non-orientable ones:  $\mathbb{R}P^2 \# \dots \# \mathbb{R}P^2$ .  
 $g=0$  called  $\Sigma_g$

Exercise (Homework) For  $M, N$  compact connected

By MV,  $H_*(M \# N) \cong H_*(M) \oplus H_*(N)$  for  $1 \leq * \leq n-2$

If  $M$  or  $N$  orientable:  $* = n-1$  also works

If both non-orientable:  $* = n-1$  one of  $\mathbb{Z}/2$  summands becomes  $\mathbb{Z}$

Cor 1)  $\chi(M \# N) = \chi(M) + \chi(N) - \begin{cases} 2 & \text{for } n \text{ even} \\ 0 & \text{" " odd} \end{cases}$   
 2)  $H_*(\Sigma_g) \leftarrow \text{genus } g \cong \begin{cases} \mathbb{Z}^{2g} & * = 0 \\ \mathbb{Z} & * = 1 \\ \mathbb{Z} & * = 2 \end{cases} \cong \chi(S^n)$

$H_0(M \# N) \cong \mathbb{Z}$  since connected  
 fact:  $H_n(M \# N)$  is  $\mathbb{Z}$  or  $0$   
 if  $M, N$  both orientable (see later in course)

# 7. DEGREE OF MAPS OF SPHERES

$$f: S^n \rightarrow S^n \implies f_n: H_n S^n \xrightarrow{\cong} H_n S^n \xrightarrow{\cong} \mathbb{Z} \xrightarrow{1} \mathbb{Z} \xrightarrow{\text{deg}(f)} \mathbb{Z}$$

$\implies f_*: \tilde{H}_* S^n \rightarrow \tilde{H}_* S^n$  is  $\text{deg}(f) \cdot \text{id}$

Properties

- $\text{deg}(\text{id}) = 1$
- $\text{deg}(f \circ g) = \text{deg } f \cdot \text{deg } g$
- $f \simeq g \implies \text{deg } f = \text{deg } g$
- $f \simeq \text{const} \implies \text{deg } f = 0$
- $f$  homeomorphism  $\implies \text{deg } f = \pm 1$

Pf  $\text{id}_* = \text{id}$ ,  $(f \circ g)_* = f_* \circ g_*$ ,  $f \simeq g \implies f_* = g_*$ ,  $\text{const}_* = 0$ ,  $f$  homeo  $\implies f_n$  iso.  $\square$   
 since  $S^n \xrightarrow{\text{pt}} S^n$  factors so  $H_n S^n \xrightarrow{H_n(\text{pt})} H_n S^n$

Examples

- $S^n = \Delta^n \times 1 \cup \Delta^n \times 0 \leftarrow \text{call this } \Delta_0$   
 $(b, 1) \sim (b, 0)$  if  $b \in \partial \Delta$   
 recall  $H_n S^n = \mathbb{Z} \cdot (\Delta_1, -\Delta_0)$   
 reflection:  $r: S^n \rightarrow S^n$ ,  $r(x, t) = (x, 1-t)$   
 so  $\Delta_0 \leftrightarrow \Delta_1$  swapped by  $r$ , so  $r_*(\Delta_1, -\Delta_0) = -(\Delta_1, -\Delta_0)$   
 $\implies \text{deg}(r) = -1$

2) antipodal map  $-\text{id}: S^n \rightarrow S^n$  viewing  $S^n \subseteq \mathbb{R}^{n+1}$   
 $\implies \text{deg}(-\text{id}) = (-1)^{n+1}$


Pf  $-\text{id} = \begin{pmatrix} -1 & & \\ & \ddots & \\ & & -1 \end{pmatrix} \circ \dots \circ \begin{pmatrix} 1 & & \\ & \ddots & \\ & & -1 \end{pmatrix}$  composition of  $n+1$  reflections each homotopic to  $r$ .  $\square$

3)  $A \in O(n) \implies A: S^{n-1} \rightarrow S^{n-1} \implies \text{deg } A = \det A = \det A = \pm 1$   
Pf  $\text{fact}$   $SO(n)$  is path-connected so  $A \in SO(n)$  is  $\simeq \text{id}$  so  $\text{deg } A = \det A = +1$   
 The other path-component of  $O(n)$  is  $r \circ O(n)$  where  $r$  is any reflection.  $\square$

4)  $f$  not surjective  $\implies \text{deg } f = 0$   
Pf If  $y \notin \text{Im } f \implies H_n(S^n) \xrightarrow{f_*} H_n(S^n) \rightarrow H_n(\mathbb{R}^n) = 0$

# Application to vector fields on $S^n$

$v: S^n \rightarrow \mathbb{R}^{n+1}$  tangent vector field on  $S^n$   
 so  $v(x) \perp x$



Cor Hairy ball theorem  $\exists$  nowhere zero v.f. on  $S^n \iff n$  odd

(case  $n=2$ : "you cannot comb a ball of hair without creating a tuft")

Pf Suppose  $v(x) \neq 0 \forall x$

$\implies \text{hpy } F: S^1 \times [0, 1] \rightarrow S^n$

$$F(x, t) = \cos(\pi t)x + \sin(\pi t) \frac{v(x)}{\|v(x)\|}$$

$\implies F_0 = \text{id}, F_1 = -\text{id}$

$\implies 1 = \text{deg } F_0 = \text{deg } F_1 = (-1)^{n+1}$

$\implies n$  odd

For  $n$  odd  $\exists v(x) = (-x_2, x_1, \dots, -x_{2k}, x_{2k-1}) \square$

Cultural Remark Adams in 1962 proved using alg. topology:

(max # pointwise linearly independent vector fields on  $S^n = 2^b + 8a - 1$

where  $n+1 = 2^{4a+b}$ . (odd number),  $0 \leq b \leq 3, a, b \in \mathbb{N}, n \geq 1$ .  $\leftarrow$  get 0 if  $n$  even  $\implies \text{cor } \checkmark$

# Local degree

$f: S^n \rightarrow S^n$   
 $x \rightarrow y = f(x)$

$\star$  Suppose points  $x$  near  $x$  do not map to  $y$ :


$\exists$  nbhds  $x \in U, y \in V$  s.t.  $(U, U \setminus x) \xrightarrow{f} (V, V \setminus y)$

$\implies (f|_x)_*: H_n(U, U \setminus x) \xrightarrow{f_*} H_n(V, V \setminus y)$   
 $\xrightarrow{\cong} H_n(S^n, S^n \setminus x) \xrightarrow{\cong} H_n(S^n, S^n \setminus \text{pt}) \xrightarrow{\cong} \mathbb{Z} \xrightarrow{1} \mathbb{Z} \xrightarrow{\text{deg } f} \mathbb{Z}$

call this  $f|_x$   
local map at  $x$

Lemma  $f: S^n \rightarrow S^n, f^{-1}(y) = \{x_1, \dots, x_k\}$

$$\Rightarrow \deg f = \sum \deg_{x_i} f$$



map to each summand is exc. of  $S^n \setminus U_i$  so iso.

$\text{Rmk}$  (can use same  $V$  for all  $i$  by taking  $\tilde{V} = \cup V_i, \tilde{U}_i = f^{-1}(V) \cup U_i$ )  
 (the 2 squares commute: 1st: quotient is natural, 2nd: excision is natural)

Example  $p: \mathbb{C} \rightarrow \mathbb{C}$  polynomial  $a_n z^n + a_{n-1} z^{n-1} + \dots + a_0, a_i \in \mathbb{C}$   
 $\Rightarrow f: S^2 = \mathbb{CP}^1 \rightarrow \mathbb{CP}^1 = S^2$  (where view  $\mathbb{CP}^1 = \mathbb{C} \cup \infty \cong S^2$ )  
 $\infty \mapsto \infty$  stereographic projection

$\Rightarrow \text{hpfy } F(z, t) = a_n z^n + t(a_{n-1} z^{n-1} + \dots + a_0)$   
 $F_0 = a_n z^n$  and  $F_1 = f$

$\Rightarrow \text{deg } f = \deg(a_n z^n) \stackrel{\text{Lemma}}{=} \sum_{k=0}^{n-1} \deg a_k z^k \leftarrow \omega_k = e^{\frac{2\pi i k}{n}}$   
 $\stackrel{\text{Lemma}}{=} \sum_{k=0}^{n-1} \deg a_k z^k \stackrel{\text{Lemma}}{=} \sum_{k=0}^{n-1} \deg a_k z^k \leftarrow \omega_k = e^{\frac{2\pi i k}{n}}$

$\Rightarrow \text{deg } f = \deg(a_n z^n) \stackrel{\text{Lemma}}{=} \sum_{k=0}^{n-1} \deg a_k z^k \leftarrow \omega_k = e^{\frac{2\pi i k}{n}}$   
 $\stackrel{\text{Lemma}}{=} \sum_{k=0}^{n-1} \deg a_k z^k \leftarrow \omega_k = e^{\frac{2\pi i k}{n}}$

Cor (Fundamental Thm of Algebra)  $n \geq 1 \Rightarrow p$  has a root

PF  $p^{-1}(0) = \emptyset \Rightarrow f^{-1}(0) = \emptyset \Rightarrow \deg f = 0 \neq \mathbb{Z} \square$


Cultural Rmk For smooth  $f: S^n \rightarrow S^n$

$\deg f =$  (the number of preimages of a generic point.)  
 (i.e. almost any point works)

$\Rightarrow \text{deg } f = d = \# \text{ preimages of a point}$   
 except if pick North/South pole

## 8. CELLULAR HOMOLOGY

Def CW complex  $X$  is sequence  $\phi = X^{-1} \subseteq X^0 \subseteq X^1 \subseteq X^2 \subseteq \dots$  s.t.  $X^0$  is any set



$\Rightarrow X = \bigcup_{n \geq 0} X^n$  top space with weak topology:  $U \subseteq X$  open  $\Leftrightarrow U \cap X^n \subseteq X^n$  open  $\forall n$

Call  $X$  n-dimensional if  $X = X^n$  and this is the least such  $n$ .

Example  $S^n = (D^0 \sqcup D^1) / (D^0 \sim \partial D^1) \rightarrow S^1 = S^2$

Example  $X = \mathbb{RP}^2 = (D^0 \sqcup D^1) / (\text{wrap } \partial \text{ of } D^1 \text{ twice around } D^0) \rightarrow X^0 = \bullet$

$X^0 = \bullet = D^0$

$X^1 = (D^0 \sqcup D^1) / (\bullet \sim \varphi_1(x)), \partial D^1 = S^0 = \{0, 1\} \xrightarrow{\varphi_1} X^0 = \bullet$

$X^2 = (D^0 \sqcup D^1) / (\text{wrap } \partial \text{ of } D^1 \text{ twice around } D^0) \rightarrow X^1 = S^1$

$\partial D^2 = S^1 \xrightarrow{\varphi_2} S^1, \varphi_2(z) = z^2$

Fact If we homotope  $\varphi_\alpha$ , we get a homotopy equivalent space

Example If we use another degree 2 map  $\varphi_2$  above, get  $X \simeq \mathbb{RP}^2$ .

$X$  is partitioned as a set by interiors of n-cells

$$X^n = X^{n-1} \cup \bigcup_{\alpha \in I_n} e_\alpha^n$$

$$= \left( \bigcup_{\alpha \in I_{n-1}} e_\alpha^{n-1} \right) \cup \left( \bigcup_{\alpha \in I_n} e_\alpha^n \right) \cup \dots$$

$\leftarrow \text{Rmk}$   
 interior  $D^n = \mathbb{D}^n$   
 so  $e_\alpha^n = \mathbb{D}^n$

$e_\alpha^n = \text{Image}(D_\alpha^n \rightarrow X)$

Examples real projective space  $\mathbb{R}P^n = S^n / (\mathbb{Z}/2\text{-action by } \pm \text{id})$   
 $X^k = \mathbb{R}P^k$  inductively  
 $X^n = X^{n-1} \cup e^n$  with  $\varphi: S^{n-1} \rightarrow X^{n-1} = \mathbb{R}P^{n-1} = S^{n-1} / \pm \text{id}$   
 $x \mapsto [x] = [-x]$



Complex projective space

$\mathbb{C}P^n = (S^{2n+1} \subseteq \mathbb{C}^n) / (S^1\text{-action by } \lambda \cdot \text{Id})$   
 $X^0 = X^1 = pt = \mathbb{C}P^0$   
 $X^2 = X^3 = X^0 \cup e^2 = \mathbb{C}P^1$   $\varphi: S^1 \rightarrow pt \leftarrow \mathbb{C}P^1 \cong S^2$   
 $X^4 = X^5 = X^2 \cup e^4 = \mathbb{C}P^2$ ,  $\varphi: S^3 \rightarrow \mathbb{C}P^1 = S^3 / (S^1\text{-action})$   
 $X^{2n} = X^{2n-2} \cup e^{2n} = \mathbb{C}P^n$ ,  $\varphi: S^{2n-1} \rightarrow X^{2n-2} = \mathbb{C}P^{n-1}$   
 $x \mapsto [x] = [\lambda x]$ ,  $\forall \lambda \in S^1$

In coordinates:  $\mathbb{C}P^n = \{ [z_0 : \dots : z_n] : \text{not all } z_i \in \mathbb{C} \text{ are } 0 \}$  and  $[z] \sim [\lambda z]$ ,  $\forall \lambda \in \mathbb{C}^*$   
 Can rescale so that  $\sum |z_i|^2 = 1$  so  $z \in S^{2n-1}$  and left with rescaling by  $\lambda \in S^1 \subseteq \mathbb{C}^*$ .  
 $\mathbb{C}P^{n-1} \cong X^{n-2} = \{ [z_0 : \dots : z_{n-1} : 0] \} \subseteq \mathbb{C}P^n = X^n$  and  $\leftarrow$  notice this = 0 if  $w \in S^{2n-1} \cong \partial \mathbb{D}^{2n}$   
 $e^{2n}: \mathbb{D}^{2n} = \{ (w_0, \dots, w_{n-1}) : \sum |w_j|^2 \leq 1 \} \rightarrow X^n$  via  $[w_0 : \dots : w_{n-1} : \sqrt{1 - \sum |w_j|^2}]$

Observe: For  $X$  CW complex, for  $n \geq 1$ :  
 •  $(X^n, X^{n-1})$  is a good pair  $\leftarrow$  (since  $\exists$  nbhd of  $\partial \mathbb{D}^n$  that deformation retracts to  $\partial \mathbb{D}^n$ )

•  $X^n / X^{n-1} \cong \bigvee_{\alpha \in I_n} S^n$   
  
 $\leftarrow$   $S^n = \mathbb{D}^n / \partial \mathbb{D}^n$   
 $X^{n-1}$  identified to a point

Def Cellular complex for  $X$  a CW cx,

$$C_n^{CW}(X) = H_n(X^n, X^{n-1}) \cong H_n(X^n / X^{n-1})$$


= free abelian gp gen. by the  $n$ -cells  $e_\alpha^n$

since  $\Delta^n \cong \mathbb{D}^n \rightarrow (e^n \subseteq X^n) \rightarrow \mathbb{D}^n / \partial \mathbb{D}^n = S^n$  generate  
 as usual we use the standard orientations of  $\Delta^n, \mathbb{D}^n, S^n$ .

Will build cellular differential  $d$ , prove  $d \circ d = 0$ ,

$$\Rightarrow \text{get } H_*^{CW}(X) = H_*(C_*^{CW}(X), d)$$

$d e_\alpha^n = \sum_{\beta \in I_{n-1}} d_{\alpha\beta}^n \cdot e_\beta^{n-1}$   
 now describe the coefficients  $d_{\alpha\beta}^n \in \mathbb{Z}$  and why that is a finite sum.



Recall LES  $H_*(X^n, X^{n-1}) \rightarrow H_*(X^{n-1})$   
 $\leftarrow$   $S^n \xrightarrow{[-1]} S^n \xrightarrow{q_*} H_*(X^n, X^{n-1})$   
 $\leftarrow$   $\mathbb{D}^n \rightarrow \mathcal{U}_\alpha(S^{n-1}) \rightarrow \sum d_{\alpha\beta} e_\beta^{n-1}$   
 $\leftarrow$   $\mathbb{D}_\alpha^n \rightarrow e_\alpha^n \subseteq X^n$   
 $\leftarrow$   $\partial \mathbb{D}_\alpha^n \subseteq X^{n-1}$

here it is important that we chose identifications  $\Delta^n \cong \mathbb{D}^n$ ,  $S^n \cong \mathbb{D}^n / \partial \mathbb{D}^n$  compatibly with orientations.  
 Quotient by  $V_{\mathbb{R}^{n-1} \times \mathbb{R}}$   
 Therefore: 
$$d_{\alpha\beta}^n = \text{deg} \left( \begin{array}{ccc} S^{n-1} \xrightarrow{q_\alpha} X^{n-1} \xrightarrow{q} X^{n-1} / X^{n-2} \cong V S^{n-1} & \rightarrow & S^{n-1} \\ \parallel \mathbb{D}_\alpha^n & & \parallel \mathbb{D}_\beta^{n-1} / \partial \mathbb{D}_\beta^{n-1} \end{array} \right)$$

Rmk Only finitely many  $d_{\alpha\beta}^n \neq 0$  (for fixed  $\alpha$ ) because  $\mathcal{U}_\alpha, q$  are continuous and  $S^{n-1}$  compact, so get a compact image in  $V S^{n-1}$ , therefore cannot surject onto  $\infty$  many  $S_\beta^{n-1}$ .

Lemma  $d \circ d = 0$   
 $\leftarrow$  recall if don't surject then deg = 0



pf  $d_n = q_{n-1}^{n-1} \circ S_n^n$   
 $d_{n-1} \circ d_n = q_{n-2}^{n-2} \circ S_{n-1}^{n-1} \circ q_{n-1}^{n-1} \circ S_n^n = 0$  by LES

Cor  $\text{rank } H_n^{CW}(X) \leq \# \text{ n-cells}$

pf  $\# \text{ n-cells} = \text{rank } C_n^{CW}(X) \geq \text{rank } \ker d_n^{CW} \geq \text{rank } H_n^{CW}(X) \square$

Example  $I \times I$   $I = [0,1]$   $D^1 = [-1,1]$

arrows here tell us how we map  $[-1,1] \rightarrow$  edge (so orientation)



$e^2 : D^2 \cong \square \rightarrow X^1$

$\partial e^2 : S^1 \cong \square \rightarrow X^1/X^0 =$

$\Rightarrow \partial e^2 = +e^1 x 0 + 1x e^1 - e^1 x 1 - 0x e^1$   
 $(= (\partial e^1) x e^1 - e^1 x (\partial e^1) \leftarrow$  we come back to this later)

Example  $RP^n$  recall: 1 cell in each dim,  $\varphi: S^k \rightarrow X^k = RP^k$

$S^{k-1} \xrightarrow{\varphi} X^{k-1}/X^{k-2} = RP^{k-1}/RP^{k-2} \xrightarrow{\partial/\partial \Delta} S^{k-1}$   
 $\Delta_1 \xrightarrow{\varphi} \Delta_2 \xrightarrow{-id(\Delta_1)} -id(\Delta_1)$   
 $\deg = +1$   
 $\deg = (-1)^k$

$\Rightarrow d_{\alpha\beta} = \begin{cases} 2 & k \text{ even} \\ 0 & k \text{ odd} \end{cases}$


$C_*^{CW}(RP^n) \rightarrow \mathbb{Z} \xrightarrow{k=n} \mathbb{Z} \xrightarrow{k=n-1} \mathbb{Z} \xrightarrow{k=n-2} \mathbb{Z} \xrightarrow{k=n-3} \mathbb{Z} \rightarrow 0$

$H_*^{CW}(RP^n) \cong \begin{cases} \mathbb{Z} & * = 0, \text{ also } * = n \text{ if odd} \\ \mathbb{Z}/2 & * = 1, 3, 5, \dots < n \\ 0 & \text{else} \end{cases}$

Example  $S^n$ :  $C_*^{CW}(S^n): n > 2: 0 \rightarrow \mathbb{Z} \cdot D^0 \xrightarrow{0} 0 \rightarrow \dots \rightarrow 0 \rightarrow \mathbb{Z} \cdot D^n \rightarrow 0$

$n=1: 0 \rightarrow \mathbb{Z} \cdot D^1 \xrightarrow{0} \mathbb{Z} \cdot D^0 \rightarrow 0$   
 $\# = 0, n$   
 $\# = 0, n$  else

Example  $\Sigma_g =$  genus  $g$  surface  
 $H_1(S^1, \mathbb{Z}) \cong \mathbb{Z} \xrightarrow{H_0(p,t)} H_0(p,t) \xrightarrow{\partial} H_0(p,t) \cong \mathbb{Z}$   
 $(\Delta^1 \cong [0,1] \rightarrow S^1)$   
 if you work with degrees, need to remember orientations:  $\partial D^1 \cong \partial[0,1] = [1] - [0] \rightarrow$  point  
 so degree =  $+1 - 1 = 0$



$0 \rightarrow \mathbb{Z} \xrightarrow{\partial} \mathbb{Z} \xrightarrow{\partial} \mathbb{Z} \xrightarrow{\partial} 0$   
 $\mathbb{Z} \cdot D \cong \mathbb{Z} \cdot D < a_1, b_1, a_2, b_2 > \mathbb{Z} \cdot v$

$D \mapsto -a_1 - b_1 + a_1 + b_1 + \dots - a_g - b_g + a_g + b_g = 0$

$H_k(\Sigma_g) \cong \begin{cases} \mathbb{Z} & * = 0, 2 \\ \mathbb{Z}^{2g} & * = 1 \\ 0 & \text{else} \end{cases}$

Lemma  $X \Delta$ -cx structure  $\Rightarrow$  induces CW-cx structure on  $X$  and  $(C_*^{CW}(X), d^{CW}) \cong (C_*^{\Delta}(X), d^{\Delta})$

$\Rightarrow H_*^{CW}(X) \cong H_*^{\Delta}(X)$

Pf  $X^n = \cup$  n-simplices of  $X$  and degrees are  $\pm 1$  depending on orient

Example  $X =$  triangle  $= \Delta^2$   
 $\Rightarrow$  so can identify  $d^{CW}$  and  $d^{\Delta}$ .  $\square$



$d_{\alpha\beta_2} = d_{\alpha\beta_0} = +1, d_{\alpha\beta_1} = -1$   
 $\Rightarrow d^{CW} \alpha = \beta_0 - \beta_1 + \beta_2 \checkmark \square$

signs: compare edge orientation with anticlockwise orientation of  $\partial D$



Theorem  $X$  CW  $cx$  (or  $\Delta$ - $cx$ )  $\implies$   $H_*^{CW}(X) \cong H_*(X)$

$\implies H_*^{\Delta}, H_*^{CW}$  independent of choice of CW- $cx$ / $\Delta$ - $cx$  structure.

Pf ①  $H_*(X^n, X^{n-1}) \cong \tilde{H}_*(X^n/X^{n-1}) \cong \tilde{H}_*(V_{\alpha} S^n) \cong \bigoplus_{\alpha} \tilde{H}_* S^n$   
 $= 0 \iff * \neq n$  lives in  $\mathbb{Z}$

LES for  $(X^n, X^{n-1}) \implies H_*(X^{n-1}) \rightarrow H_*(X^n) \rightarrow H_*(X^n/X^{n-1}) \rightarrow 0$  iso for  $* \leq n-1$

② for  $* < n$ :  $H_*(X^n) \cong H_*(X^{n+1}) \cong H_*(X^{n+2}) \cong \dots \cong H_*(X)$

by compactness each sing. chain lands in  $X^N$  some  $N$

③ for  $* > n$ :  $H_*(X^n) \cong H_*(X^{n-1}) \cong H_*(X^{n-2}) \cong \dots \cong H_*(X^{n-1}) = 0$

④ LES:  $\dots \rightarrow H_n(X^{n-1}) \rightarrow H_n(X^n) \xrightarrow{q_n} H_n(X^n/X^{n-1}) \rightarrow \dots$

$\implies q_n$  injective  $\forall n$

⑤ LES:  $\dots \xrightarrow{q_{n+1}} H_{n+1}(X^{n+1}, X^n) \xrightarrow{\delta_{n+1}} H_n(X^n) \rightarrow H_n(X^{n+1}) \rightarrow 0$

UPSHOT  $H_n(X) \cong H_n(X^{n+1})$

$H_n(X^n) / \text{im } \delta_{n+1} \cong (q_n^n H_n(X^n)) / \text{im } q_n^n \circ \delta_{n+1} \cong H_n^{CW}(X)$

exactness LES  $\text{im } q_n^n \xrightarrow{\cong} \text{Ker } \delta_n^n = \text{Ker } q_{n-1}^{n-1} \circ \delta_n^n \xrightarrow{\cong} \text{Ker } q_n^{n-1} \circ \delta_n^n$

Rmk by ①  $H_k$  not affected if attach  $(k+2)$ -cells or higher

by ② Inclusion  $X^n \rightarrow X$  induces iso  $H_*(X^n) \rightarrow H_*(X)$  for  $* < n$

Cor  $X$   $n$ -dimensional cell  $cx \implies H_*(X) = 0$  for  $* > n$

Axioms for homology: Eilenberg-Steenrod axioms

It was no accident that  $H_*^{\Delta}, H_*^{CW}, H_*^*$  all agreed.

Def A generalised homology theory (GHT)

is a functor  $F$ : Top Pairs = (Category of pairs of spaces and maps of pairs)  $\rightarrow$  Graded Abelian Gps

with a natural transformation  $\delta : F_*(X, A) \rightarrow F_{*-1}(X, \emptyset)$  satisfying:

1) homology invariance:  $f \simeq g \implies F(f) = F(g)$  abbreviated:  $F_{*-1}(X)$

2) exactness:  $\exists$  LES  $\dots \rightarrow F_*(A) \xrightarrow{f} F_*(X) \xrightarrow{F(f)} F_{*-1}(X, A) \xrightarrow{\delta} F_{*-1}(A) \rightarrow \dots$

3) additivity:  $(X, A) = \sqcup (X_i, A_i)$ ,  $\text{incl}_i : (X_i, A_i) \rightarrow (X, A)$   
 $F(\text{incl}_i : A \rightarrow X) \quad F(\text{incl}_i : X_i, \emptyset) \rightarrow (X, A)$

then  $\Sigma F(\text{incl}_i) : \bigoplus F(X_i, A_i) \cong F(X, A)$

4) excision:  $\bar{E} \subseteq A^{\circ} \subseteq X \implies F(X \setminus E, A \setminus E) \xrightarrow{\cong} F(X, A) \xrightarrow{\cong} F(X, A)$

Remark (4)  $\iff X = A^{\circ} \cup B^{\circ}$ ,  $\text{incl} : (B, A \cap B) \rightarrow (X, A)$   
 $F(\text{incl}) : F(B, A \cap B) \cong F(X, A)$

Pf  $B = X \setminus E$ ,  $E = X \setminus B$  noticing that  $(X \setminus E)^{\circ} \cup A^{\circ} = X$   
 $E = A \setminus B$  noticing that  $\bar{E} \subseteq \bar{A} \subseteq A^{\circ} \setminus B^{\circ} \subseteq A^{\circ}$ .  $\square$   
 $X = A^{\circ} \cup B^{\circ}$   
 $\text{so } \partial B \subseteq A^{\circ}$

Rmk In (3), the topology on the disjoint union  $\sqcup (X_i, A_i)$  is defined by:  $U \subseteq \sqcup (X_i, A_i)$  open  $\iff U \cap X_i \subseteq X_i$  open  $\forall i$

FACT Theorem

a)  $(F, \delta_F), (G, \delta_G)$  GHTs,  $\alpha : F \rightarrow G$  a natural transformation commuting with  $\delta_F, \delta_G$  such that  $\alpha_{\text{point}} : F(\text{point}) \rightarrow G(\text{point})$  is an iso, then  $\alpha$  is an iso.

b) If  $(F, \delta_F)$  GHT satisfies (5) dimension:  $F_*(\text{point}) = \begin{cases} \mathbb{Z} & * = 0 \\ 0 & \text{else} \end{cases}$

Then  $\exists$  natural iso  $F \cong H_*$  (such an  $F$  is called a homology theory)

Rmk In (b) if require  $F_0(\text{point}) = \mathbb{G}$  an abelian group (instead of  $\mathbb{Z}$ )  $\implies F(X, A) \cong H_*(X, A; \mathbb{G})$  = (homology with coefficients in  $\mathbb{G}$ )  $\leftarrow$  later in course

## 9. COHOMOLOGY

$(C_*, \partial_*)$  chain complex s.t.  $C_n$  free  $\mathbb{Z}$ -module  $\leftarrow C_* \cong \bigoplus_{\mathbb{Z}}$

Def **n-cochains**

$$C^n = \text{Hom}(C_n, \mathbb{Z})$$

boundary map

$$\partial^n : C^n \rightarrow C^{n+1}$$

(this is the dual of  $\partial$ )

$$\partial^n(\phi) = \phi \circ \partial_{n+1}$$

Notice  $\partial^*$  is degree +1 map (not -1)

$$H^m(C_*, \partial_*) = \text{Ker} \frac{\partial^m}{\text{Im } \partial^{m-1}} \leftarrow \begin{matrix} \text{cocycles} \\ \text{coboundaries} \end{matrix}$$

Remark If we use negative grading,  $(C^{-*}, \partial^{-*})$  is a chain complex with homology so many results from  $H_*$  carry over to  $H^*$ . So abusively write

$$H^*(C_*, \partial_*) = H_*(\text{Hom}(C_*, \mathbb{Z}), \partial^*)$$

Warning A cochain  $\varphi \in C^*$  takes values  $\varphi(c) \in \mathbb{Z}$  on chains  $c \in C_*$ . However the cohomology class  $\alpha = [\varphi] \in H^*$  does not have a well-defined value on  $c$ :  $[\varphi] = [\varphi + \partial^*(\psi)]$  and  $(\varphi + \partial^*(\psi))(c) = \varphi(c) + \psi(\partial_* c)$ . If  $c$  is a cycle, so  $\partial_* c = 0$  then  $\alpha(c) = \varphi(c)$  is well-defined, so  $\exists$  pairing  $H^* \times H_* \rightarrow \mathbb{Z}$

Remarks about dualisation

$\text{Hom}(\mathbb{Z}^n, \mathbb{Z}) \cong \mathbb{Z}^n$  generated by projection maps  $\pi_i(x_1, \dots, x_n) = x_i$

$$\alpha : \mathbb{Z}^n \rightarrow \mathbb{Z}^m \Rightarrow \text{Hom}(\mathbb{Z}^n, \mathbb{Z}) \xleftarrow{\text{dual}} \text{Hom}(\mathbb{Z}^m, \mathbb{Z}) \quad \alpha^* \phi = \phi \circ \alpha$$

$$\begin{matrix} \mathbb{Z}^n & \xrightarrow{\parallel} & \mathbb{Z}^m \\ \uparrow & \text{transpose (A)} & \downarrow \\ \text{m} \times \text{n matrix} & & \end{matrix}$$

Def  $X$  space  $\Rightarrow$  **singular cohomology**  $H^*(X) = H^*(C^*(X), \partial^*)$

similarly define  $H_{\Delta}^*$ ,  $H_{CW}^*$

Example  $\mathbb{RP}^3 : C_*^{CW}(\mathbb{RP}^3) : 0 \rightarrow \mathbb{Z} \xrightarrow{\partial} \mathbb{Z} \xrightarrow{\partial} \mathbb{Z} \xrightarrow{\partial} \mathbb{Z} \rightarrow 0$

dualise :  $C_*^{CW}(\mathbb{RP}^3) : 0 \leftarrow \mathbb{Z} \xleftarrow{\partial} \mathbb{Z} \xleftarrow{\partial} \mathbb{Z} \xleftarrow{\partial} \mathbb{Z} \rightarrow 0$

$$H^*(\mathbb{RP}^3) \cong H_{CW}^*(\mathbb{RP}^3) = \begin{cases} \mathbb{Z} & * = 0, 3 \\ \mathbb{Z}/2 & * = 2 \\ 0 & \text{else} \end{cases} \leftarrow \text{notice } H_1(\mathbb{RP}^3) \cong \mathbb{Z}/2 \text{ has moved to grading 2.}$$

## Functionality

$$f : X \rightarrow Y \Rightarrow f_* : C_* X \rightarrow C_* Y \quad \leftarrow \text{called pull-back}$$

$$\Rightarrow C^* X \xleftarrow{f^*} C^* Y \text{ is dual: } \boxed{f^* \phi = \phi \circ f_*}$$

Lemma  $f^*$  is a **cochain map** (meaning  $\partial^* \circ f^* = f^* \circ \partial^*$ )

$$\Rightarrow \boxed{f^* : H^* Y \rightarrow H^* X}$$

Pf  $(\partial^* \circ f^*)(\phi) = \partial^*(\phi \circ f_*)$

$$= (\phi \circ f_*) \circ \partial$$

$$= (\phi \circ \partial) \circ f_* \quad \text{as } f_* \text{ chain map}$$

$$= f_* \circ (\phi \circ \partial)$$

$$= f_* \circ (\partial^* \phi)$$

$$= (f_* \circ \partial^*)(\phi)$$

Properties  $\cdot \text{id}^* = \text{id}$

$\cdot (f \circ g)^* = g^* \circ f^*$  notice order!

$$\Rightarrow \boxed{H^* : \text{Top} \rightarrow \text{Graded AbGps}}$$

Contravariant functor

Exercise  $H^0(X) = \prod_{\text{Path } X} \mathbb{Z}$  where  $\text{Path } X = \{\text{path-components of } X\}$

## Homotopy invariance

Lemma  $f_*, g_* : C_* \xrightarrow{\text{free}} C_*$  chain hpic  $\Rightarrow f^* = g^* : H^* \tilde{C} \rightarrow H^* \tilde{C}$

Pf  $f_* - g_* = \tilde{\partial} \circ h + h \circ \partial$  same  $h : C_* \rightarrow \tilde{C}_*[1]$

$f^* - g^* = h^* \circ \tilde{\partial}^* + \partial^* \circ h^*$  for dual  $h^* : \tilde{C}^* \rightarrow C^*[-1]$

(notice degree -1, not +1)  $\square$

Def  $h^*$  called **cochain homotopy**

Cor  $f \simeq g : X \rightarrow Y \Rightarrow f^* = g^* : H^* Y \rightarrow H^* X \quad \square$

### Algebra: dual of SES

**Lemma**  $0 \rightarrow A \xrightarrow{i} B \xrightarrow{j} C \rightarrow 0$  exact,  $A, B, C$  free  
 $\Rightarrow 0 \leftarrow A^* \xleftarrow{i^*} B^* \xleftarrow{j^*} C^* \leftarrow 0$  exact  
**Pf**  $C$  free  $\Rightarrow \exists$  splitting  $B \xrightarrow{j} C \xleftarrow{s} B$   $j \circ s = \text{id}$   
 Pick preimages  $b_i$  for basis  $e_i$  of  $C$ , then  $s(e_i) = b_i$   
 $\Rightarrow A \oplus C \xrightarrow{i \oplus s} B$

**dual**  
 $\Rightarrow A^* \oplus C^* \xrightarrow{i^* \oplus s^*} B^*$  and  $s^* \circ j^* = \text{id}$   
 $\Rightarrow 0 \leftarrow A^* \xleftarrow{i^*} B^* \xleftarrow{j^*} C^* \leftarrow 0$

where  $0 = (j \circ i)^* = i^* \circ j^*$  so  $\text{Im } j^* \subseteq \text{Ker } i^*$   
 prove  $\supseteq$ :  $i^* b = 0 \Rightarrow b - j^* s^* b \in \text{Ker } i^* \cap \text{Ker } s^* = \{0\}$   
 $\Rightarrow b = j^* s^* b \in \text{Im } j^*$   
 $\Rightarrow \text{Ker } i^* = \text{Im } j^* \quad \square$

**Relative cohomology**  
 $H^*(X, A) = H^*(\text{Hom}(C_*(X, A), \mathbb{Z}))$   
 recall  $C_*(X, A) = C_*(X)/C_*(A)$  and  $\text{homs } C_*(X)/C_*(A) \rightarrow \mathbb{Z}$  correspond precisely to  $\text{homs } C_*(X) \rightarrow \mathbb{Z}$  which vanish on  $C_*(A)$ . So relative cocycles are cocycles on  $X$  which vanish on chains in  $A$ .

**Excision, LES, Mayer-Vietoris**  
 By previous Lemma get dual results:

**Excision**  $\bar{E} \subseteq A^0 \subseteq X \Rightarrow H^*(X \setminus E, A) \xleftarrow{\cong} H^*(X, A)$   
**LES for pair**  $(X, A) \quad \dots \xleftarrow{q^{[+1]}} H^{*+1}(X, A) \xleftarrow{\delta} H^*(A) \xleftarrow{i^*} H^*(X) \xleftarrow{q^*} H^*(X, A) \leftarrow \dots$

**M.V.**  $X = A \cup B \Rightarrow \dots \leftarrow H^{*+1}(X) \leftarrow H^*(A \cap B) \leftarrow H^*(A) \oplus H^*(B) \leftarrow H^*(X) \leftarrow \dots$   
 where  $A \cap B \xrightarrow{i_A^*} A \xrightarrow{j_A} X$  and  $A \cap B \xrightarrow{i_B^*} B \xrightarrow{j_B} X$  are the obvious maps

**Axioms for cohomology** These are analogous to the axioms for homology except we reverse all arrows, and we change axiom (3):  $\prod$  instead of  $\oplus$   
**additivity**:  $(X, A) = \sqcup (X_i, A_i)$ ,  $\text{incl}_i: (X_i, A_i) \rightarrow (X, A)$

then  $\prod F(\text{incl}_i): \prod F(X_i, A_i) \xleftarrow{\cong} F(X, A)$

### 10. CUP PRODUCT

**Theorem**  $H^*(X)$  is **unital graded-commutative** ring via  $\cup: H^k(X) \times H^l(X) \rightarrow H^{k+l}(X)$  determined by

$$\cup: C^k(X) \times C^l(X) \rightarrow C^{k+l}(X)$$

$$(\phi \cup \psi)(\sigma) = \phi(\sigma|_{[e_0, \dots, e_k]}) \cdot \psi(\sigma|_{[e_{k+1}, \dots, e_{k+l}]})$$

- ①  $1 \in C^0(X)$  constant function  $\Rightarrow 1 \cup \phi = \phi \cup 1 = \phi$
- ②  $\phi \cup \psi = (-1)^{\text{deg } \phi \cdot \text{deg } \psi} \psi \cup \phi$

**Useful trick** If  $X$  is  $\Delta$ -cx,  $C_*^\Delta(X) \xrightarrow{\text{inclusion}} C_*(X)$ , so  $C_*^\Delta(X) \xleftarrow{\text{restriction}} C^*(X)$  and can define cup product on  $C_*^\Delta(X)$  so that:

$$H_*^\Delta(X) \times H_*^\Delta(X) \xrightarrow{\cong} H_*^\Delta(X) \xleftarrow{\text{at chain level}} H_*(X) \times H_*(X) \xrightarrow{\cong} H_*(X)$$

$$(\phi \cup \psi)([v_0, \dots, v_n]) = \phi([v_0, \dots, v_k]) \cdot \psi([v_{k+1}, \dots, v_n])$$

So you can compute cup products on  $H^*(X)$  by picking simplicial cocycle representatives: so define values on the simplicial chains defining the  $\Delta$ -cx structure, and use

**Proof of Theorem**  
 $\partial^*(\phi \cup \psi)(\sigma) = (\phi \cup \psi)(\partial\sigma)$   
 $= (\phi \cup \psi) \sum (-1)^i \sigma|_{[e_0, \dots, \hat{e}_i, \dots, e_n]}$   
 $= \sum_{i \leq k} (-1)^i \phi(\sigma|_{[e_0, \dots, \hat{e}_i, \dots, e_{k+1}]}) \cdot \psi(\sigma|_{[e_{k+1}, \dots, e_n]})$   
 $+ \sum_{i > k} (-1)^i \phi(\sigma|_{[e_0, \dots, e_k]}) \cdot \psi(\sigma|_{[e_k, \dots, \hat{e}_i, \dots, e_n]}) \cdot (-1)^{i-k} \underbrace{(-1)^{k-i}}_1$   
 $= ((\partial^* \phi) \cup \psi)(\sigma) + (-1)^k \phi \cup \partial^* \psi$   
 induces  $[\phi] \cup [\psi] = [\partial^* \phi \cup \psi] + (-1)^k [\phi \cup \partial^* \psi]$

well-defined:  $\bullet$  cycles  $\rightarrow$  cycle:  $\partial(\phi \cup \psi) = (\partial\phi) \cup \psi \pm \phi \cup (\partial\psi) = 0$   
 $\bullet [\phi] = [\phi + \partial\alpha] \cup \psi = \partial(\alpha \cup \psi) + \phi \cup \psi = 0$  (using  $\partial\psi = 0$ )  
 $\bullet$  Similarly  $[\phi] \cup [\partial\beta] = 0$  (using  $\partial\psi = 0$ )

bilinear, associative, distributive: true at chain level

unital:  $(\partial 1)(\sigma) = 1(\sigma|_{[e_0]}) - 1(\sigma|_{[e_1]}) = 1 - 1 = 0$

$(1 \cup \psi)(\sigma) = 1(\sigma|_{[e_0]}) + \psi(\sigma|_{[e_0, \dots, e_n]}) = \psi(\sigma)$  ( $\psi|_{\partial} = \phi$  similar)

graded-comm. sketch proof:  $\leftarrow$  **non-examinable**

Let  $r: C_n(X) \rightarrow C_n(X)$ ,  $r(\sigma) = \varepsilon_n \bar{\sigma}$  where:  $\varepsilon_n = (-1)^{\frac{n(n+1)}{2}}$

and  $\bar{\sigma} |_{[v_0, \dots, v_n]} = \sigma |_{[v_n, \dots, v_0]}$   $\leftarrow$  reverse order of vertices:

(idea: each transposition is a reflection in hyperplane, so reverses orientation of simplex, so insert  $\varepsilon_n$  to compensate)

one checks:  $\bullet$   $r^*$  chain map

$\bullet$   $\frac{r^* \psi \cup r^* \psi}{\varepsilon_k \varepsilon_l} = r^*(\psi \cup \psi)$

$\leftarrow$  differ by  $(-1)^{kl}$

$\bullet$   $r \simeq \text{id}$  so can drop  $r^* = \text{id}$  on cohomology

$(r - \text{id}) = \partial \partial + \partial \partial$  with  $v_i, w_i$  like for prism operator

$(P\sigma) = \sum (-1)^i \varepsilon_{n-i} (\sigma \circ \pi_i) |_{[v_0, \dots, v_i, \dots, v_n, \dots, v_n, \dots, v_i]}$   $\square$

Naturality of cup product

Lemma  $f: X \rightarrow Y \implies f^*: H^* Y \rightarrow H^* X$  hom of unital rings

Pf  $f^*(\psi \cup \psi)(\sigma) = (\psi \cup \psi)(f_* \sigma)$

$= \psi(f_* \sigma|_{[e_0, \dots, e_k]}) \cup \psi(f_* \sigma|_{[e_{k+1}, \dots, e_n]})$

$= ((\psi \circ f_*) \cup (\psi \circ f_*))(\sigma)$

$= (f^* \psi \cup f^* \psi)(\sigma)$

unital:  $f^*(1) = 1 \circ f_* = 1 \quad \square$

UPSHOT  $H^*: \text{Top} \rightarrow \left\{ \begin{array}{l} \text{Graded-commutative unital rings} \\ \text{with graded unital ring homs} \end{array} \right\}$  contravariant functor.


Warning An (iso)morphism  $H^*(Y) \rightarrow H^*(X)$  of groups will also preserve the ring structure if  $f^*$  is induced by a map of spaces  $X \rightarrow Y$  (by above Lemma).

$\implies$  Cor The excision theorem iso on cohomology is an iso of rings. However the connecting hom in M.V. or LES cannot possibly be a ring hom since it drops gradings by 1 ( $\implies \delta(a \cup b)$  and  $\delta(a) \cup \delta(b)$  have different gradings!)

Example  $H^1(T^2) \times H^1(T^2) \xrightarrow{\cup} H^2(T^2) \rightarrow \mathbb{Z}$  with matrix  $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$

Pf recall:

*	$H_*(T^2)$	$H^*(T^2)$
0	$\mathbb{Z}$ -pt	$\mathbb{Z} \cdot 1$
1	$\mathbb{Z} \oplus \mathbb{Z} b$	$\mathbb{Z} a + \mathbb{Z} b^*$
2	$\mathbb{Z} \cdot D$	$\mathbb{Z} \cdot D^*$



$T^2$




$1, a^*, b^*, D^*$  are dual basis in  $H^*$

Identify  $H^*(T^2) \cong H^*(\mathbb{Z}^2)$  so at chain level:


$a^*: C_1^{CW}(X) \rightarrow \mathbb{Z} \quad b^*: C_2^{CW}(X) \rightarrow \mathbb{Z}$   
 $a \mapsto 1 \quad b \mapsto 0$   
 $a^* \mapsto 0 \quad b^* \mapsto 1$

$\implies b^*(c) = \#$  a intersects c counted with orientation signs




$a^*(c) = -\#$  b intersects c counted with signs.

Fact Same holds for smooth singular 1-chains  $c: \Delta^1 \cong I \rightarrow T^2$  which intersect a transversely: velocity vectors  $c', c^*$



Otherwise ill-defined:  $\int_{c \text{ not smooth}} \omega$  and  $\int_{c \text{ not transverse (tangency)}} \omega$  are bad.

trick need first pick homologous representative which is smooth & transverse, by continuously deforming the chain (continuous map  $\simeq \text{id}$  so id on  $H_*$ )



Example  $\int_{\tilde{c}} \omega$  deform  $\int_{\tilde{c}'} \omega$  both cases:  $a^*(\tilde{c}) = 0$


Claim  $\int_{\Delta^2} (a^* \cup b^*)(D_1 + D_2) = a^*(D_1|_{[e_0, e_1]}) \cdot b^*(D_2|_{[e_1, e_2]}) + \text{same for } D_2$

Notice we are using the "Useful Trick" (start of Sec 10) We view  $D$  as the simplicial cycle  $D_1 + D_2$ .


Graded-comm.  $\implies b^* a^* = -D^* a^* \cup a^* = (-1)^{|a^*|} a^* \cup a^*$  so  $= 0$ , similarly  $b^* b^* = 0$ .

Idea  $\cup$  just counts (signed) geometric intersection # of corresponding curves. Why " $a \cap a = 0$ "? Can deform  $a$  to make it disjoint from  $a$ :



Exercise  $\Sigma_2$



remove balls & glue babies



Make life simpler: deform generators:



$H_*(\Sigma_2)$	$H^*(\Sigma_2)$
$\mathbb{Z}$	$\mathbb{Z} \cdot 1$
$\mathbb{Z}^4$	$\mathbb{Z} \langle a_1^*, b_1^*, a_2^*, b_2^* \rangle \leftarrow$ dual basis
$\mathbb{Z} \cdot D$	$\mathbb{Z} \cdot D^*$

Notice on  $C_1^{CW}(\Sigma_2)$ :  
 $a_i^*(c) = -\#(b_i \text{ intersects } c)$   
 $b_i^*(c) = \#(a_i \text{ intersects } c)$

Exercise  $a_i^* \cup b_j^* = \delta_{ij}$   
 $a_i^* \cup a_i^* = b_i^* \cup b_i^* = 0$   
 hint:  $D$  is homologous to the sum of triangles in last picture (orientation signs)  
 so same as geometric intersection numbers of corresponding curves.

Cultural Rmk on general theory (Intersection Theory/Differential Topology)

$M^m$  oriented m-mfd  $\Rightarrow H_n(N) \xrightarrow{\text{incl}} H_n(M) \xrightarrow{\text{see later in course}}$   
 $N^n \subseteq M^m$  oriented n-dim submfd  $\Rightarrow [N] \xrightarrow{\text{with signs}}$

$N_1, N_2$  also smooth (see Differential Geometry course)  $\Rightarrow \omega_N \in H^{m-n}(M)$  counts # intersections with  $N$   
 (can always homotope  $N_1, N_2$  to achieve transversality, and class  $\omega_N$  does not change if homotope)  
 $N_1, N_2 \subseteq M$  compact submfd  $\Rightarrow \omega_{N_1} \cup \omega_{N_2} = \omega_{N_1 \cap N_2} \in H^{2m-n_1-n_2}(M)$   
 and transverse (= at every pt  $N_1, N_2$  the tangent spaces to  $N_1, N_2$  at  $p$  span the tangent space to  $M$  at  $p$ )  
 (vector space approximation at  $p$  in the local smooth coordinates)  
 Fact (Thom 1954)  $\Rightarrow$  Not all  $a \in H^j(M)$  arise as  $\omega_N$  for connected compact oriented codim=j smooth submfd  $N$ . But  $\exists N \in \mathbb{N}$  s.t.  $N \cdot a$  does arise. They do arise for  $H^*(M; \mathbb{Q}), H^*(M; \mathbb{R}), H^*(M; \mathbb{Z}/2)$

II. KÜNNETH FORMULA AND PRODUCT SPACES

Algebra: tensor products

$R$  ring (comm. with 1) e.g. abelian groups =  $\mathbb{Z}$ -mods  
 vector spaces/ $\mathbb{F}$  =  $\mathbb{F}$ -mods  
 Def  $A, B$   $R$ -modules  $\Rightarrow$  Tensor product is  $R$ -module

$A \otimes_R B = \langle (a, b) : a \in A, b \in B \rangle$  / relations of bilinearity & rescaling  
 (or  $A \otimes B$ )  $R$ -mod generated write  $a \otimes b$  for its class

bilinearity:  $(a_1 + a_2) \otimes b = a_1 \otimes b + a_2 \otimes b$   
 $a \otimes (b_1 + b_2) = a \otimes b_1 + a \otimes b_2$   
rescaling:  $r \cdot (a \otimes b) = (ra) \otimes b = a \otimes (rb)$   $\forall r \in R$   
 ("can move  $r \in R$  across the  $\otimes$  symbol")

So general element looks like  $\sum a_k \otimes b_k$  (finite sum)  $\leftarrow$  NOT UNIQUE!  
 Don't confuse with  $A \times B$ : e.g.  $0 \otimes b = 0 \forall b$

Rmk Can define  $A \otimes_R B$  also by a universal property: for all  $R$ -mods  $C$ ,

$\text{Hom}_R(A \otimes_R B; C) \xrightarrow{\text{natural}} \{R\text{-bilinear maps } A \times B \rightarrow C\}$

Using above description of  $A \otimes B: \varphi \mapsto (A \times B \rightarrow C, (a, b) \mapsto \varphi(a \otimes b))$

Example  $(R = \mathbb{F})$   $V, W$  v.s./ $\mathbb{F}$   $\Rightarrow V \otimes W$  v.s./ $\mathbb{F}$  basis  $v_i \otimes w_j$   
 basis:  $\dim V \otimes W = \dim V \cdot \dim W$

Exercise  $V, W$  finite dim/ $\mathbb{F} \Rightarrow V^* \otimes W \cong \text{Hom}_{\mathbb{F}}(V, W)$

Hint  $f: V \rightarrow W, v \in W, f \otimes w \mapsto (V \rightarrow W, v \mapsto f(v) \cdot w)$

Examples  $(R = \mathbb{Z})$   
 $\mathbb{Z}^n \otimes \mathbb{Z}^m \cong \mathbb{Z}^{n \cdot m}$  e.g.  $(\mathbb{R}^n)^* \otimes \mathbb{R}^m \cong \text{Mat}_{m \times n}(\mathbb{R}) \cong \text{Hom}(\mathbb{R}^n, \mathbb{R}^m)$   
 $\mathbb{Z}/n \otimes \mathbb{Z} \cong \mathbb{Z}/n$   $\leftarrow \sum a_i \otimes b_i = (\sum a_i b_i) \otimes 1$   
 $\mathbb{Z}/2 \otimes \mathbb{Z}/3 \cong 0$   $\leftarrow 1 \otimes x = 3 \otimes x = 1 \otimes 3x = 0$   
 $\mathbb{Z}/2 \otimes \mathbb{Z}/4 \cong \mathbb{Z}/2$   $\leftarrow \{1 \otimes 1 = 3 \otimes 1 = 1 \otimes 3 = 0\}$   
 $A \otimes B \cong B \otimes A$   $\leftarrow 1 \otimes 2 = 2 \otimes 1 = 0$

$(\bigoplus_i A_i) \otimes (\bigoplus_j B_j) \cong \bigoplus_{i,j} (A_i \otimes B_j)$  hence now know  $A \otimes B$  for any f.g.  $R$ -mods  $A, B$ .  
 $A \otimes R \cong A$  (so " $\otimes_R$  does nothing")  
 $A \otimes R/d \cong A/d \cdot A$

for example  $\mathbb{Z}/2 \otimes \mathbb{Z}/4 \cong (\mathbb{Z}/4)/2 \cdot \mathbb{Z}/4 \cong \mathbb{Z}/2$   $\leftarrow \left( \frac{\text{Rmk}}{\cong} \mathbb{Z}/m / m \cdot \mathbb{Z}/n \right)$   
 More generally:  $\left\{ \begin{array}{l} R/I \otimes R/J \cong R/(I+J) \text{ for ideals } I, J \subseteq R. \\ A \otimes R/J \cong A/J \cdot A \end{array} \right.$

Warning  $\otimes A$  often not an exact functor, i.e. does not preserve exact sequences  
indeed it can ruin injectivity:  $0 \rightarrow \mathbb{Z} \xrightarrow{\cdot 2} \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$

Fact  $\cdot \otimes \mathbb{Z}$  and  $\otimes \mathbb{R}$  are exact functors on  $\mathbb{Z}$ -mods  
 ← More generally  $\otimes \text{Frac}(R)$  is exact on  $R$ -mods where  $\text{Frac}(R)$  is fraction field, and  $R$  is an integral domain  
 // Localization is an exact functor"

example  $A$  f.g.  $\mathbb{Z}$ -mod  $\Rightarrow A \cong \mathbb{Z} \oplus \mathbb{Z}/d_1 \oplus \dots \oplus \mathbb{Z}/d_k$  some  $d_i \neq 0$   
 $\Rightarrow A \otimes \mathbb{Q} \cong \mathbb{Q}^r \Rightarrow r = \text{rank } A = \dim_{\mathbb{Q}}(A \otimes \mathbb{Q})$

Corollary Rank-nullity thm holds for  $\mathbb{Z}$ -modules if use rank instead of dim.  
 PF  $0 \rightarrow A \rightarrow B \xrightarrow{f} C \rightarrow 0$  exact  $\Rightarrow 0 \rightarrow A \otimes \mathbb{Q} \rightarrow B \otimes \mathbb{Q} \rightarrow C \otimes \mathbb{Q} \rightarrow 0$  exact  
 here  $\xrightarrow{\text{im } f} \Rightarrow \dim(\text{im } f) + \dim(A \otimes \mathbb{Q}) = \dim(B \otimes \mathbb{Q})$ .  $\square$   
 rank-nullity for  $\mathbb{Q}$ -vector spaces.

Tensor product of chain cxes  
 $C_*, \tilde{C}_*$  chain cxes  $\Rightarrow (C_* \otimes \tilde{C}_*)_n = \bigoplus_{i+j=n} C_i \otimes \tilde{C}_j$   
 of  $R$ -mods

$\partial(x \otimes y) = (\partial x) \otimes y + (-1)^{\deg x} x \otimes \partial y$   
 Think of  $\partial$  as an operator of  $\deg = -1$  acting from left  
 since  $\partial$  "jumps over  $x$ "  
 get  $(-1)^{\deg x} \cdot \deg x$

Exercise  $\partial \circ \partial = 0$  ← would fail without sign  
 recall  $\mathbb{Z}_k = \ker \partial_k = \text{im } \partial_{k+1}$   
 $\mathbb{B}_k = \text{im } \partial_k = \text{boundaries}$   
 $\mathbb{Z}_i \otimes \tilde{\mathbb{Z}}_j \subseteq \mathbb{Z}_{i+j}(C_* \otimes \tilde{C}_*)$  and  $\mathbb{Z}_i \otimes \tilde{\mathbb{B}}_j \subseteq \mathbb{B}_{i+j}(C_* \otimes \tilde{C}_*)$

Cor  $\exists$  natural maps  

$$H_i(C_*) \otimes H_j(\tilde{C}_*) \rightarrow H_{i+j}(C_* \otimes \tilde{C}_*)$$

$$\sum [c_k] \otimes [\tilde{c}_k] \mapsto \sum [c_k \otimes \tilde{c}_k]$$

FACT:  
Algebraic Künneth Thm  
 $C_*, H_*(C_*)$  f.g. free  $R$ -mods (no assumption on  $\tilde{C}_*$ )  
 $\Rightarrow \bigoplus_{i+j=n} H_i(C_*) \otimes H_j(\tilde{C}_*) \cong H_n(C_* \otimes \tilde{C}_*)$  via

Algebra: Euler characteristic  
 $C$  finitely generated graded abelian gr (so  $\mathbb{Z}$ -mod) | more generally:  $R$ -mod for PID  $R$   
Def Euler characteristic  $\chi(C) = \sum (-1)^i \text{rank } C_i$   
Example/Motivation  $X$  finite CW-cx then take  $C = C_*^{CW}(X)$  to get

$$\chi(X) = \#(0\text{-cells}) - \#(1\text{-cells}) + \#(2\text{-cells}) - \dots$$
  
Lemma If  $C_*$  f.g. chain cx  $\Rightarrow \chi(C_*) = \chi(H_*(C_*)) = \chi(\sum (-1)^i \text{rank } H_i(C_*))$

PF Abbreviate  $|C_i| = \text{rank } C_i = (\dim_{\mathbb{Q}}(C_i \otimes \mathbb{Q}))$   
 By previous Corollary about rank-nullity:  
 $0 \rightarrow \mathbb{Z}_i \rightarrow C_i \xrightarrow{\partial} B_{i-1} \rightarrow 0 \Rightarrow |C_i| = |\mathbb{Z}_i| + |B_{i-1}|$   
 $0 \rightarrow B_i \rightarrow \mathbb{Z}_i \rightarrow H_i \rightarrow 0 \Rightarrow |H_i| = |\mathbb{Z}_i| - |B_i|$   
 $\Rightarrow \chi(C_*) - \chi(H_*) = \sum (-1)^i |B_{i-1}| - \sum (-1)^i |B_i| = \sum (-1)^i (|B_{i-1}| - |B_i|) = 0. \square$

Cor  $X$  space  $\Rightarrow \chi(X) = \sum (-1)^i \text{rank } H_i(X)$   
 $= \sum (-1)^i \text{rank } C_i(X)$   
 ← if finite rank  $H_*(X)$   
 ← if finite rank  $C_*(X)$

So  $\chi(X)$  is invariant up to htpy equivalence! Example  $\chi(\text{platonic solid}) = \chi(S^2) = 2$


Product spaces  
 $X, Y$  CW-cxes  $\Rightarrow X \times Y$  CW-cx with cells  $e_\alpha \times e_\beta$  attaching maps  
 $\partial(D_\alpha^i \times D_\beta^j) = (\partial D_\alpha^i) \times D_\beta^j \cup D_\alpha^i \times (\partial D_\beta^j)$   
 $\downarrow \text{id} \times \partial_\beta$   
 $\downarrow \text{id} \times \partial_\alpha$   
 $X^{i-1} \times Y^j \cup X^i \times Y^{j-1}$   
 $(X \times Y)^{i+j-1}$

Cor  $\chi(X \times Y) = \chi(X) \cdot \chi(Y)$   
 $\forall$  finite CW-cxes  $X, Y$   
PF  $\sum (-1)^k \text{rank } H_k^{CW}(X \times Y) = \sum_k \sum_{i+j=k} (-1)^{i+j} \text{rank } C_i^{CW}(X) \cdot \text{rank } C_j^{CW}(Y)$   
 $= \sum (-1)^k \text{rank } C_k^{CW}(X \times Y) = (d e_\alpha^i \times e_\beta^j + (-1)^i e_\alpha^i \times (d e_\beta^j))$

Lemma  $d(e_\alpha^i \times e_\beta^j) = (d e_\alpha^i) \times e_\beta^j + (-1)^i e_\alpha^i \times (d e_\beta^j)$   
 hence  $C_*^{CW}(X \times Y) \cong C_*^{CW}(X) \otimes C_*^{CW}(Y)$   
 (proof later)

Hence if  $H_*(Y)$  free then by Künneth  $H_*(X \times Y) \cong H_*(X) \otimes H_*(Y)$ .  
Example

$H_*(S^1)$	$A \cong \mathbb{Z}$	$A \otimes A$	$H_*(S^1 \otimes H_*(S^1)) \cong H_*(S^1 \times S^1) \leftarrow \text{torus}$
$A \cong \mathbb{Z}$	$B \cong \mathbb{Z}$	$(A \otimes B) \oplus (B \otimes A)$	$\cong \mathbb{Z}^2$
$0$	$0$	$B \otimes B$	$\cong \mathbb{Z}$




for  $R$ -mods, do  $\dim_{\mathbb{F}}(C_i \otimes \mathbb{F})$  with  $\mathbb{F} = \text{Frac}(R)$  ( $R$  integral domain).  
 [Corollary still holds, same proof]

Pf  $(\partial D_\alpha^i) \times D_\beta^j \xrightarrow{\varphi_\alpha \times \text{id}} X^{i-1} \times Y^j \rightarrow X^{i-1} \times Y^j$   
 This proof is Non-examinable

$(X \times Y)^{i+j-2} \cap (X^{i-1} \times Y^j)$   
 || easy check  
 $X^{i-2} \times Y^j \cup X^{i-1} \times Y^{j-1}$   
 get ~ from attaching maps

$X^{i-1} \times Y^j = X^{i-2} \cup (D_\beta^{i-1} \times D_\gamma^j) \cup (D_\beta^{i-1} \times D_\gamma^{j-1})$   
 $\Rightarrow \textcircled{\star} = (D_\beta^{i-1} \times D_\gamma^j \cup \dots) / \text{boundaries}$   
 $= D_\beta^{i-1} \times D_\gamma^j / \partial(D_\beta^{i-1} \times D_\gamma^j) \vee \dots$   
 $(\partial D_\alpha^i) \times D_\beta^j \xrightarrow{\varphi_\alpha \times \text{id}} D_\beta^{i-1} \times D_\gamma^j / \text{bdry} \vee \dots$




By considering local degrees now we see we get degree =  $d_\alpha d_\beta$  for this.  
 $\Rightarrow$  get contribution  $(d_\alpha^i) \times e_\beta^j$  ✓

similarly  $D_\alpha^i \times \partial D_\beta^j \xrightarrow{\text{id} \times \varphi_\beta} D_\alpha^i \times D_\beta^{j-1} / \text{bdry}$   
 $\Rightarrow$  degree  $(-1)^i d_\alpha d_\beta$   
 so get  $(-1)^i e_\alpha^i \times d_\beta^j$

$(-1)^i$  caused by orientations:  
 could reorder factors:  $D_\alpha^i \times D_\beta^j \cong D_\beta^j \times D_\alpha^i$  by  $(\circ \text{Id}_i \circ)$   
 whose det =  $(-1)^i$ . Then  $\partial D_\beta^j \times D_\alpha^i \rightarrow D_\beta^{j-1} \times D_\alpha^i / \text{bdry}$  gives degree  $d_\alpha d_\beta$ .  
 Swap factors  $D_\beta^{j-1} \times D_\alpha^i / \text{bdry}$  by  $(\circ \text{Id}_i \circ)$ , det =  $(-1)^{i(j-1)}$ . Total sign =  $(-1)^i$ .

Example Recall after definition of  $H_*^{CW}$  we had example IX I:  
 arrows here tell us how we map  $[-1, 1] \rightarrow \text{edge}$  (so orientation)



$\partial e^2 = +e^1 \times 0 + 1 \times e^1 - e^1 \times 1 - 0 \times e^1$   
 $= (\partial e^1) \times e^1 - e^1 \times (\partial e^1)$  ✓  
 (-1) dim e^1

A further comment on orientation sign  $(-1)^i$


$D^i \times D^j \cong \Delta^i \times \Delta^j \cong [v_0, \dots, v_i] \times [w_0, \dots, w_j]$   
 viewed in  $\mathbb{R}^i, \mathbb{R}^j$   
 project  $\mathbb{R}^{i+j} \rightarrow \mathbb{R}^i$   
 $(t_0, \dots, t_i) \mapsto (t_0, \dots, t_i)$

$\partial(D^i \times D^j) \cong \partial \Delta^i \times \Delta^j \cup \Delta^i \times \partial \Delta^j$

$\sum_k (-1)^k [v_0, \dots, \hat{v}_k, \dots, v_i] \times [w_0, \dots, w_j]$   
 $\cong \sum_k (-1)^k [w_0, \dots, \hat{w}_k, \dots, w_j]$

would be correct orientation sign for basis  $w_1 - w_0, \dots, w_k - w_{k-1}, \dots, w_j - w_0$  but actually we have  $[w_0, \dots, w_k, \dots, w_j] \times [v_0, \dots, v_i] \subseteq \mathbb{R}^i \times \mathbb{R}^j$   
 and  $(-1)^{i+k}$  is the orientation sign for the basis  
 $v_1 - v_0, \dots, v_i - v_0, w_1 - w_0, \dots, w_k - w_0, \dots, w_j - w_0$   
 for the hyperplane in  $\mathbb{R}^{i+j+1}$  containing  
 $\Rightarrow$  need  $(-1)^i$  to fix orientation sign.

Example  $\Delta^1 \times \Delta^2 \subseteq \mathbb{R}^3$



$[v_0, v_1] \times [w_0, w_1, w_2]$   
 out  $w_2 - w_1$  is positive  $\mathbb{R}^2$ -basis  
 out  $v_1 - v_0$  is negative  $\mathbb{R}^2$ -basis  
 differ due to  $(-1)^i, i=1$ .

Projections  $X \times Y \xrightarrow{p_X} X$   
 $\xrightarrow{p_Y} Y$

FACT: no conditions on  $X$

**Künneth Theorem** If  $H_n(Y)$  finitely generated, free  $\forall n$

$$H_n(X \times Y) \cong \bigoplus_{i+j=n} H_i(X) \otimes H_j(Y)$$

$$H^n(X \times Y) \cong \bigoplus_{i+j=n} H^i(X) \otimes H^j(Y)$$

eg.  $Y \cong$  finite CW cplx  
 automatic if use field coefficients

Recall for cellular homology this on generators is: (chain level)

$$e_i \times e_j \mapsto e_i \otimes e_j$$

This is hom of rings if use following product

$$(a \otimes b) \cdot (\tilde{a} \otimes \tilde{b}) = (-1)^{|b| \cdot |\tilde{a}|} (a \cup \tilde{a}) \otimes (b \cup \tilde{b})$$

think of it as 'exchanging order of  $b, \tilde{a}$ '

An indirect proof the Thm is to write down two generalised cohomology theories  $F(X,A) = H^*(X,A) \otimes H^*(Y)$  and  $G(X,A) = H^*(X \times Y, A \times Y)$ , and consider the natural transformation  $\alpha: F \rightarrow G$  given by  $\otimes$ , notice for  $X = pt$  both  $F, G$  give  $H^*(Y)$ .

Example  $X = S^n, Y = S^m, n \neq m$

$$H_*(S^n \times S^m) \cong \begin{cases} \mathbb{Z} & * = 0, n, m, n+m \\ 0 & \text{else} \end{cases}$$

$$H_*(S^n \times S^n) \cong \begin{cases} \mathbb{Z} & * = 0, 2n \\ \mathbb{Z}^2 & * = n \leftarrow \text{gens: } a_n^{(1)}, a_n^{(2)} \\ 0 & \text{else} \end{cases}$$

**exterior algebra**

Cor  $H^k(T^n) \cong \wedge^k [x_1, \dots, x_n]$  = free abelian gp. on gens.

$$\{x_i, \wedge x_i, \dots, \wedge x_k : i_1 < \dots < i_k\}$$

so rank =  $\binom{n}{k}$   
 product is " $\wedge$ " using the rule  $x_i \wedge x_j = -x_j \wedge x_i$   
 (compare graded-commutativity at cup product)

FACT cup product equals composition

$$\cup : H^i(X) \otimes H^j(X) \rightarrow H^{i+j}(X \times X)$$

$$\left( \Delta_{\sigma_1}^i \rightarrow X \right) \otimes \left( \Delta_{\sigma_2}^j \rightarrow X \right) \mapsto \left( \Delta_{\sigma_1 \times \sigma_2}^{i+j} \rightarrow X \times X \right)$$

**12. UNIVERSAL COEFFICIENTS THEOREM**

Proof is non-examinable. For  $(C_*, \partial_*)$  chain cplx:

$$\Rightarrow 0 \rightarrow Z_* = \ker \partial_* \xrightarrow{\text{incl}} C_* \xrightarrow{\partial_*} B_{*-1} \rightarrow 0 \text{ is SES}$$

FACT: Submodules of a free  $\mathbb{Z}$ -module are free

Rmk The same holds for R-mods if R is PID

Assume  $C_*$  free  $\mathbb{Z}$ -mod

FACT  $\Rightarrow Z_*, B_*$  free (as  $\ker \partial, \text{Im } \partial$  are submods of  $C_*$ )

$\Rightarrow$  SES splits, choose splitting  $C_* \xrightarrow{\partial_*} B_{*-1}$  so  $\partial_* \circ s = \text{id}$

dual SES  $0 \leftarrow Z^* \xleftarrow{\text{incl}^*} C^* \xleftarrow{\partial^*} B^{*-1} \leftarrow 0$

$0 \leftarrow Z^n \xleftarrow{\partial} C^n \xleftarrow{\partial} B^{n-1} \leftarrow 0$

$0 \leftarrow Z^{n-1} \xleftarrow{\partial} C^{n-1} \xleftarrow{\partial} B^{n-2} \leftarrow 0$

note:  $\text{incl}^* = \text{restrict to } Z_*$  since  $\text{incl}^* \circ \partial = \partial \circ \text{incl}^*$

Connecting map  $\delta: Z^{n-1} \rightarrow B^n$

of LES:  $\varphi|_{Z^*} = \phi \leftarrow \exists \varphi$

LES  $\Rightarrow \dots \leftarrow \delta^n \leftarrow Z^n \leftarrow H^n C \leftarrow H^{n-1} C \leftarrow \dots$

$(H^* B = B^*, H^* C = C^* \text{ since } \partial^* = 0)$

$\Rightarrow 0 \leftarrow \ker \delta^n \leftarrow H^n C \leftarrow B^{n-1} / \text{Im } \delta^{n-1} \leftarrow 0$

$\ker \delta^n = \{ \phi \in Z^n : \phi(B_n) = 0 \} \Rightarrow$  so:  $\phi: Z_n \rightarrow Z$

$= \text{Hom}(H_n(C_*), \mathbb{Z})$

Universal Coefficients Thm:

$$0 \rightarrow B^n / \text{Im } \delta^{n-1} \rightarrow H^n(C) \rightarrow \text{Hom}(H_n(C_*), \mathbb{Z}) \rightarrow 0 \text{ is SES}$$

$$\text{Ext}^1(H_{n-1}(C_*), \mathbb{Z}) \rightarrow [ \varphi ] \rightarrow (\varphi: H_n(C_*) \rightarrow \mathbb{Z}) \text{ and natural}$$

and SES splits (but not naturally):  $B^{n-1} / \text{Im } \delta^{n-1} \xrightarrow{\partial^*} H^n(C)$

MOTIVATION: What is difference between  $H^*(\text{Hom}(C_*, \mathbb{Z}))$  and  $\text{Hom}(H_n(C_*), \mathbb{Z})$  Similarly:  $H_n(C_* \otimes \mathbb{Q})$  vs.  $H_n(C_*) \otimes \mathbb{Q}$ .

$(\mathbb{Z}$ -module  $\equiv$  abelian gp free means:  $\bigoplus_{\text{indexing set}} \mathbb{Z}$ )

PID = principal ideal domain = integral domain R s.t. every ideal =  $R \cdot a$  some  $a$

recall just pick preimages under  $\partial_*$  of a basis for  $B_*$

note:  $\text{incl}^* = \text{restrict to } Z_*$  since  $\text{incl}^* \circ \partial = \partial \circ \text{incl}^*$

Rmk Although  $\partial^* = 0: B^n \rightarrow B^{n+1}$  the map  $\partial^*: B^{n-1} \rightarrow C^n$  need not = 0

$\varphi: B_{n-1} \rightarrow \mathbb{Z}$

$\Rightarrow \partial^* \varphi = \varphi \circ \partial: C_n \rightarrow B_{n-1} \xrightarrow{\partial} B_n$

$$\Rightarrow \delta(\phi) = \phi|_{B_*}$$

see next Lemma

and  $H^n(C) \cong \text{Hom}(H_n(C), \mathbb{Z}) \oplus \text{Ext}^1(H_{n-1}(C), \mathbb{Z})$

$S^* \circ \partial^* = \text{id}$  (since  $\partial \circ S = \text{id} \Rightarrow \text{id} = (\partial \circ S)^* = S^* \circ \partial^*$ )



Lemma  $\text{Ext}^1(H_{n-1}(C); \mathbb{Z}) \cong B^{n-1}/\text{Im } \delta^{n-1}$  canonically

Algebra background on Extension groups  $\text{Ext}^i(M; \mathbb{Z})$

general case

$M$   $R$ -module,  $R$  ring (comm. with 1)

$\Rightarrow \exists$  free resolution:

$$\dots \rightarrow P_2 \xrightarrow{\psi_2} P_1 \xrightarrow{\psi_1} P_0 \xrightarrow{\psi_0} M \rightarrow 0 \quad \text{exact, } P_i \text{ free } R\text{-mods}$$

(pick gens  $x_\alpha$  for  $M \Rightarrow P_0 = \bigoplus_{\alpha} R \xrightarrow{\psi_0} M, x_\alpha \mapsto x_\alpha$   
 " "  $y_\beta$  for  $\text{Ker } \psi_0 \Rightarrow P_1 = \bigoplus_{\beta} R \xrightarrow{\psi_1} \text{Ker } \psi_0, e_\beta \mapsto y_\beta$   
 continue inductively)

Take  $\text{Hom}(\cdot; \mathbb{Z})$  and drop  $\text{Hom}(M; \mathbb{Z})$

$$0 \rightarrow \text{Hom}(P_0; \mathbb{Z}) \xrightarrow{\psi_1^*} \text{Hom}(P_1; \mathbb{Z}) \xrightarrow{\psi_2^*} \dots$$

Is cochain complex but not exact

$\Rightarrow$  take cohomology groups:

Def  $\text{Ext}^0(M; \mathbb{Z}) = \text{Ker } \psi_1^*$

$\text{Ext}^1(M; \mathbb{Z}) = \text{Ker } \psi_2^* / \text{Im } \psi_1^*$

Fact independent of choices  $P_i, \psi_i$

Example 1  $\text{Ext}^0(M; \mathbb{Z}) \cong \text{Hom}(M; \mathbb{Z})$

$$P_1 \xrightarrow{\psi_1} P_0 \xrightarrow{\psi_0} M$$

$\downarrow \phi$  descends:  $m \mapsto \phi(\psi_0^{-1}m)$  will be defined since  $\phi(\text{Ker } \psi_0) = 0$

Example 2  $\text{Ext}^1(M; \mathbb{Z}) =$

$$\left\{ \begin{array}{l} \phi : P_2 \xrightarrow{\psi_2} P_1 \rightarrow P_0 \\ \downarrow \phi \quad \downarrow \psi_1 \quad \downarrow \psi_0 \\ \mathbb{Z} \quad \mathbb{Z} \quad \mathbb{Z} \end{array} \right\} / \left\{ \phi = \psi_0 \psi_1 : P_1 \xrightarrow{\psi_1} P_0 \right\}$$

Rmk If  $R$  PID, then  $\text{Ker}(P_0 \rightarrow M)$  is free (since submod of free mod  $P_0$ )

$\Rightarrow$  can pick  $P_1 = \text{Ker}(P_0 \rightarrow M)$ ,  $P_k = 0$  for  $k \geq 2 \Rightarrow \text{Ext}^k(M; \mathbb{Z}) = 0 \quad k \geq 2$

our case  
 $H_{n-1}(C_*) \quad \mathbb{Z}$ -mod

$$0 \rightarrow B_{n-1} \hookrightarrow Z_{n-1} \rightarrow H_{n-1}(C) \rightarrow 0$$

$$\parallel \quad \parallel \quad \parallel$$

$$P_1 \quad P_0 \quad M$$

$$0 \rightarrow Z^{n-1} \rightarrow B^{n-1} \rightarrow 0$$

Proof of Lemma

By Example 2,

$$\text{Ext}^1(H_{n-1}(C_*); \mathbb{Z}) =$$

$$= \left\{ \begin{array}{l} 0 \rightarrow B_{n-1} \rightarrow Z_{n-1} \\ \downarrow \phi \quad \downarrow \psi \\ \mathbb{Z} \quad \mathbb{Z} \end{array} \right\} \text{ modulo those arising from restriction}$$

those arising from restriction

$$\left\{ \begin{array}{l} 0 \rightarrow B_{n-1} \rightarrow Z_{n-1} \\ \downarrow \phi \quad \downarrow \psi \\ \mathbb{Z} \quad \mathbb{Z} \end{array} \right\}$$

Thus  $B^{n-1}/\text{Im } \delta^{n-1} \quad \square$

(Co)homology with coefficients in a ring/field/module

Motivation

So far we had  $(C_*, \partial_*)$  chain cx of abelian groups } in graded sense

$$\Rightarrow H_*(C_*) = \text{Ker } \partial_* / \text{Im } \partial_* \text{ abelian group (since } \text{Ker } \partial, \text{Im } \partial \text{ are)}$$

We cannot use a chain cx of (non-abelian) groups, because  $\text{Im } \partial_*$  need not be a normal subgroup of  $\text{Ker } \partial_*$ .

However, abelian groups can be thought of as  $\mathbb{Z}$ -modules,

then given any abelian group  $G$ , define homology with coeffs in  $G$  with differential  $\partial_* \otimes \text{id}$


$$H_*(C_*; G) = H_*(C_* \otimes_{\mathbb{Z}} G)$$

$$\text{Def } X \text{ space} \Rightarrow H_*(X; G) = H_*(C_*(X) \otimes_{\mathbb{Z}} G)$$

Explanation:

$C_k(X)$  free  $\mathbb{Z}$ -mod  $\cong \bigoplus_{I_k} \mathbb{Z} \Rightarrow C_k(X) \otimes_{\mathbb{Z}} G \cong \bigoplus_{I_k} G$ : just replace  $\mathbb{Z}$  by  $G$  (as  $\mathbb{Z} \otimes \cong \cdot$ )

Why care? We hope to get more/new invariants of spaces

Example  $X = \mathbb{R}P^2 =$  

$$C_*(\mathbb{R}P^2; G) = \begin{array}{c|ccc} * & 2 & 1 & 0 \\ \hline & G \otimes G \oplus G & G \otimes G \oplus G & G \otimes G \oplus G \end{array}$$

$$\text{for } G = \mathbb{Z}/2: 0 \rightarrow \mathbb{Z}/2 \oplus \mathbb{Z}/2 \xrightarrow{\partial_2} \mathbb{Z}/2 \oplus \mathbb{Z}/2 \xrightarrow{\partial_1} \mathbb{Z}/2 \oplus \mathbb{Z}/2 \rightarrow 0$$

$$\left( \begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array} \right) \quad \left( \begin{array}{c} 110 \\ 110 \end{array} \right)$$

$$\Rightarrow H_*(\mathbb{R}P^2; \mathbb{Z}/2) \cong \begin{cases} \mathbb{Z}/2 & * = 0 \\ \mathbb{Z}/2 & * = 1 \\ \mathbb{Z}/2 & * = 2 \\ 0 & \text{else} \end{cases} \quad \text{compare: } H_*(\mathbb{R}P^2) \cong \begin{cases} \mathbb{Z} & * = 0 \\ \mathbb{Z}/2 & * = 1 \\ 0 & \text{else} \end{cases}$$

Form cochain complex using  $\text{Hom}_{\mathbb{Z}}(\cdot, G)$  (= group homs in place of  $\text{Hom}(\cdot, \mathbb{Z})$ )

$$H^*(C_*; G) = H^*(\text{Hom}_{\mathbb{Z}}(C_*, G))$$

$$H^*(X; G) = H^*(\text{Hom}_{\mathbb{Z}}(C_*(X); G))$$

so:  $\partial^* \phi = \phi \circ \partial_*$

Universal coefficients thm (same proof using  $\text{Hom}_{\mathbb{Z}}(\cdot, G)$ )

$$0 \rightarrow \text{Ext}_{\mathbb{Z}}^1(H_{n-1}(C_*; G) \rightarrow H^n(C_*; G) \rightarrow \text{Hom}_{\mathbb{Z}}(H_n(C_*), G) \rightarrow 0$$

$$[\varphi] \mapsto (\varphi : H_n(C_*) \rightarrow G)$$

Example  $X = \mathbb{R}P^2$ ,  $G = \mathbb{Z}/2$ , apply  $\text{Hom}_{\mathbb{Z}}(\cdot, G)$  to the complex in previous example

$$0 \leftarrow \mathbb{Z}/2 \oplus \mathbb{Z}/2 \xrightarrow{\partial^1} \mathbb{Z}/2 \oplus \mathbb{Z}/2 \oplus \mathbb{Z}/2 \xrightarrow{\partial^0} \mathbb{Z}/2 \oplus \mathbb{Z}/2 \leftarrow 0$$

$$\begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 0 & 0 & 0 \end{pmatrix} \begin{matrix} \leftarrow \partial^0 \\ \leftarrow \partial^1 \\ \leftarrow \partial^2 \end{matrix}$$

$\Rightarrow H^*(\mathbb{R}P^2; \mathbb{Z}/2) \cong \begin{cases} \mathbb{Z}/2 & * = 0 \\ \mathbb{Z}/2 & * = 1 \\ \mathbb{Z}/2 & * = 2 \\ 0 & \text{else} \end{cases}$  compare:  $H^*(\mathbb{R}P^2) \cong \begin{cases} \mathbb{Z} & * = 0 \\ 0 & * = 1 \\ \mathbb{Z}/2 & * = 2 \\ 0 & \text{else} \end{cases}$  ( $G = \mathbb{Z}$  case)

Can generalise further:

$C_* =$ chain cx of ...	coefficients in:
abelian gps ( $\mathbb{Z}$ -mods)	$H_*(C_*; G) = H_*(C_* \otimes_{\mathbb{Z}} G)$
$R$ -modules $\leftarrow$ ring (comm. with 1)	$H_*(C_*; M) = H_*(C_* \otimes_R M)$

Rmk  $H_*(C; M)$  will be an  $R$ -module since  $\ker \partial, \text{Im } \partial$  are ( $\partial_*$  is  $R$ -linear hom by assumption)

$X$  space  $\Rightarrow C_k(X; R) = C_k(X) \otimes_{\mathbb{Z}} R \cong \bigoplus_{\mathbb{Z}} R$ : just replace  $\mathbb{Z}$  by  $R$  (as  $\mathbb{Z} \otimes R \cong R$ )

$$\Rightarrow H_*(X; M) = H_*(C_*(X; R); M) = H_*(C_*(X; R) \otimes_R M)$$

so just replace each  $\mathbb{Z}$  by  $M$  in  $C_*(X)$

Form cochain complex using  $\text{Hom}_R(\cdot, M)$  ( $= R$ -linear homs to  $M$ ) in place of  $\text{Hom}(\cdot, \mathbb{Z})$

$$\begin{matrix} H^*(C_*; M) = H^*(\text{Hom}_R(C_*, M)) \\ H^*(X; M) = H^*(\text{Hom}_R(C_*(X; R); M)) \end{matrix} \leftarrow \begin{matrix} \text{with differential } \partial^* \\ \partial^* \phi = \phi \circ \partial \end{matrix}$$

so:  $H^*(C_*(X; R); M) \leftarrow H^*(C_*(X; R), M)$

Rmk These are  $R$ -mods. If we use  $M=R$ , then they are also rings via cup product

Universal Coefficient Thm For  $R$  any PID,  $C_*$  chain cx of  $R$ -mods,

$$0 \rightarrow \text{Ext}_R^1(H_{n-1}(C_*), M) \rightarrow H^n(C_*; M) \rightarrow \text{Hom}_R(H_n(C_*), M) \rightarrow 0$$

$\text{Ext}_R^1$  is SES and natural.

$B^{n-1}/\text{Im } \delta^{n-1}$  working over  $R$  using homs to  $M$

$[\varphi] \mapsto (\varphi: H_n(C_*) \rightarrow M)$

and the SES splits but the splitting is not natural.

Example  $R = \mathbb{F}$  field  $\Rightarrow C_*, H_*, H^*$  are vector spaces/ $\mathbb{F}$ .

Rmk all  $\mathbb{F}$ -mods (i.e. vector spaces/ $\mathbb{F}$ ) are free  $\mathbb{F}$ -mods  $\cong \bigoplus \mathbb{F}b_i$  up to iso they are determined by  $\dim_{\mathbb{F}} =$  cardinality of basis.

Cor  $C_* =$  chain cx of  $\mathbb{F}$ -vector spaces  $\Rightarrow H^n(C_*; \mathbb{F}) \cong (H_n(C_*))^*$  dual v.s.:  $\text{Hom}_{\mathbb{F}}(H_n(C_*), \mathbb{F})$

Pf Pick any basis  $v_i$  for  $\mathbb{F}$ -v.s.  $B_{n-1}$ , extend it to a basis  $v_i, w_j$  of  $Z_{n-1}$  (also works in  $\infty$  dim case).

$\Rightarrow$  can extend any  $\mathbb{F}$ -linear map  $\psi: B_{n-1} \rightarrow \mathbb{F}$  to  $\phi: Z_{n-1} \rightarrow \mathbb{F}$  just pick any values  $\phi(w_j) \in \mathbb{F}$  e.g.  $\phi(w_j) = 0$ .

$\Rightarrow B^{n-1}/\text{Im } \delta^{n-1} = 0$  so  $H^n(C_*; \mathbb{F}) \rightarrow \text{Hom}(H_n(C_*), \mathbb{F})$  iso  $\square$

Cor  $H^n(X; \mathbb{F}) \cong H_n(X; \mathbb{F})^*$  for any field  $\mathbb{F}$ .

$$\text{Cor } H^n(X; M) \cong H_{CW}^n(X; M) \cong H_{\Delta}^n(X; M)$$

if  $X \cong CW$ -cx  $\uparrow$  if  $X \cong \Delta$ -cx

Pf Cor holds for homology and the isos are natural.  $\leftarrow$  i.e. functional w.r.t. maps

The universal coeff. thm SES is natural. So result holds by 5-Lemma.  $\square$

Algebra: structure thm for f.g. abelian groups and R-mods

Fact 1 A f.g. abelian gp  $\Rightarrow A \cong \mathbb{Z}^r \oplus \mathbb{Z}/p_1 \oplus \dots \oplus \mathbb{Z}/p_a$

where  $p_i \in \mathbb{Z}$  prime (need not be distinct)  $\leftarrow$  free part  $\mathbb{F}$

Also  $r, k, p_i, n_i$  are unique (up to reordering)  $\leftarrow$  torsion part  $T$

Example  $\mathbb{Z}/4 = \mathbb{Z}/2 \neq \mathbb{Z}/2 \oplus \mathbb{Z}/2$  (Chinese Remainder Thm)

$\mathbb{Z}/6 \cong \mathbb{Z}/2 \oplus \mathbb{Z}/3$  with  $d_1 | d_2 | \dots | d_k$  ( $d_i \in \mathbb{N}$  unique)

Fact 2  $T \cong \mathbb{Z}/d_1 \oplus \dots \oplus \mathbb{Z}/d_k$  with  $d_1 | d_2 | \dots | d_k$  ( $d_i \in \mathbb{N}$  unique)

Example  $\mathbb{Z}/2 \oplus \mathbb{Z}/2 \oplus \mathbb{Z}/3 \cong \mathbb{Z}/2 \oplus \mathbb{Z}/3 \cdot 2^2$   $d_1=2, d_2=12$

Fact 3  $M$  f.g.  $R$ -mod,  $R$  PID, then:

$$\begin{matrix} M \cong \mathbb{F} \oplus T \\ \mathbb{F} \cong R^r \\ T \cong R/p_1 \oplus \dots \oplus R/p_a \end{matrix} \leftarrow \begin{matrix} r \in \mathbb{N} \text{ unique, called rank of } M \\ p_i \in R \text{ primes, } p_i \text{ unique up to ordering \& mult by } \\ d_i \text{ called invariant factors} \\ \text{unique up to mult}^n \text{ by invertible elements} \\ \text{if } R = \mathbb{Z} \end{matrix}$$

$$\begin{matrix} \cong R/d_1 \oplus R/d_2 \oplus \dots \oplus R/d_k \end{matrix}$$

Rmk  $T = \{m \in M : rm = 0 \text{ some } r \neq 0 \in R\} =$  torsion elements  $\leftarrow$  e.g.  $\pm 1$  if  $R = \mathbb{Z}$

$\mathbb{F} \cong M/T$

### Torsion shift

Easy Exercise

$$\text{Ext}_R^*(\bigoplus_i M_i; \bigcap_j N_j) \cong \prod_i \text{Ext}_R^*(M_i; N_j)$$

any R-mods  $M_i, N_j$

Upshot To compute  $\text{Ext}_R^i(M, R)$  for  $M = R \oplus R/d \oplus \dots$  just need:

$$\begin{aligned} \text{Ext}_R^1(R; R) &= 0 \\ \text{Ext}_R^1(R/d; R) &\cong R/d \end{aligned}$$

since  $0 \rightarrow R \xrightarrow{d} R \rightarrow R/d \rightarrow 0$

so choice of  $\phi(1) \in R$  module  $\phi$  coming from  $R \xrightarrow{d} R$  so  $\phi(1) = d \cdot \varphi(1) \in d \cdot R$

$$\Rightarrow \text{Ext}_R^1(M; R) \cong \text{Torsion}(M)$$

Exercises

- $\text{Ext}_{\mathbb{Z}}^1(\mathbb{Z}_m; \mathbb{Z}) \cong \mathbb{Z}/\text{gcd}(m, n)$
- Gabelian gp  $\Rightarrow \text{Ext}_{\mathbb{Z}}^1(\mathbb{Z}; G) = 0$  and  $\text{Ext}_{\mathbb{Z}}^1(\mathbb{Z}/d; G) \cong G/d \cdot G$
- $R$  any ring (comm. with 1)  
 $x \in R$  not zero divisor  $\Rightarrow \text{Ext}_R^*(R/(x); N) \cong \begin{cases} \{n \in N : x \cdot n = 0\} & * = 0 \\ N/x \cdot N & * = 1 \\ 0 & \text{else} \end{cases}$

Cor If  $H_n(X; R)$  f.g. R-mod  $\forall n$ ,  $R$  PID,  
 $\Rightarrow H_n(X; R) = R^n \oplus T_n$  (free & torsion parts)

$$\Rightarrow H^n(X; R) \cong R^n \oplus T_{n-1}$$

not natural

$$\text{Pf } 0 \rightarrow T_{n-1} \rightarrow H^n(X; R) \rightarrow \text{Hom}(R^n \oplus T_{n-1}, R) \rightarrow 0$$

$$\text{Hom}(R^n \oplus T_{n-1}, R) \cong (\text{Hom}(R; R))^n \oplus \text{Hom}(T_{n-1}, R)$$

$R \rightarrow R \xrightarrow{1} x \rightarrow R^n$   
 $x$  determines the hom

$\Rightarrow 0 \rightarrow T_{n-1} \rightarrow H^n(X; R) \rightarrow R^n \rightarrow 0$   
 free, so can split the SES (pick lifts of basis).  $\square$

Example

*	$H_*(\mathbb{R}P^3)$	$H^*(\mathbb{R}P^3)$
0	$\mathbb{Z}$	$\mathbb{Z}$
1	$\mathbb{Z}/2$	$0$
2	$0$	$\mathbb{Z}/2$
3	$\mathbb{Z}$	$\mathbb{Z}$

torsion moves up

### Universal coefficients Theorem in homology

FACT Theorem  $C_*$  chain cx of free  $R$ -mods,  $M$   $R$ -module

$$\Rightarrow \text{SES } 0 \rightarrow H_*(C_*) \otimes_R M \rightarrow H_*(C_* \otimes_R M) \rightarrow \text{Tor}_1^R(H_{*+1}(C_*), M) \rightarrow 0$$

$[C] \otimes m \mapsto [C \otimes m]$

The SES splits, but the splitting is not natural.

Torsion groups:  $A, B$   $R$ -mods ( $R$  comm. ring with 1)

pick  $\dots \rightarrow P_2 \xrightarrow{\psi_2} P_1 \xrightarrow{\psi_1} P_0 \xrightarrow{\psi_0} A \rightarrow 0$  free resolution  
 $\Rightarrow \dots \rightarrow P_2 \otimes B \xrightarrow{\psi_2 \otimes \text{id}} P_1 \otimes B \xrightarrow{\psi_1 \otimes \text{id}} P_0 \otimes B \rightarrow 0$  not exact  
 but is chain cx

$\text{Tor}_k^R(A, B) = H_k$  (this complex)  $\leftarrow$  fact independent of choices of  $P_i, \psi_i$   
 Rmk  $R$  PID  $\Rightarrow \ker \psi_0$  free  $\Rightarrow$  can pick  $P_i = \ker \psi_i, P_k = 0$  for  $k > 2$   
 $\Rightarrow$  only  $\text{Tor}_0^R, \text{Tor}_1^R$  can be non-zero

Example  $\text{Tor}_*^{\mathbb{Z}}(\mathbb{Z}/a, \mathbb{Z}/b) = ?$

$$0 \rightarrow \mathbb{Z} \xrightarrow{a} \mathbb{Z} \xrightarrow{b} \mathbb{Z} \xrightarrow{a} \mathbb{Z} \xrightarrow{b} \mathbb{Z} \rightarrow 0$$

free resolution

take  $\otimes \mathbb{Z}/b \Rightarrow 0 \rightarrow \mathbb{Z}/b \xrightarrow{a} \mathbb{Z}/b \rightarrow 0$  (since  $\mathbb{Z} \otimes \mathbb{Z} \cong \mathbb{Z}$  any  $G$ )

drop  $\mathbb{Z}/a \otimes \mathbb{Z}/b \Rightarrow \text{Tor}_0^{\mathbb{Z}}(\mathbb{Z}/a, \mathbb{Z}/b) = (\mathbb{Z}/b)/a \cdot \mathbb{Z}/b \cong \mathbb{Z}/\langle a, b \rangle \cong \mathbb{Z}/\text{gcd}(a, b) \cong \mathbb{Z}/a \otimes \mathbb{Z}/b$

$\text{Tor}_1^{\mathbb{Z}}(\mathbb{Z}/a, \mathbb{Z}/b) = \{x \in \mathbb{Z}/b : a \cdot x = 0\} \cong \mathbb{Z}/\text{gcd}(a, b)$

Facts  $\text{Tor}_0^R(A, B) = P_0 \otimes B / \text{Im}(\psi_0 \otimes \text{id}) \cong A \otimes B$   
 $\text{Tor}_*^R(A, B) \cong \text{Tor}_*^R(B, A)$

Exercise  $\text{Tor}_*^R(\bigoplus_i A_i; \bigoplus_j B_j) \cong \bigoplus_i \text{Tor}_*^R(A_i; B_j)$

$\text{Tor}_*^R(A, B) = 0$  for  $* > 1$  if  $A$  or  $B$  is free (use  $M \otimes_R N \cong M$ )

$\text{Tor}_*^R(R/u, M) \cong \begin{cases} M/u \cdot M & * = 0 \\ 0 & \text{else} \end{cases}$   
 deduce  $\text{Tor}_*^R(A, M)$  for f.g.  $R$ -mods  $A$   
 $u \in R$  not zero divisor any ring (comm. with 1)  $u$ -torsion  $(M) = \{x \in M : u \cdot x = 0\} \cong \begin{cases} M/u \cdot M & * = 0 \\ 0 & \text{else} \end{cases}$

Example  $H_*(\mathbb{R}P^2; \mathbb{Z}_2) \cong \begin{cases} \mathbb{Z}_2 & * = 0 \\ \mathbb{Z}_2 & * = 1 \\ \mathbb{Z}_2 & * = 2 \end{cases}$   
 $H_*(\mathbb{R}P^2) \otimes \mathbb{Z}_2 \cong \begin{cases} \mathbb{Z} \otimes \mathbb{Z}_2 & * = 0 \\ \mathbb{Z}_2 \otimes \mathbb{Z}_2 & * = 1 \\ \mathbb{Z} \otimes \mathbb{Z}_2 & * = 2 \end{cases}$   
 $\cong \begin{cases} \mathbb{Z}_2 & * = 0 \\ \mathbb{Z}_2 & * = 1 \\ \mathbb{Z}_2 & * = 2 \end{cases}$


Künneth Thm

$R$  PID  $\Rightarrow$  natural SES:  $0 \rightarrow \bigoplus_{i+j=n} H_i(C_*) \otimes H_j(D_*) \rightarrow H_n(C_* \otimes D_*) \rightarrow \bigoplus_{i+j=n-1} \text{Tor}_1^R(H_i(C_*), H_j(D_*)) \rightarrow 0$   
 $(C_*$  free ch. cx.  $R$ -mods  $\rightarrow$  free  $R$ -mods)  
 $(D_*$  any ch. cx.  $R$ -mods)

and the SES splits but the splitting is not natural. Example  $R = \text{field}$ , then this  $= 0$ .

### 13. MANIFOLDS: POINCARÉ-LÉFSCHETZ DUALITY

- $M$   $n$ -mfd is Hausdorff topological space s.t.  $\forall p \in M$   $\exists$  open neighbourhood  $U_p \subseteq M$  homeomorphic to  $\mathbb{R}^n$




(equivalently: to an open ball, or any open set in  $\mathbb{R}^n$ )

One also requires  $M$  **second countable** i.e.  $\exists$  countable basis of open sets

$\Leftrightarrow M$  is covered by countably many such  $U_p$ :  
← exercise

A **submanifold**  $N \subseteq M$  is a mfd s.t. inclusion  $N \rightarrow M$  is an embedding (i.e. a homeomorphism onto its image)

- $M$   $n$ -mfd with **boundary** if also allow  $U_p \cong$  upper half space  $\mathbb{H}^n$  which they form the **boundary**  $\partial M$  which is an  $(n-1)$ -mfd without boundary.



equivalently: any open nbhd of  $o \in \mathbb{H}^n$

**FACT** (Collar nbhd thm)  $\partial M \subseteq M$  has an open neighbourhood  $\cong \partial M \times (0,1]$



$M$  is **closed** if compact without boundary.

**Rmk** For manifolds, connected components = path components. (since locally  $\cong$  disc, so locally path-connected, so conn.  $\Leftrightarrow$  path-con.)

**Examples**

closed mfd:  $S^n, \mathbb{R}P^n, T^n = S^1 \times \dots \times S^1, \mathbb{C}P^n, O(n), SU(n)$

non-compact mfd:  $\mathbb{R}^n, \text{Mat}_{m \times n} \cong \mathbb{R}^{mn}, GL(n, \mathbb{R})$

mfd with bdr:  $D^n, D^1 \times S^1 =$   $, T^2 \setminus \text{open disc} =$

**FACT** (Milnor 1959) Any mfd is homotopy equivalent to a CW-complex

**fact** If  $M$  is a compact manifold then  $H_*(M)$  are finitely generated

**Rmk**  $M$  **triangulable** if  $M \cong$  simplicial cx.

Not all mfd are triangulable, but most of those we encounter are.

### Compact manifolds have f.g. homology

← Non-examinable proof

①  $X$  space is a **Euclidean neighbourhood retract** if

$\exists$  **embedding**  $j: X \rightarrow \mathbb{R}^m$  some  $N$ , s.t.  $i(X)$  is a retract of a nbhd  $V \subseteq \mathbb{R}^m$  (homeo onto image)


②  $X$  is **weakly locally contractible** if  $\forall$  nbhd  $x \in U \subseteq X, \exists$  nbhd  $x \in V \subseteq U$  s.t.  $V$  is contractible inside  $U$ .

**FACT** compact  $X \subseteq \mathbb{R}^n$  is ①  $\Leftrightarrow$  ②

**Rmk** If we find nbhd  $V$  as in ① with retraction  $V \xrightarrow{f} X$  then any smaller nbhd  $V'$  also retracts using  $f|_{V'}: V' \rightarrow X$ . Similarly in ②  $V' \subseteq V$  is contractible: restrict the hpy.

**Lemma A**  $X$  compact & ①  $\Rightarrow X$  is the retract of a finite simplicial cx

**pf**  $i(X) \subseteq \mathbb{R}^n$  compact  $\Rightarrow$  lies inside some large  $n$ -simplex  $\Delta^n \rightarrow \mathbb{R}^n$   
 Apply barycentric subdivision until simplices have diameter  $< \text{dist}(X, \partial V)$ .



**Simpl. cx.** =  $\cup$  {simplices which intersect  $X$ } using the restriction of retraction  $V \rightarrow X$ .

**Rmk** Also deduce  $X$  has f.g. homology since retractions are surjective on  $H_k$ .

(①  $\Rightarrow$  ②)  $\rightarrow H_k(\text{finite simpl. cx}) \xrightarrow{\text{retract}} H_k(X)$  so get surjection from free  $\mathbb{Z}$ -mod, so f.g.)

**Lemma B**  $M$  compact mfd  $\Rightarrow M$  embeds into  $\mathbb{R}^n$ , some  $n$ .

**pf** "Just do it proof":

$\forall p \in M, \exists$  homeo  $D^n \xrightarrow{\psi_p} \text{nbhd}(p \in M)$

Pick finite subcover of  $\psi_p$ :  $M = \cup_{p \in M} \psi_p(D^n)$ . Say  $i = 1, \dots, k$

$\psi_i: M \xrightarrow{\psi_i^{-1}} D^n \rightarrow D^n / \partial D^n \cong S^n \subseteq \mathbb{R}^{n+1}$  define embedding  $(\psi_{i_1}, \dots, \psi_{i_k}): M \rightarrow \mathbb{R}^{k \cdot (n+1)}$

Finally use: a continuous bijection from a compact space to a Hausdorff space is  $\cong$

**Rmk** Same works if  $M$  has boundary, just consider its **double**  $M \cup M$  identify along  $\partial M$  and apply the Lemma to the double.

**Cor**  $M$  compact mfd (possibly with bdr)  $\Rightarrow M$  has f.g. homology

**pf** Mfd's satisfy ② since locally ball  $\cong$  pt.  $M$  embeds in  $\mathbb{R}^n$  by Lemma B.

① holds by **FACT**. Done by Lemma A.  $\square$


### Local orientations and orientability

Def A local orientation of  $M$  at  $x \in M$  is a choice of generator

$$\mu_x \in H_n(M, M \setminus x) \cong H_n(D^n, D^n \setminus \{0\}) \cong \tilde{H}_n(S^n) \cong \mathbb{Z} \quad \begin{matrix} \text{excise complement of nbhd } V_x \cong \mathbb{D}^n \\ \text{choice of } \mu_x \text{ is not canonical!} \\ \text{choice of } \mu_x \text{ is not canonical!} \end{matrix}$$

Def An orientation of  $M$  is a locally consistent choice  $x \mapsto \mu_x$

meaning:




Def  $M$  orientable if  $\exists$  orientation on  $M$


oriented if we chose an orientation

Examples  $S^n, \mathbb{R}^n, \mathbb{C}P^n$ , orientable surfaces  $\Sigma_g, \mathbb{R}P^n \leftarrow \text{odd } n$

Non-example  $\mathbb{R}P^2 = \text{Möbius band} \cup D^2$




by local consistency can move disc continuously and preserves orientation



choice of  $\mu_x$  is choice of orientation of boundary circle of small disc containing  $x$

$\Rightarrow \mathbb{R}P^2$  not orientable



discs are differently oriented  $\Rightarrow$  contradicts local consistency.

### The fundamental class $[M]$

FACT Theorem For  $M$  closed  $n$ -mfd:

$$M \text{ orientable connected} \Rightarrow H_n(M) \cong H_n(M, M \setminus x) \cong \mathbb{Z} \text{ (choice)}$$

$$\Rightarrow \exists [M] \in H_n(M) \text{ called fundamental class}$$

once we choose an orientation  $(\mu_x)_{x \in M}$

(if swap orientation: for  $-\mu_x$  get  $-[M]$ )

$$M \text{ not orientable connected} \Rightarrow H_n(M) = 0$$

$$H_n(M; \mathbb{F}_2) = \mathbb{F}_2 \cdot [M] \cong \mathbb{F}_2$$

(or any field of characteristic 2)

Construction of  $[M]$  if  $M$  has  $\Delta$ -complex structure

$M$  compact  $\Rightarrow$  finite #  $n$ -simplices  $\delta_1, \dots, \delta_N$

$M$  oriented  $\Rightarrow$  pick orientations of  $\delta_1, \dots, \delta_N$  to agree with given orientation of  $M$ :  $\checkmark$  for  $x \in \text{Int}(\delta_i)$



$$\mathbb{Z} \cdot \mu_x = H_n(M, M \setminus x) \xrightarrow{\text{exc}} H_n(\delta_i, \delta_i \setminus x) = \mathbb{Z} \cdot \delta_i$$

$$\mu_x \mapsto \delta_i$$

$$\Rightarrow [M] := \sum \delta_i \text{ satisfies } \partial[M] = 0 \checkmark$$

(each facet arises twice with opposite signs)

$$H_n(M) \rightarrow H_n(M, M \setminus x) \cong H_n(\delta_i, \delta_i \setminus x)$$

$$[M] \xrightarrow{\mu_x} \delta_i$$

More generally:  
 $[M] := \sum \pm \delta_i$   
 where signs come from  $H_n(M, M \setminus x) \cong H_n(\delta_i, \delta_i \setminus x)$   
 $\mu_x \mapsto \pm \delta_i$   
 (so compare orientation of  $\mu_x$  with orientation of  $\delta_i$ )

Not difficult to see that  $H_n^\Delta(M) = \mathbb{Z} \cdot [M]$ , so  $\Rightarrow H_n(M) \cong H_n(M, M \setminus x)$


Also  $\mathbb{Z} \cong \mathbb{Z} \cdot [M]$  since  $C_{n+1}(M) = 0$  ( $\Delta^{(n+1)}$ -simplices since  $\dim M = n$ )

$M$  non-orientable  $\Rightarrow$  each facet of  $\delta_i$  appears twice in  $\partial \sum \delta_i$


$\Rightarrow \partial \sum \delta_i = 0$  over  $\mathbb{F}_2$  independently of choices of orientations of  $\delta_i$ .

Examples


1)  $S^n = \Delta^n \cup \Delta^n$    
 glue bodies




$[S^n] = \Delta_+ - \Delta_-$  if use canonical orientation we discussed  
 hence  $\partial[S^n] = \partial\Delta - \partial\Delta = 0$   
 $D^n \subseteq \mathbb{R}^n$  canonical orientation  
 $\Rightarrow S^{n-1} = \partial D^n$  using outward normal first rule




2)  $T^2 =$




$\Delta$ -complex structure (compatibly with side identifications!)



Want orientation induced by square  $\subseteq \mathbb{R}^2$



$v_1 - v_0, v_2 - v_0$  positive  $\mathbb{R}^2$ -basis  
 $\Rightarrow \delta_1$  agrees with orientation



$[T^2] = +\delta_1 - \delta_2$


$\delta_2$  orientation disagrees

RMK general feature: if two simplices are identified along a facet then the 2 respective outward normals are related by reflection.



So consistency  $\Rightarrow$  either simplices are compatibly oriented and the two induced orientations on facet are opposite or not compatibly oriented but facet orient<sup>n</sup> is same, then need sign like in example when build  $[T^2]$

3) Recall  $\Sigma_2 =$

$\Delta$ -cx structure (compatible with side identifications!):




Use the orientation induced by polygon  $\subseteq \mathbb{R}^2$   
 $\Rightarrow [\Sigma_2] = -\delta_1 - \delta_2 + \delta_3 + \delta_4 - \delta_5 + \delta_6 + \delta_3 - \delta_2$




$v_1 - v_0, v_2 - v_0$  is negative  $\mathbb{R}^2$ -basis

All simplices  $\delta_i$  have  $v_0 =$  centre of polygon

$\Rightarrow$  sign  $<$   $+$  if overedge clockwise  $---$  anti



3)  $\mathbb{RP}^2 =$  (non-orientable example)



Won't get  $\Delta$ -cx structure if you try (since get issue here)

Use the orientation induced by square  $\subseteq \mathbb{R}^2$

$\Rightarrow [\mathbb{RP}^2] = -\delta_1 + \delta_2$   
 $\partial[\mathbb{RP}^2] = -(b - a + c) + (a - b + c) = -2b + 2a \neq 0$  so not cycle in  $C_*^{CW}(\mathbb{RP}^2)$

However, working modulo 2:

$\partial[\mathbb{RP}^2] = 0 \in C_*^{CW}(\mathbb{RP}^2; \mathbb{F}_2)$  since  $2=0$  in  $\mathbb{F}_2$   
 $\Rightarrow [\mathbb{RP}^2] \in H^2(\mathbb{RP}^2; \mathbb{F}_2)$

### Degree

Def  $M, N$  oriented closed connected  $n$ -mfds,  $f: M \rightarrow N$

$$f_*: H_n(M) \rightarrow H_n(N)$$

$$[M] \mapsto \deg(f) \cdot [N] \in \mathbb{Z}$$

Local degree

Lemma If  $f^{-1}(y)$  finite, Local map like in chapter 7


$$\text{then } \deg(f) = \sum_{x \in f^{-1}(y)} \deg(f|_{x,*})$$

$$\begin{array}{c} \text{pf} \\ [M] \in H_n(M) \xrightarrow{f_*} H_n(N) \ni [N] \\ \downarrow \oplus_{x \in f^{-1}(y)} \downarrow (f_x)_* \\ \oplus_{x \in f^{-1}(y)} H_n(M, M \setminus x) \xrightarrow{(\sum \deg(f_x)_*) \cdot \mu_y^N} H_n(N, N \setminus y) \ni \mu_y^N \end{array}$$

### Examples

$$1) S^1 \rightarrow S^1, z \mapsto z^n, [S^1] \mapsto n \cdot [S^1] \text{ so } \deg = n$$

$$2) \Sigma_3 = \Sigma_3 / \mathbb{Z}_3 \text{-rotation action} \xrightarrow{q} \text{torus} = \Sigma_1$$



Explanation:

rotation symmetry

$$\text{Easy check: } \deg(q) = 3 \text{ (e.g. use local degrees)}$$

### Cultural Rmk

For  $M, N, f$  smooth, the  $\deg f = \#$  (preimages of a generic point of  $N$ )  
 Idea:  $\deg f$  tells you how many times you cover  $N$ . (almost all points work)


### Poincaré duality

FACT Theorem For  $M$  closed  $n$ -mfd


$$M \text{ oriented} \Rightarrow H^k(M) \cong H_{n-k}(M) \text{ s.t. } 1 \leftrightarrow [M]$$

$M$  non-oriented  $\Rightarrow$  same holds with  $\mathbb{F}_2$  coefficients


Sketch proof when  $M$  is a simplicial complex  $K$  (Non-examinable)



1) simplex  $\sigma = \sigma_v$  of  $K$  with barycentre  $v \leftrightarrow v^*$  vertex of  $S(K)$



$$2) \text{ht}(v) = (\text{height of } v) = \dim \sigma_v$$



3)  $\sigma$   $k$ -simplex of  $K$

dual simplex

$$\hat{\sigma} = \bigcup_{\tau \in S(K)} \tau$$

$\text{ht}(v_i)$  is min of heights of vertices of  $\tau$

Rmk:  $\bigcup_{v \in \sigma} \tau$  with  $\text{ht}(v_i) \max$  will give back  $\sigma$ .

Thus  $\hat{\sigma}, \sigma$  intersect transversely at  $v^*$ . One can also describe  $\hat{\sigma}$  as


$\hat{\sigma} = \bigcap_{v \in \sigma} \text{Star } S(K) \leftarrow$  (closed star is the union of simplices of  $S(K)$  having  $v$  as a face)

FACTS •  $\dim \hat{\sigma} = n - \dim \sigma$  ("polygonal" complex rather than  $\Delta$ -cx)

• dual cells  $\hat{\sigma}$  give a cell decomposition of  $M$

•  $\partial \hat{\sigma} = \sum_{\substack{\text{facet} \\ \sigma \neq \tau}} \pm \hat{\tau}$

need compare orientations of  $\sigma, \tau$  (+ if  $\sigma$  as a facet of  $\tau$  has boundary orientation)



4) dual chain complex

$D_{n-k} =$  free abelian group on dual chains  $\hat{\sigma}$

$H_*(M) \cong H_*(D_*, \partial_*)$  (since  $\hat{\sigma}$  give a cell decomp. of  $M$ )

5)  $\varphi: D_{n-k} \rightarrow C^k(M)$

•  $\varphi$  linear bijection ✓ where  $\sigma^*(\alpha) = \begin{cases} 0 & \text{for } k\text{-cells } \alpha \neq \sigma \\ 1 & \text{for } \alpha = \sigma \end{cases}$

• chain map:  $\xrightarrow{\text{see } \textcircled{*}}$

$\varphi(\partial \hat{\sigma}) = \varphi \sum \pm \hat{\tau} = \sum \pm \tau^*$

$\partial^* \varphi(\hat{\sigma}) = \partial^* \sigma^* = (\sigma^* \circ \partial : \tau \mapsto \sum \pm \sigma_i \xrightarrow{\text{facet } \sigma} \begin{cases} \pm 1 & \text{if one of } \sigma_i = \sigma \\ 0 & \text{else} \end{cases}$

UPSHOT  $\varphi$  is chain iso so get iso:


$H_*(M) \cong H_*(D_*, \partial_*) \xrightarrow{\varphi} H^{n-k}(M)$

Cor  $\chi$  (odd dimensional closed orientable mfd) = 0

Pf Betti numbers  $b_i = \text{rank } H_i(M) \stackrel{\text{universal coeff. thm.}}{=} \text{rank } H_i(M) \stackrel{\text{Poincaré duality}}{=} \text{rank } H_{n-i}(M)$

$\chi(M) = b_0 - b_1 + \dots + b_{\dim M-1} - b_{\dim M}$  equal.  $\square$

(Poincaré-)Lefschetz duality




Non-oriented  $\Rightarrow$  same holds with  $\mathbb{F}_2$  coefficients.  $\square$   
Pf basically same as Poincaré duality.  $\square$

Cor  $M$  compact, connected,  $\partial M \neq \emptyset \Rightarrow H^n(M) = H_0(M, \partial M) = 0$

Examples

- 1)  $D^n$   $\partial D^n = S^{n-1}$   
 $Z \cong H^0 D^n \cong H_n(D^n, S^{n-1})$  generator  $D_1, -D_2$
- 2)  $A = \text{annulus} \subseteq \mathbb{R}^2 \simeq S^1$   
 $Z \cong H^0 A \cong H_2(A, \partial A)$  generator  $\leftarrow D_1$   
 $Z \cong H^1 A \cong H_1(A, \partial A)$  generator  $\leftarrow D_2$   
 $0 \cong H^2 A \cong H_0(A, \partial A)$  (notice  $\partial D \rightarrow \partial A$ )  
Remark notice gen. of  $H_1(A)$  is  $\circlearrowleft$  which intersects gen. of  $H_1(A, \partial A)$  once transversely.



$\Rightarrow H_*^*(M, \partial M) \cong H^{2-*}(S^1 \vee S^1) = \begin{cases} \mathbb{Z}^2 & * = 1 \leftarrow \text{gen. by 2 loops} \\ \mathbb{Z} & * = 2 \leftarrow \text{gen. by } [M, \partial M] \\ 0 & \text{else} \end{cases}$



What happens in the non-compact case?

Locally finite homology (Borel-Moore)

$C_*^{lf}(X)$  allow infinite sums  $\sum_{i \in \mathbb{Z}} n_i \sigma_i$  generators of  $C_*(X)$   
 s.t. given any compact subset  $K \subseteq X$ ,  
 $\#\{n_i \neq 0 : K \cap \text{supp } \sigma_i \neq \emptyset\} < \infty$ .

Examples

$C_1^{lf}(\mathbb{R}) \ni \sum_{m \in \mathbb{Z}} \sigma_m$  

$\Rightarrow$  get cycle  $[\mathbb{R}] \in H_1^{lf}(\mathbb{R})$   $\sigma_m: I \cong [m, m+1] \subseteq \mathbb{R}$

$C_0^{lf}(\mathbb{R}) \ni [\text{point}] \ni \partial(\sigma_1 + \sigma_2 + \dots)$  is a boundary: 

exercise  $H_*^{lf}(\mathbb{R}) = \begin{cases} \mathbb{Z} & * = 1 \\ 0 & \text{else} \end{cases} (\cong H_*^{-1}(\mathbb{R}))$

FACT Theorem  $M$  orientable n-mfd  $\Rightarrow H_*^{lf}(M) \cong H_{n-*}^{lf}(M)$   
 (possibly not compact)  $\swarrow$  depends on  $\phi$

cohomology with compact supports  $H_c^*(X)$

$C_c^*(X)$ : only allow cochains  $\phi: C_* X \rightarrow \mathbb{Z}$  s.t.  $\exists$  compact  $K \subseteq X$  with  $\phi(C_*(X \setminus K)) = 0$  (vanish on chains in  $X \setminus K$ )

Example  $c \in C_*(X) \Rightarrow \phi(c) = \text{signed \# intersections of } c \text{ with } \alpha$   
 (geometric intersection #)  
 $\Rightarrow \phi \in C_c^*(X)$  since  $\phi(\alpha) = 0$  if  $\alpha \subseteq X \setminus \text{Im}(c)$

Thm  $M$  orientable n-mfd  $\Rightarrow H_*(M) \cong H_c^{n-*}(M)$   
 (possibly not compact)

Warning  $H_*^{lf}, H_c^*$  are not homotopy invariant (indeed non-trivial for  $\mathbb{R}^n$ )  
 Caused because they are not functorial. They are however functorial for proper maps  
 Mayer-Vietoris holds for  $H_c^*$  but not for  $H_*^{lf}$ .  
 (preimages of compact sets are compact)

Fact  $H_c^*(X) \cong \varinjlim H^*(X, X \setminus K)$  where compacts  $K_1 \subseteq K_2$  give  $H^*(M, M \setminus K_1) \rightarrow H^*(M, M \setminus K_2)$   
Direct limit  $\varinjlim G_i$  via maps  $G_i \rightarrow G_j$  means  $\sqcup G_i$  / identify  $g \in G_i$  with its images under those maps  
 (The indices are partially ordered & directed:  $\forall i, j, \exists k > i, j$  so can compare  $G_i, G_j$  inside  $G_k$  via  $G_i \rightarrow G_k, G_j \rightarrow G_k$ )  
Fact  $\varinjlim$  is an exact functor.

Cap product and Poincaré duality revisited

$X$  space,  $k \geq l$   $\swarrow$  (sometimes write)  $\emptyset \cap \sigma$

$n: C_k(X) \times C^l(X) \rightarrow C_{k-l}(X)$  cap product

$(\sigma: \Delta^k \rightarrow X) \cap (\phi: C_l(X) \rightarrow \mathbb{Z}) = \phi(\sigma|_{[e_0, \dots, e_l]}) \cdot \sigma|_{[e_{l+1}, \dots, e_k]}$   
 "bottom face"  $\in \mathbb{Z}$  "top face"  $\cong \Delta^{k-l} \in C_{k-l}(X)$

(easy) Properties

- $\cap$  bilinear
- $\partial(\sigma \cap \phi) = (-1)^l (\partial\sigma \cap \phi - \sigma \cap \partial^*\phi)$
- cycle  $\cap$  cocycle is cycle
- boundary  $\cap$  cocycle are boundaries
- cycle  $\cap$  coboundary

$\Rightarrow n: H_k(X) \times H^l(X) \rightarrow H_{k-l}(X)$  bilinear

Theorem (Poincaré duality) The map  $\phi \mapsto [M] \cap \phi$  gives following isos

- For  $M$  closed oriented n-mfd  $[M] \cap \cdot: H^*(M) \xrightarrow{\cong} H_{n-*}(M)$
- For  $M$  non-compact oriented n-mfd,  $[M] \cap \cdot: H_c^*(M) \xrightarrow{\cong} H_{n-*}(M)$   $\otimes$
- $[M] \cap \cdot: H^*(M) \xrightarrow{\cong} H_{n-*}^{lf}(M)$

Sketch Pf of  $\otimes$  for smooth mfd (Non-examinable)

If  $M$  smooth  $\Rightarrow \exists$  "good cover"  $U_i$  of  $M$  meaning open cover s.t.  
 FACT from Riemannian geometry ("convex neighbourhoods")  $U_i \cong \mathbb{R}^n$   
 $U_{i_1} \cap \dots \cap U_{i_k} \cong \mathbb{R}^n$  or  $\emptyset$   
 Then compute  $H_c^*(\mathbb{R}^n) \cong \begin{cases} \mathbb{Z} & * = n \\ 0 & \text{else} \end{cases}$  and  $\otimes$  holds for  $\mathbb{R}^n$ .  
 $\Rightarrow \otimes$  holds  $\forall U_i$   
 $\Rightarrow$  by naturality of  $\otimes$  and of Mayer-Vietoris get  $\otimes$  for  $\cup U_i$  finite  
 $\Rightarrow \otimes$  for  $M$ , which is  $\otimes$ .  $\square$   $\nwarrow$  use 5-lemma

General pf of Poincaré duality ← Non-examinable

Step 1: holds for  $\mathbb{R}^n$

pf  $H_c^k(\mathbb{R}^n) \cong H^k(D^n, S^{n-1}) \cong \begin{cases} 0 & k \neq n \\ \mathbb{Z} & k = n \end{cases} \cong H_{n-k}(\mathbb{R}^n)$

can make  $K$  larger by picking  $K = \text{large ball}$   
then  $H^k(\mathbb{R}^n, \mathbb{R}^n \setminus K) \cong H^k(K, \partial K)$

Pick  $\Delta$ -cx structure for  $\mathbb{R}^n$ . So  $[\mathbb{R}^n] = \sum \pm \sigma_i \leftarrow \text{sum over } n\text{-simplices}$ .

Say  $\exists$  simplex  $\sigma_0: \Delta^n \rightarrow \mathbb{R}^n$ . Define  $\phi: C_c^k(\mathbb{R}^n) \rightarrow \mathbb{Z}, \phi(\sigma_0) = \pm 1$  (other simplices) = 0

$[\mathbb{R}^n] \cap \phi = \sum \phi(\pm \sigma_i) = \phi(\pm \sigma_0) = 1$  (pick sign in  $\oplus$ )

Step 2 holds for  $A, B, A \cap B \Rightarrow$  holds for  $A \cup B$

pf Mayer-Vietoris for  $H_c^*$ , naturality, 5-lemma  $\checkmark$

Step 3 holds for  $A_i$ , and  $A_1 \subseteq A_2 \subseteq A_3 \subseteq \dots \Rightarrow$  holds for  $\cup A_i$

pf By applying ling: both sides of P.D. iso commute with limits  $\checkmark$

Step 4 holds for open subsets in  $\mathbb{R}^n$

pf Every such set is a union of convex open sets (e.g. balls)

By Step 3 enough to consider case of finite union.

By induction on  $\#$  convex open sets:

1 convex set  $U \cong \mathbb{R}^n$  via a proper homeomorphism,

now use Step 1  $\checkmark$

2 convex sets: KEY TRICK convex set  $\cap$  convex set is convex in  $\mathbb{R}^n$ !

$\Rightarrow$  use Step 2 & previous case

$k+1$  convex sets:  $A = \cup \{\text{first } k \text{ convex sets}\}, B = \text{last convex set} \Rightarrow$  use step 2

$\Rightarrow A \cap B \subseteq B$  is a union of  $k$  convex sets  $\Rightarrow$  inductive hypothesis  $\checkmark$

Step 5 holds for mfd  $M$

Consider open sets in  $M$  for which it holds.

By a Zorn's Lemma argument we get a maximal open subset  $U$  where holds.

If  $U \neq M$  pick  $p \in M \setminus U$  and nbhd  $V \cong \mathbb{R}^n$  of  $p$ . Then holds for  $U, V, U \cup V$

(note  $U \cup V \subseteq \mathbb{R}^n$  open, so Step 4 applies) so by Step 2 holds for  $U \cup V$

Contradicts maximality.  $\checkmark \square$

This page (Corollary of Poincaré duality) is non-examinable

Recall there is a well-defined evaluation of  $H^*$ -classes on  $H_*$ :

$\langle \cdot, \cdot \rangle: H_k(M; \mathbb{R}) \otimes H^k(M; \mathbb{R}) \rightarrow \mathbb{R}$   
 $\downarrow$  any representative cycle of form  $\varphi(c)$   
 $c \otimes \alpha \mapsto \langle c, \alpha \rangle = \varphi(c)$

Easy exercise  $\langle c, \alpha \cup \beta \rangle = \langle c \cap \alpha, \beta \rangle$  any  $\alpha, \beta \in H^*, c \in H_*$


Corollary of Poincaré duality

$M$  compact oriented  $n$ -mfd,  $F$  field.

$$\begin{array}{ccc} H^k(M; F) \otimes H^{n-k}(M; F) & \xrightarrow{\otimes} & F \\ \alpha \otimes \beta & \mapsto & \langle [M], \alpha \cup \beta \rangle \end{array}$$
  
is a non-singular bilinear form.

pf. By exercise,  $\langle [M], \alpha \cup \beta \rangle = \langle [M] \cap \alpha, \beta \rangle = \langle PD(\alpha), \beta \rangle$

So the following diagram commutes:



By universal coefficients,  $H^*(M; F) \cong \text{Hom}(H_*(M; F), F)$  via  $\beta \mapsto \langle \beta, \cdot \rangle$

Hence  $\langle \cdot, \cdot \rangle$  is a non-degenerate bilinear pairing.

Hence so is the pairing  $\otimes$  in the diagram.  $\square$

Remark For  $M$  non-orientable, the same holds for  $F$  of characteristic 2, eg.  $\mathbb{Z}/2$

For  $\mathbb{Z}$  coefficients it can fail if  $H^*(M) \neq \text{Hom}(H_*(M), \mathbb{Z})$ . So we define:

Betti group  $B^k(M) = H^k(M) / \text{torsion}(H^k(M))$   
 $B_k(M) = H_k(M) / \text{torsion}(H_k(M))$


By what we proved in the section on universal coefficients,  $B^1(M) \cong \text{Hom}(B_0(M), \mathbb{Z})$  whenever  $H_{q-1}(M)$  is finitely generated (which we know holds for compact mfd's)

The iso is given by  $\langle \cdot, \cdot \rangle$  again: this descends to quotients since  $\langle c, \alpha \rangle = 0 \in \mathbb{Z}$  if  $c$  or  $\alpha$  has finite order (i.e. torsion). The same proof as above yields:

$M$  compact oriented  $n$ -mfd  $\Rightarrow B^k(M) \otimes B^{n-k}(M) \rightarrow \mathbb{Z}, \alpha \otimes \beta \mapsto \langle [M], \alpha \cup \beta \rangle$   
is non-degenerate bilinear form.

Also the Remark holds.  
Example Use this to prove ex. 4(c) sheet 3. (Hint:  $H^{2k}(\mathbb{C}P^n) \cup H^{2n-2k}(\mathbb{C}P^n) = H^{2n}(\mathbb{C}P^n)$ )

# Alexander duality



(in fact, enough to assume \$X\$ is locally contractible)

\$\emptyset \neq X \subseteq S^n\$ compact subset s.t.

\$\exists\$ open neighbourhood \$N(X)\$ which deformation retracts to \$X\$

such that \$\overline{N(X)} \subseteq S^n\$ is an \$n\$-mfd with boundary.

Theorem  $\widetilde{H}_*(X) \cong \widetilde{H}^{n-*}(\overline{S^n \setminus X})$

Pf later

Example \$X \subseteq S^3\$ knot (i.e. \$X = \text{image}(S^1 \xrightarrow{\text{homeomorphism}} S^3)\$ onto the image)

\$\Rightarrow N(X) \cong \text{solid torus} \cong S^1\$

\$\Rightarrow \widetilde{H}\_0(X) = 0 = \widetilde{H}^2(S^3 \setminus X)\$

\$\widetilde{H}\_1(X) = \mathbb{Z} = \widetilde{H}^1(S^3 \setminus X)\$

\$\widetilde{H}\_2(X) = 0 = \widetilde{H}^0(S^3 \setminus X)\$


so the homology of a knot complement does not tell knots apart (always same)

## Theorem (Jordan curve Theorem)

\$C \cong S^1\$ closed curve in \$\mathbb{R}^2 \subseteq S^2\$

\$\Rightarrow \mathbb{R}^2 \setminus C\$ has 2 path-components (= connected components)

Similarly for \$S^n \cong C \subseteq \mathbb{R}^{n+1} \subseteq S^{n+1}\$.



"inside" "outside"

Pf \$S^n \cong C \subseteq \mathbb{R}^{n+1} \subseteq S^{n+1} \Rightarrow \mathbb{Z} \cong \widetilde{H}\_n(S^n) \cong \widetilde{H}^0(S^{n+1} \setminus C)\$

\$\Rightarrow H^0(S^{n+1} \setminus C) \cong \mathbb{Z}^2\$

\$\Rightarrow S^{n+1} \setminus C\$ has 2 path components. \$\square\$

e.g. by stereographic projection \$S^2 \cong \mathbb{C} \cup \infty\$

Alexander duality

# Proof Alexander duality Abbreviate \$N = N(X)\$ (mfd of \$X\$ which is \$\cong X\$)

\$Y := S^n \setminus N \cong S^n \setminus X\$

for \$\* \leq n-1\$

\$\widetilde{H}^{n-\*}(\overline{Y}) = H^{n-\*}(\overline{Y})\$

\$\cong H\_{\*+1}(Y, \partial Y)\$

Lefschetz

\$\cong\_{\text{exc.}} H\_{\*+1}(S^n, \overline{N})\$

\$\cong\_{\text{LES}} \widetilde{H}\_\*(\overline{N})\$  
using \$\* \leq n-1\$

for \$\* = n-1\$

\$\widetilde{H}^0(Y) \oplus \mathbb{Z} \cong H^0(Y)\$

\$\cong\_{\text{Lef.}} H\_n(Y, \partial Y)\$

\$\cong\_{\text{exc.}} H\_n(S^n, \overline{N})\$

\$\cong \widetilde{H}\_{n-1}(\overline{N}) \oplus \mathbb{Z}\$

Explanation of \$\oplus\$:

LES: \$0 \rightarrow \widetilde{H}\_n(S^n) \rightarrow H\_n(S^n, \overline{N}) \rightarrow \widetilde{H}\_{n-1}(\overline{N}) \rightarrow 0\$ is SES

\$\downarrow\$ quotient

\$\cong H\_n(S^n, S^n \setminus \infty) \cong \mathbb{Z}\$

\$\otimes\$ since each (path-)connected component of \$N\$ has non-empty boundary



\$\Rightarrow\$ Hence that quotient map gives a splitting of the SES.

for \$\* = n\$ \$H^{n-\*}(\overline{Y}) = H^{-1}(Y) = 0\$

\$H\_n(X) \cong H\_n(N) \cong H^0(N, \partial N) = 0 \cdot \square\$

\$\uparrow\$ Lefschetz duality

see \$\otimes\$