

## Problem sheet 3

## General Relativity II, Hilary Term 2020

Questions marked with a star have lowest priority to be discussed during class. Any comments or corrections please to Jan.Sbierski@maths.ox.ac.uk.

- 1) Consider the linearised Einstein equations in the wave gauge

$$\begin{aligned}\square \bar{h}_{\mu\nu} &= -16\pi T_{\mu\nu}^{(1)} \\ \partial^\mu \bar{h}_{\mu\nu} &= 0,\end{aligned}$$

where we recall that that  $\bar{h}_{\mu\nu} = \tilde{h}_{\mu\nu} - \frac{1}{2}\eta_{\mu\nu}\tilde{h}$ , where  $\eta$  is the Minkowski metric. Now consider a point mass (modelling a spherically symmetric body) of mass  $\varepsilon M$  which is at rest at the coordinate origin  $x = y = z = 0$ . The corresponding stress energy tensor is given by  $T_{\mu\nu} = \varepsilon T_{\mu\nu}^{(1)} = \varepsilon M \delta^3(\underline{x}) U_\mu U_\nu$ , where  $U = \partial_t$  is the four velocity of the particle. Derive the external gravitational field  $g_{\mu\nu} = \eta_{\mu\nu} + \varepsilon \tilde{h}_{\mu\nu}$  under the assumption that it is stationary and compare it to the Schwarzschild metric with mass  $\varepsilon M$  in the region where  $\frac{\varepsilon M}{\sqrt{x^2+y^2+z^2}}$  is small.

- 2) According to the Quadrupole formula the asymptotically leading order terms of the metric corrections  $\tilde{h}_{\mu\nu}$  of a radiating system in wave gauge take the form

$$\begin{aligned}\bar{h}_{00}(t, \underline{x}) &\simeq \frac{4M}{r} + \frac{2x_i x_k}{r^3} \frac{d}{dt} Q_{ik}(t-r) \\ \bar{h}_{0i}(t, \underline{x}) &\simeq -\frac{2x_k}{r^2} \frac{d^2}{dt^2} Q_{ik}(t-r) \\ \bar{h}_{ij}(t, \underline{x}) &\simeq \frac{2}{r} \frac{d^2}{dt^2} Q_{ij}(t-r).\end{aligned}$$

Recall that  $\bar{h}_{\mu\nu} = \tilde{h}_{\mu\nu} - \frac{1}{2}\eta_{\mu\nu}\tilde{h}$ , where  $\eta$  is the Minkowski metric. Compute the metric corrections  $\tilde{h}_{\mu\nu}$  and show that the curvature component  $R^i{}_{00j}$  of  $g_{\mu\nu} = \eta_{\mu\nu} + \varepsilon \tilde{h}_{\mu\nu}$  to leading order in  $0 < \varepsilon \ll 1$  and to leading order in  $\frac{1}{r}$  is given by

$$R^i{}_{00j}(t, \underline{x}) \simeq \frac{\varepsilon}{r} \left[ \Pi^m{}_i \Pi^n{}_j - \frac{1}{2} \Pi^{mn} \Pi_{ij} \right] \frac{d^4}{dt^4} Q_{mn}(t-r),$$

where  $\Pi^{mn} = \delta^{mn} - \frac{x_n x_m}{r^2}$ . What is the interpretation of  $\Pi^n{}_n$ ?

- 3) Consider two stars, each of mass  $m$  and modelled as point particles, moving in a circular *Newtonian* orbit of radius  $R$  in the  $x, y$ -plane centred at the origin.

- (a) Show that their trajectories may be taken to be  $\gamma_\pm(t) = \pm(t, R \cos(\omega t), R \sin(\omega t), 0)$ , where  $\omega^2 = \frac{m}{4R^3}$ .  
 (b) Consider the corresponding stress-energy tensor

$$T^{\mu\nu} = \sum_{a=\pm} m \int \dot{\gamma}_a^\mu \dot{\gamma}_a^\nu \delta^4(x^\alpha - \gamma_a^\alpha(\tau)) d\tau,$$

where  $\tau$  is proper time of the particles and  $\dot{\gamma}_a^\mu = \frac{d}{d\tau} \gamma_a^\mu$ . Compute the stress-energy tensor and the quadrupole moment in the slow motion approximation.

- (c) Recall that our derivation of the quadrupole formula required the system to be non-self-gravitating, which is violated in this scenario. However, assume now that it still serves as a good approximation even in the weakly-self-gravitating case if all other assumptions made in its derivation are met. What restrictions does this impose on the parameters  $m$  and  $R$ ?

Consider the stars to be each of one solar mass  $\approx 2 \cdot 10^{30} kg$ . Give an order of magnitude estimate on  $R$  for which we might expect the quadrupole formula to be a good approximation.

- (d) (\*) Compute the metric corrections  $\bar{\bar{h}}_{\mu\nu}$  according to the quadrupole formula.

- 4) This question recalls the derivation of electromagnetic radiation and thus makes explicit the differences and similarities between the gravitational and electromagnetic cases. Let  $(M, g)$  be the 3 + 1-dimensional Minkowski spacetime with canonical coordinates  $x^\mu$ . Maxwell's equation read  $dF = 0$  and  $\partial_\mu F^{\mu\nu} = 4\pi J^\nu$ , where  $J$  is the source 4-vector and  $F$  is the Faraday tensor, an antisymmetric 2-form.

- (a) Show that  $J$  satisfies the conservation law  $\partial_\mu J^\mu = 0$  by virtue of the Maxwell equations.  
 (b) We now consider a source  $J$  that is compactly supported in space. Show that the charge  $q(t) := \int_{\mathbb{R}^3} J^0(t, \underline{x}) d\underline{x}$  is independent of time.  
 (c) Consider now the Maxwell equations in the Lorentz gauge

$$\begin{aligned}\square \tilde{A}_\mu &= 4\pi J_\mu \\ \partial^\mu \tilde{A}_\mu &= 0\end{aligned}$$

where  $\tilde{A}$  is a one-form such that  $d\tilde{A} = F$ . Follow the approach taken in the gravitational case in the lectures to derive the following expression for the far field (i.e. for large  $r = |\underline{x}|$ )

$$\begin{aligned}\tilde{A}_0(t, \underline{x}) &\simeq \frac{q}{r} + \frac{x_i}{r^2} \frac{d}{dt} D_i(t-r) \\ \tilde{A}_i(t, \underline{x}) &\simeq -\frac{1}{r} \frac{d}{dt} D_i(t-r),\end{aligned}$$

where  $D_i(t) := \int_{\mathbb{R}^3} J^0(t, \underline{x}) x_i d\underline{x}$  is the *dipole moment*. Spell out the assumptions under which the above approximation is valid.

- (d) (\*) Compute the leading order of the electric and magnetic field in the far region and show that they (the leading orders) are orthogonal.  
 (e) (\*) **(Distributional part is optional)** Consider a point particle with charge  $e$  oscillating according to

$$t \mapsto (t, 0, 0, L \sin(\omega t)).$$

The associated current 4-vector is given by  $J^\mu = e \int \delta^4(x^\nu - \gamma^\nu(\tau)) \dot{\gamma}^\mu(\tau) d\tau$ , where  $\tau$  is the proper time of the particle and  $\dot{\gamma}^\mu = \frac{d}{d\tau} \gamma^\mu$ . Show that  $\partial_\mu J^\mu = 0$  in the sense of distributions.

Assume now that the particle is moving slowly compared to the speed of light, i.e.  $0 < \omega \ll 1$ . Compute in this approximation the dipole moment  $D_i(t)$ .

- 5) (\*) Let  $(M, g)$  be a  $(d + 1)$ -dimensional Lorentzian manifold and let  $f : M \rightarrow \mathbb{R}$  be a function such that for  $f^{-1}(0) =: \Sigma$  we have  $df \neq 0$  on  $\Sigma$ , i.e.,  $\Sigma$  is a hypersurface. Let now  $\Sigma$  be a *null* hypersurface, i.e., we have in addition  $g^{-1}(df, df) = 0$  on  $\Sigma$ .

Show/recall that one can locally introduce coordinates  $\{x_0, \dots, x_d\}$  such that  $x_0 = f$ . Let

$$g_{\mu\nu} = \begin{pmatrix} g_{00} & g_{01} & \cdots & g_{0d} \\ g_{10} & & & \\ \vdots & & g_{ij} & \\ g_{d0} & & & \end{pmatrix}$$

be the components of  $g$  in these coordinates. Show that  $\det(\{g_{ij}\}_{1 \leq i, j \leq d}) = 0$ .

---

- 6) Let  $M = \mathbb{R} \times (r_+, \infty) \times \mathbb{S}^2$  with the standard  $\{t, r, \theta, \varphi\}$  coordinates where  $r_+ = M + \sqrt{M^2 - a^2}$ ,  $M > 0$ , and  $0 < a < M$ . We define the Kerr metric  $g$  on  $M$  by

$$g = -\left(1 - \frac{2Mr}{\rho^2}\right) dt^2 - \frac{2Mra \sin^2 \theta}{\rho^2} (dt \otimes d\varphi + d\varphi \otimes dt) + \frac{\rho^2}{\Delta} dr^2 \\ + \rho^2 d\theta^2 + \left(r^2 + a^2 + \frac{2Mra^2 \sin^2 \theta}{\rho^2}\right) \sin^2 \theta d\varphi^2,$$

where  $\rho^2 = r^2 + a^2 \cos^2 \theta$  and  $\Delta = r^2 - 2Mr + a^2$ . Show that the vector field  $\partial_t$  is a Killing vector field and that it is timelike for  $r > M + \sqrt{M^2 - a^2 \cos^2 \theta}$ . Also show that  $\partial_t$  is not hypersurface orthogonal. Thus  $(M, g)$  is stationary but not static.