

# String Theory II: Assignment 3

## 1. Spin Connections and Killing Spinors

[Conventions for this questions are as in BLT Section 14.8].

The spin connection  $\omega$  is a connection for the local Lorentz symmetry in a given representation and can be expanded in terms of 1-forms

$$\omega = \omega_\mu(x) dx^\mu. \quad (1)$$

As for gauge connections in Yang-Mills theory we can define the curvature 2-form as

$$\mathcal{R} = d\omega + \omega \wedge \omega. \quad (2)$$

Infinitesimal local Lorentz transformations map

$$\delta_\Lambda \omega = d\Lambda + [\omega, \Lambda] \quad (3)$$

and so  $\delta_\Lambda \mathcal{R} = [\mathcal{R}, \Lambda]$ .

Let  $e_\mu^a$  be the viel-bein where  $a, b, \dots$  are the flat indices and  $\mu, \nu, \dots$  the curved indices. Let  $\nabla_\mu$  be the covariant derivative with Christoffel symbols

$$\Gamma_{\mu\nu}^\rho = \frac{1}{2} g^{\rho\sigma} (\partial_\mu g_{\sigma\nu} + \partial_\nu g_{\mu\sigma} - \partial_\sigma g_{\mu\nu}).$$

The spin connection is then the 1-form valued in the local Lorentz algebra ( $\omega$  has a  $\mu$  curved index, and is a matrix valued object with  $a, b$  flat indices) that satisfies

$$\nabla_\mu e_\nu^a = \partial_\mu e_\nu^a - \Gamma_{\mu\nu}^\rho e_\rho^a + \omega_{\mu b}^a e_\nu^b = 0 \quad (4)$$

In components it is given by

$$\omega_\mu^{ab} = \frac{1}{2} (\Omega_{\mu\nu\rho} - \Omega_{\nu\rho\mu} + \Omega_{\rho\mu\nu}) e^{\nu a} e^{\rho b} \quad \text{where} \quad \Omega_{\mu\nu\rho} = (\partial_\mu e_\nu^a - \partial_\nu e_\mu^a) e_{a\rho}. \quad (5)$$

- (a) Determine the components of the curvature 2-tensor  $R_{\mu\nu}{}^{ab}$  in terms of  $\omega$ .
- (b) Consider 10d spacetime  $\mathbb{R}^{1,3} \times M_6$ , with local Lorentz group  $SO(1,3) \times SO(6)$  in the spinor representation as in the lecture (i.e. the generators are  $\Gamma^{ab}$ ). Let  $\epsilon$  be a **16** component spinor of  $SO(1,9)$ . Compute  $[\nabla_\mu, \nabla_\nu]\epsilon$ .

## 2. Kähler Manifolds, Projective Space

- (a) Let  $X_n$  be a complex  $n$ -dimensional Kähler manifold. Determine the non-trivial Christoffel symbols and the Ricci tensor for  $X_n$ .
- (b) Consider  $\mathbb{P}^n = \mathbb{C}^{n+1} / \sim$ ,  $n$ -dimensional projective space, where the equivalence relation is  $(z^0, \dots, z^n) \sim (w^0, \dots, w^n)$  if  $\exists \lambda \neq 0$  such that  $(z^0, \dots, z^n) = \lambda(w^0, \dots, w^n)$ . An open cover is given by  $U_r = \{z^r \neq 0\}$ , with local coordinates in  $U_r$  given by  $z_{(r)}^i$ ,  $i = 1, \dots, n$
- Determine charts  $\phi_r$  and transition functions  $\phi_r \circ \phi_s^{-1}$  which make this into a complex manifold.
  - Determine the metric that follows from the Kähler potential in the open patch  $U_r$

$$K_{(r)} = \log\left(1 + \sum_{i=1}^n |z_{(r)}^i|^2\right)$$

This is the Fubini–Study metric for  $\mathbb{P}^n$ .

- Compute the Ricci form for this metric.
- Show that for  $n = 1$  the space is (as a real manifold) a 2-sphere  $S^2$ .

## 3. Calabi-Yau Manifolds

- (a) Consider the Type IIB supergravity in 10d compactified on a Calabi-Yau three-fold  $W_6$  – with Hodge numbers  $h^{p,q}(W_6)$ , in particular  $h^{1,1}(W_6)$ ,  $h^{1,2}(W_6)$ . Determine the bosonic field content by expanding the 10d fields into harmonic forms along  $W_6$ . Confirm that the massless spectrum agrees with this of IIA on the mirror Calabi-Yau  $M_6$ , where  $h^{1,1}(W_6) = h^{1,2}(M_6)$  and  $h^{1,2}(W_6) = h^{1,1}(M_6)$ .
- (b) Consider now a Calabi-Yau 2-fold (real 4d space), also called a K3-surface. The Hodge diamond is completely fixed in this case and has non-trivial entries

$$h^{2,2} = h^{0,0} = h^{2,0} = h^{0,2} = 1, \quad h^{1,1} = 20. \quad (6)$$

By Hodge duality, the (1,1) forms are self-dual.

- Determine the degrees of freedom that the following bosonic fields have in 6d: scalar, anti-symmetric tensor, graviton, vector.
- IIB has two spinors  $\epsilon, \epsilon'$  in the **16**. Decompose the spinors appropriate for a compactification to  $\mathbb{R}^{1,5} \times M_4$ , where  $M_4$  is (1) a generic 4d manifold and (2) a K3 surface. What is the supersymmetry of the 6d theory obtained from IIB on K3?
- By expanding the IIB bosonic supergravity fields determine the massless spectrum of IIB on K3 (note: be careful about the self-duality of  $F_5$ .)