

Quantum Theory

Sheet 4 — MT20

1. A particle of mass m and charge e moving in a plane perpendicular to a magnetic field B has Hamiltonian

$$H = \frac{1}{2m} \left((P_1 + \frac{1}{2}eBX_2)^2 + (P_2 - \frac{1}{2}eBX_1)^2 \right) ,$$

where we suppose that $eB \neq 0$ is constant. Show that the energy levels have the form

$$E = E_n = \frac{|eB|\hbar}{m} \left(n + \frac{1}{2} \right) ,$$

where n is a non-negative integer.

[Hint: Introduce new operators P and X , proportional to $P_1 + \frac{1}{2}eBX_2$ and $P_2 - \frac{1}{2}eBX_1$, and show that the given Hamiltonian takes the same form as the harmonic oscillator Hamiltonian for a suitable choice of angular frequency ω .]

2. The *spin representation* of angular momentum has $j = 1/2$, with angular momentum matrices $J_i = S_i$ given by

$$S_1 = \frac{1}{2}\hbar \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} , \quad S_2 = \frac{1}{2}\hbar \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} , \quad S_3 = \frac{1}{2}\hbar \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} .$$

Here $J_3 = S_3$ has eigenvalues $\pm \frac{1}{2}\hbar$, with corresponding eigenstates $\psi_+ = (1, 0)^T$, $\psi_- = (0, 1)^T$, which are called *spin up* and *spin down* states. Introducing the unit vector $\mathbf{n} = (\sin \theta \cos \phi, \sin \theta \sin \phi, \cos \theta)$ in spherical polar coordinates, we define the operator $S_{\mathbf{n}} \equiv \sum_{i=1}^3 n_i S_i$, which is the component of spin angular momentum in the direction $\mathbf{n}$.

- (a) Show that

$$S_{\mathbf{n}} = \frac{\hbar}{2} \begin{pmatrix} \cos \theta & \sin \theta e^{-i\phi} \\ \sin \theta e^{i\phi} & -\cos \theta \end{pmatrix} .$$

- (b) Hence, or otherwise, show that $(S_{\mathbf{n}})^2 = \frac{\hbar^2}{4} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$, and deduce that $S_{\mathbf{n}}$ has eigenvalues $\pm \frac{\hbar}{2}$. Verify that the corresponding normalized eigenstates of $S_{\mathbf{n}}$ are

$$\begin{aligned} \psi_{\mathbf{n},+} &= \cos \frac{\theta}{2} \psi_+ + \sin \frac{\theta}{2} e^{i\phi} \psi_- , \\ \psi_{\mathbf{n},-} &= \sin \frac{\theta}{2} \psi_+ - \cos \frac{\theta}{2} e^{i\phi} \psi_- . \end{aligned}$$

- (c) The spin angular momentum of S_3 is measured for a quantum system, obtaining the value $\frac{1}{2}\hbar$ (so that this is “spin up” along the z -axis). The spin angular momentum of $S_{\mathbf{n}}$ is now measured. Find the probabilities for obtaining the values $\frac{1}{2}\hbar$ and $-\frac{1}{2}\hbar$.

3. Recall that the orbital angular momentum operators L_i are in spherical polar coordinates given by

$$L_{\pm} = \pm \hbar e^{\pm i\phi} \left(\frac{\partial}{\partial \theta} \pm i \cot \theta \frac{\partial}{\partial \phi} \right), \quad L_3 = -i\hbar \frac{\partial}{\partial \phi},$$

where $L_{\pm} = L_1 \pm iL_2$ are raising and lowering operators.

- (a) Using the fact that $L_+ Y_{\ell,\ell}(\theta, \phi) = 0$, hence show that $Y_{\ell,\ell}(\theta, \phi) = a_{\ell}(\sin \theta e^{i\phi})^{\ell}$, where a_{ℓ} is a normalization constant.
- (b) By applying the lowering operator L_- appropriately, hence find $Y_{\ell,m}(\theta, \phi)$ for $(\ell, m) = (1, 1), (1, 0), (1, -1), (2, 2), (2, 1)$ and $(2, 0)$, up to overall normalization constants that you may ignore.
4. In a two-dimensional model of the hydrogen atom, the stationary state Schrödinger equation takes the form

$$-\frac{\hbar^2}{2m} \left[\frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial \psi}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 \psi}{\partial \phi^2} \right] - \frac{e^2}{4\pi\epsilon_0 r} \psi = E\psi,$$

where (r, ϕ) are polar coordinates.

- (a) By separating the equation via $\psi(r, \phi) = R(r)\Phi(\phi)$, show that $\Phi(\phi)$ is a constant linear combination of $e^{i\ell\phi}$ and $e^{-i\ell\phi}$, where ℓ is a non-negative integer. [*Hint: use the fact that $\Phi(\phi + 2\pi) = \Phi(\phi)$.*]
- (b) By further substituting $R(r) = f(r)e^{-\kappa r}$, where $\kappa = \sqrt{-2mE}/\hbar$, show that the radial equation becomes

$$f'' + \left(\frac{1}{r} - 2\kappa \right) f' - \left(\frac{\ell^2}{r^2} + \frac{\kappa - \beta}{r} \right) f = 0,$$

where β is a constant you should identify.

- (c) By substituting a generalized power series expansion for f , of the form $f(r) = \sum_{k=0}^{\infty} a_k r^{k+c}$, argue that $c = \ell$ for a non-singular wave function, and in this case hence deduce the recurrence relation

$$a_k = \frac{2\kappa(k + \ell) - \kappa - \beta}{(k + \ell)^2 - \ell^2} a_{k-1}.$$

- (d) Hence or otherwise show that the energy levels are of the form $E_n = -\nu/(2n+1)^2$, where ν is a positive constant and n is a non-negative integer. [*Hint: appeal to normalizability to make the series terminate.*] What is the degeneracy of each energy level?

5. (a) (i) Making use of any results you need from the lecture notes, show that the $(N, \ell, m) = (2, 1, 0)$ state of the hydrogen atom has wave function

$$\psi(r, \theta, \phi) = B r e^{-r/2a} \cos \theta ,$$

where a is the Bohr radius and B is a normalization constant.

- (ii) By normalizing $\psi = \psi(r, \theta, \phi)$, show that we may take $B = \sqrt{\frac{1}{32\pi a^5}}$.
 (iii) Compute $\mathbb{E}_\psi(r)$, the expected value of the distance of the electron from the nucleus in this excited state.
- (b) Recall that the normalized ground state wave function for an electron in a hydrogen-like atom with Z protons and any number A neutrons in the nucleus is

$$\psi = \sqrt{\frac{Z^3}{\pi a^3}} e^{-Zr/a} ,$$

where a is the Bohr radius and r is distance from the nucleus.

An electron is in the ground state of tritium ($Z = 1, A = 2$). A nuclear reaction (β -decay) instantaneously changes the nucleus into a helium-3 ion ${}^3\text{He}^+$ ($Z = 2, A = 1$). Show that the probability of measuring the electron to be in the ground state of ${}^3\text{He}^+$ is $\frac{512}{729}$.

[In both parts you may use the integral $\int_0^\infty r^n e^{-r/b} dr = b^{n+1} n!$, where $b > 0$ and $n \in \mathbb{N}$.]