

A2: Metric Spaces and Complex Analysis

Sheet 6, sections 7.2-9 from the notes — MT20

1. Suppose that $f: U \rightarrow \mathbb{C}$ is a holomorphic function on a domain U .

1. Show that, if $\bar{B}(a, r) \subseteq U$ then

$$f(a) = \frac{1}{2\pi} \int_0^{2\pi} f(a + re^{it}) dt.$$

2. Suppose that $a \in U$ is such that $|f(a)|$ is a maximum for $|f|: U \rightarrow \mathbb{R}$. Show that $|f|$ must in fact be constant near a .

3. Deduce that f is constant on all of U .

[Hint: Consider the set $S = \{z \in U : |f(z)| = |f(a)|\}$.

2. 1. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be the function defined by

$$f(x) = \begin{cases} \exp(-\frac{1}{x}), & x > 0; \\ 0, & x \leq 0. \end{cases}$$

Show that f is infinitely differentiable everywhere on \mathbb{R} but is not equal to its Taylor series in any open interval centred at 0. (Hint: Show by induction that the n -th derivative for $x > 0$ (on the right at $x = 0$) is of the form $f^{(n)}(x) = p_n(1/x) \exp(-1/x)$ where $p_n(x)$ is a polynomial of degree $2n$ and is identically zero for $x \leq 0$.)

2. Show that

$$g(z) = \begin{cases} \exp(-z^{-4}), & z \neq 0; \\ 0, & z = 0. \end{cases}$$

satisfies the Cauchy-Riemann equations at $z = 0$ and hence in every point in the complex plane, but is not holomorphic on \mathbb{C} .

3. Let f be holomorphic on \mathbb{C} .

1. Prove that f is a polynomial of degree at most k if and only if there exist real constants $M, R > 0$ and an integer k such that

$$|f(z)| \leq M |z|^k \quad \text{for } |z| > R.$$

2. What holomorphic functions f satisfy $|f(z)| \leq |z|^k$ for all $z \in \mathbb{C}$?

3. Let $p(z)$ be a polynomial. What holomorphic functions f satisfy $|f(z)| \leq |p(z)|$ for all $z \in \mathbb{C}$?

4. Suppose that $f: \mathbb{C} \rightarrow \mathbb{C}$ is holomorphic on the whole complex plane (*i.e.* f is an entire function).

1. If $f(1/n) = 1/n$ for all $n \in \mathbb{N}$ must $f(z) = z$ for all $z \in \mathbb{C}$?
2. If $f(n) = n$ for all $n \in \mathbb{N}$ must $f(z) = z$ for all $z \in \mathbb{C}$?
3. Show that there must be some $n \in \mathbb{N}$ such that $f(1/n) \neq 1/(n+1)$.

5. Suppose that $f: \mathbb{C} \rightarrow \mathbb{C}$ is an entire function, and for each $z_0 \in \mathbb{C}$ the power series expansion $f(z) = \sum_{n=0}^{\infty} c_n(z - z_0)^n$ has at least one c_n equals zero. Prove that $f(z)$ is a polynomial.

[Hint: Note that $n!c_n = f^{(n)}(z_0)$ and that \mathbb{C} is uncountable.]

6. Identify the singularities of the following functions. Classify any singularities which are isolated.

$$\frac{1}{e^z - 1}, \quad \frac{\sin 2\pi z}{z^3(2z - 1)}, \quad \sin\left(\frac{1}{z}\right), \quad \bar{z}, \quad \frac{1}{\exp\left(\frac{1}{z}\right) + 2}.$$

7. Let

$$F(z) = \frac{1}{(z - 1)^2(z + 2)}.$$

Find Laurent expansions for F in

$$A_1 = D(0, 1), \quad A_2 = \{z : 1 < |z| < 2\}; \quad A_3 = \left\{z : \sqrt{2} < |z - i| < \sqrt{5}\right\}.$$

8. (*Optional:*) A famous theorem of Weierstrass shows that any continuous function $f: [0, 1] \rightarrow \mathbb{R}$ can be uniformly approximated arbitrarily closely by a polynomial, in that given any $\epsilon > 0$ there is a polynomial $p_\epsilon: [0, 1] \rightarrow \mathbb{R}$ such that $|f(t) - p_\epsilon(t)| < \epsilon$ for all $t \in [0, 1]$. If we let $B = \bar{B}(0, 1)$ be the closed unit disk in \mathbb{C} , is it true that any continuous function $f: B \rightarrow \mathbb{C}$ on B can be uniformly approximated by polynomials?