

Questions marked as optional are not expected to be handed in and will not be marked.

1. Suppose that $\phi(x, t; \tau)$ is the solution of

$$\phi_{tt} - \phi_{xx} = 0, \quad \text{with} \quad \phi(x, \tau; \tau) = 0, \quad \phi_t(x, \tau; \tau) = g(x, \tau).$$

Construct a solution for ϕ involving an integral of g and verify that

$$u(x, t) = \int_0^t \phi(x, t; \tau) d\tau$$

satisfies the inhomogeneous problem

$$u_{tt} - u_{xx} = g(x, t) \quad \text{with} \quad u(x, 0) = u_t(x, 0) = 0.$$

2. Consider the problem

$$\nabla^2 \phi = 0 \quad \text{for } 1 \leq r \leq 2, \quad \text{with} \quad \alpha \phi + \frac{\partial \phi}{\partial r} = \begin{cases} k \cos \theta & \text{on } r = 1, \\ 0 & \text{on } r = 2. \end{cases}$$

for constant α .

- (i) If $\alpha = 0$, show that the solvability condition for existence of a solution is satisfied.
(Use an extension of Green's theorem for a non-simply-connected domain.)
(ii) Show that the homogeneous problem (i.e. $k = 0$) has a solution $\phi = \phi(r)$ if

$$\alpha = 0 \quad \text{or} \quad \alpha = \frac{1}{2 \log 2}.$$

- (iii) By seeking a solution of the homogeneous problem of the form $\phi = f(r)g(\theta)$, show that that there are countably infinitely many such solutions (i.e. countably infinitely many α for which such solutions exist).
(iv) What can you say about existence and uniqueness of the inhomogeneous problem as a function of α ?
3. Construct the Green's function for Laplace's equation in the domain $x > 0, y > 0$ with Neumann boundary data. Hence, assuming suitable behaviour at infinity, give a solution to the problem

$$\begin{aligned} \nabla^2 u &= f & \text{in } x > 0, y > 0, \\ \frac{\partial u}{\partial x} &= g(y) & \text{on } x = 0, \\ \frac{\partial u}{\partial y} &= h(x) & \text{on } y = 0. \end{aligned}$$

4. [This question is optional.] Let w be the difference between two solutions of

$$\begin{aligned} \frac{\partial u}{\partial t} &= \nabla^2 u + au + f(\mathbf{x}, t) & \mathbf{x} \in \Omega, t > 0 \\ \nabla u \cdot \mathbf{n} + \alpha u &= g(\mathbf{x}, t) & \mathbf{x} \in \partial\Omega \\ u &= h(\mathbf{x}) & \text{at } t = 0 \end{aligned}$$

where a and α are constants with $\alpha > 0$, and \mathbf{n} is the unit normal on $\partial\Omega$. Derive the relation

$$\frac{d}{dt} \int_{\Omega} w^2 d\mathbf{x} + 2 \int_{\Omega} (|\nabla w|^2 - aw^2) d\mathbf{x} + 2\alpha \int_{\partial\Omega} w^2 dS = 0$$

Deduce that

$$\left(\frac{d}{dt} + \text{constant} \right) \int_{\Omega} w^2 d\mathbf{x} \leq 0$$

and thus that $w \equiv 0$.

5. Consider the equation

$$u_t = u_{xx}, \quad x > 0, \quad t > 0$$

with

$$u(x, 0) = 0, \quad u(0, t) = f(t), \quad u \rightarrow 0 \quad \text{as } x \rightarrow \infty.$$

- (i) Explain why this admits a *similarity solution* when f is constant. Thus obtain the solution $u = u_0(x, t)$ corresponding to $f \equiv 1$.

You may find useful the formula

$$\int_0^\infty e^{-s^2/4} ds = \sqrt{\pi}.$$

- (ii) Use the Green's function approach to show that the solution for arbitrary $f(t)$ may be written in the form

$$u = \int_0^t f(t-s) \frac{\partial u_0}{\partial t}(x, s) ds.$$

6. Find a similarity solution of the equation, for constant $\alpha \in (0, 1)$,

$$u_t = x^\alpha u_{xx}, \quad \text{for } x, t > 0,$$

which also satisfies the boundary conditions

$$u(0, t) = 0, \quad u(x, 0) = T_0 > 0, \quad \text{and } u \rightarrow T_0 > 0 \quad \text{as } x \rightarrow \infty.$$

7. Find a non-trivial similarity solution of the equation

$$u_t = (uu_x)_x \quad \text{in } 0 < x < t^{1/3}, \quad t > 0,$$

where

$$u_x(0, t) = 0 = u(t^{1/3}, t) \quad \text{for } t > 0.$$

Show that

$$\int_0^{t^{1/3}} u(x, t) dx = \text{constant for } t > 0.$$