

STOCHASTIC DIFFERENTIAL EQUATIONS
MATH C8.1 - 2019 - SHEET 4

1. SHEET 4

- (i) Let $(M_t)_{t \in [0, \infty)}$ be a continuous local martingale that vanishes at zero. Let $(\mathcal{E}(M)_t)_{t \in [0, \infty)}$ denote the stochastic exponential.
- (a) Show that $(\mathcal{E}(M)_t)_{t \in [0, \infty)}$ is a nonnegative continuous local martingale.
 - (b) Show that $(\mathcal{E}(M)_t)_{t \in [0, \infty)}$ is a supermartingale with $\mathbb{E}[\mathcal{E}(M)_t] \leq 1$ for every $t \in [0, \infty)$.
 - (c) Show that $(\mathcal{E}(M)_t)_{t \in [0, \infty)}$ is a continuous martingale if and only if $\mathbb{E}[\mathcal{E}(M)_t] = 1$ for every $t \in [0, \infty)$.

- (ii) The following is Kazamaki's criterion. Let $(L_t)_{t \in [0, \infty)}$ be a continuous local martingale. Prove that if $(\exp(\frac{1}{2}L_t))_{t \in [0, \infty)}$ is a uniformly integrable submartingale, then the stochastic exponential $(\mathcal{E}(L)_t)_{t \in [0, \infty)}$ is a uniformly integrable martingale.

Hint: Show for every $\alpha \in (0, 1)$ that

$$\mathcal{E}(\alpha L_t) = (\mathcal{E}(L)_t)^{\alpha^2} (Z_t^\alpha)^{1-\alpha^2},$$

for $Z_t^\alpha = \exp(\frac{\alpha}{1+\alpha}L_t)$. Use Hölder's inequality, Question (i), and the optional stopping theorem to prove that, for every stopping time T , for every $A \in \mathcal{F}$,

$$\mathbb{E}[\mathcal{E}(\alpha L_T)\mathbf{1}_A] \leq \mathbb{E}[Z_T^\alpha \mathbf{1}_A]^{1-\alpha^2}.$$

Conclude using the assumption and $\alpha \in (0, 1)$ that $(Z_t^\alpha)_{t \in [0, \infty)}$ is a uniformly integrable submartingale, and therefore that the family

$$\{\mathcal{E}(\alpha L_T) : T \text{ is a stopping time}\} \text{ is a uniformly integrable,}$$

and therefore that $(\mathcal{E}(\alpha L_t))_{t \in [0, \infty)}$ is a uniformly integrable martingale. Hence, for every $\alpha \in (0, 1)$,

$$1 = \mathbb{E}[\mathcal{E}(\alpha L_\infty)] \leq \mathbb{E}[Z_\infty^\alpha]^{1-\alpha^2}.$$

Use the assumption and the dominated convergence theorem to pass to the limit $\alpha \rightarrow 1$ to conclude that

$$\mathbb{E}[\mathcal{E}(L)_\infty] = 1,$$

and therefore using Question (i) conclude that $(\mathcal{E}(L)_t)_{t \in [0, \infty)}$ is a martingale.

- (iii) The following is Novikov's criterion. Prove that if $(L_t)_{t \in [0, \infty)}$ is a continuous local martingale which satisfies

$$\mathbb{E} \left[\exp \left(\frac{1}{2} \langle L \rangle_\infty \right) \right] < \infty,$$

then $(\mathcal{E}(L))_{t \in [0, \infty)}$ is a uniformly integrable martingale.

(Hint: Show that Novikov's criterion implies Kazamaki's criterion. First use the fact that the exponential integrability implies for every $p \in (0, \infty)$ that

$$\mathbb{E} \left[\langle L \rangle_\infty^{\frac{p}{2}} \right] < \infty.$$

Use this fact and the Burkholder-Davis-Gundy inequalities to prove that $(L_t)_{t \in [0, \infty)}$ is a uniformly integrable martingale. Then use the equality

$$\exp\left(\frac{1}{2}L_\infty\right) = \mathcal{E}(L)_\infty^{\frac{1}{2}} \exp\left(\frac{1}{4}\langle L \rangle_t\right),$$

Hölder's inequality, Question (i), and the assumptions to conclude that

$$\mathbb{E}\left[\exp\left(\frac{1}{2}L_\infty\right)\right] < \infty.$$

Use this fact to conclude that $(\exp(\frac{1}{2}L_t))_{t \in [0, \infty)}$ is a uniformly integrable submartingale, and then apply Kazamaki's criterion.

- (iv) Let $(B_t)_{t \in [0, \infty)}$ be a standard one-dimensional Brownian motion defined on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ with respect to a filtration $(\mathcal{F}_t)_{t \in [0, \infty)}$. Suppose that $b: \mathbb{R} \rightarrow \mathbb{R}$ is a bounded, measurable function. Define measures $\{\mathbb{Q}_T\}_{T \in (0, \infty)}$ on $\{(\Omega, \mathcal{F}_T, \mathbb{P})\}_{T \in (0, \infty)}$ which satisfy the following three properties.

- (a) The measures are compatible in the sense that $\mathbb{Q}_{T_1} = \mathbb{Q}_{T_2}$ on \mathcal{F}_{T_1} for every $T_1 \leq T_2 \in [0, \infty)$.
- (b) For every $T \in [0, \infty)$ the measure \mathbb{Q}_T is mutually absolutely continuous with respect to \mathbb{P} on $(\Omega, \mathcal{F}_T, \mathbb{P})$.
- (c) The process $(\tilde{B}_t)_{t \in [0, \infty)}$ defined by

$$\tilde{B}_t = B_t - \int_0^t b(B_s) ds,$$

is for every $T \in (0, \infty)$ a standard Brownian motion with respect to $(\mathbb{Q}_T, \mathcal{F}_T)$ on $[0, T]$. Prove that for every $T \in (0, \infty)$ the pair $(B_t, \tilde{B}_t)_{t \in [0, T]}$ is a weak solution to the equation

$$(1.1) \quad \begin{cases} dB_t = b(B_t) dt + d\tilde{B}_t & \text{in } (0, T), \\ B_0 = 0, \end{cases}$$

with respect to $(\mathbb{Q}_T, \mathcal{F}_T)$. Deduce that uniqueness in law holds for (1.1).

- (v) (Skorokhod's Lemma) Let $y: [0, \infty) \rightarrow \mathbb{R}$ be a continuous real-valued function that satisfies $y(0) \geq 0$. Prove that there exist unique functions $a, z: [0, \infty) \rightarrow \mathbb{R}$ which satisfy the following three properties.

- (a) We have the decomposition

$$z = y + a.$$

- (b) The function z is nonnegative

$$z \geq 0.$$

- (c) The function a is increasing, continuous, and vanishes at zero and the corresponding Riemann-Stieltjes measure da is supported on the set $\{s: z(s) = 0\}$.

Furthermore, the function a is given by

$$a(t) = \left[\sup_{s \in [0, t]} (-y(s)) \right] \vee 0.$$

- (vi) Let $(B_t)_{t \in [0, \infty)}$ be a standard one-dimensional Brownian motion. Define the process $(X_t)_{t \in [0, \infty)}$ by

$$X_t = \int_0^t \operatorname{sgn}(B_s) dB_s.$$

- (a) Show that $(X_t)_{t \in [0, \infty)}$ is a standard Brownian motion with respect to the filtration

$$\mathcal{F}_t^{|B|} = \sigma(|B_s| : s \in [0, t]).$$

Furthermore, observe that $\mathcal{F}_t^{|B|} \subsetneq \mathcal{F}_t^B$.

- (b) Let $(L_t^0)_{t \in [0, \infty)}$ denote the local time of $(B_t)_{t \in [0, \infty)}$ at zero. Prove that

$$L_t^0 = \sup_{s \in [0, t]} (-X_s).$$

- (c) Let $(S_t)_{t \in [0, \infty)}$ be defined by $S_t = \sup_{s \in [0, t]} B_s$. Show that the two-dimensional processes $(S_t - B_t, S_t)$ and $(|B_t|, L_t^0)$ have the same law.
- (d) For every $a \in \mathbb{R}$ let $(L_t^a)_{t \in [0, \infty)}$ denote the local time of $(B_t)_{t \in [0, \infty)}$ at $a \in \mathbb{R}$. Deduce for every $a \in \mathbb{R}$ that $\mathbb{P}[L_\infty^a = \infty] = 1$.

- (vii) Let $(B_t)_{t \in [0, \infty)}$ be a standard one-dimensional Brownian motion. Prove that the stochastic differential equation

$$(1.2) \quad \begin{cases} dX_t = \operatorname{sgn}(X_t) dB_t & \text{in } (0, \infty), \\ X_0 = 0, \end{cases}$$

had a weak solution but no strong solution. Deduce that uniqueness in law holds for equation (1.2) but that pathwise uniqueness does not hold.

- (viii) Let $\alpha \in (0, 1/2)$, let $\sigma: \mathbb{R} \rightarrow \mathbb{R}$ be defined by

$$\sigma(x) = |x|^\alpha \wedge 1,$$

and let $(B_t)_{t \in [0, \infty)}$ be a standard one-dimensional Brownian motion. Show that the map

$$t \in [0, \infty) \rightarrow \int_0^t \sigma^{-2}(B_s) ds,$$

is almost surely finite. Let $(\tau_t)_{t \in [0, \infty)}$ denote the associated time-changes

$$\tau_t = \inf \left\{ s \in [0, \infty) : \int_0^s \sigma^{-2}(B_r) dr = t \right\}.$$

Show that $X_t = B_{\tau_t}$ and $X_t = 0$ are two solutions of the equation

$$\begin{cases} dX_t = \sigma(X_t) dB_t & \text{in } (0, \infty), \\ X_0 = 0. \end{cases}$$

Conclude that uniqueness in law does not hold for this equation, despite the fact that the second of these solutions is a strong solution.