Tensor products

Wednesday, 7 October 2020 18:13

let k be a field and Vec_k the category of k-vector spaces.

recall: given $V, W, L \in Vec_k$ denote by $Bilin(V,W;L)$ the set of bilinear transformations $V \times W \to L$

Can we write this using category theory language?

Note that if $V, W \in Vec_k$ then $Hom_{Vec_k}(V,W)$ $(= Hom_k(V,W))$ is actually a k-vector space. let's denote this k-v.s. by $\underline{Hom}(V,W) \in Vec_k$

Note: the category Vec_k is enriched over itself.

$$Bilin(V,W;L) \cong Hom_{Vec}(V, \underline{Hom}(W,L)) \cong Hom_{Vec_k}(W, \underline{Hom}(V,L))$$

$\underline{Hom}(-,-)$ inner Hom.

Question: Is $Bilin(V,W;-): Vec_k \to Set$ representable? In other words, is there a v.s. $V \otimes_k W$ with a natural iso. of functors $Bilin(V,W;-) \simeq Hom_{Vec_k}(V \otimes_k W, -)$?

This is equivalent to asking for the universal bilinear transformation: a v.s. $V \otimes_k W$ with a bilinear map $V \times W \to V \otimes_k W$ s.t. for any other bilinear map

$$\begin{tikzcd} & V \otimes_k W \arrow[d, "\exists!"] \\ V \times W \arrow[ur] \arrow[r] & L \end{tikzcd}$$

commutes.

This is equivalent to: asking that the functor $\underline{Hom}(W,-): Vec_k \to Vec_k$ has a left adjoint $- \otimes_k W: Vec_k \to Vec_k$

$$Hom_{Vec_k}(- \otimes_k W, -) \cong Hom_{Vec_k}(-, \underline{Hom}(W,-)) \cong Bilin(-,W;-)$$

Such a v.s. exists: look at the v.s. spanned by $V \times W$ $k[V \times W]$ and look at the subspace spanned by

$(v+v', w) - (v,w) - (v',w)$
$(v, w+w') - (v,w) - (v,w')$
$(\lambda v, w) - \lambda(v,w)$ $\quad (v, \lambda w) - \lambda(v,w)$

call it M

$$\underbrace{V \times W \to k[V \times W] \to k[V \times W]/M}_{bilinear}$$

$$Hom_{Vec_k}(k[V \times W]/M, L) = Bilin(V,W;L)$$

so $V \otimes_k W$ exists.

Note: $V \otimes_k W$ is given up to a unique iso.

$$\begin{tikzcd} & V \otimes_k W \arrow[d, shift left, "\exists!"] \\ V \times W \arrow[ur] \arrow[r] & (V \otimes_k W)' \arrow[u, shift left, "\exists!"] \end{tikzcd}$$

iso.

Since $Bilin(V,W;L) \simeq Bilin(W,V;L)$ we get $V \otimes_k W \simeq W \otimes_k V$

$k \otimes_k W \simeq W \otimes_k k \simeq W$

$Trilin(V_1,V_2,V_3;L) \simeq$
$Bilin(V_1 \otimes_k V_2, V_3; L) \simeq$
$Bilin(V_1, V_2 \otimes_k V_3; L)$

so there is a canonical iso.

$$(V_1 \otimes_k V_2) \otimes_k V_3 \simeq V_1 \otimes_k (V_2 \otimes_k V_3)$$

we get a closed symmetric monoidal category

$$(Vec_k, - \otimes_k -, \underline{Hom}(-,-))$$

Note: a k-algebra is a monoid object in $(Vec_k, \otimes_k)$: an object $A \in Vec_k$ with mult. $m: A \otimes_k A \to A$ and unit $i: k \to A$ s.t. the axioms of a monoid hold ass. unit.

$$\begin{tikzcd} A \otimes_k A \otimes_k A \arrow[r, "m \otimes id"] \arrow[d, "id \otimes m"'] & A \otimes A \arrow[d, "m"] \\ A \otimes A \arrow[r, "m"'] & A \end{tikzcd}$$

commutes.

$(ab)c = a(bc)$

Since $- \otimes_k W$ is a left adjoint it commutes with arbitrary colimits.

$$\left(\bigoplus_{i \in I} V_i\right) \otimes_k W \simeq \bigoplus_{i \in I} (V_i \otimes_k W)$$

Note: $- \otimes_k W$ doesn't commute with infinite products in general.

Non-linear version: Set

the functor $Hom_{Set}(Y,-): Set \to Set$ has a left adjoint $- \times Y$

$$Hom_{Set}(X \times Y, Z) \cong Hom_{Set}(X, Hom_{Set}(Y,Z))$$

this is an example of a Cartesian closed category

In Vec_k $\overset{binary}{product} = \overset{binary}{coproduct} = biproduct$ is $V \oplus W$, very different from $V \otimes_k W$

$dim(V \oplus W) = dim V + dim W$

$dim(V \otimes_k W) = dim V \cdot dim W.$

Exactly the same works for any commutative ring R: the category R-mod is a closed symmetric monoidal category.

$$(R\text{-mod}, - \otimes_R -, \underline{Hom}_R(-,-))$$

$$Hom_{R\text{-mod}}(M \otimes_R N, L) \simeq Hom_{R\text{-mod}}(M, \underline{Hom}(N,L))$$

$- \otimes_R M$ is a left adjoint so commutes with arbitrary (small) colimits

$R^J \to R^I \to N \to 0$ free presentation: generators and relations for the module N.

$$R^J \otimes_R M \to R^I \otimes_R M \to N \otimes_R M \to 0$$

$R^I = \bigoplus_{i \in I} R$

$R^J = \bigoplus_{j \in J} R$

$$R^I \otimes_R M = \left(\bigoplus_I R\right) \otimes_R M = \bigoplus_I (R \otimes_R M) = \bigoplus_I M = M^I$$

$$M^J \to M^I \to N \otimes_R M \to 0$$

Suppose $I \subset R$ is an ideal generated by $\{x_i\}_{i \in J}$

$$R^J \to R \to R/I \to 0$$

$$\begin{tikzcd} R^J \otimes_R M \arrow[r] \arrow[d, phantom, "\wr"] & R \otimes_R M \arrow[r] \arrow[d, phantom, "\wr"] & R/I \otimes_R M \arrow[r] & 0 \\ M^J & M \arrow[ur] & & \end{tikzcd}$$

$$R/I \otimes_R M \simeq M/IM$$

---

if R is not commutative we have

$$- \otimes_R M: \text{mod-}R \to Ab$$

$M \in R$-mod

this has a right adjoint

$$Hom_{Ab}(M,-): Ab \to \text{mod-}R$$

prop: if B is an R-S bimodule and C is a right S-module then $Hom_S(B,C)$ is a right R module

$(fr)(b) \overset{def}{=} f(rb)$

$Hom_S(B,-): \text{mod-}S \to \text{mod-}R$ is a right adjoint to $- \otimes_R B: \text{mod-}R \to \text{mod-}S$

$$Hom_S(A \otimes_R B, C) \simeq Hom_R(A, Hom_S(B,C))$$

Remark: a way to prove this is by noticing / defining

$$M \otimes_R N \overset{def}{=} Coeq(M \otimes_{\mathbb{Z}} R \otimes_{\mathbb{Z}} N \rightrightarrows M \otimes_{\mathbb{Z}} N)$$

using properties of $- \otimes_{\mathbb{Z}} -$ and Coeq.