

C2.6 Introduction to Schemes

Prof. Alexander F. Ritter
University of Oxford
HT 2021
ritter@maths.ox.ac.uk

Feedback and corrections are welcome!

References

2018-2019 Course Lecture Notes by Prof. Damian Rössler ← on course page

Ravi Vakil, The Rising Sea, Foundations of Algebraic Geometry ← online

http://stacks.math.columbia.edu ← search defines theorems/proofs in algebra & alg-geometry

Eisenbud & Harris, The Geometry of Schemes, Springer GTM 197

George R. Kempf, Algebraic Varieties, LMS Lecture notes 172

Classic books by: Mumford (Red Book of Varieties & Schemes)

Hartshorne (Algebraic Geometry)

Shafarevich (Basic Algebraic Geometry 2)

✓ or my website

My C3.4 Algebraic geometry notes (see C2.6 course webpage) try to

fill the gap between classical algebraic geometry (C3.4) and C2.6

Prerequisites

Commutative algebra (e.g. Atiyah - MacDonald, Introduction to Comm. Alg.)

Category theory — or willingness to read things up as necessary

Homological algebra — or willingness to read things up as necessary

Expectations

That you read the notes regularly after each class.

(This is a 16-lecture course, 2 lectures/week across 8 weeks.)

Not everything can be covered in detail in class, so you need to be willing to look things up as necessary.

Conventions

Diagrams commute unless we say otherwise

Ring means commutative ring with unit 1.

CONTENTS

0. INTRODUCTION

- 0.1 Classical Algebraic Geometry : Affine varieties
- 0.2 Why schemes?
- 0.3 What is a point? (reducible, irreducible)

1. DEFINITION OF SCHEMES

- 1.1 Examples of affine schemes (Spec R , $V(I)$, generic/closed point, covering trick, quasi-compact)
- 1.2 Definition of a scheme (ringed space, locally ringed space, affine scheme, scheme)
- 1.3 Pre-sheaves (pre-sheaf, morph of presheaves, sub-presheaf)
- 1.4 Sheaves (sheaf, local-to-global condition, skyscraper sheaf, $\mathcal{A}_k(X)$)
- 1.5 Stalks (stalk, direct limits, checking in/surj at stalk level)
- 1.6 Sheafification (sheafification $F \rightarrow F^+$, universal property of F^+)
- 1.7 Kernels, cokernels, images (abelian categories, additive categories, additive functor)
- 1.8 Exactness (cochain complex / cohomology in abelian cats, left/right exact)
- 1.9 Push-forward (direct image) and inverse image ($f_*F, f^{-1}F, f_!F, f^*F$)
- 1.10 Morphisms of ringed spaces (closed chain complex / cohomology in abelian cats, left/right exact)
- 1.11 A sheaf defined on a topological basis (Using $B = \{D_f\}$ for Spec R , structure sheaf $\mathcal{O}_X$, classical alg. geom.)
- 1.12 Construction of $\mathcal{O}_{\text{Spec } R}$ (Spec: Rings to equivalent Aff. fully faithful locally ringed spaces)
- 1.13 Morphisms between Specs (ideal sheaf for $I \subseteq R$ on Spec R , quasi-coherence)
- 1.14 Closed affine subschemes (sheaf of ideals on a scheme, quasi-coherence, support of a sheaf)
- 1.15 Closed subschemes (sheaf of ideals on a scheme, quasi-coherence, support of a sheaf)

2. GLOBAL SECTIONS AND THE FUNCTOR OF POINTS

- 2.0 Points of Spec R (not necessarily closed) (max ideals in local rings $\longleftrightarrow$ points)
- 2.1 Global sections and basic open sets for locally ringed spaces (X canonical, Spec $\Gamma(X, \mathcal{O}_X)$, D_f)
- 2.2 What it means to be affine (Yoneda lemma / embedding, $\text{Mor}(X, \text{Spec } R) \cong \text{Hom}(R, \Gamma(X, \mathcal{O}_X))$)
- 2.3 Functor of points $\mathbf{h}_Y$

3. PROPERTIES OF SCHEMES

- 3.0 Useful facts from commutative algebra : localisation (localisation of modules exactness)
- 3.1 Noetherian (locally Noetherian schemes, useful trick: basis $\subseteq$ overlap of affines)
- 3.2 Properties that are affine-local (locality of finite type, reduced, Noetherian)
- 3.3 Reduced schemes (stalk-local property, extending morphisms onto closures)
- 3.4 Irreducible schemes (irreducible as generic point, connectedness, irreducible components, primary decomp.)
- 3.5 Integral schemes (integral $\iff$ reduced & irreducible, injectivity of restriction, function field $K(X)$)
- 3.6 Properties of morphisms (affine quasi-compact, locally finite type, finite type, closed/open immersion, closed/open subschemes, flat, flatness & deformations, closures in Spec R)

4. GLUING THEOREMS

- 4.1 Gluing sheaves (gluing data, compatibility conditions, morphisms defined by local data)
- 4.2 Gluing schemes (gluing conditions, gluing lemma, functor of points is a sheaf of sets)
- 4.3 Affine n -space by gluing (see Homework for projective space) ($\mathbb{A}^n$ and $\mathbb{P}^n$ as representable functors)

5. PRODUCTS

- 5.0 Products in category theory (product, coproduct, category $\mathcal{C}/B$, fiber product, pushout)
- 5.1 Fiber products exist in Schemes / B (A -algebras, tensor products, fiber products in Aff & Sch)
- 5.2 Fibers and preimages (Mumford's picture, underlying topological space of products)
- 5.3 Base change (separated, universally closed, proper, projective morphism)
- 5.4 More properties of schemes (abstract varieties, complete, affine and quasi-projective vars)
- 5.5 Varieties (abstract variety, complete, affine and quasi-projective vars)
- 5.6 Scheme structure on subsets (induced scheme structure, locally closed subsets)

6. SHEAVES OF MODULES

- 6.1 $\mathcal{O}_X$ -modules
- 6.2 Modules generated by sections
- 6.3 Vector bundles and coherent modules (locally free, invertible sheaf, coherent, loc. finitely presented)
- 6.4 $\mathcal{O}_X$ -module $\tilde{M}$ on $X = \text{Spec } R$, for $R\text{-Mod}$ (locally free, invertible sheaf, coherent, loc. finitely presented)
- 6.5 Direct image and inverse image
- 6.6 Operations on $\mathcal{O}_X$ -modules
- 6.7 Pullback
- 6.8 $\tilde{M}$ on any scheme
- 6.9 Classification of $\mathcal{O}_X$ -homs $\tilde{M} \rightarrow F$ ($\text{Hom}_R(\tilde{M}, F) = \text{Hom}_R(\tilde{M}, \Gamma(X, F))$ on $X = \text{Spec } R$)
- 6.10 Flatness ($f: X \rightarrow Y$ flat $\Rightarrow f^*: \mathcal{O}_Y\text{-Mod} \rightarrow \mathcal{O}_X\text{-Mod}$ exact, flat resolutions)

7. (QUASI-) COHERENT SHEAVES

- 7.1 $\text{QCoh}(X)$ (locally finitely presented vs. coherence, coherent modules)
- 7.2 Overview of general properties of $\text{QCoh}(X)$ and $\text{Coh}(X)$ for X scheme
- 7.3 Pull-back preserves quasi-coherence
- 7.4 Pushforwards for X Noetherian
- 7.5 Gluing modules
- 7.6 $\text{QCoh}(X)$, $\text{Coh}(X)$, $\text{Vect}(X)$ for $X = \text{Spec } R$ ($R\text{-Mod} \simeq \text{QCoh}(\text{Spec } R)$, $\text{Coh } R\text{-Mod} \simeq \text{Coh}(\text{Spec } R)$)

8. ČECH COHOMOLOGY

- 8.1 Čech complex
- 8.2 Čech complex with ordering (Serre's trick)
- 8.3 Affines have no cohomology except H^0 ($H^i(\text{Spec } R, F) = 0 \forall i \geq 1$ for $F \in \text{QCoh}$)
- 8.4 Independence of cover (X separated & quasi-compact $\Rightarrow H^i_{\text{Čech}} \cong H^i_{\text{étale}}$ indep. of cover for QCoh)
- 8.5 Induced LES on H ($\Gamma(U, \cdot)$ exact on QCoh for affine U)
- 8.6 Dealing with infinite covers (refinements of covers, $H^i_{\text{Čech}}$ vs. singular cohomology)
- 8.7 Application: line bundles and $H^1(X, \mathcal{O}_X^*)$ (trivialization, vector bundle, sheaf $\mathcal{O}_X^*$ of invertible $\mathcal{O}_X$)
- 8.8 Product on Čech cohomology (Picard group, $\text{Pic}(P^1)$, $\text{Pic}(P^n)$)

9. SHEAF COHOMOLOGY

- 9.1 Resolutions (injective/projective, left/right-derived functors, "enough injectives")
- 9.2 Acyclic resolutions
- 9.3 Čech cohomology vs. Sheaf cohomology (characterization of H^i (separated quasi-compact schemes) for QCoh , separated Noeth. $\Rightarrow H^i = H^i_{\text{Čech}}$ on QCoh)
- 9.4 Product on sheaf cohomology (Serre's Theorem)

10. $\text{QCoh}(P^n)$, GRADED MODULES, $\text{PROJ}(R)$

- 10.1 Graded modules and $\text{QCoh}(P^n) \leftarrow \text{graded rings/modules}$ ($\text{Proj } R[x_0, \dots, x_n] \text{ Mod} \xrightarrow{\text{pullback}} \text{QCoh}(P^n)$)
- 10.2 $\text{Proj}(R)$ and $\text{QCoh}(\text{Proj } R)$ (line bundles via graded mods)

$$\left(\text{Proj } R, \text{ irrelevant ideal}, V(\text{graded ideal}), \mathcal{O}_{\text{Proj}(R)}, \frac{\text{exact} \& \text{faithful}}{\text{full}} \text{QCoh}(\text{Proj } R) \right)$$

0.1 Classical Algebraic Geometry: Affine varieties

$R = k[x_1, \dots, x_n]$ polynomial ring over algebraically closed field k
 $I \subseteq R$ ideal

$X = V(I) = \{a \in k^n : f(a) = 0 \forall f \in I\}$ affine variety

The topological space

Affine space: $\mathbb{A}^n = k^n$ with Zariski topology: $\left\{ \begin{array}{l} \text{closed sets: } V(I) \\ \text{open sets: } U_I = \mathbb{A}^n \setminus V(I) \\ X \subseteq \mathbb{A}^n \text{ subspace topology: } X \cap U_I \end{array} \right.$

The functions on it

$R \cong \text{Hom}(\mathbb{A}^n, \mathbb{A}^1)$, $f \mapsto (a \mapsto f(a))$
 $\Gamma(X) = \{f \in R : f(X) = 0\}$
 Remark $V(\Gamma(X)) = X$ for affine varieties X

Coordinate ring: $k[X] = R/\Gamma(X)$

Key facts: 1) Hilbert's basis theorem: R Noetherian, so $k[X]$ Noetherian
 2) Hilbert's weak nullstellensatz: Maximal ideals of R (and of $k[X]$) are $m_a = I(\{a\}) = \langle x_1 - a_1, \dots, x_n - a_n \rangle$, so correspond to points: $\{a\} = V(m_a)$
 3) Hilbert's Nullstellensatz: $\Gamma(V(I)) = \sqrt{I}$ (Hence: $\Gamma(V(I)) = I$ if I is radical)
 Lemma There are enough functions to separate points

pf $a \neq b \in X \subseteq \mathbb{A}^n \Rightarrow$ some coordinate $a_i \neq b_i \Rightarrow x_i \in k[X]$ separates a, b

Morphisms between affine varieties

$\text{Hom}(\mathbb{A}^n, \mathbb{A}^m) \cong R^m \leftarrow$ polynomial maps $a \mapsto (f_1(a), \dots, f_m(a))$
 $\text{Hom}(X, Y) = \{\text{restriction of a polynomial map } \mathbb{A}^n \rightarrow \mathbb{A}^m \text{ s.t. } X \rightarrow Y\}$
 Facts: 1) $k[X] \cong \text{Hom}(X, \mathbb{A}^1) \leftarrow$ "values of functions are enough to determine the abstract function"

2) $\text{Hom}(X, Y) \cong \text{Hom}_{k\text{-alg}}(k[Y], k[X])$

$$(f: X \rightarrow Y) \mapsto (f^*: \text{Hom}(Y, \mathbb{A}^1) \rightarrow \text{Hom}(X, \mathbb{A}^1)) \leftarrow \begin{array}{c} \text{"pullback"} \\ \begin{array}{ccc} X & \xrightarrow{f} & Y \\ f^* \downarrow & & \downarrow \\ \mathbb{A}^1 & & \mathbb{A}^1 \end{array} \end{array}$$

Equivalence of categories


$\{\text{affine varieties}\} \longleftrightarrow \{\text{finitely generated reduced } k\text{-algebras} \& \text{homs of } k\text{-algs.}\}$
 $X \mapsto k[X]$
 $(f: X \rightarrow Y) \mapsto f^*$

Remark The "same" (up to isomorphism) X can be embedded in various $\mathbb{A}^n$.
 E.g. cuspidal cubic $V(y^2 - x^3) = \sqrt{(y^2 - x^3)} \subseteq \mathbb{A}^2_{x,y} \cong V(y^2 - x^3, z - x) \subseteq \mathbb{A}^3_{x,y,z}$

0.2 Why schemes?

Some reasons:

- 1) Why always have spaces embedded in $\mathbb{A}^n$? (extrinsic)
Can you make sense of X without reference to $\mathbb{A}^n$? (intrinsic)
- 2) Why not let R be any ring?
- 3) When you deform varieties, nilpotents arise naturally and should not be ignored:



We lost information: classically you cannot tell $x=0$ apart from $x^2=0$ in the theory of schemes, the key role is not played by the topological space. The key role is played by the ring of functions, or rather, the sheaf of functions $\mathcal{O}$: on each open set $U \subseteq X$ get a ring of functions $\mathcal{O}(U)$.

Example above: $\mathcal{O}(X) = k[x]/(x^2) \leftarrow$ we do not reduce the ring of functions

At what cost? Values of functions need not determine the abstract function:

$$\mathcal{O}(X) \ni \alpha + \beta x \mapsto (\alpha + \beta x : X = \{0\} \rightarrow \mathbb{A}^1) \in \text{Hom}(X, \mathbb{A}^1)$$

Idea: the abstract " β " remembers that X arose from the collision of

two points, so β records tangential information: $\frac{\partial}{\partial x}|_{x=0} (\alpha + \beta x) = \beta$.

0.3 What is a point?

X topological space is reducible if $X = X_1 \cup X_2$ for proper closed $X_i \subsetneq X$.

Euclidean world (more generally if X Hausdorff): $Y \subseteq X$ irreducible $\Leftrightarrow Y = \text{point}$ or $Y = \emptyset$.

Classical Alg. Geom. $\leftarrow$ point $a \in X \leftrightarrow$ max ideal $m_a \subseteq k[X]$

closed $\emptyset \neq Y \subseteq X$ irreducible $\Leftrightarrow \Pi(Y) \subseteq k[X]$ prime ideal

R ring $\Rightarrow$ "points" of R are $\text{Spec}(R) = \{\text{prime ideals of } R\}$ not just max ideals

Categorically a good choice since functorial:

$$\varphi: R \rightarrow S \text{ hom of rings} \Rightarrow \varphi^{-1}(\text{prime}) = \text{prime ideal} \Rightarrow \text{Spec } S \xrightarrow{\varphi^{-1}} \text{Spec } R$$

1. DEFINITION OF SCHEMES

1.1 Examples of affine schemes

$\text{Spec}(R)$ some ring R (always: comm. ring with 1)

- As a set: $\text{Spec}(R) = \{\text{prime ideals } P \subseteq R\} \leftarrow (\text{prime}) \text{ spectrum}$
- Zariski topology: closed sets: $V(I) = \{ \text{prime ideals containing } I \} \subseteq \text{Spec } R$
- sheaf $\mathcal{O}_{\text{Spec } R}$ which we construct later. $\leftarrow$ spaces of functions

Remark The global functions are: $\mathcal{O}_{\text{Spec } R}(\text{Spec } R) = R$. $\leftarrow$ so spaces of fns can recover the space!

Key exercise $V(I) \cup V(J) = V(I \cdot J) = V(I \cap J)$ so $V(I \cdot J) = V(I \cap J)$ but $I \cdot J$ and $I \cap J$ may be $\neq$

Key $V(I) = \emptyset \Leftrightarrow I = R \Leftrightarrow 1 \in I$, since any proper ideal $\subseteq$ some maximal

Topological: open sets: $U_I = \text{Spec } R \setminus V(I) = \bigcup_{f \in I} D_f$

basis of open sets: $D_f = \{P \in \text{Spec } R : f \notin P\} = \{P \in \text{Spec } R : f(P) \neq 0\}$

"value of $f \in R$ at P ": $R/P \xrightarrow{f} K(P) = \text{Frac}(R/P) \cong R_P/P_P$

Remark $f(P) = 0 \Leftrightarrow f \in P$

Examples 1) $R = k[X] \leftarrow$ affine variety $X \subseteq \mathbb{A}^n$

$\text{Spec } R \xrightarrow{\text{bijection}} \text{irreducible subvarieties } Y \subseteq X$

Value of $f \in R$ at m_a : $m_a \rightarrow R/m_a \cong k$ $\leftarrow$ in this case the target field does not depend on the point

2) $\text{Spec } \mathbb{Z} = \{0\} \cup \{P : P \in \mathbb{N} \text{ prime}\}$

$V(0) = \{\text{prime ideals containing } 0\} = \text{Spec } \mathbb{Z}$ so the point $\{0\}$ is dense!

$V(P) = \{P\}$ are "closed points". Value of $f \in \mathbb{Z}$: $f(P) = (f \in \mathbb{Z}/P) = (f \bmod P)$

In general Prime ideals P with $V(P) = \text{Spec } R$ are called generic points

Prime ideals P with $V(P) = \{P\}$ are called closed points

Exercise $\{\text{closed points}\} = \{\text{max ideals of } R\}$

Exercises • a prime ideal $\Rightarrow$ a radical ($a = \sqrt{a}$)
 • For a, b radical, $a \subseteq b \Leftrightarrow V(a) \supseteq V(b)$ ← order reversing!

Cor $V(I) \subseteq V(J) \Leftrightarrow \sqrt{I} \supseteq \sqrt{J}$
Pf $V(I) = V(\sqrt{I})$, so: $\Leftrightarrow V(\sqrt{I}) \subseteq V(\sqrt{J}) \Leftrightarrow \sqrt{I} \supseteq \sqrt{J}$ by exercise. $\square$

Cor $V(a) = V(b) \Leftrightarrow \sqrt{a} = \sqrt{b}$
 $\Rightarrow$ [closed sets of $\text{Spec } R$] $\xleftrightarrow{\text{order-reversing correspondence}} \{\text{radical ideals of } R\}$

Proposition $f \in R$ vanishes at all $p \in \text{Spec } R \Leftrightarrow f$ nilpotent

Covering Trick $\text{Spec } R = \bigcup D_{f_i} \Leftrightarrow 1 \in \langle \text{all } f_i \rangle \Leftrightarrow \langle \text{all } f_i \rangle = R$

Pf $\text{Spec } R \setminus \bigcup D_{f_i} = \bigcap V(f_i) = V(\langle \text{all } f_i \rangle)$, now use previous key. $\square$

Theorem $\text{Spec } R$ is quasi-compact $\leftarrow$ (quasi-compact = compact = open covers have finite subcovers)

Pf $\text{Spec } R = \bigcup U_i$. As $U_i = \bigcup_j D_{f_{ij}}$, wlog $U_i = D_{f_i}$.
 Trick $\Rightarrow 1 = \sum_{\text{finite}} r_i f_i \leftarrow$ so finitely many f_i generate R , so those D_{f_i} cover. $\square$

Basic Exercises

1) $\varphi: R \rightarrow S$ ring hom $\Rightarrow \alpha: \text{Spec } S \rightarrow \text{Spec } R$, $p \mapsto \varphi^{-1}(p)$ is continuous
 indeed $\alpha^{-1}(D_f) = D_{\varphi(f)} \leftarrow$ (Hint: $f \notin p \Leftrightarrow \varphi(f) \notin \varphi(p)$)

2) Show that $\text{Spec}(R/I)$ "is" the subspace $V(I) \subseteq \text{Spec } R$ and the quotient map $\pi: R \rightarrow R/I$ induces via (1) the inclusion map on Specs.

Example $\text{Spec}(R/(f)) = \{\text{prime ideals of } R \text{ containing } f\}$
 = the points of $\text{Spec } R$ where f vanishes
 = $V(f)$

3) Show that $\text{Spec}(S^{-1}R)$ "is" a subspace of $\text{Spec } R$, where $S^{-1}R$ is localization of R at a multiplicative set $S \subseteq R$, and $R \rightarrow S^{-1}R$, $r \mapsto \frac{r}{1}$ induces via (1) the inclusion

Example $S = \{1, f, f^2, f^3, \dots\}$, so $S^{-1}R = R_f$, then:

$\text{Spec } R_f = \{\text{prime ideals of } R \text{ not containing } f\}$
 = the points of $\text{Spec } R$ where f does not vanish
 = D_f

4) $D_f \cap D_g = D_{fg}$, so $\text{Spec } R_f \cap \text{Spec } R_g = \text{Spec } R_{fg}$

5) $D_f \subseteq D_g \Leftrightarrow V(f) \supseteq V(g) \Leftrightarrow \sqrt{f} \subseteq \sqrt{g} \Leftrightarrow f \in (g)$ some $N \Leftrightarrow g \in R_f$ invertible

6) $p \subseteq R$ prime ideal $\Rightarrow R_p = S^{-1}R$ for $S = R \setminus p$, then $\exists!$ closed point $m_p = p \cdot R_p \in \text{Spec } R_p$
 so local ring: $\exists!$ max ideal m ($\Leftrightarrow$ elts outside m are invertible)

Also: $m_p \in U \subseteq \text{Spec } R_p$ open $\Rightarrow U = \text{Spec } R_p$.

1.2 Definition of a scheme

Def A ringed space is

- a topological space X
- with a sheaf of rings $\mathcal{O}_X$ on X

Locally ringed space if also:

- all stalks $\mathcal{O}_{X,x}$ are local rings (so $\exists$ unique maximal ideal $\mathfrak{m}_{X,x} \subseteq \mathcal{O}_{X,x}$ (and $\exists$ residue field at $x: k(x) = \mathcal{O}_{X,x} / \mathfrak{m}_{X,x}$)

Def An affine scheme is a locally ringed space

isomorphic to $(\text{Spec } R, \mathcal{O}_{\text{Spec } R})$ for some ring R .

Def A scheme is a locally ringed space which is

locally isomorphic to an affine scheme.

means:

$\forall x \in X \exists$ some open neighbourhood $U \ni x$ st. $(U, \mathcal{O}_X|_U) \cong (\text{Spec } R, \mathcal{O}_{\text{Spec } R})$
 $\exists$ some ring R depending on x

1.3 Pre-sheaves

$\mathcal{A}b$ = category of abelian groups and group homs

X = any topological space

$\text{Top } X$ = category with objects: open sets $U \subseteq X$
 morphs: inclusion maps

Def A presheaf (of abelian groups) on X is a contravariant functor

$F: \text{Top } X \rightarrow \mathcal{A}b$

So: $\forall$ open $U \subseteq X$ have an abelian group $F(U)$ ← elements called sections (over U)

• $\forall$ inclusion $U \hookrightarrow V$ have a "restriction" group hom $F(V) \rightarrow F(U)$

• $F(\text{id}: U \rightarrow U): F(U) \xrightarrow{\text{id}} F(U)$ so $s|_U = s$ for $s \in F(U)$.

• $U \subseteq V \subseteq W \Rightarrow F(W) \rightarrow F(V) \rightarrow F(U)$ so: $(s|_V)|_U = s|_U$ for $s \in F(W)$.

Example X topological space, $F(U) = \{\text{continuous functions } U \rightarrow \mathbb{R}\}$ with obvious restrictions

Morphism of pre-sheaves = natural transformation of such functors: $\varphi: F \rightarrow G$

So: $\forall$ open $U \subseteq X$ have $\varphi_U: F(U) \rightarrow G(U)$ group hom

$\forall$ inclusion $U \hookrightarrow V$ have $F(U) \xrightarrow{\varphi_U} G(U)$
 $F(V) \xrightarrow{\varphi_V} G(V)$
 $\uparrow$ restriction homs $\leftarrow$ i.e. this diagram with $\varphi_U = \text{inclusion}$

Sub pre-sheaf $F \subseteq G$ means $F(U) \subseteq G(U)$ subgr, compatibly with restrictions

RED: WORDS TO BE DEFINED LATER

IDEA

- ← the points
- ← the functions
- ← the germs of functions near point x
- ← the "value" of a function at x lives here

if use category $\mathcal{C}$
 get (pre)sheaves with values in $\mathcal{C}$
 e.g. $\mathcal{C} = \text{Rings}$
 get presheaf of rings

$\leftarrow \text{Mor}(U, V) = \{\emptyset \text{ if } U \not\subseteq V, \text{ incl. if } U \subseteq V\}$

So the homs are compatible with restrictions

i.e. this diagram with $\varphi_U = \text{inclusion}$

1.4 Sheaves

Def Pre-sheaf F is a sheaf on X if it satisfies the local-to-global condition:

If U_i open, $s_i \in F(U_i)$ agreeing on overlaps:
 $s_i|_{U_i \cap U_j} = s_j|_{U_i \cap U_j} \in F(U_i \cap U_j)$

Then $\exists$ unique $s \in F(\bigcup U_i)$ with $s|_{U_i} = s_i$.
 Consequences

- two sections $s, t \in F(U)$ equal $\Leftrightarrow$ they equal locally: $s|_{U_i} = t|_{U_i}$, $U = \bigcup U_i$
- you can build sections by defining local sections, compatibly on overlaps.
- exact sequence: $0 \rightarrow F(U) \rightarrow \prod_i F(U_i) \xrightarrow{\pi} \prod_i F(U_i \cap U_j) \rightarrow \dots$
- $F(\emptyset) = 0$ (Hint: consider empty covering of $\emptyset$)

Examples

- 1) Sheaf of continuous real functions: $F(U) = \{\text{continuous maps } U \rightarrow \mathbb{R}\}$
- 2) Skyscraper sheaf at $p \in X$ for group A : $F(U) = \begin{cases} A & \text{if } p \in U \\ 0 & \text{if } p \notin U \end{cases}$
- 3) Presheaf of constant functions for group A :

$$F(U) = \begin{cases} A & \text{if } U \neq \emptyset \\ 0 & \text{if } U = \emptyset \end{cases}$$

(so $f \in F(U)$ is a constant function $f: U \rightarrow A$, $f \in A \in A$)
 (only want one function on $\emptyset$)
- 4) Sheaf of locally constant functions for group A . So $f \in F(U)$ means $f: U \rightarrow A$ such that $\forall x \in U, \exists$ open $x \in V \subseteq U$ with $f|_V: V \rightarrow A$ constant.
 Warning: it implies f constant on connected components but converse can fail (e.g. consider $\mathbb{Q}$ with usual topology).

Exercise (3) is not a sheaf if $X = 2$ points with discrete topology, $\mathbb{R} \neq 0$.

Write $\text{Ab}(X) = \text{category of sheaves on } X \text{ and morphisms of sheaves}$

Sh(X) if work with category of Sets instead of Ab (morphisms of presheaves)

Def stalk at x of presheaf F is the abelian group

$$F_x = \varinjlim_{x \in U} F(U)$$

← direct limit over restriction maps induced by inclusions.

Explicitly:

An element of F_x is determined by $s \in F(U)$ some $U \ni x$ open, identify $s \sim t$ for $t \in F(V) \Leftrightarrow s|_W = t|_W$ some $U \cap V \ni W \ni x$ open

Rmk. natural map $F(U) \rightarrow F_x$, $s \mapsto s_x = \text{equivalence class of } s$. (for $x \in U$)
 or write: $s|_x$

- morph $\varphi: F \rightarrow G$ then get $\varphi_x: F_x \rightarrow G_x$
 or write: $\varphi|_x$
 $(\varphi_x(s_x) = \varphi(s)|_x \text{ if } s \in F(U))$

Exercise $\varphi, \psi: F \rightarrow G$ morphs of sheaves,

if all $\varphi_x = \psi_x: F_x \rightarrow G_x$ then $\varphi = \psi$.

Facts For sheaves F, G in category $\text{Ab}(X)$

- $F \rightarrow G$ monomorphism $\Leftrightarrow F_x \rightarrow G_x$ injective $\forall x$
- $F \rightarrow G$ epimorphism $\Leftrightarrow F_x \rightarrow G_x$ surjective $\forall x$
- $F \rightarrow G$ isomorphism $\Leftrightarrow F_x \rightarrow G_x$ iso $\forall x$

Warning $\text{mono} \Leftrightarrow F(U) \rightarrow G(U) \text{ inj } \forall U$, but fails for epi: $F(U) \rightarrow G(U)$ need not be surj.
 Exercise $F_x \xrightarrow{\varphi_x} G_x \text{ surj} \Leftrightarrow \forall t \in G(U), \exists s \in F(U): \varphi(s) = t|_U \in G(U)$ (but V can depend on t)

1.6 Sheafification

F pre-sheaf $\Rightarrow F^+$ sheaf (ification):

$$F^+(U) = \left\{ s: U \rightarrow \bigsqcup_x F_x : \text{locally } s \text{ is a section of } F \right\}$$

(in fact by definition $s(x) \in F_x$ so $s: U \rightarrow \bigsqcup_{x \in U} F_x \subseteq \bigsqcup_{x \in X} F_x$)

comes with natural morph $F \rightarrow F^+ \leftarrow (s \in F(U) \mapsto (x \mapsto s_x) \in F^+(U))$

Exercise: F^+ is a sheaf, $F_x^+ = F_x$ and it satisfies:

Universal property: $\forall$ sheaf G on X , $F^+ \dashrightarrow G$
 (determines F^+ uniquely up to unique isomorphism)

Hint. In our construction:

$F_x^+ = F_x \rightarrow G_x$ so we know locally how sections map but we need to glue them...

Trick: $F \rightarrow F^+ \rightarrow G$ finally G is sheaf so $G_x = G^+$
 $G \rightarrow G^+$ (natural iso, using $G_x = G_x^+$ and Facts)

Example (pre-sheaf of constant functions) $^+ =$ (sheaf of locally constant functions)

Exercise 1) $F \subseteq G$ sub pre-sheaf, G sheaf $\Rightarrow \exists$ smallest subsheaf $H \subseteq G$ s.t. $F \subseteq H$
 Moreover, $H_x = F_x$.

- 2) $(DF)(U) = \varinjlim_{x \in U} F_x$ with obvious restriction maps is a sheaf (sheaf of discontinuous sections)
 Hint: mimic definition of F^+
- 3) $i: F \rightarrow DF$ obvious morph, let $F^b = \text{presheaf image so } F^b(U) = i(U)$
 then $F^b \subseteq DF$ is a sub pre-sheaf and construction (1) gives $H = F^+$.

1.7 Kernels, Cokernels, Images

$\varphi: F \rightarrow G$ morph of sh.

- $(\text{Ker } \varphi)(U) = \text{Ker } \varphi_U$ is sheaf
- $\text{Coker } \varphi = (\text{pre-coker } \varphi)^+$ where $(\text{pre-coker } \varphi)(U) = \text{Coker } \varphi_U$
- $\text{Im } \varphi = (\text{pre-Im } \varphi)^+$ where $(\text{pre-Im } \varphi)(U) = \text{Im } \varphi_U$

Books in additive cat:

$\text{mono} \Leftrightarrow H \xrightarrow{\circ} G \text{ then } H \xrightarrow{\circ} F$

$\text{epi} \Leftrightarrow F \xrightarrow{\circ} H \text{ then } G \xrightarrow{\circ} H$

category ker & cok, see below

$\mathcal{A} = \text{additive category}$ such that morphisms have Ker, Coker and i) $\varphi: F \rightarrow G$ monomorph is the Ker of its Coker

ii) " epimorph " Coker " Ker
core means $\text{Mor}(A, B)$ abelian gp (so often write $\text{Hom}(A, B)$) s.t.
 Composition of morphisms distributes over addition
 $\exists$ products $A \times B$ $(\exists) (x, y) \in A \times B$
 $\exists$ zero object 0 (an object that is both initial & terminal)

ve/abelian cats is additive if $\text{Hom}(A, B) \rightarrow \text{Hom}(FA, FB)$ is gp. hom.

$\begin{array}{ccc} \text{Objects} & & \\ \text{Ker } \varphi & \longrightarrow & A \end{array}$	$\begin{array}{ccc} & & A \\ & \swarrow \varphi & \searrow \varphi \\ \text{Ker } \varphi & \xleftarrow{\varphi} & B \end{array}$ $\exists ! \uparrow$ Fact: $\text{Ker } \varphi$ is an epimorph.	$\text{Ker } \varphi$ is $B \rightarrow \text{Coker } \varphi \in \text{Ob } \mathcal{B}$ s.t. $\exists ! \uparrow$	$\text{Im } \varphi = \text{Ker}(\text{Coker } \varphi)$ which is a morph $\text{Im } \varphi \rightarrow B$ Facts: $\exists !$ factorization of φ $A \rightarrow \text{Im } \varphi \rightarrow B$ Abelian cat $\Rightarrow A \rightarrow \text{Im } \varphi \text{ epi}$ and $= \text{Coker}(\text{Ker } \varphi)$
--	--	--	---

an gps, (i) says:

$$\text{Ker } \pi \xrightarrow{\psi} B \xrightarrow{\pi} B/A$$

is $\text{Ker } \pi$ $\xrightarrow{\psi}$ $\uparrow$ $\text{Coker } \psi$

underlining Ker , Coker , Im .

as expected!

Freyd-Mitchell Thm

logical definitions can be cumbersome to work with. It turns out: in an abelian category $\mathcal{A}$, $\exists$ a possibly non-commutative ring R with 1 such that $\mathcal{A}$ is a faithful exact functor $\mathcal{A} \rightarrow \{\text{left } R\text{-modules}\}$ (in particular preserves kernels, cokernels, and is additive). $\Rightarrow$ can "pretend" you work with modules. (Example you just apply the theorem to the small abelian subcategory involved in your diagram/sequence of maps - don't need to use the whole category)

lex $F^\bullet = (\dots \rightarrow F^{i-1} \xrightarrow{d^{i-1}} F^i \xrightarrow{d^i} F^{i+1} \rightarrow \dots)$ in an abelian cat

composite of two consecutive morphs is zero: $d^{i+1} \circ d^i = 0 \quad \forall i$
($\exists$ mono $\text{Im } d^i \subset \text{Ker } d^{i+1}$ and $\text{Im } d^i$ is color)

$H^i(F^\bullet) = \text{Ker } d^{i+1} / \text{Im } d^i$

$\text{Im } d^i = \text{Ker } d^{i+1} \iff F^\bullet \text{ is exact sequence of abelian grps}$
 $\iff F^\bullet \text{ is complex with zero homology } H^\bullet = 0$
 by **Facts** on previous page

(short exact sequences) $0 \rightarrow F \xrightarrow{\beta} H \rightarrow 0$ of sheaves exactness at level of stalks, but can equivalently check: $F(U) \rightarrow G(U) \rightarrow H(U) \rightarrow 0$ exact $\forall$ open U smallest subsheaf containing $\text{pre-Im } \beta$, meaning every of H can be obtained by gluing local sections of type $\beta(\text{local section of } F)$

A functor of abelian cats is left exact if:

$$0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0 \text{ exact} \\ \Rightarrow 0 \rightarrow FA \rightarrow FB \rightarrow FC \text{ exact}$$

right exact if:

$$FA \rightarrow FB \rightarrow FC \rightarrow 0 \text{ exact}$$

Example $\text{Hom}_R(M, \cdot)$ is left exact, $\cdot \otimes_R M$ is right exact, as functors on $R\text{-mods}$ (any $R\text{-mod } M$)

1.9 Push-forward (direct image) and inverse image

 $f: X \rightarrow Y$ continuous
$$\Rightarrow \text{additive functor } f_* : \mathcal{A} \rightarrow \mathcal{B}$$

Def $F \in \text{Ab}(X)$ gives $f_F \in \text{Ab}(Y) :$

$$(f_*F)(V) = F(f^{-1}(V))$$

Exercise $(g \circ f)_* F = g_*(f_* F)$ for $X \xrightarrow{f} Y \xrightarrow{g} Z$.

$$\Rightarrow \text{additive functor } f^{-1}: \mathbf{Ab} \mathcal{Y} \rightarrow \mathbf{Ab} \mathcal{X}$$

Def $F \in \text{Ab}(Y)$ gives $f^{-1}F \in \text{Ab}(X)$ is $(\text{pre-}f^{-1}F)^+$ where

$$(\text{pre-}f^{-1}F)(u) = \varinjlim_{V \ni f(u)} F(V)$$

Exercise $(f^{-1}F)_x = F_{f(x)}$ and $(g \circ f)^{-1} \cong f^{-1} \circ g^{-1}$

Examples 1) $i: S \rightarrow X$ inclusion of an open subset:

$$F \in \text{Ab}(S) \quad i_* F: V \longrightarrow F(V \cap S)$$
$$F \in \text{Ab}(X) \quad \quad \quad i^{-1}F := \mathcal{U} \mapsto F(\mathcal{U}) \quad \quad \quad \downarrow$$

open $\leq S \leq X$ called restriction

$$2) \quad i :: \text{point} \rightarrow X$$
[illegible]
$$F \in A_b(X)$$

3) $\pi: X \rightarrow \text{point}$

$$F \in A^b(X)$$

Proposition 1) f_* is left exact

2) f^{-1} is exact

For f^* : exercise

Proof for f^{-1} :

$$= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$
[illegible]

KMK } would follow by category theory for

PICTURE IN MIND

$$\begin{array}{ccccc} f_* F & & Y & \hookrightarrow & U & V \\ & & \downarrow f & & & \\ F & \hookrightarrow & X & \hookrightarrow & U & V \\ & & & & f^{-1}V & \end{array}$$

PICTURE IN MIND

$$\begin{array}{ccccc} & & F & & \\ & & | & & \\ f^{-1}F & & & & Y \geq V \\ & & | & & | \\ & & X & \xrightarrow{f} & U \\ & & & & | \\ & & & & U \end{array}$$

also follows by uniqueness of adjoint functors, see next page.

$F|_S$
restriction of F

if $U = \{\text{point}\}$
if $U = \emptyset$

$\rightarrow$ is left exact

assumption is exact

Proposition f^{-1} is the left adjoint functor of f_* , meaning $\exists$ natural iso $\text{Mor}(f^{-1}F, G) \cong \text{Mor}(F, f_*G)$ which is natural in F and G

Sketch pf
 $\text{In} \rightarrow \text{direction:}$ since view is allowed $\xrightarrow{\text{given}} F(V) \xrightarrow{f_*} G(U)$
 $\text{II} \leftarrow \text{pick } U = f^{-1}V$
 $G(f^{-1}V) = f_*G(V)$

$\text{In} \leftarrow \text{direction:}$ $F(V) \xrightarrow{\text{given}} G(f^{-1}V)$
 $\downarrow$ assume $V \ni fU$
 $\text{take } \lim_{V \ni fU}$ over such V
 $\lim_{V \ni fU} F(V) \rightarrow \lim_{V \ni fU} G(f^{-1}V)$
 $\downarrow$ restriction $\leftarrow$ notice $f^{-1}V \supseteq U$
 $G(U)$

Now check these two are natural transformations, inverse to each other, and natural in F, G .
Rmk Another example of adjoint functors, for R -modules, are $\text{Hom}(M, ?)$ and $\otimes M$:
 $\text{Hom}(F \otimes M, G) \cong \text{Hom}(F, \text{Hom}(M, G))$ for R -mods F, G .

1.10 Morphisms of ringed spaces

Def $(f, \varphi): (X, \mathcal{O}_X) \rightarrow (Y, \mathcal{O}_Y)$ morph of ringed spaces means
 $X \xrightarrow{f} Y$ continuous map of topological spaces
 $f_* \mathcal{O}_X \xleftarrow{\varphi} \mathcal{O}_Y$ morph of sheaves of rings (on Y)
 often write $\varphi = f^\#$
 (So: $\mathcal{O}_X(f^{-1}V) \xleftarrow{f^\#} \mathcal{O}_Y(V)$ for $V \subseteq Y$, compatibly with restrictions)

For a morphism of locally ringed spaces want in addition:
 $\mathcal{O}_{X,x} \xleftarrow{\varphi_x} \mathcal{O}_{Y,f(x)}$ is local ring hom
 (Explanation: $\varphi_x(\frac{s}{t}) \in \mathcal{O}_x(f^{-1}V)$ is a representative for $\varphi_x(\frac{s}{t})$)

Rmk Can compose: $(X, \mathcal{O}_X) \xrightarrow{f} (Y, \mathcal{O}_Y) \xrightarrow{g} (Z, \mathcal{O}_Z)$:
 $(g \circ f)_* \mathcal{O}_X = g_* f_* \mathcal{O}_X \xleftarrow{g^\# \circ f^\#} g_* \mathcal{O}_Y \xleftarrow{g^\#} \mathcal{O}_Z$
 Notice in the definition we cannot just talk about a morphism $\mathcal{O}_X \leftarrow \mathcal{O}_Y$ because the sheaves are not defined over the same topological space.
 $\Rightarrow$ either need a morph $f_* \mathcal{O}_X \leftarrow \mathcal{O}_Y$ of sheaves on Y or a morph $\mathcal{O}_X \leftarrow f^{-1} \mathcal{O}_Y$ of sheaves on X

By the proposition, this is the same information since $\text{Mor}(f^{-1} \mathcal{O}_Y, \mathcal{O}_X) \cong \text{Mor}(\mathcal{O}_Y, f_* \mathcal{O}_X)$
 (Notice also the map on stalks $\mathcal{O}_{X,x} \leftarrow (\mathcal{O}_Y)_x \leftarrow (f^{-1} \mathcal{O}_Y)_x = \mathcal{O}_{Y,f(x)}$ is the φ_x above)
Rmk φ local $\Rightarrow$ also get hom on residue fields: $\varphi_x: K(f(x)) = \mathcal{O}_{Y,f(x)} / \mathfrak{m}_{Y,f(x)} \hookrightarrow \mathcal{O}_{X,x} / \mathfrak{m}_{X,x} = K(x)$
 $\Rightarrow$ field extension $\varphi_x: K(f(x)) \hookrightarrow K(x)$ (in classical algebraic geometry: K is alg. closed and x closed point)
 (get $\text{id}: K \rightarrow K$, $p(f(x)) \mapsto (f^*p)(x)$ where $\{x \in X\}$)

1.11 A sheaf defined on a topological basis

X top. space with a basis B of open subsets $\leftarrow$ means: basic sets cover X , and:
 $\forall$ basic $B_1, B_2, x \in B_1 \cap B_2$
 $\exists$ basic B with $x \in B \subseteq B_1 \cap B_2$

Def B -sheaf F means

$F(U) \in \text{Ab}, \forall$ basic U with $\text{homs } F(U) \rightarrow F(V), s \mapsto s|_V \quad \forall \text{ basic } V \subseteq U$
 and as usual: $F(U) \xrightarrow{\text{id}} F(U)$ and $F(U) \rightarrow F(W) \rightarrow F(W)$ for $W \subseteq V \subseteq U$

$\bullet$ local-to-global condition:

$\forall$ basic U with $U = \bigcup U_i \leftarrow$ basic

$\forall s_i \in F(U_i)$ "agreeing locally on overlaps":
 $\forall x \in U_i \cap U_j \exists$ basic $x \in U_k \subseteq U_i \cap U_j$ with

$$s_i|_{U_k} = s_j|_{U_k} \in F(U_k)$$



$\Rightarrow \exists$ unique $s \in F(U)$ with $s|_{U_i} = s_i$:
Rmk stalk $F_x = \varinjlim_{x \in \text{basic } U} F(U)$

(Hence also the stalk is $\mathbb{F}_x$ up to canonical iso.)

Theorem 1) B -sheaf F extends uniquely (up to unique iso) to a sheaf $\tilde{F}$ on X .
 (so $F(\text{basic } U)$ and restrictions for basic sets are same up to canonical isomorphisms.)

2) B -sheaves F, G then morph $F \rightarrow G$ on the extended sheaves is uniquely defined by data:

$\bullet$ $\text{homs } F(U) \rightarrow G(U)$ for basic U , commuting with restrictions (for basic opens)
Uniqueness Such an extension $\tilde{F}$ is unique (if it exists) because we can canonically identify $\tilde{F}(U)$ for any open U in terms of the B -sheaf data:

$\tilde{F}(U) \xrightarrow{\text{bijection}} \{s_v \in F(V) \text{ for } (\text{basic } V) \subseteq U: s_v|_W = s_v|_W \in F(W) \text{ for basic } W \subseteq V \cap V'\}$
 $s \mapsto (s_v := s|_V \in \tilde{F}(V) = F(V))$

Explanation: given s , notice that this holds: $s_v|_W = (s|_W)|_V = s|_{V \cap V'} = s_v|_{V \cap V'}$
 Conversely, given such $s_v \in F(V) = \tilde{F}(V)$, then $s_v|_{V \cap V'} \in \tilde{F}(V \cap V')$ and $s_v|_{V \cap V'} \in F(V \cap V')$ must equal because their restrictions to a covering of $V \cap V'$ by basic W agree ($= s_w$).
 (and then use sheaf property of $\tilde{F}$)

Existence

$$F(U) = \varinjlim_{(\text{basic } V) \subseteq U} F(V)$$

$$= \left\{ (s_v) \in \prod_{(\text{basic } V) \subseteq U} F(V) : s_v|_W = s_w \quad \forall W \subseteq V \subseteq U \right\}$$

 (compatible families of local sections on basic open sets)
 (inverse limit over restrictions for basics)



With obvious restriction maps (for $U' \subseteq U$ a subset of the (basic V) $\subseteq U$ are $\subseteq U'$)

Notice: $F(\text{basic } U)$ has not changed up to canonical identification:

$$F(U) \xrightarrow{\cong} \varprojlim_{(\text{basic } V) \subseteq U} F(V) \xrightarrow{s} (s|_V) \text{ which includes } s|_U = s.$$

and for stalks:

$$\varprojlim_{x \in (\text{basic } V)} F(V) \xrightarrow{\cong} \varprojlim_{x \in U} F(U) \leftarrow \begin{array}{l} \text{easy check:} \\ \text{if sections} \\ \text{agree on } x \in W \\ \text{then agree on} \\ x \in V \subseteq W \\ \text{some basic } V. \end{array}$$

Proof (2): by functoriality of $\varprojlim$:

$$\varprojlim_{(\text{basic } V) \subseteq U} F(V) \longrightarrow \varprojlim_{(\text{basic } V) \subseteq U} G(V).$$

Remark Equivalently, it is enough to remember germs around each point:

$$F(U) = \left(\varprojlim_{(\text{basic } V) \subseteq U} F(V) \right) \xrightarrow{\cong} \left\{ s: U \rightarrow \bigsqcup_{x \in X} F_x : s(x) \in F_x \text{ which } \right. \\ \left. \begin{array}{l} \text{are "locally compatible":} \\ \forall x \in U, \exists x \in (\text{basic } V) \subseteq U \\ \exists t \in F(V) \text{ with } \\ \exists \text{ open } x \in W \subseteq V \text{ } t_y = s(y) \forall y \in W \end{array} \right\}$$

with obvious restriction maps for these (just restrict the map $U \rightarrow \bigsqcup F_x$).

Remark Can simplify: $\text{WLOG } W$ also basic (just pick $x \in (\text{basic } W)$ so: $\forall x \in U \exists x \in (\text{basic } W) \subseteq U$ replace V by W , so $V = W$ basic. $\exists t \in F(V)$ with $t_y = s(y) \forall y \in V$

Inverse: have cover $U = \bigcup (\text{basic } x \in V^*)$ and $t^* \in F(V^*)$ s.t. t^* agree locally (since germs agree) so $\star$ holds so can extend to unique global section.

1.12 Construction of $\mathcal{O}_{\text{Spec } R}$

$X = \text{Spec } R$, we define $\mathcal{O}_X$ first on basic open sets: $\downarrow$

$$\mathcal{O}_X(D_f) = R \text{ localised at multiplicative set } \{g: g \text{ does not vanish on } D_f\} \\ \cong R_f \\ \uparrow \text{ natural}$$

(Recall exercise: $V(g) \subseteq V(f) \Leftrightarrow D_f \subseteq D_g \Leftrightarrow f^m \in (g) \Leftrightarrow g \in R_f$ invertible)

For $D_f \subseteq D_g$ define natural restriction homs: (which are compatible under composition)

$$\mathcal{O}_X(D_g) \longrightarrow \mathcal{O}_X(D_f) \xleftarrow{\text{"localise further"}} \\ \cong R_g \xrightarrow{\text{explicitly: } f^n = rg \text{ so}} R_f \\ x \mapsto \frac{x \cdot f^m}{(r \cdot g)^m} = \frac{x \cdot f^m}{r^m}$$

Lemma 1 This is a B -sheaf on X for $B = \{\text{basic open sets } D_f, f \in R\}$

PF Uniqueness: $\alpha, \beta \in R_f = \mathcal{O}_X(D_f)$ and $D_f = \bigcup D_{f_i}$

$\uparrow$ if $\alpha|_{D_{f_i}} = \beta|_{D_{f_i}} \forall i$ then $\alpha = \beta$

Proof By redefining X, R by D_f, R_f we can assume $f=1, R_f=R, D_f=X$.

$\alpha - \beta = 0 \in R_f \Rightarrow f_i \cdot (\alpha - \beta) = 0$ some $N \in \mathbb{N} \leftarrow N$ may depend on i , but WLOG finite subcover D_{f_i}

$\Rightarrow < \text{all } f_i^N > \cdot (\alpha - \beta) = 0$ (quasi-compactness) so pick maximal N

recall "Covering Trick" $\rightarrow R$ since $X = D_{f_1} \cup \dots \cup D_{f_N} = D_{f_1^N} \cup \dots \cup D_{f_N^N} \leftarrow (\text{recall } D_f = D_{f^N})$

$\Rightarrow 1 \cdot (f_1 - \beta) = 0$ so $\alpha = \beta$ $\square$

Existence in $\star$: as before $\text{WLOG } U = D_f, R_f$ become X, R .

Uniqueness $\Rightarrow$ in $\star$ can assume sections $s \in \mathcal{O}_X(D_{f_i})$ agree on overlaps $D_{f_i} \cap D_{f_j} = D_{f_i f_j}$

(apply Uniqueness) $s_i|_{D_{f_i f_j}} = s_j|_{D_{f_i f_j}} \in R_{f_i f_j}$

$\text{WLOG } X = D_{f_1} \cup \dots \cup D_{f_N}$ finite cover, $s_i = \frac{g_i}{f_i^m}$ since $D_{f_i} = D_{f_i^m}$, $\text{WLOG } m=1$, so $s_i = \frac{g_i}{f_i}$

$s_i = s_j$ on $D_{f_i f_j} \Rightarrow (f_i f_j)^m (f_i g_j - f_j g_i) = 0 \in R \leftarrow N$ depends on i, j but can pick largest N over finitely many i, j so N works $\forall i, j$

rewrite: $(f_j^{N+1}) \cdot \underbrace{(f_i^N g_j)}_{a_i} - \underbrace{(f_i^{N+1}) \cdot (f_j^N g_i)}_{b_j} = 0$ notice $s_i = \frac{g_i}{f_i}, D_{f_i} = D_{f_i^N}$ so $\text{WLOG } N=0!$

"Covering Trick": $X = D_{f_1} \cup \dots \cup D_{f_N}$ so $1 = \sum r_i f_i \leftarrow$ ("partition of unity" trick)

$1 \cdot g_j = (\sum_i r_i f_i) g_j = \sum_i r_i (f_i g_j) = \sum_i r_i (f_i g_i) = f_j (\sum_i r_i g_i)$

$\Rightarrow s_j = \frac{g_j}{f_j} = \sum_i \frac{r_i g_i}{f_j} \in R_{f_j} \forall j$ so we globalised the $s_j \in \mathcal{O}_X(D_{f_j})$ to $\sum_i r_i g_i \in \mathcal{O}_X(X) = R$ $\square$

Corollary $\mathcal{O}_X$ extends uniquely to a sheaf on $X = \text{Spec } R$ called structure sheaf (or sheaf of regular functions)

stalk $\mathcal{O}_{X, P} := \varprojlim_{D_f \ni P} \mathcal{O}_X(D_f) \leftarrow$ Messy unpacking of definitions: we identify $\frac{f}{g} \in R_f \cong \mathcal{O}_X(D_f)$ and $\frac{s}{g} \in R_g \cong \mathcal{O}_X(D_g)$ iff $\frac{f}{g} = \frac{s}{g} \cdot \frac{g}{g} \in R_g$ some $h \in R$ with $h \in D_g \subseteq D_f \cap D_g$ (iff $R^N (rg^N - sf^N) = 0 \in R$ some N) D_{fg}

Lemma 2 $\mathcal{O}_{X, P} \cong R_P \xleftarrow{\text{rest.}} \mathcal{O}_X(X) \cong R \xleftarrow{\text{localise}} \mathcal{O}_X(X) \cong R$

PF $\varprojlim_{D_f \ni P} \mathcal{O}_X(D_f) \cong \varprojlim_{f \notin P} R_f \cong R_P$ $\leftarrow$ straight forward algebra exercise $\leftarrow$ Recall in R_P you invert all elements $f \notin P$

$\Rightarrow \theta_X(U) = \{s : U \rightarrow \bigsqcup_{p \in X} R_p : s(p) \in R_p \text{ which are locally compatible}\}$

$\forall p \in U, \exists \text{ open nbhd } p \in D \subseteq U \text{ with } s(x) = t_x$
 $\forall t \in \theta_X(D_f) \uparrow \text{ some } f \in R$
 $\Downarrow \text{ some } f \in R$
 $\Downarrow \text{ some } f \in R$
 $\Downarrow \text{ some } f \in R$

Rmk. could assume $t = \frac{f}{g}$ since can replace D_f with D_{fg} ($= D_f$).
 • could just ask $s(x) = t_x$ on a smaller open $p \in V \subseteq D_f$.

Comparison with classical algebraic geometry

- X affine variety, $p \in U \subseteq X$ open nbhd
 $f : U \rightarrow k$ is regular at p if $\exists$ open nbhd $p \in W \subseteq U$ with $f|_W = g/h$ for $g, h \in k[X], h(p) \neq 0$

Rmk In fact can assume $W = D_g$ basic open (if $f = \frac{g}{h}$, replace D_g by $D_{gh} = D_g$)
 $\theta_X(U) = k$ -algebra of functions $U \rightarrow k$ regular at all $p \in U$
 $\theta_{X,p} = k$ -algebra of germs of functions near p , regular at p

(so pairs (U, f) with $p \in U \subseteq X$ open, $f : U \rightarrow k$ regular at p
 (and identify $(U, f) \sim (V, g) \Leftrightarrow f|_W = g|_W$ on some open $p \in W \subseteq U \cap V$)
 Theorem $\theta_X(X) \cong k[X] \leftarrow \text{Rmk This theorem is not obvious in C3.4 course.}$
 $X = \text{Spec } k[X]$ so by Lemma 1 get $\theta_X(X) = k[X]$

- $X \subseteq \mathbb{A}^n$ affine variety
 $f \in R = k[x_1, \dots, x_n]$ polynomial
 $V(f) = \{f = 0\} \subseteq X$ hypersurface
 $D_f = \{f \neq 0\} \subseteq X$ open, but identifiable
 with affine variety $Y = V(zf - 1) \subseteq \mathbb{A}^{n+1} (D_f \rightarrow Y, a \mapsto (a, \frac{1}{a}))$

and $k[Y] = k[X]/(zf - 1) \cong k[X]_f$ via $\cong \xrightarrow{f}$
 fact $\theta_X(D_f) \cong k[X]_f$
 $\theta_{X,p} \cong k[X]_m$ $\leftarrow$ where $m_p = \mathbb{I}(p) = \{f \in k[X] : f(p) = 0\}$
 is max ideal corresponding to p .

local ring $m_{X,p} = m_p \cdot k[X]_m$ = germs of functions near p vanishing at p
 $m_{X,p} = \theta_{X,p} / m_{X,p} \cong k, \frac{g}{h} \mapsto \frac{g(p)}{h(p)}$
 residue field $K(p) = \theta_{X,p} / m_{X,p} \cong k$
 Morphisms:
 $\alpha : X \rightarrow Y \Rightarrow \alpha^* : \theta_Y(U) \rightarrow \theta_X(\alpha^{-1}(U)), \alpha^*(f : U \rightarrow k) = (\alpha^* f) : \alpha^{-1}(U) \rightarrow k$
 (morph of aff. vars.)

1.13 Morphisms between Specs

$\varphi : R \rightarrow S$ hom of rings $\Rightarrow \text{Spec}(\varphi) : \text{Spec } S \rightarrow \text{Spec } R$
 $p \mapsto \varphi^{-1}(p)$

Example $\varphi : R \rightarrow R_f, r \mapsto \frac{r}{f}$ localisation

$\text{Spec } R \leftarrow \text{Spec } R_f$ is an "inclusion" with image $= D_f$.

$\alpha = \text{Spec}(\varphi) : Y \rightarrow X, p \mapsto \varphi^{-1}(p)$

Lemma $\alpha^{-1}(D_f) = D(\varphi(f))$

$\text{pf } \alpha^{-1}\{q \in X : f \notin q\} = \{p \in Y : \varphi^{-1}(p) = q \text{ some } q \in X, f \notin \varphi^{-1}(p)\}$
 $= \{p \in Y : \varphi(f) \notin p\}.$ $\square$

Claim $\exists \varphi^* : \theta_X \rightarrow \alpha_* \theta_Y$ such that $\varphi^* : \theta_X(X) = R \xrightarrow{\varphi} S = \alpha_* \theta_Y(X)$

pf Enough to build φ^* on basic opens, compatibly with restrictions

$$\varphi^* : \theta_X(D_f) \rightarrow \alpha_* \theta_Y(D_f) = \theta_Y(\alpha^{-1}(D_f)) \xrightarrow{\cong} \theta_Y(D_{\varphi(f)})$$

$$R_f \xrightarrow{\text{natural hom}} S_{\varphi(f)} \xrightarrow{\text{natural map}} \frac{\varphi(r)}{\varphi(f)}$$

Easy check: compatible with restriction maps for $D_g \subseteq D_f$. $\square$

(By theorem on B-sheaves)

Claim $\theta_{X,p}$ is local and φ_p^* is local

pf Lemma 2: $\theta_{X,p} \cong R_p$ so local with max ideal $m_p = p \cdot R_p$.
 For $p \in Y, \varphi_p^* : \theta_{X,p} \rightarrow \theta_{Y,p}$
 is local.
 Hint: $\varphi(r) \notin p \Rightarrow r \notin \varphi^{-1}(p)$
 natural map: $\frac{r}{t} \mapsto \frac{\varphi(r)}{\varphi(t)}$
 $t \notin \varphi^{-1}(p) \Rightarrow \varphi(t) \notin p$
 is direct limit of maps hence:

$\Rightarrow$ Theorem (ring $R \rightarrow S$) $\rightarrow$ locally ringed space $(\text{Spec } R, \theta_{\text{Spec } R}) \rightarrow (\text{Spec } S, \theta_{\text{Spec } S})$

(ring hom $R \xrightarrow{\varphi} S \rightarrow (\text{Spec } \varphi, \varphi^*) : (\text{Spec } R, \theta_{\text{Spec } R}) \rightarrow (\text{Spec } S, \theta_{\text{Spec } S})$)

Contravariant functor $\text{Spec} : \text{Rings} \rightarrow \text{Locally Ringed Spaces}$ (easy to check)

Claim The functor is fully faithful $\leftarrow$ i.e. surj & inj (so iso) on morphism spaces

pf Given a hom of loc. ringed spaces $(f, f^*) : (Y, \theta_Y) \rightarrow (X, \theta_X)$ $X = \text{Spec } R, Y = \text{Spec } S$

Let $\varphi := f^* : R \cong \theta_X(X) \xrightarrow{f^*} \theta_Y(X) = \theta_Y(Y) \cong S$ ring hom.

$$R \cong \theta_{X,p} \xrightarrow{f_p^*} \theta_{Y,p} \cong S_p = p \cdot S_p$$

$\Rightarrow \varphi^{-1}(p) = \varphi^{-1}(p \cdot S_p) = \varphi^{-1}(f_p^*(m_p)) = \varphi^{-1}(f_p^*(m_p)) = f_p^{-1}(m_p) = f_p^{-1}(m_p)$

$\Rightarrow \varphi^{-1}(p) = \varphi^{-1}(p \cdot S_p) = \varphi^{-1}(f_p^*(m_p)) = \varphi^{-1}(f_p^*(m_p)) = f_p^{-1}(m_p) = f_p^{-1}(m_p)$
 diagram $m_{X,p} \xrightarrow{f_p^*} m_{Y,p}$ since $f_p^* \text{ local ring hom}$

$\Rightarrow f(p) = \varphi^{-1}(p)$ so $f = \text{Spec}(\varphi)$ is the map on Specs induced by $\varphi: R \rightarrow S$.

Upshot: have two morphs of sheaves $f^\#, \varphi^\# : \mathcal{O}_X \rightarrow \text{Spec}(\varphi)_* \mathcal{O}_Y$ and $f^\# = \varphi^\#$ since equal on stalks (by the diagram have $f^\# = \varphi^\#$)

Def Aff = category of affine schemes (and morphs of locally ringed spaces)

$\Rightarrow \text{Spec} : \text{Rings}^{\text{op}} \rightarrow \text{Aff}$ is an equivalence of categories.
(op = opposite category = reverse arrows so artificially make Spec covariant)

1.14 Closed affine subschemes
 $X = \text{Spec } R, I \subseteq R$ ideal (think same as specifying a subscheme) is iso to an object in image $Y = V(I) \cong \text{Spec}(R/I)$ are called closed (affine) subschemes of X
(as top space, $V(I) = V(\sqrt{I})$ but sheaf remembers $I : \mathcal{O}_Y(Y) = R/I$)

Example $I = m$ max ideal $\Rightarrow$ get a closed point $\{m\} = \text{Spec } R/m \hookrightarrow X$.

Rmk $\text{Spec}(R/I)$ is closed subscheme of $\text{Spec}(R/I)$ means $J \supseteq I \Rightarrow \text{Spec } R/I \cap \text{Spec } R/J = \text{Spec } R/I$

Def $\text{Spec } R/I \cap \text{Spec } R/J = \text{Spec}(R/I+J)$, $\text{Spec } R/I \cup \text{Spec } R/J = \text{Spec } R_{I \cap J}$

Define sheaf of ideals $J = J_{X/Y}$ on X :
(also: ideal sheaf) $J(D_f) = I \cdot R_f \subseteq R_f = \mathcal{O}_X(D_f)$ ideal

Notice $\mathcal{O}_Y(D_f) = (R/I)_f \cong R_f/I \cdot R_f = \mathcal{O}_X(D_f)/J(D_f)$

$\Rightarrow J = \text{Ker}(\mathcal{O}_X \rightarrow j_* \mathcal{O}_Y)$
 $\mathcal{O}_Y = \mathcal{O}_X/J$ where $j : Y \rightarrow X$ inclusion.
more precisely this is $j_* \mathcal{O}_Y$

Def A sheaf of ideals on $X = \text{Spec } R$ is quasi-coherent if it arises as J as above, some ideal $I \in R$

Rmk Later will consider more generally sheaves of R -modules and quasi-coherence.

1.15 Closed subschemes
 $(X, \mathcal{O}_X)$ scheme, sheaf of ideals J means $J(U) \subseteq \mathcal{O}_X(U)$ ideal compatibly with restrictions.

Quasi-coherent means: $\forall$ affine open $U, J|_U$ is quasi-coherent.
closed subscheme means $\cdot Y \subseteq X$ closed topological space

$\cdot \mathcal{O}_Y = \mathcal{O}_X/J$ some quasi-coherent sheaf of ideals J on X ,
s.t. $Y \cap (\text{affine open } U) \subseteq U$ is closed affine subscheme for the ideal $J(U) \subseteq \mathcal{O}_X(U)$.

Rmk $\exists$ 1:1 correspondence $\{\text{closed subschemes of } X\} \leftrightarrow \{\text{quasi-coh. sheaves of ideals on } X\}$

Can recover $Y \subseteq X$ from the support of $\mathcal{O}_X/J$:
 $Y = \text{Supp } \mathcal{O}_X/J = \{x \in X : (\mathcal{O}_X/J)_x \neq 0\} = \{x \in X : J_x \neq \mathcal{O}_{X,x}\}$

Example closed point $p \in X$ (so $\overline{\{p\}} = \{p\}$) $\Rightarrow$ pick affine $p \in \text{Spec } R \hookrightarrow X$ then $p \leftrightarrow (\text{max}) \subseteq R$
 $\Rightarrow$ sheaf J on $\text{Spec } R \Rightarrow$ extend J to X by $J(V) = \mathcal{O}_X(V)$ if $p \notin V$ (so $\mathcal{O}_Y(V) = 0$)

2. GLOBAL SECTIONS AND THE FUNCTOR OF POINTS

2.0 Points of Spec R (not necessarily closed)

$R \xrightarrow{\text{loc}} R_p \xrightarrow{\text{quotient}} K(p) = R_p/m_p \Rightarrow \text{Spec } K(p) \hookrightarrow \text{Spec } R_p \hookrightarrow \text{Spec } R$
 $\text{loc}^{-1}(m_p) = p \leftarrow p \cdot R_p = m_p \leftarrow (0) \xrightarrow{\{0\}} \{0\}$

So points of $\text{Spec } R$ correspond to the max ideals in the local rings.

2.1 Global sections and basic open sets for locally ringed spaces

$(X, \mathcal{O}_X)$ locally ringed space $\Gamma(\cdot, \mathcal{O}_X) : \text{Top}(X)^{\text{op}} \rightarrow \text{Rings}$, include $\uparrow$ sections functor $\downarrow$ restrict $V \hookrightarrow X \hookrightarrow \mathcal{O}_X(V)$

global sections functor: Locally Ringed Spaces $\text{op} \rightarrow \text{Rings}$, $(X, \mathcal{O}_X) \mapsto \Gamma(X, \mathcal{O}_X) = \mathcal{O}_X(X)$
 $\exists$ canonical map $X \rightarrow \text{Spec } \mathcal{O}_X(X)$, $x \mapsto \text{res}_x^{-1}(m_{x,x})$ where $\text{res}_x : \mathcal{O}_X(x) \rightarrow \mathcal{O}_{x,x}$ restricts.

Trick $f \in \mathcal{O}_X(X)$ then $f_x \in \mathcal{O}_{X,x}$ invertible $\Leftrightarrow f(x) \neq 0 \in K(x) = \mathcal{O}_{X,x}/m_x$
Pf $f_x \in \mathcal{O}_{X,x} \setminus m_x = \{\text{invertibles of } \mathcal{O}_{X,x}\} \Leftrightarrow f_x \notin m_x \square$

image of f via $\mathcal{O}_X(X) \rightarrow \mathcal{O}_{x,x} \rightarrow K(x) \hookrightarrow f(x)$
 $\Leftrightarrow f \notin m_x \Leftrightarrow (f_x \in \mathcal{O}_{X,x})$ is open in X .

Lemma $f \in \mathcal{O}_X(X) \Rightarrow D_f = \{x \in X : f(x) \neq 0 \in K(x)\}$ is open in X .

Pf Trick $\Rightarrow \exists g \in \mathcal{O}_{X,x} : f \cdot g = 1$ so $\exists$ open $U \subseteq X$ s.t. $f, g \in \mathcal{O}_X(U)$, $f \cdot g = 1 \in \mathcal{O}_X(U)$
 $\Rightarrow x \in U \subseteq D_f$ since $\forall g \in U, f \cdot g = 1 \in \mathcal{O}_{X,g} = (f \cdot g)_g = 1 \in \mathcal{O}_{X,g}$ so $f_g \in \text{invertibles of } \mathcal{O}_{X,g}$ so $f(x) \neq 0$, so $y \in D_f \square$

Lemma $f|_{D_f} \in \mathcal{O}_X(D_f)$ is invertible
Pf Lemma $\Rightarrow f$ is locally invertible. If $f \cdot h = 1$ on U then $h = g$ on $U \cap V$. So can globalize. $\square$

2.2 What it means to be affine
 $\hookrightarrow$ locally ringed space

$(X, \mathcal{O}_X)$ affine $\Leftrightarrow \exists$ ring $R : \exists X \xrightarrow{\alpha} Y = \text{Spec } R$ homeomorph, and $\exists \mathcal{O}_Y \xrightarrow{\cong} \alpha_* \mathcal{O}_X$

But $\mathcal{O}_Y(Y) = R$ so $R \xrightarrow{\cong} \mathcal{O}_X(X)$ so $\text{Spec } \mathcal{O}_X(X) \xrightarrow{\cong} Y$.

$\varphi_x \text{ local} \Rightarrow \mathcal{O}_X \xrightarrow{\cong} \mathcal{O}_X(X) \xrightarrow{\cong} R \xrightarrow{\cong} \mathcal{O}_X(x) \xrightarrow{\cong} \mathcal{O}_{X,x} \xrightarrow{\cong} R_{\alpha(x)} \xrightarrow{\cong} \mathcal{O}_{X,x}$

So a locally ringed space $(X, \mathcal{O}_X)$ is affine precisely if:

- the canonical map $X \rightarrow \text{Spec } \Gamma(X, \mathcal{O}_X)$ is homeomorph
- $\mathcal{O}_X(D_f) \cong \Gamma(X, \mathcal{O}_X)_f \forall f \in \Gamma(X, \mathcal{O}_X)$ and restrictions are localizations $\leftarrow$ (by Sec. 1.12)

2.3 Functor of points by

MOTIVATION Y set, you recover set Y from $\text{Mor}(\text{point}, Y)$
 Y group, " " " $\text{Mor}(\mathbb{Z}, Y)$

Functor of points $h_Y : \text{Sch}^{\text{op}} \rightarrow \text{Sets}$, $h_Y(X) = \text{Mor}(X, Y)$
 $\xrightarrow{f \leftarrow \mathbb{Z}} \rightarrow$ on morphisms: $h_Y(X \xleftarrow{f} \mathbb{Z}) = (\text{Mor}(X, Y) \xrightarrow{\circ f} \text{Mor}(\mathbb{Z}, Y))$
 MOTIVATION

$Y = \text{Spec } \mathbb{Z}[X]/(x^2+1)$. $\mathbb{C}$ -valued points of Y ?
 $\mathbb{Z}[X]/(x^2+1) \rightarrow \mathbb{C}, x \mapsto i \Rightarrow \text{morph } X = \text{Spec } \mathbb{C} \rightarrow Y$ so $i \in h_Y(X) \Leftarrow \text{often write } Y(\mathbb{C})$

HWK 1 natural transformations $\text{functor } F \leftarrow \text{functor } G$ take image of $\text{id}_Y \in \text{Mor}(Y, Y) = h_Y(Y)$ given $F(Y) \xrightarrow{F(\text{id}_Y)} F(Y)$
Yoneda Lemma $\text{Nat}(h_Y, F) \cong F(Y)$ $\leftarrow$ Conversely given $\alpha \in F(Y)$, $\varphi \in h_Y(Y)$ get $F(\varphi)(\alpha) \in F(Y)$

Yoneda embedding $h_Y : \text{Sch} \rightarrow \text{Sets}^{\text{Sch}^{\text{op}}}$ $Y \mapsto h_Y$ is fully faithful (iso on morphisms: $\text{Nat}(h_Y, h_W) \cong \text{Mor}(Y, W)$)
UPSHOT $\text{Nat}(h_Y, h_W) \cong Y \cong W$

② Can now ask which functors $\text{Sch}^{\text{op}} \rightarrow \text{Sets}$ are $\cong h_Y$, i.e. represented by a scheme Y .
 Example Will show that $\mathbb{A}^n = \text{Spec } \mathbb{Z}[x_1, \dots, x_n]$ represents $\leftarrow$ "tell me who your friends are" and I will tell you who you are $\leftarrow Y$
 $\text{Sch}^{\text{op}} \rightarrow \text{Sets}, X \mapsto \{\text{morphs } \bigoplus_{i=1}^n \theta_X \rightarrow \theta_X \text{ which are } \theta_X\text{-linear}\}$

Example 1 $h_{\text{Spec } R} : \text{Sch} \rightarrow \text{Sets}^{\text{Sch}^{\text{op}}}$ $Y \mapsto h_Y$ is fully faithful (iso on morphisms: $\text{Nat}(h_Y, h_W) \cong \text{Mor}(Y, W)$)
UPSHOT $\text{Nat}(h_Y, h_W) \cong Y \cong W$

KEY EXAMPLE $Y = \mathbb{A}^1 = \text{Spec } \mathbb{Z}[X]$
 $\text{Mor}(X, \mathbb{A}^1) \cong \text{Mor}(X, \text{Spec } R) \rightarrow \text{Hom}(R, \Gamma(X, \theta_X))$ bijective
 $\text{Spec } R \xrightarrow{\vartheta} \Gamma(X, \theta_X)$

$\text{pf } \text{pf}_Y(Y) \xrightarrow{\vartheta} \theta_X(X) \rightarrow \theta_{X,X}$ preimage of m_X gives $p \in \text{Spec } R = Y$
 $\text{R} \xleftarrow{\vartheta} Y = \text{Spec } R$ defines $g : X \rightarrow Y$, $g(x) = p$
 $\cdot g$ is continuous (check $g^{-1}(D_f) = D_{\vartheta f}$) $\leftarrow$ (see 2.1 for basic)

$\cdot \theta_Y(D_f) = R_f \xrightarrow{\vartheta_f} \theta_X(X) \rightarrow \theta_X(D_{\vartheta f}) = \theta_X(g^{-1}D_f) = g_X(D_f)$
 $\uparrow$ localise ϑ
 natural map induced by restriction $\theta_X(X) \rightarrow \theta_X(D_{\vartheta f})$
 since ϑ_f invertible in $\theta_X(D_{\vartheta f})$ see 2.1

These are compatible with restrictions $\square$
 Universal property of localisation: $R \xrightarrow{\vartheta} R_2$ and $\vartheta(S) \subseteq \text{invertibles of } R_2 \Rightarrow \exists ! R_1 \rightarrow S^{-1}R_1 \rightarrow R_2$.
 Obvious: $\frac{1}{s} \mapsto \vartheta(s)$

Cor 1 (X, θ_X) scheme $\Rightarrow$ canonical morph $X \rightarrow \text{Spec } \Gamma(X, \theta_X)$
 Explicitly: on sets $x \mapsto \text{res}^{-1}(m_{x,x}) \in \theta_X(X)$
 On sheaves over $D_f \subseteq X : \theta_X(X)_f \text{ res } \theta_X(D_f) \xrightarrow{\text{now localise at } f} \theta_X(D_f)$
 sheaves over $D_f \subseteq X : \theta_X(X)_f \text{ res } \theta_X(D_f) \xrightarrow{\text{now localise at } f} \theta_X(D_f)$

Rmk Canonical morph is injective if global sections separate points meaning:
 $x \neq y \in X \Rightarrow \exists f \in \Gamma(X, \theta_X), f(x) \neq f(y)$ (equivalently $\exists f : f(x) = 0, f(y) \neq 0$)
 $f \in m_{x,x} \not\subseteq m_{y,y}$

Classical algebraic geom. $X \subseteq \mathbb{A}^n$ affine variety $(X = \mathbb{V}(I), I \subseteq k[x_1, \dots, x_n])$
 so $\Gamma(X, \theta_X) = k[X]$, $\theta_X(D_f) = k[X]_f$, $\theta_X(U) = \{ \text{regular functions} \}$, $\theta_{X,a} = k[X]_{m_a}$
 separates points, and $X \xrightarrow{\text{ini}}$ $\{ \text{closed points} \} \subseteq \text{Spec } k[X]$

$a \mapsto \text{max ideal } m_a \subseteq k[X]$ $\leftarrow$ max ideal of $\theta_{X,a}$
 in fact get embedding $\{ \text{Category of Affine Varieties} \} \hookrightarrow \text{Sch}$

Example 2 $h_Y(\text{Spec } R) \Rightarrow X = \text{Spec } R \Rightarrow \{ f \in \text{Mor}(\text{Spec } R, Y) \}$ $\xleftarrow{!} \text{Hom}_{\text{local rings}}(\theta_{Y,Y}, R)$
 $m \subseteq R = \text{local ring}$ with $f(m) = y$

Pf $\text{Spec } R \xrightarrow{f} Y$
 $\downarrow \quad \downarrow$
 $m \mapsto y$

Affine case $Y = \text{Spec } S$
 $\varphi : S_y \rightarrow R \Rightarrow S \otimes_S S_y \rightarrow R \Rightarrow \text{Spec } R \rightarrow \text{Spec } S$
 $\varphi^*(m) = y \cdot S_y \quad m \mapsto (\text{preimage of } y^m) = y$

General case
 $y \in U \subseteq Y$ open affine, then $\theta_{U,y} = \theta_{Y,y} \xrightarrow{\vartheta} R$ gives $\text{Spec } R \rightarrow U \subseteq Y$
 uniqueness: Suppose $f : \text{Spec } R \rightarrow Y$ gives same ϑ
 $m \mapsto y$

pick $y \in V \subseteq Y$ affine open $\Rightarrow f^{-1}(V)$ open $\ni m = (\text{unique closed point of Spec } R) \Rightarrow f^{-1}(m) = \text{Spec } R$
 so $f : \text{Spec } R \rightarrow V \subseteq Y$ so reduce to affine case. $\square$

Cor 2 $x \in X \Rightarrow \exists$ canonical morph $\text{Spec } \theta_{X,x} \rightarrow X$.
 (By Example 2 for $(\text{id} : \theta_{X,x} \rightarrow \theta_{X,x})$) Any $\text{Spec } R \rightarrow X$ factors as $\text{Spec } R \rightarrow \text{Spec } \theta_{X,x} \rightarrow X$ some $x \in X$.

induced by a local ring hom

Any $f : X \rightarrow Y$ of schemes get $\text{Spec } \theta_{X,x} \rightarrow X \xrightarrow{f} Y$
 $\downarrow \quad \downarrow$
 $m_{X,x} \mapsto z$
 $\downarrow \quad \downarrow$
 $m_{Y,y} \mapsto y$

induced by $f_x^{\#}$

Example Case $X = \text{Spec } k$ for field k .
 $R \text{ local} \Rightarrow \text{residue field } k = R/m$
 A local hom $R \xrightarrow{\vartheta} k = \text{field}$ factors $R \xrightarrow{\text{quot}} k \rightarrow k$
 (since $\text{Ker } \vartheta = \vartheta^{-1}(0) = m$) (since local hom)

Thus: $\{ f \in \text{Mor}(\text{Spec } k, Y) \} \xleftarrow{!} \text{Hom}(k(y), k)$ and any $\text{Spec } k \rightarrow Y$ factors: $\text{Spec } k \rightarrow \text{Spec } k(y) \rightarrow Y$

UPSHOT : Morphs from local rings or fields don't give more information than already know from $\text{Spec } \theta_{X,x} \rightarrow X$ and $\text{Spec } k(x) \rightarrow X$.

Non-separable: $\text{Rmk } y \in Y$ called k -valued point if $k(y) \cong k$, then id_k defines a morph $\text{Spec } k \rightarrow Y$
 (or k -point, or k -rational point)

If Y comes with a morph $Y \xrightarrow{\pi} \text{Spec } k$ (hence $\theta_{Y,U}$ are k -algebras) and above require morphs to commute with π , then get $\text{Hom}_k(k(y), k)$, and if $k(y) \cong k$ then $\text{Hom}_k(k(y), k) = \{ \text{id}_k \}$. E.g. $\text{Spec } (\mathbb{C})$ has many $\mathbb{C}$ -points, one for each automorphism of $\mathbb{C}$ (e.g. $\sigma \in \mathbb{C} \rightarrow \mathbb{C}, z \mapsto \bar{z}$) but if work over $\mathbb{R}$ get two $\mathbb{C}$ -points.

3. PROPERTIES OF SCHEMES

3.0 Useful facts from commutative algebra: localisation

R ring, M R -mod, $S \subseteq R$ multiplicative set $\xleftarrow{\text{write "M"}}$ $1 \in S, S \cdot S \subseteq S$
 $\Rightarrow$ localisation $S^{-1}M = M \otimes_R S^{-1}R$ relation $(m, s) \sim (n, t) \Leftrightarrow u \cdot (tm - sn) = 0$ $\swarrow$ some $u \in S$

which is an $S^{-1}R$ -mod and have R -mod hom $M \rightarrow S^{-1}M$ localisation map.

Fact $S^{-1}M \cong M \otimes_R S^{-1}R$ canonically $\leftarrow$ (via $\frac{m}{s} \mapsto m \otimes \frac{1}{s}$ and $\sum \frac{r_i m_i}{s_i} \mapsto \sum m_i \otimes \frac{r_i}{s_i}$)

Exercise $\alpha: M \rightarrow N$ hom (of R -mods) $\Rightarrow \exists$ natural $S^{-1}\alpha: S^{-1}M \rightarrow S^{-1}N$

Fact Localisation is an exact functor.

Cor $S^{-1}(M/N) \cong S^{-1}M/S^{-1}N$

Pf apply S^{-1} to exact sequence $0 \rightarrow N \rightarrow M \rightarrow M/N \rightarrow 0$ $\swarrow$ (indeed take $N = \text{preimage}$ via $M \rightarrow S^{-1}M$)

Fact Submods of $S^{-1}M$ have form $S^{-1}N$ for submod $N \subseteq M$

Fact $S^{-1}M = \varinjlim M_f$ via localisation maps $M_f \rightarrow M_g$ whenever $g = fh$

(e.g. proof: $\varinjlim M \otimes_R f = M \otimes_R S^{-1}R$) $\swarrow$ (induced by $R \xrightarrow{f} R_f$ via $M \otimes_R f \rightarrow M \otimes_R R_f$)

Local algebra theorem $\swarrow$ multiplicative set $S = R \setminus \mathfrak{p}$

① $x \in M: x = 0 \Leftrightarrow x_{\mathfrak{p}} = 0 \in M_{\mathfrak{p}} \forall \mathfrak{p} \in \text{Spec } R$

② $M = 0 \Leftrightarrow M_{\mathfrak{p}} = 0 \forall \mathfrak{p} \in \text{Spec } R$

③ $M \xrightarrow{\alpha} M' \xrightarrow{\beta} M''$ exact $\Leftrightarrow M_{\mathfrak{p}} \xrightarrow{\alpha_{\mathfrak{p}}} M'_{\mathfrak{p}} \xrightarrow{\beta_{\mathfrak{p}}} M''_{\mathfrak{p}}$ exact $\forall \mathfrak{p} \in \text{Spec } R$

④ $f: M \rightarrow N$ inj. $\Leftrightarrow f_{\mathfrak{p}}: M_{\mathfrak{p}} \rightarrow N_{\mathfrak{p}}$ inj. $\forall \mathfrak{p} \in \text{Spec } R$

" " surj. " " iso. " " $\swarrow$ (if $\text{Ann } x \neq 0$ unless $x=0$)

Pf ① $\Leftrightarrow \text{Ann}(x) = \{r \in R: rx=0\}$ ideal $\subseteq$ max ideal $\mathfrak{m}$ (unless $x=0$)

$x_{\mathfrak{m}} = 0 \in R_{\mathfrak{m}} \Rightarrow \exists r \in R \setminus \mathfrak{m}$ s.t. $rx = 0 \in R$ $\Rightarrow$ (since $r \notin \text{Ann}(x)$)

by ① $\Rightarrow H := \text{Ker } \beta / \text{Im } \alpha \Rightarrow H_{\mathfrak{p}} \cong (\text{Ker } \beta)_{\mathfrak{p}} / (\text{Im } \alpha)_{\mathfrak{p}} = \text{Ker } \beta_{\mathfrak{p}} = 0$ now use ②

(holds since localisation is exact) $\left(\begin{array}{l} \text{since } 0 \rightarrow \text{Ker } \beta \xrightarrow{\text{inj}} M' \xrightarrow{\beta} \text{Im } \beta \rightarrow 0 \text{ exact} \\ \Rightarrow 0 \rightarrow (\text{Ker } \beta)_{\mathfrak{p}} \xrightarrow{\text{inj}} M'_{\mathfrak{p}} \xrightarrow{\beta_{\mathfrak{p}}} (\text{Im } \beta)_{\mathfrak{p}} \rightarrow 0 \text{ exact, so} \end{array} \right)$

④ by ③ $\leftarrow$ (e.g. inj means $0 \rightarrow M \xrightarrow{f} N$ exact) $\square$

Rmk $\text{Spec } R = \bigcup D_{f_i}$ then above results hold $\Leftrightarrow$ hold when localise at each f_i

Pf $x_i = 0 \in M_{f_i} = M \otimes_R R_{f_i} \Rightarrow$ localise further at $\mathfrak{p} \in \text{Spec } R_{f_i}: M_{f_i} = M \otimes_R R_{f_i} \rightarrow M \otimes_R R_{\mathfrak{p}} = M_{\mathfrak{p}}$

(Note every $\mathfrak{p} \in \text{Spec } R$ is in some $D_{f_i} = \text{Spec } R_{f_i}$) $\swarrow$ $0 = x_i \mapsto \frac{x_i}{1} \in R_{f_i} \xrightarrow{R \rightarrow R_{\mathfrak{p}}} R_{\mathfrak{p}} \rightarrow 0$ $\square$

3.1 Noetherian

Recall: ring R is $\Leftrightarrow$ ideals of $R \Leftrightarrow$ submods of f.g. R -mods are f.g. Noetherian

Rmk localisation and quotients preserve Noetherian property

Def An affine open (for the ring R) means an open subset $U \subseteq X$ admitting an isomorphism

$(U, \theta_X|_U) \cong (\text{Spec } R, \theta_{\text{Spec } R})$ for some ring R . [Note: $\theta_X(U) \in R$

Def scheme (X, θ_X) is Noetherian if quasi-compact and locally Noetherian:

Claim The following are equivalent definitions for (X, θ_X) to be locally Noetherian

1) every point has an affine open neighbourhood U with $\theta_X(U)$ Noetherian

2) $X = \bigcup U_i$ for open affines U_i with $\theta_X(U_i)$ Noetherian

3) given any open affine for a ring R , R must be Noetherian

Pf (1) $\Leftrightarrow$ (2) and (3) $\Rightarrow$ (1) since schemes are locally affine.

(1) $\Rightarrow$ (2) $\Rightarrow$ (3): consider $\text{Spec } R \cong U \subseteq X$

$\forall \mathfrak{p} \in U, \exists$ affine open $\mathfrak{p} \in V = \text{Spec } S \subseteq X$ with S Noetherian (by (1))

$\Rightarrow \exists$ basic open $\mathfrak{p} \in D_g \subseteq U$ for $\text{Spec } S$, some $g \in S$

By the useful trick, $\text{Wlog } D_g$ is basic also for $\text{Spec } R$, say $\text{Spec } R_f$.

Since $\text{Spec } S_g \subseteq \text{Spec } R_f$ get $S_g \cong R_f$ so Noetherian. Get over for U ,

so need: Algebra Lemma R_{f_i} Noeth. $\forall i \Rightarrow R$ Noeth.

proof $I \subseteq R$ ideal (aims: I is f.g.) $\leftarrow$ by "Covering Trick"

$\Rightarrow I_{f_i} := I \cdot R_{f_i} \subseteq R_{f_i}$ ideal, f.g. since R_{f_i} Noeth., say generators $g_{ij} = \frac{h_{ij}}{f_i^n}$

$\Rightarrow f_i^n \cdot g_{ij} = h_{ij}$ also generate (since $\frac{h_{ij}}{f_i^n} \in R_{f_i}$) (localisation at f_i) $\swarrow$ (some $h_{ij} \in I$) $\swarrow$ f.i.

$\Rightarrow \bigoplus_i R \xrightarrow{q} I, e_{ij} \mapsto h_{ij}$ satisfies q_{f_i} surjective $\forall f_i$ so q surj. $\square$

Exercise give an alternative proof of algebra lemma by proving the ACC for R

(Key trick: $I = \bigcap \varphi_i^{-1}(I_{f_i})$ where $\varphi_i: R \rightarrow R_{f_i}$ is localisation.

You may need the famous Trick: $\text{Spec } R = D_{f_1} \cup \dots \cup D_{f_n}$ so $\sum r_i f_i^n = 1$)

3.2 Properties that are affine-local

Above we had a property $\star$ of affine opens ("ring is Noetherian") satisfying Affine-local conditions

1) $\text{Spec } R \hookrightarrow X \star \Rightarrow \text{Spec } R_f \hookrightarrow X \star \forall f \in R$ $\swarrow$ so property is preserved by localisation

2) $\text{Spec } R = \bigcup D_{f_i}, \text{Spec } R_{f_i} \hookrightarrow X \star \Rightarrow \text{Spec } R \hookrightarrow X \star$ $\swarrow$ can globalise from basic affines to affine

3.6 Properties of morphisms ← all properties we list are preserved when compose such morphs

A morph of schemes $f: X \rightarrow Y$ is: (will suppress f^*, θ_X, θ_Y from notation)

- ① affine: equivalent conditions:
 - f^{-1} (affine open) is affine
 - $\exists$ affine open cover V_i of Y , $f^{-1}(V_i)$ affine
 - $\forall$ affine open cover V_i of Y , $f^{-1}(V_i)$ affine

- ② quasi-compact: replace affine by quasi-compact
- ③ locally of finite type:
 - $\forall$ affine opens $U \subseteq X, V \subseteq Y$ with $f(U) \subseteq V$, $f^*: \theta_Y(V) \rightarrow \theta_X(U)$ finite type
 - (Rings: $A \rightarrow B$ finite type means B f.g. as A -alg., i.e. $\exists$ surj $A[x_1, \dots, x_n] \rightarrow B$ of A -algs.)
 - (meaning: $\theta_Y(V) \xrightarrow{f^*} \theta_X(f^{-1}(V)) \xrightarrow{\text{surj}} \theta_X(U)$)
 - $\exists$ open affine covers $Y = \cup V_i$, $f^{-1}(V_i) = \cup U_{ij}$ finite type
 - $f^*: \theta_Y(V_i) \rightarrow \theta_X(U_{ij})$ finite type

- ④ finite type: ② + ③: quasi-compact & locally finite type
- ⑤ closed immersion: iso onto a closed subscheme.

Explicitly: $f: X \xrightarrow{\text{homeo}} f(X) \subseteq Y$ (see 1.14.1.15)
 $f^*: \theta_Y \rightarrow f_* \theta_X$ surjective (so ideal sheaf $J = \ker f^*$)
 • $\forall$ aff. open $U = \text{Spec } R \subseteq Y \exists$ ideal $I \subseteq R$ s.t. $f^{-1}(U) \cong \text{Spec}(R/I)$
 $f|_U: \text{aff. cover } Y = \cup \text{Spec } R_i$, ideals $I_i \subseteq R_i$, $f^{-1}(\text{Spec } R_i) = \text{Spec}(R_i/I_i)$
 • $\exists$ aff. cover $Y = \cup \text{Spec } R_i$, ideals $I_i \subseteq R_i$, $f^{-1}(\text{Spec } R_i) = \text{Spec}(R_i/I_i)$
 Example $X = Y_{\text{red}} \subseteq Y$ closed subscheme: $X = Y$ as topological space and (reduction of Y : it's reduced) sheaf of ideals $J(U) = \{s \in \theta_Y(U) : s(p) = 0 \in \kappa(p) \forall p \in U\}$ (so $\theta_X = \theta_Y/J$)
 Note locally: on $U = \text{Spec } R$, $J(U) = \{s \in R : s \in \mathfrak{p} \forall \mathfrak{p} \in \text{Nil}(R)\} = \text{Nil}(R)$, so locally J agrees with $\text{Nil}(\theta_Y)$, indeed J is the sheafification of $\text{Nil}(\theta_Y)$ ← need not be sheaf e.g. $Y = \mathbb{A}^1$, $X_0 = \text{Spec}(\mathbb{Z}[t])$, $2 \in \theta_Y(Y)$, $2 \notin \text{Nil}(\theta_Y(Y))$ but $2 \in \text{Nil}(\theta_Y(Y_n))$, $2 \in J(Y)$

- ⑥ open immersion: iso onto an open subscheme
 Explicitly: $f: X \xrightarrow{\text{homeo}} f(X) \subseteq Y$ (idea: functions on X are the same as "n" Y locally)
 $f^*: \theta_Y \rightarrow f_* \theta_X$ iso ($\Leftrightarrow$ iso on stalks $f_x^*: \theta_{Y,x} \rightarrow \theta_{X,x}$)

- ⑦ flat: all $\theta_{Y,x} \rightarrow \theta_{X,x}$ are flat ring homs

Not intuitively clear, but ensures that fibers of f vary in a controlled way:
 Many invariants of fibers like dimension, do not change unless you "expect" it!
 It is weaker than saying the fibers are locally iso e.g. it allows two points to collide as vary fiber.

Algebra: R -mod M is flat if $M \otimes_R \cdot$ is exact functor on R -mods

$\varphi: R \rightarrow S$ flat ring hom means S flat R -mod (using $r \cdot s = \varphi(r) \cdot s$)

Basic facts

- 1) $M \otimes_R \cdot$ always right exact, so M flat R -mod $\Leftrightarrow N_1 \hookrightarrow N_2$ implies $M \otimes_R N_1 \hookrightarrow M \otimes_R N_2$
- 2) M free $\Rightarrow M$ flat (pf. $M \cong \bigoplus_{i \in I} R \Rightarrow M \otimes_R N \cong \bigoplus_{i \in I} N$)

Example: $\prod_{\text{finite}} \mathbb{Z}$ is not free $\mathbb{Z}$ -mod, but it is flat. An abelian gp is flat $\mathbb{Z}$ -mod $\Leftrightarrow$ torsion-free

Non-example: $\mathbb{Z}_n$ is not flat $\mathbb{Z}$ -mod: $\mathbb{Z} \xrightarrow{\cdot n} \mathbb{Z}$ then $\cdot \theta_{\mathbb{Z}_n}$ get $\mathbb{Z}_n \xrightarrow{\cdot n} \mathbb{Z}_n$ not inj.
 Fact (Lazard) R -mod M is flat $\Leftrightarrow M = \varinjlim M_i$ some f.g. free R -mods M_i

- 3) R local, M finite R -mod (so $M = \sum_{\text{finite}} R m_i$): M flat $\Leftrightarrow M$ free

- 4) $A \rightarrow B$ flat, $B \rightarrow C$ flat $\Rightarrow A \rightarrow C$ flat

pf $N_1 \hookrightarrow N_2$ A -mods $\Rightarrow B \otimes_A N_1 \hookrightarrow B \otimes_A N_2$ B -mods $\Rightarrow C \otimes_B B \otimes_A N_1 \hookrightarrow C \otimes_B B \otimes_A N_2$ C -mods

- 5) $A \rightarrow B$ flat $\Rightarrow A_p \rightarrow B_p = B \otimes_A A_p$ flat $\forall p \in \text{Spec } A$

pf $N_1 \hookrightarrow N_2$ A_p -mods $\Rightarrow N_1 \hookrightarrow N_2$ A -mods (via $A \rightarrow A_p$) $\Rightarrow B \otimes_A N_1 \hookrightarrow B \otimes_A N_2$ B_p -mods

- 6) Ring hom $\varphi: A \rightarrow B$, multiplicative sets $S \subseteq A$, $T \subseteq B$ with $\varphi(S) \subseteq T$, then $\psi: S^{-1}B \rightarrow S^{-1}A \otimes_A B \rightarrow T^{-1}B$, $\varphi \otimes b \mapsto \varphi(a)b \mapsto \varphi(a)b$

Since isos of rings and localization are exact functors, get ψ flat.

Example: $P \subseteq B$ prime ideal, $q = \varphi^{-1}(P) \subseteq A$ prime ideal, $S = A \setminus q$, $T = B \setminus P \Rightarrow B_q = B \otimes_A A_q \rightarrow B_p$ flat

Theorem $\varphi: A \rightarrow B$ flat ring hom $\Leftrightarrow \varphi^*: \text{Spec } B \rightarrow \text{Spec } A$ flat

- pf $\Rightarrow$ $A \rightarrow B$ flat $\Rightarrow A_q \rightarrow B_q$ flat for $q = \varphi^{-1}(p)$ by (5), $B_q \rightarrow B_p$ flat by (6) $\Rightarrow A_q \rightarrow B_p$ flat.

- $\Leftarrow$ Recall $\ker(B \otimes_A N_1 \rightarrow B \otimes_A N_2) \neq 0 \Leftrightarrow \ker \psi \neq 0 \forall p \in \text{Spec } B$.

$\ker(N_1 \rightarrow N_2) = 0 \Rightarrow \ker(A_q \otimes_A N_1 \rightarrow A_q \otimes_A N_2) = 0 \Rightarrow \ker(B_p \otimes_{A_q} A_q \otimes_A N_1 \rightarrow B_p \otimes_{A_q} A_q \otimes_A N_2) = 0$

Fact: $B = \text{Spec } k[t]$
 $B^* = B \setminus 0 = \text{Spec } k[t, t^{-1}]$
 $X \subseteq A^*_g$ closed subscheme

Fact: $B = \text{Spec } k[t]$
 $B^* = B \setminus 0 = \text{Spec } k[t, t^{-1}]$
 $X \subseteq A^*_g$ closed subscheme

Fact: $B = \text{Spec } k[t]$
 $B^* = B \setminus 0 = \text{Spec } k[t, t^{-1}]$
 $X \subseteq A^*_g$ closed subscheme

Fact: $B = \text{Spec } k[t]$
 $B^* = B \setminus 0 = \text{Spec } k[t, t^{-1}]$
 $X \subseteq A^*_g$ closed subscheme

Fact: $B = \text{Spec } k[t]$
 $B^* = B \setminus 0 = \text{Spec } k[t, t^{-1}]$
 $X \subseteq A^*_g$ closed subscheme

Fact: $B = \text{Spec } k[t]$
 $B^* = B \setminus 0 = \text{Spec } k[t, t^{-1}]$
 $X \subseteq A^*_g$ closed subscheme

Fact: $B = \text{Spec } k[t]$
 $B^* = B \setminus 0 = \text{Spec } k[t, t^{-1}]$
 $X \subseteq A^*_g$ closed subscheme

Fact: $B = \text{Spec } k[t]$
 $B^* = B \setminus 0 = \text{Spec } k[t, t^{-1}]$
 $X \subseteq A^*_g$ closed subscheme

Fact: $B = \text{Spec } k[t]$
 $B^* = B \setminus 0 = \text{Spec } k[t, t^{-1}]$
 $X \subseteq A^*_g$ closed subscheme

Fact: $B = \text{Spec } k[t]$
 $B^* = B \setminus 0 = \text{Spec } k[t, t^{-1}]$
 $X \subseteq A^*_g$ closed subscheme

Fact: $B = \text{Spec } k[t]$
 $B^* = B \setminus 0 = \text{Spec } k[t, t^{-1}]$
 $X \subseteq A^*_g$ closed subscheme

Fact: $B = \text{Spec } k[t]$
 $B^* = B \setminus 0 = \text{Spec } k[t, t^{-1}]$
 $X \subseteq A^*_g$ closed subscheme

Fact: $B = \text{Spec } k[t]$
 $B^* = B \setminus 0 = \text{Spec } k[t, t^{-1}]$
 $X \subseteq A^*_g$ closed subscheme

Fact: $B = \text{Spec } k[t]$
 $B^* = B \setminus 0 = \text{Spec } k[t, t^{-1}]$
 $X \subseteq A^*_g$ closed subscheme

Fact: $B = \text{Spec } k[t]$
 $B^* = B \setminus 0 = \text{Spec } k[t, t^{-1}]$
 $X \subseteq A^*_g$ closed subscheme

Fact: $B = \text{Spec } k[t]$
 $B^* = B \setminus 0 = \text{Spec } k[t, t^{-1}]$
 $X \subseteq A^*_g$ closed subscheme

Fact: $B = \text{Spec } k[t]$
 $B^* = B \setminus 0 = \text{Spec } k[t, t^{-1}]$
 $X \subseteq A^*_g$ closed subscheme

Fact: $B = \text{Spec } k[t]$
 $B^* = B \setminus 0 = \text{Spec } k[t, t^{-1}]$
 $X \subseteq A^*_g$ closed subscheme

Fact: $B = \text{Spec } k[t]$
 $B^* = B \setminus 0 = \text{Spec } k[t, t^{-1}]$
 $X \subseteq A^*_g$ closed subscheme

Fact: $B = \text{Spec } k[t]$
 $B^* = B \setminus 0 = \text{Spec } k[t, t^{-1}]$
 $X \subseteq A^*_g$ closed subscheme

Fact: $B = \text{Spec } k[t]$
 $B^* = B \setminus 0 = \text{Spec } k[t, t^{-1}]$
 $X \subseteq A^*_g$ closed subscheme

Fact: $B = \text{Spec } k[t]$
 $B^* = B \setminus 0 = \text{Spec } k[t, t^{-1}]$
 $X \subseteq A^*_g$ closed subscheme

Fact: $B = \text{Spec } k[t]$
 $B^* = B \setminus 0 = \text{Spec } k[t, t^{-1}]$
 $X \subseteq A^*_g$ closed subscheme

Fact: $B = \text{Spec } k[t]$
 $B^* = B \setminus 0 = \text{Spec } k[t, t^{-1}]$
 $X \subseteq A^*_g$ closed subscheme

Fact: $B = \text{Spec } k[t]$
 $B^* = B \setminus 0 = \text{Spec } k[t, t^{-1}]$
 $X \subseteq A^*_g$ closed subscheme

Fact: $B = \text{Spec } k[t]$
 $B^* = B \setminus 0 = \text{Spec } k[t, t^{-1}]$
 $X \subseteq A^*_g$ closed subscheme

Fact: $B = \text{Spec } k[t]$
 $B^* = B \setminus 0 = \text{Spec } k[t, t^{-1}]$
 $X \subseteq A^*_g$ closed subscheme

Fact: $B = \text{Spec } k[t]$
 $B^* = B \setminus 0 = \text{Spec } k[t, t^{-1}]$
 $X \subseteq A^*_g$ closed subscheme

Fact: $B = \text{Spec } k[t]$
 $B^* = B \setminus 0 = \text{Spec } k[t, t^{-1}]$
 $X \subseteq A^*_g$ closed subscheme

Fact: $B = \text{Spec } k[t]$
 $B^* = B \setminus 0 = \text{Spec } k[t, t^{-1}]$
 $X \subseteq A^*_g$ closed subscheme

Fact: $B = \text{Spec } k[t]$
 $B^* = B \setminus 0 = \text{Spec } k[t, t^{-1}]$
 $X \subseteq A^*_g$ closed subscheme

Remarks about calculating closures of sets in $X = \text{Spec } R$

1) $p \in \text{Spec } R \Rightarrow \overline{\{p\}} = V(p)$
 $\underline{\text{pf}}$ $p \in V(p) \Rightarrow \overline{\{p\}} \subseteq V(p)$ (since $V(p)$ closed)
 converse: $p \in \overline{\{p\}} \subseteq V(I) \Rightarrow I \subseteq p \Rightarrow p \in V(I) \square$
 $q \in V(p) \Rightarrow p \subseteq q$
Example $X^* = V(p_1, p_2, \dots, p_k) \subseteq \mathbb{A}_k^n \xleftarrow{\text{say}} B^* = \text{Spec } R[t, t^{-1}]$
 $= V(p_1, p_2, \dots, p_k) \subseteq \mathbb{A}_k^n$ where $V_k(\cdot)$ is $V(\cdot)$ calculated in $\mathbb{A}_k^n$
 $\Rightarrow \overline{X^*} = V(p_1, p_2, \dots, p_k) \subseteq \mathbb{A}_k^n \xleftarrow{\text{say}} B^* = \text{Spec } R[t, t^{-1}]$
 $= V(p_1, p_2, \dots, p_k) \subseteq \mathbb{A}_k^n$ and $p_i \in X^* \subseteq X^*$
 $= V(p_1, p_2, \dots, p_k)$

2) For $\varphi: R \rightarrow S$ ring hom, $\alpha: \text{Spec } S \rightarrow \text{Spec } R$, $\alpha(p) = \varphi^{-1}p$:

Given $C = V(J) \subseteq \text{Spec } S$, $\overline{\alpha(C)} = V(\varphi^{-1}J)$

$\underline{\text{pf}}$ $J = \sqrt{J} \Rightarrow \varphi^{-1}J = \bigcap_{p \in J} \varphi^{-1}p$ since $\alpha(C) \subseteq \overline{\alpha(C)} = V(I)$, $I \subseteq \varphi^{-1}p$
 $\varphi^{-1}p \in \text{Spec } S$ so $\varphi^{-1}p \in V(\varphi^{-1}J)$
 $\alpha(p) = \varphi^{-1}p \in V(\varphi^{-1}J)$
 $\alpha(C) \subseteq V(\varphi^{-1}J)$ so $\overline{\alpha(C)} \subseteq V(\varphi^{-1}J)$
 $\varphi^{-1}J \subseteq \overline{\alpha(C)}$ so $\overline{\alpha(C)} = V(\varphi^{-1}J)$

Example $S = R_f$ localisation, $f \in R$, if $\varphi: R \hookrightarrow R_f$ injection then $\varphi^{-1}J = R \cap J$

e.g. $X^* = V(J) \subseteq \mathbb{A}_k^n$ for $B = \text{Spec } R[t]$, $B^* = \text{Spec } R[t, t^{-1}]$
 so $\mathbb{A}_k^n = \text{Spec } R[x_1, \dots, x_n, t]$, $\mathbb{A}_k^n = R[x_1, \dots, x_n, t, t^{-1}]$

$\Rightarrow \overline{X^*} = V(R[x_1, \dots, x_n, t] \cap J) \subseteq \mathbb{A}_k^n$ is the closure

Rmk Also know inverse images of closed sets: $\alpha^{-1}(V(I)) = V(\varphi^{-1}I)$

$\underline{\text{pf}}$ $I = \langle f_i \rangle$, $\text{Spec } R \setminus V(I) = \cup D_{f_i}$,

$\cup D_{\varphi f_i} = \alpha^{-1}(\cup D_{f_i}) = \alpha^{-1}(\text{Spec } R \setminus V(I)) = \text{Spec } S \setminus \alpha^{-1}V(I)$
 $\Rightarrow \alpha^{-1}V(I) = \text{Spec } S \setminus \cup D_{\varphi f_i} = V(\langle \varphi f_i \rangle) \square$
 (recall $\alpha^{-1}D_f = D_{\varphi f}$)


4- GLUING THEOREMS

4.1 Gluing sheaves

$X = \cup U_i$ open cover, abbreviate $U_{ij} = U_i \cap U_j$, $U_{ijk} = U_i \cap U_j \cap U_k$

F_i sheaf on U_i

$\varphi_{ij}: F_i|_{U_{ij}} \xrightarrow{\sim} F_j|_{U_{ij}}$
 Compatibility conditions
 1) $\varphi_{ii} = \text{id}$
 2) $\varphi_{ji} = \varphi_{ij}^{-1}$
 3) $\varphi_{ik}|_{U_{ijk}} = \varphi_{jk} \circ \varphi_{ij}|_{U_{ijk}}$



Example F sheaf on X , $F_i := F|_{U_i}$ (so $F_i(V) = F(U_i \cap V) = F(U_i \cap V)$, $\forall \text{ open } V \subseteq U_i$)

$\varphi_{ij} = \text{isos}$ induced by double restrictions (iso of functors $\cdot|_{U_{ij}} \cdot|_{U_{ij}}$)

Theorem $\exists$, up to unique iso, a sheaf F on X with isos

$$\psi_i: F|_{U_i} \xrightarrow{\sim} F_i$$

s.t. $\psi_j^{-1} \circ \varphi_{ij} \circ \psi_i|_{U_{ij}} = F|_{U_{ij}} \cong F|_{U_{ij}}|_{U_{ij}} \cong F|_{U_{ij}}$

$$\begin{array}{c} F|_{U_i}|_{U_{ij}} \xrightarrow{\psi_i} F_i|_{U_{ij}} \\ \cong \downarrow \quad \downarrow \varphi_{ij} \\ F|_{U_j}|_{U_{ij}} \xrightarrow{\psi_j} F_j|_{U_{ij}} \end{array}$$

pf Let $E = \bigsqcup_i (F_i)_x / \sim$ equivalence relation $(F_i)_x \sim (F_j)_x$ for $x \in U_{ij}$

(using conditions)

$F(U) = \{s: U \rightarrow E : s \text{ is locally a section of some } F_i\}$
 ($\forall x \in U, \exists i, \exists \text{ open } V \ni x \subseteq U_i, \exists t \in F_i(V), s|_V = t|_V$)

Theorem Given sheaves F, G constructed as above from local data F_i, φ_{ij} on U_i

a morph $f: F \rightarrow G$ can be uniquely defined from data:

- morphs $f_i: F_i \rightarrow G_i$
- compatibility condition: $\psi_j \circ f_i|_{U_{ij}} = f_j \circ \varphi_{ij}|_{U_{ij}}$

$$\begin{array}{c} \varphi_{ij} \downarrow \quad \downarrow f_j \\ F_i|_{U_{ij}} \xrightarrow{\psi_i} F_i|_{U_{ij}} \\ \downarrow f_i \quad \downarrow \psi_j \\ G_i|_{U_{ij}} \xrightarrow{\psi_j} G_j|_{U_{ij}} \end{array}$$

s.t. via identifications $F|_{U_i} \cong F_i$, $G|_{U_i} \cong G_i$ recover $f|_{U_i} = f_i$

4.2 Gluing schemes


U_i schemes, $U_{ij} \subseteq U_i$ open subschemes ($U_{ii} = U_i$)

$\varphi_{ij}: U_{ij} \xrightarrow{\sim} U_{ji}$ isos $\leftarrow$ (think "go from U_i to U_j ")


gluing conditions 1) $\varphi_{ii} = \text{id}$

2) $\varphi_{ij}(U_{ij} \cap U_{ik}) \subseteq U_{jk} \cap U_{ik}$

3) $\varphi_{ik} = \varphi_{jk} \circ \varphi_{ij}$ when restricted as maps $U_{ij} \cap U_{ik} \rightarrow U_{ik}$




Example if $U_i \subseteq X$ open subschemes, can take $U_{ij} = U_i \cap U_j \subseteq X$ with $\varphi_{ij} = id$
Claim (exercise) $\exists$ unique (up to iso) scheme X with open cover $X = \cup X_i$



Gluing Lemma

$\Rightarrow f: X \rightarrow Y$ morph can be uniquely defined from morphs $f_i: X_i \rightarrow Y$ st.



Pf Continuous map: $f: X \rightarrow Y$ defined by $f|_{X_i} = f_i$ (Compatibility)

on sheaves need $f^{-1}\theta_Y \rightarrow \theta_X \leftarrow$ (recall get $\theta_Y \rightarrow f_*\theta_X$ by adjunction)
 $(f^{-1}\theta_Y)|_{X_i} = f_i^{-1}\theta_Y = f_i^{-1}\theta_{X_i} \leftarrow (X_i \xrightarrow{\varphi_i} X \text{ inclusion, then } \varphi_i^* f^{-1}\theta_Y = (f \circ \varphi_i)^* \theta_Y$
 $f_i^* \in \text{Mor}(\theta_Y, f_i^* \theta_{X_i}) \cong \text{Mor}(f_i^* \theta_Y, \theta_{X_i})$ and $\theta_{X_i} = \theta_X|_{X_i}$ since open subs.
 Finally we can glue the $f_i^*: f_i^* \theta_Y \rightarrow \theta_{X_i}$ by $\oplus$ to get $f^{-1}\theta_Y \rightarrow \theta_X$. $\square$

Consequence $h_Y|_{\text{Top}(X)^{\text{op}}} : \text{Top}(X)^{\text{op}} \rightarrow \text{Sets}$ is a sheaf of sets.

$(X, Y \text{ schemes}) \quad U \mapsto h_Y(U) = \text{Mor}(U, Y)$

4.3 Affine space by gluing (see Homework for projective space)

Affine n-space over $\text{Spec } R: \mathbb{A}_R^n := \text{Spec } R[x_1, \dots, x_n] (= \mathbb{A}_R^n \text{ Spec } R) \xrightarrow{\text{Spec}} \text{Spec } R$

$\text{Rmk } R \rightarrow S \text{ ring hom} \Rightarrow \text{hom on poly} (so: R[x_1, \dots, x_n] \rightarrow S[x_1, \dots, x_n]) \Rightarrow \mathbb{A}_S^n \rightarrow \mathbb{A}_R^n$

Example $R \rightarrow R_f \Rightarrow \mathbb{A}_R^n \rightarrow \mathbb{A}_R^n$ is the basic open set of $\mathbb{A}_R^n$ for $f \in R$

If $U \subseteq \text{Spec } R$ open $\Rightarrow U = \cup D_{f_i} \Rightarrow \mathbb{A}_U^n = \cup \mathbb{A}_{R_{f_i}}^n \subseteq \mathbb{A}_R^n \xleftarrow{\text{glued along}} \mathbb{A}_R^n$

X scheme, affine n-space over $X: \mathbb{A}_X^n := \cup \mathbb{A}_{X_i}^n$ where $X = \cup X_i$ affine open cover

$(\mathbb{A}_X^n = \cup \mathbb{A}_{X_i}^n) \xrightarrow{\downarrow} \mathbb{A}_X^n$ (notice $\mathbb{A}_{X_i}^n = \mathbb{A}_{X_i \cap X_j}^n$, then identify these copies) $\xrightarrow{\text{glued along}} \mathbb{A}_X^n$ open in affine X_i

Claim $\mathbb{A}^n = \text{Spec } \mathbb{Z}[x_1, \dots, x_n]$ represents functor $F: \text{Set}^{\text{op}} \rightarrow \text{Sets}$ $X \mapsto \{\text{morphs } \mathbb{A}^n \rightarrow X \text{ st. } \forall U_i \subseteq X, \theta_{X_i}(U_i) \cap \theta_X(U_i) \text{ is hom of } \theta_X(U_i)\}$

Pf $F|_{\text{Top}(X)^{\text{op}}}$ is a sheaf of sets (easy to check: can glue morphs since θ_X sheaf)

$h_{\mathbb{A}^n|_{\text{Top}(X)^{\text{op}}}}$ by consequence above. Thus if the two functors agree on affines then

by sheaf property they agree everywhere. For affine $X = \text{Spec } R$ just need compare global sections

$F(\text{Spec } R) = \text{Hom}_R(R^n, R) \leftarrow$ (here: R -mod homs!) in both cases just need specify $\{e_i = (0, \dots, 1, \dots, 0) \mapsto r_i\}$


$h_{\mathbb{A}^n}(\text{Spec } R) = \text{Mor}(\text{Spec } R, \mathbb{A}^n) \cong \text{Hom}(\mathbb{Z}[x_1, \dots, x_n], R)$ where generators go $\{x_i \mapsto r_i\}$ $\square$

5. PRODUCTS


5.0 Products in category theory

Category theory: $\mathcal{C}$ Cat., $C_i \in \mathcal{C}$

product $C_1 \times \dots \times C_n$ (if exists) is an object with morphs π_i to C_i st.



coproduct $C_1 \sqcup \dots \sqcup C_n$:



Examples Sets / Top spaces: $X = \text{product}$, $\pi_i = \text{projections}$, $\sqcup = \text{disjoint union}$, π_i are inclusions.
 Vector spaces / algebras/modules: $\sqcup = \text{direct sum}$, π_i are inclusions.
 Rings: $\sqcup = \text{tensor product}$, $\pi_i(r) = 1 \otimes \dots \otimes r \otimes \dots \otimes 1$

Fix $B \in \mathcal{C}$ ("base")


Category of B -objects: $\mathcal{C}/B$

obj: morphs $C \rightarrow B$, morphs: $C \rightarrow D$ in $\mathcal{C}$

(think of B as a parameter space and $\mathcal{C}$ as a family parametrized by B)

fiber product $C \times_B D$ is the product in $\mathcal{C}/B$ of $C \xrightarrow{f} B$, $D \xrightarrow{g} B$ (if exists)

(or pullback, or Cartesian square)



Example for Sets or Top spaces: $C \times_B D = \{(c, d) \in C \times D : f(c) = g(d)\} \in C \times D = C \times D$

for example if f, g are inclusions of subsets (subspaces) then $C \times_B D = C \cap D$

Pushout The opposite diagram (reverse arrows)

Example: for Rings the pushout of $B \rightarrow C, B \rightarrow D$ is the tensor product $C \otimes_B D$ sec 4.2

Exercise: $B \xrightarrow{f} C, B \xrightarrow{g} D$ inclusions of open subschemes, then pushout $C \sqcup_B D$ is the gluing!

Exercise: $(co)product$, fiber product, pushout are unique up to unique iso if they exist.

(Hint: compose unique maps between them (s.t. diagram commutes) then composites = id by uniqueness of self-map)

Examples of fiber products in cat. of Sets or Top spaces: $C \times_B D = \{(c, d) : f(c) = g(d)\} \subseteq C \times D$

$B = \text{point} \Rightarrow C \times_B D = C \times D$

$C \xrightarrow{f} B, D \xrightarrow{g} B \Rightarrow C \times_B D \cong C \cap D$

$D \xrightarrow{g} B \Rightarrow C \times_B D \cong f^{-1}(D) \subseteq C$ for example $D = \text{point} = b \in B$ get fiber $f^{-1}(b)$

$C = D \Rightarrow C \times_B D = \{(x, y) : f(x) = g(y)\} \subseteq C \times D$ ("equalizer")

IMPORTANT EXAMPLES:

All schemes X have canonical $X \rightarrow \text{Spec } \mathbb{Z}$

by giving canonical maps on affines:

$\text{Spec } R \rightarrow \text{Spec } \mathbb{Z}$ from $\mathbb{Z} \rightarrow R, 1 \mapsto 1$

Schemes over field k means have

$X \rightarrow \text{Spec } k$, same as saying all $\theta_X(U)$

are k -algebras and restrictions are k -algebra homs

Functor of points interpretation:

$\text{Hom}(\mathbb{A}^n_C, C \times_B D) \cong \text{Hom}(\mathbb{A}^n_C \times \text{Hom}(\mathbb{A}^n_D), \text{Hom}(\mathbb{A}^n_B))$

So we are asking whether $h_C \times_{h_B} h_D$ is representable

5.1 Fiber products exist in Schemes/B

Fix scheme B, consider category Schemes/B

Theorem fiber products $X_1 \times_B \dots \times_B X_n$ exist

Inductively suffices to do case $n=2$. First need some algebraic preliminaries

An A-algebra R is a ring R together with a ring hom $A \rightarrow R$
(A ring) $(\Rightarrow R \text{ is } A\text{-mod via } a \cdot r = \psi(a)r)$

R, S A-algebras $\Rightarrow (R \otimes_A S) = \text{free } R\text{-alg. on } R \times S$
relations

relations: i) $\otimes$ is bilinear

ii) $a \cdot (r \otimes s) = (\psi(a)r) \otimes s = r \otimes (\psi(a)s)$

In particular $A \rightarrow R \otimes_A S$ is $a \mapsto a \cdot (1 \otimes 1) = \psi(a) \otimes 1 = 1 \otimes \psi(a)$

The product on generators: $(r \otimes s) \cdot (r' \otimes s') = rr' \otimes ss'$

Rmk R, S rings $\Rightarrow R \otimes S = R \otimes_{\mathbb{Z}} S$

Facts 1) $R \otimes_R S \cong S$ (via $\sum r_i \otimes s_i \mapsto \sum r_i s_i$)

2) $R[x_1, \dots, x_n] \otimes_R R[y_1, \dots, y_m] \cong R[x_1, \dots, x_n, y_1, \dots, y_m]$

3) $(S/I) \otimes_R T \cong (S \otimes_R T) / (I \otimes 1) \cdot (S \otimes_R T)$ where S, T are R-algebras

4) k field, A k-alg, for A-algs R, S get: $R \otimes_A S \cong (R \otimes_k S) / \langle \psi_R(a) \otimes 1 - 1 \otimes \psi_S(a) : a \in A \rangle$

Affine case: $\text{Spec } R \times_{\text{Spec } A} \text{Spec } S = \text{Spec } (R \otimes_A S)$ exists in Aff/Spec A:

have pushout: $R \otimes_A S \leftarrow S$
(in category of A-algs) $\uparrow \psi_S$
 $R \xleftarrow{\psi_R} A$

Now apply Spec. $\square$
(property of pushout for A-algs gives you the univ. prop. for fiber prod. for Spec A)

Claim: this is fiber product also in Sch/Spec A: let $X = \text{Spec } R$
 $Y = \text{Spec } S$
 $B = \text{Spec } A$
 $F = \text{Spec } (R \otimes_A S)$

affine cover: U_i of scheme Z

ind

scheme Z

V_{ψ_1}

V_{ψ_2}

F

X

Y

B

f

g

Rmk $B = \text{Spec } \mathbb{Z}$ gives $X \times_B Y = X \times Y$

Recall fiber products are unique up to unique iso if they exist.

By construction (as U_i affine) $\exists! U_i \rightarrow F$ making diagram commute

It can show these affine on overlaps $U_{ij} = U_i \cap U_j$, then glue to unique $Z \rightarrow F$.

If U_{ij} were affine, this would have been immediate.

$U_{ij} \subseteq \text{affine } U_i$, so running same argument with Z replaced by U_{ij} ,

we can cover U_{ij} by basic open affines $D_{f_k} \subseteq U_i$ and now $D_{f_k} \cap D_{f_l} = D_{f_k f_l}$ affine

$\Rightarrow$ glue uniquely to give $U_{ij} \rightarrow F$

Recall trick that can pick open cover of U_{ij} that are basic opens simultaneously for U_i, U_j

$\Rightarrow U_{ij} \rightarrow F$ and $U_{ij} \rightarrow F$ agree.

General case build schemes/morphs by 3 gluing procedures (tedious!)

1) case $U_i \times_B Y$ with B, Y affine, $X = U_i$: affine open cover $\Rightarrow \exists X \times_B Y$ affine

2) case $X \times_B V_j$ with B affine, $Y = U_j$: " " $\Rightarrow \exists X \times_B Y$ affine

3) case $X \times_{U_k} Y$ with $B = U_k$: " " $\Rightarrow \exists X \times_B Y$

Gluing work because agreement on overlaps is ensured by uniqueness up to iso of fiber products. Sketch:

① if know $U_i \times_B Y$ exist, then $\pi_i^{-1}(U_{ij})$ is fiber product

$U_{ij} \times_B Y$ so by uniqueness $\exists$ iso $\pi_i^{-1}(U_{ij}) \rightarrow \pi_j^{-1}(U_{ij})$, so glue & get $X \times_B Y$

(indeed a natural identification since $U_{ij} = U_i$ with sheaf $\mathcal{O}_{U_{ij}}$)

② as in ①, swapping roles X, Y .

③ let $X_k = f^{-1}(W_k)$, $Y_k = g^{-1}(W_k) \Rightarrow X_k \times_{U_k} Y_k$ exists by ② (we affine)

Key trick: notice $X_k \times_{U_k} Y_k = X_k \times_B Y$

"because images are trapped in W_k, Y_k anyway"

Then use argument in ① to glue the $X_k \times_B Y$. $\square$

Rmk Proof shows that $X \times_B Y$ has affine open cover by $U(U_i \times_{U_k} V_j)$

where $X = U_i, Y = U_j$ are " " $B = U_k$ with $U_i \rightarrow W_k \subseteq B, V_j \rightarrow W_k \subseteq B$

Examples 1) $A_R^n \times_{\text{Spec } R} A_R^m = \text{Spec } R[x_1, \dots, x_n, y_1, \dots, y_m] = A_R^{n+m}$

2) $\text{Spec } \mathbb{Z}_2 \times_{\text{Spec } \mathbb{Z}} \text{Spec } \mathbb{Z}_3 = \text{Spec } (\mathbb{Z}_2 \otimes_{\mathbb{Z}} \mathbb{Z}_3) = \text{Spec } (0) = \emptyset$

Exercise $X \times_B Y \cong X, X \times_B Y \cong Y \times_B X, (X \times_B Y) \times_B Z \cong X \times_B (Y \times_B Z), X \times_B B \times_B Y \cong X \times_B Y$

5.2 Fibers and preimages

$f: X \rightarrow B$ morph of schemes

fiber over point $b \in B: f^{-1}(b) = \text{Spec } k(b) \times_B X$

preimage of closed subscheme $Y \subseteq B: f^{-1}(Y) = Y \times_B X$

Diagram showing the relationship between schemes X, Y, B and their fiber products and preimages.

Diagram showing the relationship between schemes X, Y, B and their fiber products and preimages.

Diagram showing the relationship between schemes X, Y, B and their fiber products and preimages.

Diagram showing the relationship between schemes X, Y, B and their fiber products and preimages.

Diagram showing the relationship between schemes X, Y, B and their fiber products and preimages.

Diagram showing the relationship between schemes X, Y, B and their fiber products and preimages.

Diagram showing the relationship between schemes X, Y, B and their fiber products and preimages.

Diagram showing the relationship between schemes X, Y, B and their fiber products and preimages.

Diagram showing the relationship between schemes X, Y, B and their fiber products and preimages.

Diagram showing the relationship between schemes X, Y, B and their fiber products and preimages.

Diagram showing the relationship between schemes X, Y, B and their fiber products and preimages.

Diagram showing the relationship between schemes X, Y, B and their fiber products and preimages.

Diagram showing the relationship between schemes X, Y, B and their fiber products and preimages.

Diagram showing the relationship between schemes X, Y, B and their fiber products and preimages.

Diagram showing the relationship between schemes X, Y, B and their fiber products and preimages.

Diagram showing the relationship between schemes X, Y, B and their fiber products and preimages.

Diagram showing the relationship between schemes X, Y, B and their fiber products and preimages.

Diagram showing the relationship between schemes X, Y, B and their fiber products and preimages.

Diagram showing the relationship between schemes X, Y, B and their fiber products and preimages.

Diagram showing the relationship between schemes X, Y, B and their fiber products and preimages.

Diagram showing the relationship between schemes X, Y, B and their fiber products and preimages.

Diagram showing the relationship between schemes X, Y, B and their fiber products and preimages.

Diagram showing the relationship between schemes X, Y, B and their fiber products and preimages.

Diagram showing the relationship between schemes X, Y, B and their fiber products and preimages.

Diagram showing the relationship between schemes X, Y, B and their fiber products and preimages.

Diagram showing the relationship between schemes X, Y, B and their fiber products and preimages.

Diagram showing the relationship between schemes X, Y, B and their fiber products and preimages.

Diagram showing the relationship between schemes X, Y, B and their fiber products and preimages.

Diagram showing the relationship between schemes X, Y, B and their fiber products and preimages.

Diagram showing the relationship between schemes X, Y, B and their fiber products and preimages.

Examples

3) k = algebraically closed field $\leftarrow$ (so classical alg. geometry)

$f: A_k^1 \rightarrow A_k^1$ induced by $f^\#: k[x] \rightarrow k[y], x \mapsto y^2$

fiber over 0: (view point 0 as $\text{Spec } k \rightarrow \text{Spec } k \rightarrow A_k^1$ so $k \cong k[x]$)


fiber = $\text{Spec } k[x^2] \subset \text{Spec } k[x] = \text{Spec}(k \otimes_k k[x]) = \text{Spec}(k[y])$

$= \text{Spec}(k[x^2]) \otimes_{(y)} k[y] \cong \text{Spec}(k[y]/(y^2))$ where $f(x) = y^2$

(e.g. use facts about $\otimes$ from 5.1)

Rmk Notice how a product of affine varieties gave a scheme that was not an affine variety.

4) Mumford's picture of $\text{Spec } \mathbb{Z}[x]$:



$X = \text{Spec } \mathbb{Z}[x] = \{(2, x), (3, x), (5, x), (7, x), (0, x)\}$

$B = \text{Spec } \mathbb{Z} = \{(2), (3), (5), (7), (0)\}$

π is induced by inclusion $\mathbb{Z} \rightarrow \mathbb{Z}[x]$

$\Rightarrow \pi^{-1}((p)) = V((p)) = \{(p, f(x)) : f(x) \text{ mod } p \text{ is irreducible in } \mathbb{F}_p[x]\}$

(so (p) is a dense) $\rightarrow$ if $p \in \mathbb{Z}$ then $\mathbb{Z}[x]/I \cong \mathbb{F}_p[x]/I$ where $\mathbb{F}_p = \mathbb{Z}/p$

(point in $\pi^{-1}((p))$).

Rmk curve $V(x^2+1)$ passes through $(p, x+i)$ iff x^2+1 vanishes at that

point, so iff $x^2+1=0$ in $\mathbb{F}_p[x]/(x+i) \cong \mathbb{F}_p, x \mapsto -i$, so iff $j^2 = -1$.

(classical number theory says a square root of -1 exists in $\mathbb{F}_p \Leftrightarrow (p \equiv 1 \text{ mod } 4)$ or $p=2$)

fiber over $(p) : K(p) = \mathbb{Z}(p)/p \cdot \mathbb{Z}(p) = (\mathbb{Z}/p)[p] = \mathbb{F}_p = \mathbb{Z}/p$

$\Rightarrow \pi^{-1}(p) = \text{Spec}(K(p) \otimes_{\mathbb{Z}} \mathbb{Z}[x]) = \text{Spec } \mathbb{F}_p[x] = \{(0), (f(x))\}$ irreducible in $\mathbb{F}_p[x]$

fiber over $(0) : K(0) = \mathbb{Z}(0) = \mathbb{Q}$

$\Rightarrow \pi^{-1}(0) = \text{Spec}(K(0) \otimes_{\mathbb{Z}} \mathbb{Z}[x]) = \text{Spec } \mathbb{Q}[x] = \{(0), (f(x))\}$

[Gauss's Lemma: for $f \in \mathbb{Z}[x]$ primitive (gcd(coeffs)=1) irreducible in $\mathbb{Q}[x]$, so WLOG irreducible in $\mathbb{Z}[x]$, f irreducible in $\mathbb{Z}[x] \Leftrightarrow f$ irreducible in $\mathbb{Q}[x]$]

Consequence $\text{Spec } \mathbb{Z}[x] = \{(0), (p), (f), (p, f)\}$ $f \in \mathbb{Z}[x]$ irreducible mod p

$\leftarrow p \in \mathbb{Z}$ prime $f \in \mathbb{Z}[x]$ irreducible, nonconstant

$\leftarrow p \in \mathbb{Z}$ prime $f \in \mathbb{Z}[x]$ irreducible, nonconstant

Forgetful functor $|\cdot|: \text{Sch} \rightarrow \text{TopSpaces}, X \mapsto |X| = \text{underlying topological space}.$

Claim $f: X \rightarrow B$ morph schemes $\Rightarrow |f^{-1}(b)| = |f|^{-1}(b)$ $\leftarrow$ fiber is homeomorphic to topological fiber

Pf WLOG B affine = $\text{Spec } S$ and b is prime ideal $p \subseteq S$

$f^{-1}(B) = \cup \text{Spec } R_i$ given by $\varphi_i: S \rightarrow R_i$

WLOG just consider one affine, so $R = R_i$, so $\text{WLOG } X = \text{Spec } R$

$\Rightarrow \text{Spec } K(b) \times_B X = \text{Spec}(K(b) \otimes_S R)$

$K(b) = (S/p)_p \Rightarrow K(b) \otimes_S R = (S/p)_p \otimes_S R = S_p \otimes_S R = R_{(p)} / (p)_{(p)}$

$\Rightarrow \text{Spec}(K(b) \otimes_S R) \xrightarrow{1:1} \{q \subseteq R \text{ prime ideal containing } q(p) \text{ but not intersecting } q(S \setminus p)\}$

$q \cdot R_p \hookrightarrow q \quad (= \text{preimage of } qR_p \text{ via localization } R \rightarrow R_p = S_p \otimes_S R) \quad q \cap p = p$

$q \subseteq R \setminus q(S \setminus p) \Rightarrow q^{-1}q \subseteq S \setminus (S \setminus p) = p$ so get $\{q \in \text{Spec } R : q^{-1}q = p\}.$

$q \supseteq q(p) \Rightarrow q^{-1}q \supseteq p$ (apply 1.1 to diagram defining $x \times_B y$ then by universal property in category of topological spaces get unique map $\otimes$)

Cor Given $f: X \rightarrow B, g: Y \rightarrow B,$ where $f(x) = g(y) = b$

fiber of $|X \times_B Y| \xrightarrow{\otimes} |X| \times |B| \times |Y|$ over (x, y) is $|\text{Spec}(K(x) \otimes_{K(b)} K(y))|$

Pf fiber of $X \times_B Y \rightarrow X$ over $x: \text{Spec } K(x) \times_X (X \times_B Y) = \text{Spec } K(x) \times_B Y$

fiber of $\text{Spec } K(x) \times_B Y \rightarrow Y$ over $y: \text{Spec } K(x) \times_Y \text{Spec } K(y) = \text{Spec } K(x) \times_{\text{Spec } K(b)} \text{Spec } K(y)$

fiber of $\text{Spec } K(x) \times_{\text{Spec } K(b)} \text{Spec } K(y) \rightarrow \text{Spec } K(b): \text{Spec } K(x) \times_{\text{Spec } K(b)} \text{Spec } K(y) = \text{Spec } K(x) \otimes_{K(b)} K(y).$

$\square$

or at category level, with abuse of notation:

at algebra level: if A_1, A_2 are modules over $S = R/pR_p$ then $S \otimes (A_1 \otimes_R A_2) \cong A_1 \otimes_S A_2$

$R_p \otimes (S/p)_p \otimes \frac{R}{p} \otimes A_1 \otimes A_2 \xrightarrow{f} \frac{R}{p} \otimes A_1 \otimes A_2 \xrightarrow{f} \frac{R}{p} \otimes A_1 \otimes A_2$

namely: $\frac{R}{p} \otimes (S/p)_p \otimes \frac{R}{p} \otimes A_1 \otimes A_2 \xrightarrow{f} \frac{R}{p} \otimes A_1 \otimes A_2$

lands in $\{b\} \subseteq B$

by claim can work with fiber in Sch before applying 1.1

here $\xrightarrow{f} \exists! x \times_B y$ hence $\xrightarrow{f} \exists! x \times_B y$

so $|A_k| \neq |A'_k| \times |A''_k|$: the fiber over $(0, (0))$ is complicated.

note $\text{Spec } k = \text{point} = \{(0)\}$ so often omit "Spec k " from notation.

Examples $|\text{Spec } \mathbb{Z}_k \times_{\text{Spec } \mathbb{Z}} \text{Spec } \mathbb{Z}_k| = |\text{Spec } \mathbb{Z}_k| \times_{|\text{Spec } \mathbb{Z}|} |\text{Spec } \mathbb{Z}_k| = \emptyset$ since 1^{st} factor $\mapsto (2) \in \text{Spec } \mathbb{Z}$ and 2^{nd} factor $\mapsto (3) \in \text{Spec } \mathbb{Z}$

$\bullet A_k^2 = A_k^1 \times_{\text{Spec } k} A_k^1 = \text{Spec } k[x, y] \xrightarrow{f} \text{Spec } k[x, y]$ then $(x, y) \mapsto (0)$ via both projections to A_k^1 but $(x, y) \neq (0)$

(field k) so $|A_k^2| \neq |A_k^1| \times |A_k^1|$: the fiber over $(0, (0))$ is complicated.

Rmk If x, y closed points of schemes X, Y finite type over k, k algebraically closed,

then fiber over (x, y) of $X \times_{\text{Spec } k} Y$ is $\text{Spec}(K(x) \otimes_k K(y)) = \text{Spec}(K \otimes_k K) = \text{Spec } k = (0)$

so over closed points you get the product of sets. $\leftarrow$ (so classical alg. geom.)

Warning $A_k^2 = A_k^1 \times A_k^1$ does not have the product topology, e.g. consider $V(x, y)$

Non-examinable Rmk Working over an algebraically closed field k , the stalk of $X \times_{\text{Spec } k} Y$

at (x, y) is $\mathcal{O}_{X, x} \otimes_k \mathcal{O}_{Y, y}$ localised at max ideal $\mathfrak{m}_{X, x} \otimes \mathfrak{m}_{Y, y} + \mathcal{O}_{X, x} \otimes \mathfrak{m}_{Y, y}$

5.6 Scheme structure on subsets

Motivation: classically, a projective variety is a closed subset of $\mathbb{P}^n$. A quasi-proj. var. is an open $\subseteq$ proj. var., so $\Leftrightarrow$ locally closed subset of $\mathbb{P}^n$.

Claim Any closed subset $C \subseteq X$ of a scheme $\Rightarrow \exists!$ closed reduced subscheme $(C, \mathcal{O}_C) \rightarrow X$
 $\mathcal{P}_C(U) := \{s \in \mathcal{O}_X(U) : s(p) = 0 \in k(p) \ \forall p \in C \cap U\}$ is sheaf of ideals
 Locally: $U = \text{Spec } R$, $C \cap U = V(I)$ for unique radical ideal I

then $s(p) = 0 \in k(p) = (R/p)$ $\forall p \in V(I) \Leftrightarrow s \in \bigcap_{p \in V(I)} p = \sqrt{I} = I$
 $\Rightarrow \mathcal{P}_C(U) = I$ for unique radical ideal I
 Same trick shows $\mathcal{P}_C(D_p) = I_f$, so $\mathcal{P}$ is the quasi-coherent ideal sheaf corresponding to I .
 Note: $C = \text{supp}(\mathcal{O}_X/\mathcal{P})$ and $C \cap U = \text{Spec } R/I$, and we define $\mathcal{O}_C = \mathcal{O}_X/\mathcal{P}$. $\square$

Def call this the induced reduced scheme structure on C .
 Example: When we consider an irreducible component $Z \subseteq X$, we use this scheme structure.
 Exercise: For $C = X \subseteq X$ get the reduced scheme X_{red} (see 3.6)

Def $Z \subseteq X$ locally closed means $\forall z \in Z, \exists$ open $U \ni z$ s.t. $Z \cap U$ is closed in U .
 Lemma: Z locally closed $\Leftrightarrow Z$ open in $\bar{Z}$ (i.e. $Z = \bar{Z} \cap U$ some open $U \subseteq X$) by Lemma, $C = \bar{Z}$ works

Def $Z \subseteq X$ locally closed $\Leftrightarrow Z$ open in $\bar{Z}$ (i.e. $Z = \bar{Z} \cap U$ some open $U \subseteq X$) by Lemma, $C = \bar{Z}$ works
 Pf: $\Leftarrow$: $Z = \bar{Z} \cap U$ for open $U \subseteq X \Rightarrow Z \cap U = Z = \bar{Z} \cap U$
 $\Rightarrow$: $Z \cap U$ closed in U so equals its closure in U which is: $\overline{Z \cap U} = \bar{Z} \cap U$
 $\Rightarrow Z \cap U = \bar{Z} \cap U \subseteq \bar{Z}$ so Z contains an open neighbourhood of z in $\bar{Z}$ $\Rightarrow Z \cap U = \bar{Z} \cap U$

Rmk $\bar{Z} \subseteq X$ closed, so $\exists!$ induced reduced scheme structure $\mathcal{O}_{\bar{Z}}$ on $\bar{Z}$
 $Z \subseteq \bar{Z}$ is open so get " " $\mathcal{O}_Z = \mathcal{O}_{\bar{Z}}|_Z$ (so $\mathcal{O}_Z(V) = \mathcal{O}_{\bar{Z}}(V)$)
 The local description is the same as above: $Z \cap U = \bar{Z} \cap U = \text{Spec}(R/I)$, $\mathcal{O}_Z|_U \cong \mathcal{O}_{\bar{Z}}|_U$

Rmk If Z irreducible ($\Rightarrow \bar{Z}$ irreducible) then $I = p \in \text{Spec } R$ where p is a generic point for both $Z, \bar{Z}$
 Hw 3.3 $\bar{Z}$ irred. locally closed $\subseteq$ variety $(X, \mathcal{O}_X) \Rightarrow (\bar{Z}, \mathcal{O}_{\bar{Z}})$ variety

Hw 3.3 $(X, \mathcal{O}_X)$ variety, $Z \subseteq X$ irreducible subspace $\Leftarrow$ (the irreducibility is not so important if allow varieties to be reducible)
 Define sheaf $\mathcal{O}_Z$ on Z : for open $V \subseteq Z$,
 $\mathcal{O}_Z(V) = \{s: V \rightarrow \coprod_{x \in V} k(x) : s(x) = t(x) \in k(x) \ \forall x \in V \cap U\}$
 such that $s(x) = t(x) \in k(x) \ \forall x \in V \cap U$

Prove that:
 $(Z, \mathcal{O}_Z)$ variety $\Rightarrow Z$ locally closed and $\mathcal{O}_Z$ is the induced reduced scheme structure
 (universal property for the above sheaf)

Lemma: With that definition, if Y reduced scheme, $f: Y \rightarrow X$ morph of sch.
 if $f(Y) \subseteq Z$ (as topological spaces) then f factorizes $f: Y \rightarrow Z \rightarrow X$

Pf Need check sheaves: $s \in \mathcal{O}_Z(U \cap \bar{Z})$ for $U \subseteq X$ open then $\exists$ open cover $U \cap \bar{Z} = \bigcup U_i \cap \bar{Z}$ and $s_i \in \mathcal{O}_X(U_i)$, $s(x) = s_i(x) \in k(x) \ \forall x \in U_i \cap \bar{Z}$
 $\Rightarrow f^\#(s_i) \in \mathcal{O}_Y(f^{-1}(U_i))$, $f^\#(s_i)(y) = f^\#(s_i)(y) \in k(y)$, $\forall y \in f^{-1}(U_i \cap \bar{Z})$ where by 1.6:
 $\Rightarrow$ by Sec. 3.3 since Y reduced: $f^\#(s_i)_y = f^\#(s)_y \in \mathcal{O}_Y(y)$, $\forall y \in f^{-1}(U_i \cap \bar{Z})$
 $\Rightarrow f^\#(s)$ glue to a unique section $r \in \mathcal{O}_Y(f^{-1}(U))$. Define $\mathcal{O}_Z(U) \rightarrow \mathcal{O}_Y(f^{-1}(U))$, $s \mapsto r|_{f^{-1}(U)}$
 and note $\mathcal{O}_X(U_i) \rightarrow \mathcal{O}_Z(U \cap \bar{Z}) \rightarrow \mathcal{O}_Y(f^{-1}(U_i))$, $s_i \mapsto s|_{U_i \cap \bar{Z}} \mapsto r|_{f^{-1}(U_i)}$

Rmk Applying the Lemma to the case $Y =$ locally closed $Z \subseteq X$ with induced reduced sheaf, implies $\mathcal{O}_Z \cong \mathcal{O}_{\bar{Z}}$.

6. SHEAVES OF MODULES

6.1 $\mathcal{O}_X$ -modules

Def $\mathcal{O}_X$ -module is: $\bullet$ sheaf $F \in \text{Ab}(X)$ (often abbreviate $\mathcal{O}_X = \mathcal{O}_{X|U}$)
 (or sheaf of $\mathcal{O}_X$ -mods) $\bullet$ $F(U)$ is an $\mathcal{O}_X(U)$ -module
 (restrictions are compatible with module structure)

Morphism $F \rightarrow G$ of $\mathcal{O}_X$ -module is: $\bullet$ morph $F \xrightarrow{\varphi} G$ of sheaves
 (if monomorph, i.e. φ_U injective, F is $\mathcal{O}_X$ -submod of G) $\bullet$ $F(U) \xrightarrow{\varphi_U} G(U)$ is hom of $\mathcal{O}_X(U)$ -mods

Rmk stalk F_x is $\mathcal{O}_{X,x}$ -mod, and for morph $F \rightarrow G$ get $F_x \rightarrow G_x$ is $\mathcal{O}_{X,x}$ -mod hom.
 Example: A sheaf of ideals is an $\mathcal{O}_X$ -submod of $\mathcal{O}_X \Leftarrow$ (just like R -submods of R are ideals)

Fact $\mathcal{O}_X$ -Mods = (category of $\mathcal{O}_X$ -mods on X) is an abelian cat $\Leftarrow$ (proof similar to $\text{Mod}_R(X)$)
 Need notions of submod, quotient mod, ker, coker, Im agree with what get in $\text{Ab}(X)$
 e.g. $F \rightarrow G \rightarrow H$ exact $\Leftrightarrow$ exact in $\text{Ab}(X) \Leftrightarrow$ exact on stalks

Will write $\text{Hom}_{\mathcal{O}_X}$ for morphisms in this category.

6.2 Modules generated by sections

$\text{Hom}_{\mathcal{O}_X}(\mathcal{O}_X, F) \xrightarrow{1:1} F(X) \quad \forall F \in \mathcal{O}_X\text{-Mods} \quad \Leftarrow$ analogue of $\text{Hom}_R(R, M) \cong M$
 $(\varphi: \mathcal{O}_X \rightarrow F) \mapsto s = \varphi(1)$ since $\varphi_U(r) = \varphi_U(r \cdot 1) = r \cdot s|_U \quad \forall r \in \mathcal{O}_X(U)$

Similarly $\text{Hom}_{\mathcal{O}_X}(\mathcal{O}_X^n, F) \xrightarrow{1:1} F(X)^{\oplus n}$ defined by n global sections $s_1, \dots, s_n \in F(X)$
Def F is generated by global sections if

$\exists$ surjection $\bigoplus_{i \in I} \mathcal{O}_X \rightarrow F$ of $\mathcal{O}_X$ -mods $(\Leftrightarrow)$ s.t. s_i generate $\mathcal{O}_{X,x}$ -mod $F_x \quad \forall x \in X$
 same as picking sections $s_i \in F(X)$
 (as $\mathcal{O}_{X,x}$ -module, $\bigoplus_{i \in I} \mathcal{O}_{X,x} \rightarrow F_x$)

Def F is locally generated by sections if $\forall x \in X \exists$ open $U \ni x$ s.t. $F|_U$ generated by global sections

Rmk Can produce $\mathcal{O}_X$ -submods from given local sections $s_i \in F(U_i) \Leftarrow$ sheafify $U \mapsto$ possible $\mathcal{O}_X(U)$ -linear combos of $s_i|_U$ (if $s_i|_U \in F(U_i)$)

Def A sheaf has finite type if locally generated by finitely many sections.
 (equivalent definitions)
 so $\mathcal{O}_X^n \rightarrow F|_U$ some open U some $n \in \mathbb{N}$ (not fixed)

6.3 Vector bundles and coherent modules

Def $\mathcal{O}_X$ -mod F is locally free $\mathcal{O}_X$ -mod of finite rank (or vector bundle) if
 $\forall x \in X \exists$ open $U \ni x$: $F|_U \cong \mathcal{O}_U^n$ (as $\mathcal{O}_U$ -mods)
 i.e. locally generated by finite # of "independent sections"

Def X invertible sheaf ("or line bundle") if $n=1$ (fixed) $\Leftarrow$ locally $\mathcal{O}_U \cong \mathcal{O}_U$ so $F(U) = F(U)$
 generated by one section $s \in F(U)$

Question Is it enough to ask $F_x \cong \mathcal{O}_{X,x}^n \quad \forall x \in X$ (as $\mathcal{O}_{X,x}$ -mods) to say F is locally free?
 (as clear) $\Leftarrow$ (can fail)

Lemma F finite type, $\mathcal{O}_X^n \rightarrow F|_U \xrightarrow{\text{surj}} F|_U \Rightarrow \exists x \in U$ with surj $\mathcal{O}_U^n \rightarrow F|_U$, $\varphi|_x = \varphi_x$.
 (see Hw 4.4)
Pf finite type $\Rightarrow \exists$ surj $\mathcal{O}_U^n \rightarrow F|_U$. Let $s_i = \varphi_x(e_i) \in F_x$ ($e_i = 1$ in i th), so $F_x = \sum \mathcal{O}_{X,x} s_i$. Now $s_i \in F(U)$ some $x_i \in U$. Replace U by $U \cap U_1 \cap \dots \cap U_n$ so wlog $s_i \in F(U)$. Let $f_j = 1 \in (j$ -th copy of $\mathcal{O}_U$) $\mapsto \varphi(f_j)|_x = \sum \varphi(f_j)_i s_i$. So $\varphi(f_j)|_x \in \sum \mathcal{O}_{X,x} s_i$. s.t. some $v_j \in U$, again wlog $v_j = U$ (replace U by $U \cap v_1 \cap \dots \cap v_n$) $\Rightarrow \varphi(f_j) \in \text{Im } \varphi$ for $\varphi: \mathcal{O}_U^n \rightarrow F|_U$ with $\varphi(e_i) = s_i$ on U . So φ hits $\mathcal{O}_{X,x}$ -mod generators $\varphi(f_j)|_x$

Continuing above Question: We know φ_x is inj at x , but we don't know if the same φ works also for y close to x , so we do not know whether φ_y inj (recall φ inj $\Leftrightarrow \varphi_y$ inj at all stalks at $y \in U$).

Lemma In previous Lemma, if $\text{Ker } \varphi$ finite type, φ_x iso $\Rightarrow \varphi: \mathcal{O}_U^{\oplus n} \rightarrow \mathcal{F}|_U$ iso, some u_x .
 Pf shrinking U , $\exists \text{ surj } \mathcal{O}_U^{\oplus m} \xrightarrow{\psi} \mathcal{O}_U^{\oplus n} \xrightarrow{\varphi} \mathcal{F}|_U \rightarrow 0$ exact.
 Apply lemma to $\text{Ker } \varphi$: using $(\text{Ker } \varphi)_x = 0$ deduce $(\text{Ker } \varphi)|_U = 0$ possibly after shrinking U further. So φ is also injective. $\square$
 This motivates the definition: $\mathcal{O}_U$ -mod boms φ open U , $\forall n \in \mathbb{N}$ locally finitely presented

Def $\mathcal{F} \in \mathcal{O}_X$ -Mod is coherent if $\{\mathcal{F}|_U \mid U \text{ open}\}$ finite type
Def $\mathcal{F} \in \mathcal{O}_X$ -Mod is coherent if $\{\mathcal{F}|_U \mid U \text{ open}\}$ finite type
Rmk $\mathcal{F} \in \text{Coh}(X) \Rightarrow \mathcal{F}$ locally finitely presented
Pf $\mathcal{F}$ finite type $\Rightarrow \exists \text{ surj } \mathcal{O}_U^{\oplus n} \rightarrow \mathcal{F}|_U$, then consider $\text{Ker } \varphi$. $\square$

$\text{Vect}(X) = \{\text{vector bundles on } X\} \subseteq \mathcal{O}_X$ -Mods, but not an abelian cat (need not be free)
 $\text{Coh}(X) = \{\text{coherent } \mathcal{O}_X\text{-mod}\} \leftarrow \text{Fact abelian category! (explains partly its importance)}$

Claim $\mathcal{F} \in \text{Coh}(X)$ and $\mathcal{F}_x \cong \mathcal{O}_{X,x}^{\oplus n} \forall x \Rightarrow \mathcal{F} \in \text{Vect}(X)$ ($\forall x \in X$, some $n \in \mathbb{N}$ depending on x unless we fix the rank)
Claim follows by Lemmas. Converse of Claim?

Cor X locally Noetherian scheme $\Rightarrow \text{Vect}(X) = \{\mathcal{F} \in \text{Coh } X: \mathcal{F}_x, \mathcal{F}_{x,x} \cong \mathcal{O}_{X,x}^{\oplus n} \text{ some } n\} \subseteq \text{Coh}(X)$
Pf $\mathcal{F} \in \text{Vect}(X) \Rightarrow \mathcal{F}$ finite type, in general $\leftarrow$ Noetherian
 $\text{Ker}(\mathcal{O}_U^{\oplus n} \xrightarrow{\varphi} \mathcal{F}|_U)$ (need show finite type) shrinking U wlog U affine = $\text{Spec } R$
 In sections below we will prove that because $\mathcal{O}_U^{\oplus n}, \mathcal{F}|_U$ are "quasi-coherent" the problem reduces to taking global sections: $\text{Ker}(R^0 \gamma_* \mathcal{F}(U))$ and this is finitely generated since R Noeth.
 So got exact sequence $R^m \rightarrow R^n \xrightarrow{\gamma_* \mathcal{F}(U)} 0$ and this implies $\mathcal{O}_U^{\oplus m} \rightarrow \mathcal{O}_U^{\oplus n} \xrightarrow{\gamma_* \mathcal{F}(U)} 0$ exact. $\square$

6.4 $\mathcal{O}_X$ -module $\tilde{M}$ on $X = \text{Spec } R$, for R -mod M
 sheaf $\tilde{M}$ on $X = \text{Spec } R$ by Sec. 1.12 method:
 • $\tilde{M}(D_f) = M_f$ (so $\tilde{M}(X) = \tilde{M}(D_1) = M$)
 • $D_g \subseteq D_f \Rightarrow M_f \rightarrow M_g$ induced by $R_f \rightarrow R_g$
 • stable $\tilde{M}_p = \lim_{D_f \ni p} \tilde{M}(D_f) = \lim_{D_f \ni p} M_f \cong M_p$
 • $\tilde{M}(U) = \{s: U \rightarrow \coprod_{p \in \text{Spec } R} M_p : s(p) \in M_p \text{ which are locally compatible: } \forall p \in U, \exists \text{ open nbhd } p \in D_f \subseteq U \text{ with } s(x) = t_x \text{ for } x \in D_f\}$
 with the obvious restriction maps.

Rmk could assume $t = \frac{m}{f}$ since can replace D_f with D_{fm} ($= D_f$).
 • could just ask $s(x) = t_x$ on a smaller open $p \in V \subseteq D_f$.
 • $\tilde{M}$ = sheafification of $U \mapsto M \otimes_R \mathcal{O}_X(U)$
 Call $\tilde{M}$ the sheaf associated to M

UPSHOT $\tilde{M}$ is $\mathcal{O}_X$ -module on $X = \text{Spec } R$
 $\varphi: M \rightarrow N$ R -mod hom $\Rightarrow \tilde{M} \rightarrow \tilde{N}$ $\mathcal{O}_X$ -mod morph by gluing $\tilde{M}(D_f) \rightarrow \tilde{N}(D_f)$
 (just need check stalks, then use sec. 3.0) $\leftarrow$ for converse take global sections
 $\Rightarrow$ fully faithful exact functor $R\text{-Mod} \rightarrow \mathcal{O}_{\text{Spec}(R)}\text{-Mod}$

EXAMPLS • $\tilde{R} = \mathcal{O}_X$ ($X = \text{Spec } R$)
 • $\bigoplus_{i \in \mathbb{I}} \tilde{M}_i \cong \tilde{\bigoplus_{i \in \mathbb{I}} M_i}$, so $\tilde{\mathcal{O}_X} \cong \bigoplus_{i \in \mathbb{I}} \mathcal{O}_X$

EXAMPLS • $\tilde{R} = \mathcal{O}_X$ ($X = \text{Spec } R$)
 • $\bigoplus_{i \in \mathbb{I}} \tilde{M}_i \cong \tilde{\bigoplus_{i \in \mathbb{I}} M_i}$, so $\tilde{\mathcal{O}_X} \cong \bigoplus_{i \in \mathbb{I}} \mathcal{O}_X$

Call $\tilde{M}$ the sheaf associated to M
UPSHOT $\tilde{M}$ is $\mathcal{O}_X$ -module on $X = \text{Spec } R$
 $\varphi: M \rightarrow N$ R -mod hom $\Rightarrow \tilde{M} \rightarrow \tilde{N}$ $\mathcal{O}_X$ -mod morph by gluing $\tilde{M}(D_f) \rightarrow \tilde{N}(D_f)$
 (just need check stalks, then use sec. 3.0) $\leftarrow$ for converse take global sections
 $\Rightarrow$ fully faithful exact functor $R\text{-Mod} \rightarrow \mathcal{O}_{\text{Spec}(R)}\text{-Mod}$

EXAMPLS • $\tilde{R} = \mathcal{O}_X$ ($X = \text{Spec } R$)
 • $\bigoplus_{i \in \mathbb{I}} \tilde{M}_i \cong \tilde{\bigoplus_{i \in \mathbb{I}} M_i}$, so $\tilde{\mathcal{O}_X} \cong \bigoplus_{i \in \mathbb{I}} \mathcal{O}_X$

6.5 Direct image and inverse image

$\mathcal{O}_X$ -mod $\rightarrow \mathcal{F} \xrightarrow{f_*} \mathcal{F}'$ $f_* \mathcal{F}$ is $f_* \mathcal{O}_X$ -mod
 $f: X \rightarrow Y$ $\leftarrow$ top.sp.
 ringed.sp.

Algebra: Recall $R \xrightarrow{f_*} S$ hom of rings, then S is R -mod via $r \cdot s = \varphi(r)s$.
 $f: X \rightarrow Y$ morph of ringed spaces, then: $\leftarrow$ (recall $\text{Mor}(\mathcal{O}_Y, f_* \mathcal{O}_X) = \text{Mor}(f^{-1} \mathcal{O}_Y, \mathcal{O}_X)$)
 $f^{-1} \mathcal{O}_Y(U) \rightarrow \mathcal{O}_X(U)$ makes $\mathcal{O}_X$ an $f^{-1} \mathcal{O}_Y$ -mod on ringed space $(X, f^{-1} \mathcal{O}_Y)$

$f^{-1} \mathcal{F} \xrightarrow{f^*} \mathcal{F}$ $f^{-1} \mathcal{F}$ is $f^{-1}(\mathcal{O}_Y)$ -mod
 $X \rightarrow Y$ $\leftarrow$ top.sp.
 top.sp. $\leftarrow$ ringed.sp.

$(f^{-1} \mathcal{F})(U) = \lim_{V \supseteq U} f^{-1} \mathcal{F}(V)$
 $\leftarrow$ (presheaf) $\leftarrow$ so can act by $\mathcal{O}_Y(V)$
 $(f^{-1} \mathcal{O}_Y)(U) = \lim_{V \supseteq U} \mathcal{O}_Y(V)$
Warning: $\text{Hom}_{\mathcal{O}_Y}(f^{-1} \mathcal{F}(U), \mathcal{G}(U))$ would not work since do not get restriction maps.

6.6 Operations on $\mathcal{O}_X$ -mod

$\text{Hom}_{\mathcal{O}_X}(\mathcal{F}, \mathcal{G}): U \mapsto \text{Hom}_{\mathcal{O}_U}(\mathcal{F}|_U, \mathcal{G}|_U)$ is a sheaf of $\mathcal{O}_X$ -mod.
Corollary in $\mathcal{O}_X$ -Mod: $\mathcal{F}_i$ $\mathcal{O}_X$ -mods, $\bigoplus \mathcal{F}_i = \text{sheafify}(U \mapsto \bigoplus \mathcal{F}_i(U))$
 (Need sheafify: could get ∞ sums when globalise, e.g. $X = \mathbb{N}$, $\mathcal{F}_i = \mathbb{Z}$ on $\{i\}$, $s_n = (1, \dots, 1, 0, \dots)$ at $\{n\}$, try globalise)
Fact $\exists$ canonical iso $\text{Mor}(\bigoplus \mathcal{F}_i, \mathcal{G}) \cong \prod \text{Mor}(\mathcal{F}_i, \mathcal{G})$ natural in $\mathcal{F}_i, \mathcal{G}$.
product in $\mathcal{O}_X$ -Mod: $\mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{G} = \text{sheafify}(U \mapsto \mathcal{F}(U) \otimes_{\mathcal{O}_X(U)} \mathcal{G}(U))$

Fact $\exists!$ $\mathcal{O}_X$ -mod structure s.t. $\mathcal{F}(U) \otimes_{\mathcal{O}_X(U)} \mathcal{G}(U) \rightarrow (\mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{G})(U)$ hom of $\mathcal{O}_X(U)$ -mod
Universal property: $\text{Hom}_{\mathcal{O}_X}(\mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{G}, \mathcal{H}) = \text{Bilinear}_{\mathcal{O}_X}(\mathcal{F} \times \mathcal{G}, \mathcal{H})$
Rmk Stalks are $\text{Hom}_{\mathcal{O}_{X,x}}(\mathcal{F}_x, \mathcal{G}_x), \bigoplus (\mathcal{F}_i)_x, \mathcal{F}_x \otimes_{\mathcal{O}_{X,x}} \mathcal{G}_x$.
Examples on $X = \text{Spec } R: \bigoplus \tilde{M}_i \cong \tilde{\bigoplus M_i}, \tilde{M} \otimes_R \tilde{N} \cong \tilde{\bigoplus M_i \otimes_R N}$ (so $\bigoplus \mathcal{O}_X$ Hom algebra $\text{Hom}_R(M \otimes_R N, P) \cong \text{Hom}_R(M, \text{Hom}_R(N, P))$ canonically, for R -mod M, N, P (are adjoint))
Fact $\text{Hom}_{\mathcal{O}_X}(\mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{G}, \mathcal{H}) \cong \text{Hom}_{\mathcal{O}_X}(\mathcal{F}, \text{Hom}_{\mathcal{O}_X}(\mathcal{G}, \mathcal{H}))$ canonically & functorial in $\mathcal{F}, \mathcal{G}, \mathcal{H}$.
Cor $\mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{G}$ adjoint, $\mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{G}$ right exact, $\text{Hom}_{\mathcal{O}_X}(\mathcal{G}, \mathcal{H})$ left exact.
Fact $f: X \rightarrow Y \Rightarrow f^{-1}(\mathcal{F} \otimes_{\mathcal{O}_Y} \mathcal{G}) \cong f^{-1} \mathcal{F} \otimes_{f^{-1} \mathcal{O}_Y} f^{-1} \mathcal{G}$ canonically ($\mathcal{F}, \mathcal{G}$ $\mathcal{O}_Y$ -mod)
6.7 Pullback $\leftarrow$ can prove this using universal property, or by hand thinking about $\mathcal{O}$
Rmk $R \rightarrow S$ rings, M R -mod, N S -mod $\Rightarrow M \otimes_R N$ is $\{R\text{-mod since } N \text{ } R\text{-mod via } R \rightarrow S \text{ (} r \cdot (m \otimes n) = (rm) \otimes n = m \otimes rn)\}$
 $\Rightarrow M \otimes_R N$ is S -mod by $s \cdot (m \otimes n) = m \otimes sn$
 similarly: $X \xrightarrow{f} Y$ $\leftarrow$ ringed $\Rightarrow f^* \mathcal{F} = f^{-1}(\mathcal{F}) \otimes_{f^{-1} \mathcal{O}_Y} \mathcal{O}_X$ is an $f^{-1} \mathcal{O}_Y$ -mod but also an $\mathcal{O}_X$ -mod!

Fact $\exists!$ $\mathcal{O}_X$ -mod : presheaf tensor = $f^{-1}(F)(U) \otimes_{f^{-1}\mathcal{O}_Y(U)} \mathcal{O}_X(U) \rightarrow f^*F(U)$ is $\mathcal{O}_X(U)$ -mod hom structure s.t.

Example $f^*\mathcal{O}_Y = \mathcal{O}_X$ (since $f^{-1}\mathcal{O}_Y \otimes_{f^{-1}\mathcal{O}_Y} \mathcal{O}_X \cong \mathcal{O}_X$ canonically)

Exercise $X \xrightarrow{f} Y \xrightarrow{g} Z \Rightarrow f^* \circ g^* = (g \circ f)^*$ (use last Fact in 6.6, using Sec.1.9)
 $f^*(F \otimes_{\mathcal{O}_Y} G) = f^*F \otimes_{\mathcal{O}_X} f^*G$ canonically & functorial

Upshot $f: X \rightarrow Y$ morph of ringed spaces $\Rightarrow \text{Mod}_{\mathcal{O}_X}(X) \xrightarrow{f^*} \text{Mod}_{\mathcal{O}_Y}(Y)$ and $\leftarrow^*$

Theorem (exercise) f^*, f_* are adjoint functors: $\text{Mod}_{\mathcal{O}_X}(f^*F, G) \cong \text{Mod}_{\mathcal{O}_Y}(F, f_*G)$

hence f_* left exact, f^* right exact

HWK 3 f_* commutes with limits $\lim$ for example $\prod$, f^* commutes with colimits $\varinjlim$ for example $\oplus$

Example $f^*(\oplus \mathcal{O}_Y) = \oplus f^*\mathcal{O}_Y = \oplus \mathcal{O}_X$

Exercise Deduce from that $f^*(\text{Vect}(Y)) \subseteq \text{Vect}(X)$.

6.8 $\tilde{M}$ on any scheme

M R -mod, $X \xrightarrow{\text{canonical}} \text{Spec } \Gamma(X, \mathcal{O}_X) \xrightarrow{\alpha} \text{Spec } R$ Then get $F_M := \alpha^* \tilde{M}$

Easier: $(X, \mathcal{O}_X) \xrightarrow{\pi} \text{ringed space (point, } R) \text{ (on sheaves } \pi_* \mathcal{O}_X = \Gamma(X) \leftarrow R)$

$F_M := \pi^* M$
 $= \text{sheafify } (U \mapsto M \otimes_{R-\pi^{-1}R} \mathcal{O}_X(U)) \leftarrow \text{(since } \pi^{-1}M \otimes_{\pi^{-1}R} \mathcal{O}_X \text{ and } (\pi^{-1}R)(U) = R)$

(get same answer since $X \xrightarrow{\alpha} \text{Spec } R \xrightarrow{\pi} \text{(point, } R)$, $\tilde{M} = \pi^* M$ by construction, $\pi^* = \alpha^* \pi_*$)

Claim $f: Y \rightarrow X$ (morph of ringed spaces) $\Rightarrow f^*F_M = F_N$ where $N = M \otimes_{\Gamma(X)} \Gamma(Y)$

Pf $M \xrightarrow{\Gamma(X)} \Gamma(X)$ -module (case $R \xrightarrow{\Gamma(X)} \Gamma(X)$)
 $\pi_Y \downarrow \quad \downarrow \pi_X$
 $Y \xrightarrow{f} X$
 $f^* \pi_X^* M = \pi_Y^* \psi^* M$
 $\psi^* M = \psi^* M \otimes_{\psi^{-1}\Gamma(X)} \Gamma(Y) = M \otimes_{\Gamma(X)} \Gamma(Y)$

Cor $\alpha: \text{Spec } S \rightarrow \text{Spec } R \Rightarrow \alpha^* \tilde{M} = \widetilde{M \otimes_R S}$ (compare Sec.2.3)

Example $D_f = \text{Spec } R_f \hookrightarrow \text{Spec } R \Rightarrow \tilde{M}|_{D_f} = \widetilde{M \otimes_R R_f} = \tilde{M}_f$ (stronger statement than saying $\tilde{M}(D_f) = M_f$)

6.9 Classification of $\mathcal{O}_X$ -homs $\tilde{M} \rightarrow F$

Lemma $X = \text{Spec } R \Rightarrow \text{Hom}_{\mathcal{O}_X}(\tilde{M}, F) \xrightarrow{\cong} \text{Hom}_R(M, \Gamma(X, F)) \quad \forall \mathcal{O}_X\text{-mod } F$ (compare Sec.2.3)

Pf $\pi: (X, \mathcal{O}_X) \rightarrow (\text{point}, R)$ morph of ringed spaces $(\pi^*: R \xrightarrow{\text{id}} \pi_* \mathcal{O}_X = \mathcal{O}_X(X) = R)$
 $\tilde{M} = \pi^* M, \quad \Gamma(X, F) = \pi_* F$

$\Rightarrow \text{Hom}_{\mathcal{O}_X}(\tilde{M}, F) = \text{Hom}_{\mathcal{O}_X}(\pi^* M, F) \xrightarrow{\pi^* \pi_* \text{ adjoint}} \text{Hom}_R(M, \pi_* F) = \text{Hom}_R(M, \Gamma(X, F)). \square$

Exercise Using 6.8: $\text{Hom}_{\mathcal{O}_X}(F, M) \xrightarrow{\cong} \text{Hom}_R(M, \Gamma(X))$ using $R \xrightarrow{\text{sheaf}} \Gamma(X, \mathcal{O}_X)$ to make $F(X)$ an R -mod.

6.10 Flatness

Def F is flat $\mathcal{O}_X$ -mod if $F \otimes_{\mathcal{O}_X} \cdot$ is exact

so $\Leftrightarrow F_x$ flat $\mathcal{O}_{X,x}$ -mod $\forall x$.

Example $U \xrightarrow{\sim} X$ open subsch. $\Rightarrow i_U \mathcal{O}_U$ is flat $\mathcal{O}_X$ -mod

Rmk Morph of schemes $f: X \rightarrow Y$ is flat $\Leftrightarrow \mathcal{O}_X$ flat $f^{-1}\mathcal{O}_Y$ -module (see Sec.3.6)

Claim $f: X \rightarrow Y$ flat $\Rightarrow f^*: \mathcal{O}_Y\text{-Mod} \rightarrow \mathcal{O}_X\text{-Mod}$ is exact (not just right exact)

Pf f^{-1} is exact $\Rightarrow \mathcal{O}_Y\text{-Mod} \xrightarrow{f^{-1}} f^{-1}\mathcal{O}_Y\text{-Mod}$ exact,

$\mathcal{O}_X$ exact by Rmk $\Rightarrow f^*F = f^{-1}F \otimes_{f^{-1}\mathcal{O}_Y} \mathcal{O}_X$ is composite of two exact functors $\square$

Facts $\cdot$ free $\Rightarrow$ flat

$\cdot (X, \mathcal{O}_X)$ ringed space

$\cdot$ Can take $\oplus$ of flat mods

$\cdot \mathcal{O} \rightarrow F_1 \rightarrow F_2 \rightarrow F_3 \rightarrow \mathcal{O}$ exact : outer two or last two flat $\Rightarrow$ all flat

$\cdot$ Combine SES's show images $(F_1 \rightarrow F_{n-1})$ flat (so "flat resolution of flat $\mathcal{O}_X\text{-mod } F"$)

$\cdot F_3$ flat $\Rightarrow$ sequence $\mathcal{O} \rightarrow \mathcal{O}_X\text{-mod } G$ is exact

$\cdot$ any $\mathcal{O}_X$ -mod G is exact

$\cdot$ Fact " $\Leftarrow$ " holds also if just assume $\mathcal{O}_X$ is coherent

7. (QUASI-)COHERENT SHEAVES

7.1 QCoh(X)

Recall F coherent $\Rightarrow F$ locally finitely presented

(Sec.6.3) and " $\Leftarrow$ " holds if X locally Noetherian scheme. (now weaken this condition by dropping finiteness)

Def F quasi-coherent $\Leftrightarrow F$ is locally presented, i.e. $\forall x, \exists$ open $U \ni x \in U$

$\exists \bigoplus_{i \in I} \mathcal{O}_U \rightarrow \bigoplus_{j \in J} \mathcal{O}_U \rightarrow F|_U \rightarrow 0$ exact.

(any ringed space $(X, \mathcal{O}_X)$) $\mathcal{O}_U$ -mods

Summary: coherent $\Rightarrow$ locally finitely presented $\Rightarrow$ quasi-coherent (= locally presented)

vector bundle $\Rightarrow$ locally generated by finitely many sections $\Rightarrow$ locally generated by sections

Lemma For $X = \text{Spec } R: \left(\bigoplus_{i \in I} \mathcal{O}_X \rightarrow \bigoplus_{j \in J} \mathcal{O}_X \rightarrow F \rightarrow 0 \right) \Leftrightarrow (F \cong \tilde{M} \text{ some } R\text{-module } M)$

Pf $\Rightarrow$ Let $M = \bigoplus_j R / \text{Im}(\bigoplus_i \mathcal{O}_R \rightarrow \mathcal{O}_R)$ (taking global sections)

by exact functor from 6.4: $\bigoplus_i \mathcal{O}_X \rightarrow \bigoplus_j \mathcal{O}_X \rightarrow F \rightarrow 0$ exact

$\bigoplus_i \tilde{R} \rightarrow \bigoplus_j \tilde{R} \rightarrow \tilde{M} \rightarrow 0$ exact

$\tilde{R} \cong \tilde{M}$ by uniqueness of kernels up to iso:

$\tilde{R} \cong \tilde{M}$

$\Rightarrow F = \tilde{M}$: pick J = set of generators m_j for $R\text{-mod } M$ (e.g. $J = M$)

pick $I = \{ \text{generators } m_j \}$ $\tilde{R} \rightarrow \tilde{M} \rightarrow 0$ exact

apply $\sim$ to $\tilde{R} \rightarrow \tilde{M} \rightarrow 0$ exact

$\tilde{R} \rightarrow \tilde{M} \rightarrow 0$ exact

$\tilde{R} \rightarrow \tilde{M} \rightarrow 0$ exact

$\tilde{R} \rightarrow \tilde{M} \rightarrow 0$ exact

$\tilde{R} \rightarrow \tilde{M} \rightarrow 0$ exact

$\tilde{R} \rightarrow \tilde{M} \rightarrow 0$ exact

$\tilde{R} \rightarrow \tilde{M} \rightarrow 0$ exact

$\tilde{R} \rightarrow \tilde{M} \rightarrow 0$ exact

$\tilde{R} \rightarrow \tilde{M} \rightarrow 0$ exact

$\tilde{R} \rightarrow \tilde{M} \rightarrow 0$ exact

$\tilde{R} \rightarrow \tilde{M} \rightarrow 0$ exact

$\text{FEQ Coh}(X) \iff \forall x \in X \exists \text{ affine open } U \ni x \in \text{Spec } R, F|_U \cong \widetilde{M}$ some R -mod M
 $\text{FE Coh}(X) \iff$ in addition require M is coherent R -mod $\swarrow$ $\begin{matrix} M = F(U) \\ R = \text{Coh}(U) \end{matrix}$ $\swarrow$ WLOG

$\{ M \text{ finitely generated} \}$
 $\{ \ker(R^n \xrightarrow{f} M) \text{ is f.g., any } n \in \mathbb{N} \}$
 $\leftarrow \text{any hom of } R\text{-mods}$
 Noeth., coherent = f.g. (since R^n f.g., so its submods are f.g. as R Noeth.)
 $\Rightarrow \partial_X$ is coherent
 loc. Noeth. scheme $\Rightarrow$ ideal sheaf of any closed subsh. is coherent.
 (as $\text{Fluc}(u) = \tilde{R}(u) = 0$)
 Idea: want $\forall$ f.g. submod of M to have finite presentation, indeed get exact sequence
 $R^m \xrightarrow{f} R^n \xrightarrow{g} \text{im } g \rightarrow 0$
 $\quad \quad \quad \uparrow \quad \quad \quad \uparrow$
 $\quad \quad \quad \text{f.g.} \quad \quad \quad \text{f.g.}$
 $\quad \quad \quad \text{f.g.} \quad \quad \quad \text{f.g.}$

Noeth., coherent = f.g. (since R^n f.g., so its submodules are f.g. as R -Noeth.)
 loc. Noeth. scheme $\Rightarrow \mathcal{O}_X$ is coherent $\Rightarrow$ ideal sheaf of any closed subsch. is coherent.
linearly exact sequence
 $R^m \rightarrow R^n \xrightarrow{\varphi} R^m \rightarrow 0$
 $\downarrow$ map to gens of ker φ

$$\begin{aligned} \text{me: } F \in \mathbf{QCoh}(X) &\Leftrightarrow \exists \text{ affine open cover } X = \bigcup U_i \text{ s.t. } F|_{U_i} \simeq \widetilde{M}_i \text{ for } R_i\text{-mods } M_i \\ &U_i = \text{Spec } R_i \\ &(\text{WLOG: } R_i = \mathcal{O}_X(U_i) \quad M_i = F(U_i)) \\ F \in \mathbf{Coh}(X) &\Leftrightarrow \text{" and } M_i \text{ coherent.} \end{aligned}$$

restriction to open $V \subseteq X$: $Q\text{Coh}(X) \rightarrow Q\text{Coh}(V)$, $\text{Coh}(X) \rightarrow \text{Coh}(V)$
 $\bigvee \mathcal{U} = \bigcup D_{f_i}$ for $f_i \in R$ then $F|_{\bigcup D_{f_i}} \approx F|_{D_{f_i}} \approx \widetilde{M}|_{D_{f_i}} \approx \widetilde{M_{f_i}}|_{D_{f_i}}$ (and use fact that localisation preserves "coherent" property)
 is an locally module. $\square$ Example in 6.8

asi-coherence a good notion?

$\rightarrow \text{Aff}, R \mapsto (\text{Spec}(R), \theta_{\text{Spec}})$ equivalence of cats

$\rightarrow \mathcal{O}_{\text{Spec}(R)}\text{-Mod}_s, \quad M \mapsto \widetilde{M}$ not equivalence of cats
 $X = \text{Spec } k[x] = \mathbb{A}^1_k$, skyscraper sheaf at 0: $F(U) = \begin{cases} k[x] & \text{if } 0 \in U \\ 0 & \text{else} \end{cases}$ $\xrightarrow{\text{FEQ-mod}}$
 $\Rightarrow$ if the above were an equivalence of cats then $F \cong \widetilde{M}$ some $b[x]\text{-mod } M$

so $k[x] = F(X) \cong \widetilde{F}(X) = M$. But $\widetilde{k[x]} = \theta_X$ is not isomorphic to F !
 restrict which θ_X -mods you allow : want them locally to look like $\widetilde{M}$,
 just like when we studied sheaves of ideals that locally look like $\widetilde{I}$

For $X = \text{Spec } R : R\text{-Mods} \rightarrow \text{QCoh}(X)$ equivalence of categories

view of general properties of $\text{QGh}(X)$ and $\text{Coh}(X)$ for X scheme

abelian category, and $\text{Coh}(X) \xrightarrow{\text{incl}} \text{Mod}$
 $\text{Coh}(X) \xrightarrow{\text{incl}} \text{Mod}$ are exact functors
 (for Coh(X properties) enough if X ringed)

Similar can take Ker, Coker , Image in both $\rightarrow F_2 \rightarrow F_3 \rightarrow 0$ exact in $\mathcal{O}_X\text{-Mods.}$

the $F_i \in \text{Coh}(X) \Rightarrow$ all three are. Same holds for $\text{Coh}(X)$ (not for $\text{Vect}(X)$)
 $0 \rightarrow F_1 \rightarrow F_2 \rightarrow F_3$ exact, and F_2, F_3 are, then F_1 is. (Pf. $F_1 \cong \text{Ker}(F_2 \rightarrow F_3)$, use (1.1))

Let $\underline{\text{finite}} \oplus, \cdot \otimes_{\theta_X}, \text{Hom}_{\theta_X}(\cdot, \cdot)$ in $\mathcal{Q}\text{Coh}(X)$, $\text{Coh}(X)$ and $\text{Vect}(X)$ Rosenberg thm $\hookrightarrow$ for $\mathcal{Q}\text{Coh}, \text{Hom}_{\theta_X}(F, G)$ need assume F locally finitely presented

-compact & separated (e.g. variety) $\Rightarrow X$ is determined up to iso by $\mathcal{Q}(\text{Coh } X)$!

type subsheaf $F \subseteq G, G \in \text{Coh}(X) \Rightarrow F \in \text{Coh}(X)$

$\varphi: G \rightarrow G$, $G \in \text{Coh } X$, F finite type, $\varphi_x: F_x \rightarrow G_x$ injective $\Rightarrow \varphi|_U: F|_U \rightarrow G|_U$ injective $\Rightarrow \varphi|_U$ is injective
 Picard group $\text{Pic}(X) = \{\text{isomorphism classes of invertible sheaves}\}$
 group operation is $\cdot$. $\otimes_{\mathcal{O}_X}$. (abelian group as $F \otimes_{\mathcal{O}_X} G \cong G \otimes_{\mathcal{O}_X} F$)
 We proved it in case F is in $\mathcal{P}_f$ claim in Sec. 6.3

Abstract

X loc. Noeth. scheme $\Rightarrow f^*: \text{Coh } Y \rightarrow \text{Coh } X$.

$$i^* (fx \in U \subseteq Y \text{ open}) \quad \text{Vect}^U Y \xrightarrow{f^*} \text{Vect}^U X$$

$\rightarrow U$, using g^* right exact & commutes with $\oplus$.
 $\circ$ exact, and $x \in f^{-1}U$ open. ← using $X_{loc, Noeth}$

$\lambda \Rightarrow f^*F$ Loc. finitely presented $\Rightarrow f^*F \in \text{Coh}(X) \square$

(above proof for I, J finite)

issue is $f^{-1}(\text{affine})$ need not be affine.

$$\text{f}_x: \mathbb{Q}G_h X \rightarrow \mathbb{Q}G_h Y$$

sheaf property, where $X = U_i$: affine open cover
 on overlaps (Sec. 14) $U_i \cap U_j = U_{i,j} = U_{j,i}$ " " "

$$\Gamma \Rightarrow 0 \rightarrow f_* F \rightarrow \Gamma f_*(F|_{U_i}) \rightarrow \Gamma f_*(F|_{U_{i,j,k}})_{\text{exact}}$$

$(i_k) \in \text{QCh}(Y)$ then $f_* F \in \text{QCh}(Y) \xleftarrow[\text{in } \mathbb{Z}_2]{\text{Trick}(2)} \text{sec. 6.5}$

$\hookrightarrow$ is $\oplus$, but $\sim$ commutes with $\oplus$ so finally done! $\square$

$$\text{h.} \Rightarrow f_x: \text{Coh } X \rightarrow \text{Coh } Y \leftarrow \begin{array}{l} \omega_{ijk} = \text{unif. amne.} \\ f_{*A_i} = k[x] \text{ not} \\ \text{finite } k\text{-mod} \end{array}$$

previous morph, $F = \widetilde{\prod k[t]}$, $f_* F = \widetilde{\prod k[t]}$ if $\text{assume } \in \text{Qcoh}$ $\leftarrow \text{by 4.6}$

$$= (\bigcup D_i) = ({}^a F)(D_i) \text{ but } \notin (\bigcap [t_i])(D_i) = (\bigcap [t_i])_t$$

$$\text{s.t. } 1 \in \text{all } f_i > * \neq \bigcap [t_i]_t.$$

$\underbrace{\text{cycle}}_s \underbrace{\text{condition}}_s (M, f, f_w) \xrightarrow{\psi_i, k} (M, f, f_i) \xrightarrow{\psi_i, k} \text{case } k = i \text{ get } \psi_i = \psi_i$

$$\psi_{ij} = (M_{ij}, f_{ij})$$

idea: local data which agrees on overlaps

$$- \phi_{j,i} \left(\frac{1}{T} \right) \Bigg|$$
$$f_i \rightarrow M_i \text{ and } \psi_{i,0}^j = \frac{1}{\pi_j(m)} = \frac{1}{\pi_j(m)} \quad \forall m \in M$$

at every prime $q \in \text{Spec } R_{\mathfrak{f}_0}$

$(\oplus_i \mathcal{M}_i) \xrightarrow{g_2} (\oplus_i \mathcal{M}_i) \oplus (\oplus_i \mathcal{M}_i)$

$M \rightsquigarrow M_P, M_i \rightsquigarrow (M_i)_P, f_0 \rightsquigarrow 1.$ localizing at f_0 is like localizing at 1 since f_0 is a unit in R_P

[illegible]

WLOG $M_i = N_{f_i}$ (identifies via ψ_{f_i}), so cocycle cond. becomes:

$$\Rightarrow 0 \rightarrow N \xrightarrow{\text{natural}} \bigoplus_{i \neq \ell} N_{f_i} \xrightarrow{\varphi_p} \bigoplus_{i,j} N_{f_i, f_j}$$

$$(N \rightarrow \bigoplus_{i \neq \ell} N_{f_i} \xrightarrow{\text{natural}} \bigoplus_{i,j} N_{f_i, f_j}) \xrightarrow{\psi_{f_i, f_j}} \bigoplus_{i,j} (M_j)_{f_i} \xrightarrow{\psi_{f_i, f_j}} \bigoplus_{i,j} (M_j)_{f_i} \xrightarrow{\psi_{f_i, f_j}} \bigoplus_{i,j} (M_j)_{f_i} \xrightarrow{\psi_{f_i, f_j}} \bigoplus_{i,j} (M_j)_{f_i}$$

Sub-claim This is exact ($\Rightarrow N = \ker \varphi_p = M$, π_ℓ iso, $\psi_{f_i} = \text{id}$ under identifications via π maps)
 Pf Enough to prove after localising at each max ideal m $\leftarrow$ see 3.0
 By $\otimes$ not all $f_i \in m$ otherwise $1 \in \langle \text{all } f_i \rangle \subseteq m$
 Say $f_k \notin m$, so WLOG replace $N \rightarrow N_m, R \rightarrow R_m, f_k \mapsto 1$ $\leftarrow$ $f_k \notin m$ so unit in localising R_m

$\Rightarrow 0 \rightarrow N \rightarrow \bigoplus_{i \neq k} N_{f_i} \xrightarrow{\varphi_p} \bigoplus_{i,j} N_{f_i, f_j}$
 clearly injective
 $n \otimes n_i \in \ker \text{then } \frac{n_i}{1} = \frac{n_i}{1} N_{f_i} f_k = N_{f_i} \quad \forall i \quad \square$
 hence $= n \otimes \frac{n_i}{1}$ so image of n via previous map

7.6 Qcoh(X), Coh(X), Vect(X) for $X = \text{Spec } R$

Theorem For $X = \text{Spec } R$, $\exists$ equivalence of categories

$$\begin{array}{ccc} R\text{-Mod} & \xrightarrow{\sim} & \text{Qcoh}(X) \\ M & \xrightarrow{\sim} & \tilde{M} \\ F(X) = \Gamma(X, F) & \xleftarrow{\sim} & F \end{array}$$

Pf. Easy direction: $M \mapsto F = \tilde{M} \mapsto F(X) = \tilde{M}(X) = M$. Converse: given F want $F \cong \tilde{F}(X)$.
 $\Rightarrow$ locally $\forall p \in X, \exists p \in D_f$ st. $F|_{D_f} \cong \tilde{N}$ some R_f -mod N $\leftarrow$ By cor in 7.1
 cover X by finitely many such, say N_i on $D_{f_i}, i=1, \dots, n$, so $1 \in \langle \text{all } f_i \rangle$ and Spec quasi-compact
 $\Rightarrow$ On overlaps: $\psi_{ij} : (N_i)_{f_j} \xrightarrow{\varphi_{f_i}} (N_j)_{f_i} \xrightarrow{\varphi_{f_j}} (N_j)_{f_i f_j} \xrightarrow{\varphi_{f_i}} (N_i)_{f_i f_j}$ satisfies cocycle condition $\leftarrow$ since $(N_i)_{f_j}$ and other two are identified with $F|_{D_{f_i f_j}}$
 $\Rightarrow$ by gluing them $\exists M$ with $M_{f_i} = N_i$ compatibly with the ψ_{ij}
 But then $\tilde{M}, F$ have isomorphic local gluing data for cover $X = D_{f_1} \cup \dots \cup D_{f_n}$ so $\tilde{M} \cong F$.
 Explicitly: $m \in M \mapsto m_i = \frac{m}{1} \in M_{f_i} = N_i \xrightarrow{\varphi_{f_i}} s_i \in F(D_{f_i})$ and $s_i|_{D_{f_i f_j}} = s_j|_{D_{f_i f_j}}$ so globalises to unique $s \in F(X)$. Recall $M \rightarrow F(X)$ determines $\tilde{M} \rightarrow F$ by sec. 6.9. $\square$

Cor $X = \text{Spec } R: F \in \text{Coh } X \Leftrightarrow F = \tilde{M}$ for coherent module M $\leftarrow$ and if R Noether, get:
 Pf $F = \tilde{F}(X)$ by Theorem. In definition of coherent take global sections $\Rightarrow F(X)$ coherent R -mod,
 and conversely if M coherent get $\tilde{M}$ coherent since $\sim$ is exact & fully faithful. $\square$
 Fact $X = \text{Spec } R: F \in \text{Vect } X \Leftrightarrow F = \tilde{M}$ for finitely presented $\tilde{M}$ $\leftarrow$ (see 11.4.4)
 means in R -mods $\text{Hom}(M_i, \cdot)$ exact $\Rightarrow M$ is a direct summand of some free R -mod

8. Čech Cohomology

8.1 Čech complex

X top. space, $X = \bigcup U_i$ open cover
 $U_I = U_{i_0} \cap \dots \cap U_{i_n}$ for $I = (i_0, \dots, i_n)$ multi-index, abbreviate $|I| = n$
 $C^n = \check{C}^n_{\{U_i\}} = \prod_{|I|=n} \Gamma(U_I, F) \xleftarrow{F \in \text{Ab}(X)}$
 $d = d^n: C^n \rightarrow C^{n+1}$
 $(ds)_I = \sum_{j=0}^{n+1} (-1)^j s_j|_{U_I}$
 $\leftarrow$ where $I_j = (i_0, \dots, i_j, \dots, i_{n+1})$
 $\leftarrow$ omit i_j if omit $i_j, i_k, \dots$
 later also use notation $I_{j,k,\dots}$ if omit $i_j, i_k, \dots$
 $\leftarrow$ so SEC^n is a collection $s_I \in F(U_I)$
 $\leftarrow$ called cochain

Example $C^0 = \prod_i \Gamma(U_i) \xrightarrow{d} \prod_{i,j} \Gamma(U_{ij}) = C^1$

$(s_i) \mapsto (s_j|_{U_{ij}} - s_i|_{U_{ij}})$

$C^1 = \prod_{i,j} \Gamma(U_{ij}) \xrightarrow{d} \prod_{i,j,k} \Gamma(U_{ijk}) = C^2$

$(s_{ij}) \mapsto (s_{jk}|_{U_{ijk}} - s_{ik}|_{U_{ijk}} + s_{ij}|_{U_{ijk}})$

Claim $d^2 = 0$, so (C^*, d) is a complex

Pf $(d^2 s)_I = \sum_{j,k=0}^{n+2} (-1)^k (ds)_{I_{j,k}} = \sum_{j,k=0}^{n+2} (-1)^k \left(\sum_{l=0}^{k+1} (-1)^l s_{I_{j,k,l}} \right)$

$= 0$. $\square$ $\leftarrow$ anti-symmetry if swap j,k (notice full sum is over all $j \neq k$)

Def $H^n(X, F) = \check{H}^n_{\{U_i\}}(X, F) = \ker d^n / \text{Im } d^{n-1}$

Lemma $H^0(X, F) = \Gamma(X, F)$

Pf $s_j|_{U_{ij}} = s_i|_{U_{ij}}$ says s glues to global section. $\square$

Terminology 1) hom of complexes $f: C^n \rightarrow C^n$ is chain map if $f \circ d = d \circ f$

2) $h: C^n \rightarrow C^{n-1}$ is chain homotopy between chain maps f, g if $f - g = d \circ h + h \circ d$

Consequences: 1) $f: H^n \rightarrow H^n$ via $f[c] = [fc]$ well-defined $\leftarrow [c] = [c + db]$

2) $f = g: H^n \rightarrow H^n \leftarrow (dc = 0 \Rightarrow [fc - gc] = [d(hc)] = 0)$

Key trick To show $H^k = 0$ can find chain homotopy between id and 0 .

Rmk If a homomorphism $d_n: C_n \rightarrow C_{n-1}$ decreases the degree by 1, and $d_{n-1} \circ d_n = 0$

then $H_n = \ker d_n / \text{Im } d_{n+1}$ is called the homology of (C, d) . In this case a chain homotopy is degree increasing: $h: C_n \rightarrow C_{n+1}$ with $f_n - g_n = d_{n+1} \circ h_n - h_n \circ d_n$.

$U_{ij} = U_i \cap U_j$
 $U_{ijk} = U_i \cap U_j \cap U_k$
 $\leftarrow$ ordered, allow repetitions
 $\leftarrow$ size is actually $n+1$

$C^n = \check{C}^n_{\{U_i\}} = \prod_{|I|=n} \Gamma(U_I, F) \xleftarrow{F \in \text{Ab}(X)}$

$d = d^n: C^n \rightarrow C^{n+1}$

$(ds)_I = \sum_{j=0}^{n+1} (-1)^j s_j|_{U_I}$

$\leftarrow$ where $I_j = (i_0, \dots, i_j, \dots, i_{n+1})$

$\leftarrow$ omit i_j if omit $i_j, i_k, \dots$

later also use notation $I_{j,k,\dots}$ if omit $i_j, i_k, \dots$

Example $C^0 = \prod_i \Gamma(U_i) \xrightarrow{d} \prod_{i,j} \Gamma(U_{ij}) = C^1$

$(s_i) \mapsto (s_j|_{U_{ij}} - s_i|_{U_{ij}})$

$C^1 = \prod_{i,j} \Gamma(U_{ij}) \xrightarrow{d} \prod_{i,j,k} \Gamma(U_{ijk}) = C^2$

$(s_{ij}) \mapsto (s_{jk}|_{U_{ijk}} - s_{ik}|_{U_{ijk}} + s_{ij}|_{U_{ijk}})$

Claim $d^2 = 0$, so (C^*, d) is a complex

Pf $(d^2 s)_I = \sum_{j,k=0}^{n+2} (-1)^k (ds)_{I_{j,k}} = \sum_{j,k=0}^{n+2} (-1)^k \left(\sum_{l=0}^{k+1} (-1)^l s_{I_{j,k,l}} \right)$

$= 0$. $\square$ $\leftarrow$ anti-symmetry if swap j,k (notice full sum is over all $j \neq k$)

Def $H^n(X, F) = \check{H}^n_{\{U_i\}}(X, F) = \ker d^n / \text{Im } d^{n-1}$

Lemma $H^0(X, F) = \Gamma(X, F)$

Pf $s_j|_{U_{ij}} = s_i|_{U_{ij}}$ says s glues to global section. $\square$

Terminology 1) hom of complexes $f: C^n \rightarrow C^n$ is chain map if $f \circ d = d \circ f$

2) $h: C^n \rightarrow C^{n-1}$ is chain homotopy between chain maps f, g if $f - g = d \circ h + h \circ d$

Consequences: 1) $f: H^n \rightarrow H^n$ via $f[c] = [fc]$ well-defined $\leftarrow [c] = [c + db]$

2) $f = g: H^n \rightarrow H^n \leftarrow (dc = 0 \Rightarrow [fc - gc] = [d(hc)] = 0)$

Key trick To show $H^k = 0$ can find chain homotopy between id and 0 .

Rmk If a homomorphism $d_n: C_n \rightarrow C_{n-1}$ decreases the degree by 1, and $d_{n-1} \circ d_n = 0$

then $H_n = \ker d_n / \text{Im } d_{n+1}$ is called the homology of (C, d) . In this case a chain homotopy is degree increasing: $h: C_n \rightarrow C_{n+1}$ with $f_n - g_n = d_{n+1} \circ h_n - h_n \circ d_n$.

8.2 Čech complex with ordering

e.g. if X quasi-compact

Repetitions of indices are annoying since $C^n \neq 0$ all $n \geq 0$ even if finite # U_i

Trick pick total ordering on indices

C_n^+ : as C^n but only allow $I = (i_0, \dots, i_n)$ if $i_0 < i_1 < \dots < i_n$, as before

$\Rightarrow C_n^+ \subseteq C^n$ subcomplex

Claim $H_n^+ \cong H^n$

Non-examinable Proof ("Serre's Trick")

Let S_* = free abelian group generated by all index sets I , so: $S_n = \langle I : |I| = n \rangle$

Differential: $\partial I = \sum (-1)^j I_j$ so $\partial: S_n \rightarrow S_{n-1}$

S_*^+ = subgroup generated by strictly ordered index sets I

$S_*^+ \subseteq S_*$, S_*^+ are acyclic

Step 1 S_*, S_*^+ are acyclic

Pf $h: S_*^+ \rightarrow S_*$, $h(I) = \begin{cases} (I, I) & \text{if } I \neq i_0 \\ 0 & \text{if } I = i_0 \end{cases}$ if $I \neq i_0: \partial h I = \partial(\partial I) = I + \sum (-1)^j (I_j)$

$\Rightarrow I = (\partial I + h \partial) I$. Exercise: check same holds if $I = i_0$.

$\Rightarrow \text{id} - 0 = \partial h + h \partial$ For S_* it is even easier: $h(I) = (I, I)$ works. $\square$

Step 2 $f(I) = \begin{cases} 0 & \text{if } \exists \text{ repeated indices in } I \\ \text{Sign}(\sigma) \cdot \sigma(I) & \text{otherwise, where } \sigma \text{ unique permutation s.t. } \sigma(I) \text{ ordered} \end{cases}$

$\Rightarrow f$ chain map, $f = \text{id}$ on S_0 , $f(S_*) \subseteq S_*^+$, f is id on S_*^+ , f is a projection to S_*^+

Pf $\sigma(I) \in S_*^+$ and if I is ordered then $\sigma = \text{id}$. On $S_0: f((i_0)) = (i_0)$.

$\partial f I = \sum (-1)^j \text{sign}(\sigma) \sigma(I_j) \xrightarrow{\sigma} \sum (-1)^j \text{sign}(\sigma \circ \tau) \tau(I_j)$ get same set, $\text{sign}(\sigma) = \text{sign}(\tau) \cdot (-1)^{k-j}$ since

$f \partial I = \sum (-1)^k \text{sign}(\tau) \tau(I_k)$ σ does an extra $k-j$ transpositions to move j to position k

Step 3 General trick: C_* free acyclic complex, a chain map $f: C_* \rightarrow C_*$ has $f_0 = \text{id}: C_0 \rightarrow C_0$

then f, id are chain homotopic: $\exists k: C_n \rightarrow C_{n+1}$ with $f - \text{id} = \partial k + k \partial$

Pf Build k inductively by equation $\partial_{n+1} \circ k_n = f_n - \text{id} - k_{n-1} \circ \partial_n$

$n=0: \partial_1 \circ k_0 = f_0 - \text{id} - k_{-1} \circ \partial_0$ so just define $k_0 = 0$.

Assume true for $n-1: \partial_n k_{n-1} = f_{n-1} - \text{id} - k_{n-2} \circ \partial_{n-1}$

$\partial_n(f_n - \text{id} - k_{n-1} \circ \partial_n) = f_{n-1} \circ \partial_n - \partial_n - (\partial_n k_{n-1}) \circ \partial_n$

$\xrightarrow{\text{C}_0 \text{ exact}} f_{n-1} \circ \partial_n - \partial_n - (f_{n-1} - \text{id} - k_{n-2} \circ \partial_{n-1}) \partial_n$

$= 0$ since $\partial \circ \partial = 0$ and f is chain map

$\Rightarrow \partial_n k_{n-1} = 0$ since $\partial \circ \partial = 0$ and f is chain map

we can pick basis elts c_n of C_n and pick such c_{n+1} , then define $k_n(c_n) := c_{n+1}$

and extend k_n linearly to get $k_n: C_n \rightarrow C_{n+1} \Rightarrow$ get required equation for n

Step 4 chain maps/homotopies on S_*, S_*^+ induce corresponding chain maps/homies on C_*^+, C_*^+

Pf If $q(z) = \sum \alpha_i z^i$, then define $(\psi(s))_I = \sum \alpha_i \text{sign}(\sigma) \cdot \sigma(I)_i$

(ψ hom on S_* or S_*^+) (ψ hom on C_*^+ or C_*^+ respectively)

Example $d = \sum (-1)^j \partial_j$ and for f of Step 2: $(\psi(s))_I = \sum \alpha_i \text{sign}(\sigma) \cdot \sigma(I)_i$ hence

Conclusion: $\psi: C_*^+ \rightarrow C_*^+$ chain map to id and surjects onto $C_*^+ \Rightarrow [f] = \text{id}: H_*^+ \rightarrow H_*^+$ equal $\square$

Cor $\bullet H_*^+$ is independent of choice of total ordering on set of indices (since $H_*^+ \cong H^*$)

$\bullet H_{\{U_i\}}^m(X, F) = 0$ for $m \geq N$ if $X = \bigcup U_i$ if finite cover with N sets (since $U_i \neq \emptyset$ in C_n^+ if $|I| \geq N$)

Example $X = \mathbb{P}_k^n$ with cover by $N=n+1$ affine sets $U_i \cong \mathbb{A}_k^n$ (Hw k 2)

8.3 Affines have no cohomology except H^0

(compare $H^*(\mathbb{A}^n) = 0$ for $n \geq 1$)
in algebraic topology

Theorem $X = \text{Spec } R$

$F \in \text{QCoh}(X)$

$X = \bigcup U_i$ finite affine open cover

Pf X separated (since affine) $\Rightarrow U_i$ all affine (Sec. 5.3, 8)

Easy case: minimal index I satisfies $U_I = X$

chain homotopy: $(h s)_I = \begin{cases} 0 & \text{if } i_0 = l \\ s_{\partial I} & \text{if } i_0 \neq l \end{cases}$

for I with $i_0 \neq l$

$(d(h s))_I = \sum (-1)^j (h s)_{I_j} = \sum (-1)^j s_{\partial I_j} \Rightarrow \text{id} = d h + h d$

$(h(d s))_I = (d s)_{\partial I} = s_I + \sum (-1)^{j+1} s_{\partial I_j}$

General case

$X = \text{Spec } R = \bigcup U_i, U_i = \text{Spec } R_i$

By easy case, know result for space U_ℓ with covering $\{U_\ell \cap U_i\}$ for minimal ℓ .

Ordering of indices does not affect H^* , so know result for $\mathcal{I}$ any ℓ by Cor of 8.2

$\Rightarrow$ Reduce to claim: if C^* exact when restrict to $U_i \forall i$, then C^* exact

$F \in \text{QCoh}(X), U_I$ affine say $\text{Spec } R_I \xrightarrow{\sim} F|_{U_I} \cong \widetilde{M_I}$ some R_I -module M_I

$C^n = \prod_{|I|=n} F(U_I, F) = \prod_{|I|=n} M_I$ finite product so $= \bigoplus$ (in particular, an R -mod)

$\Rightarrow C^0 \xrightarrow{d} C^1 \xrightarrow{d} C^2 \rightarrow \dots$ is a complex of R -mods

and by assumption of exactness on U_i have: $\leftarrow \text{using } F|_{U_i} = \widetilde{M_I}|_{U_i} \cong \widetilde{M_I \otimes R_i} \text{ by 6.8}$

$C^0 \otimes_R R_i \rightarrow C^1 \otimes_R R_i \rightarrow \dots$ exact $\forall i$

$\Rightarrow$ localising further by $\otimes_{R_i} (R_i)$ get exactness of localisation of C^* at each $p \in \text{Spec } R$.

$\Rightarrow$ by Sec. 3.0 deduce exactness of C^* . $\square$

8.4 Independence of cover

Theorem X separated, quasi-compact $\Rightarrow H^*(X, F)$ independent of choice of

Pf Will use ordered Čech cohomology.

X separated $\Rightarrow \bigcap_{\text{finite}} U_i$ is affine (Sec. 5.3, 8)

$X = \bigcup U_i, X = \bigcup_j \{U_i \cap U_j\}$ take mixed intersections: $C^{n,m} = \prod_{|I|=n} \prod_{|J|=m} F|_{U_I \cap U_J}$

$C^{n,0} \cong \prod_{|I|=n} \check{C}^n\{U_i \cap U_I\}$

$C^{0,m} \cong \prod_{|J|=m} \check{C}^m\{U_i \cap U_J\}$

$\Rightarrow$ rows & columns are exact except for degree 0:

$H^0(C^{n,0}) = \prod_{|I|=n} \Gamma(U_I, F) = \check{C}^n\{U_i\}(F)$

$H^0(C^{0,m}) = \prod_{|J|=m} \Gamma(U_J, F) = \check{C}^m\{U_i\}(F)$

$H^0(C^{n,m}) = \prod_{|I|=n} \prod_{|J|=m} \Gamma(U_I \cap U_J, F) = \check{C}^{n+m}\{U_i\}(F)$

$\Rightarrow$ rows & columns are exact except for degree 0:

$H^0(C^{n,0}) = \prod_{|I|=n} \Gamma(U_I, F) = \check{C}^n\{U_i\}(F)$

$H^0(C^{0,m}) = \prod_{|J|=m} \Gamma(U_J, F) = \check{C}^m\{U_i\}(F)$


$H^0(C^{n,m}) = \prod_{|I|=n} \prod_{|J|=m} \Gamma(U_I \cap U_J, F) = \check{C}^{n+m}\{U_i\}(F)$

Rmk $\mathcal{L}$ line bundle with transition maps α_{ij} } and $\mathcal{L} \otimes \mathcal{L}^{-1} \cong \mathcal{O}_X = \text{trivial line bundle}$
 $\Rightarrow \mathcal{L}^{-1}$ " " " α_{ij}^{-1}

FACT line bundles on A^n are always trivial
indeed vector bundles on A^n are always trivial $\leftarrow$ (Serre's Conjecture 1955) (Quillen-Suslin Theorem 1976)

EXAMPLE $\text{Pic}(P^1) \cong P^1 = A_0 \cup A_1$ $\leftarrow$ In C3.4 course: view $P^1 = k^2 / \sim$ k^n -rescaling
 $\text{Spec } k[t]$ $\cong \text{Spec } k[t^{-1}]$ Have homogeneous coordinates $[x_0 : x_1]$
and A_0 corresponds to $\{[1:t] : t \in A^1\}$ where $t = x_1/x_0$

$\mathcal{L}$ line bundle on $P^1_k \Rightarrow \mathcal{L}|_{A_i}$ trivial since $A_i \cong A^1$.
 $(\alpha_{10} : \mathcal{L}|_{A^1} \rightarrow \mathcal{L}|_{A_0}) \in k[t, t^{-1}]^* = \{a t^i : a \in k^*, i \in \mathbb{Z}\}$ $\leftarrow$ note: $A_0 \cap A_1 = \text{Spec } k[t, t^{-1}]$
 $\beta_0 \in k[t, t^{-1}]^* = k^*$, $\beta_1 \in k[t, t^{-1}]^* = k^*$ $\leftarrow$ exercise

$\Rightarrow$ $\text{Pic}(P^1) \cong \check{H}^1(P^1, \mathcal{O}_{P^1}^*) \cong \mathbb{Z}$ so define $\theta(i)$ by using A_1
 $\mathcal{O}(i) \hookrightarrow (\alpha_{10} = t^i) \hookrightarrow i$ 

Rmk $\mathcal{O}(0) = \mathcal{O}_{P^1}$ trivial line bdl.
Hwk 4 Ideal sheaf of a closed point in P^1_k is $\cong \mathcal{O}(-1)$ for disjoint union of n closed pts get $\cong \mathcal{O}(-n)$
for order n point $(t^i) \leq k[t]$ (i.e. closed subscheme $\text{Spec } k[t]/(t^n) \subseteq A_0 \subseteq P^1_k$) get $\mathcal{O}(-n)$.
Non-examinable Rmk (for differential geometry): i determines the Chern class $c_1(\mathcal{L}) = i = c_1(\mathcal{L})_{P^1}$
 $T P^1$ is $\mathcal{O}(2)$ since $2 = \chi(P^1) = \chi(S^2)$ and $c_1(T P^1) = \text{Euler class of } P^1$, and $T^* P^1 = \mathcal{O}(-2)$.
 $\mathcal{O}(-1) \rightarrow P^1$ is blow-up of $\mathbb{A}^2$ at 0 : the lines through 0 in k^2 are the fibres.

Theorem 1) $\check{H}^0(P^1, \mathcal{O}(i)) = \begin{cases} 0 & i < 0 \\ \{f \in k[t] : \deg f \leq i\} \cong k[x_0, x_1]_i & i \geq 0 \end{cases}$ $\leftarrow$ i-th graded part in x_0, x_1 of degree i
2) $\check{H}^1(P^1, \mathcal{O}(i)) = \begin{cases} 0 & i \geq -1 \\ k[t^{-1}] / k + t^i k[t^{-1}] \cong k[x_0, x_1]_{-i-2} & i < -1 \end{cases}$ $\leftarrow$ exercise
3) $\check{H}^n(P^1, \mathcal{O}(i)) = 0$ for $n \geq 2$

Pf By 8.6, since P^1 separated & quasi-compact enough to calculate $\check{H}^i_{\{A_0, A_1\}}(P^1, \mathcal{O}(i))$.
3) no triple overlaps or higher $\leftarrow$ $g \in \mathcal{O}_{A_1} \xrightarrow{\alpha_{10}} \mathcal{O}_{A_0} \ni f$ where α_{10} is defined on $A_0 \cap A_1$.

1) $\check{H}^0 = \Gamma : g(t^{-1}) \in k[t^{-1}]$ on A_1 , $f(t) \in k[t]$ on A_0 , on overlap: $f(t) = g(t^{-1}) = f(t) \in k[t, t^{-1}]$
 $\Rightarrow \deg f \leq i$ and g is determined by f from equation

2) $\mathcal{L} = \mathcal{O}(i)$ $\Gamma(A_0, \mathcal{L}) \oplus \Gamma(A_1, \mathcal{L}) \xrightarrow{d} \Gamma(A_0 \cap A_1, \mathcal{L}) \xrightarrow{d} 0$
 $\leftarrow$ strictly speaking $\Gamma(A_0 \cap A_1, \mathcal{L}) \cong k[t, t^{-1}]$ $\leftarrow$ $\mathcal{L}|_{A_0} \cong \mathcal{O}_{A_0}(i) \cong k[t]$ $\mathcal{L}|_{A_1} \cong \mathcal{O}_{A_1}(i) \cong k[t^{-1}]$
 $(f, g) \mapsto t^i \cdot g(t^{-1}) - f(t)$ 

$\check{H}^1 = k[t, t^{-1}] / k[t] + t^i k[t^{-1}]$ $\leftarrow$ need to transition from $g(t^{-1}) \in \mathcal{O}_{A_1}(A_{10})$ to $\mathcal{O}_{A_0}(A_{01})$ via $\alpha_{10} : \mathcal{O}_{A_0} \cong \mathcal{L}|_{A_0} = \mathcal{L}|_{A_{10}} \cong \mathcal{O}_{A_{01}}$
 $\leftarrow$ is all of $k[t, t^{-1}]$ if $i \geq -1$
 $\leftarrow$ does not contain $t^{-1}, t^{-2}, \dots, t^{-i-1}$ if $i < -1$

EXAMPLE: P^n

$X = P^n_k = A_0 \cup A_1 \cup \dots \cup A_n$ $A_i = \text{Spec } k[\frac{x_0}{x_i}, \dots, \frac{x_n}{x_i}]$ $\leftarrow$ omit x_i
 $\rightarrow \mathcal{O}(1) = \text{line bundle with } \alpha_{ij} = (\frac{x_j}{x_i}) : k[\frac{x_0}{x_i}, \dots, \frac{x_n}{x_i}] \rightarrow k[\frac{x_0}{x_j}, \dots, \frac{x_n}{x_j}]$ $\leftarrow$ P^1 case: $t = x_1/x_0$
 $\mathcal{O}(m) = \mathcal{O}(1)^{\otimes m}$ so $\alpha_{ij} = (\frac{x_j}{x_i})^m$ both equal to $\Gamma(A_i \cap A_j; \mathcal{O}_X)$ $\leftarrow$ it multiplies by $\frac{x_j}{x_i} = t^{-1}$ $\checkmark$
 $\leftarrow$ tensor m times

Rmk $\mathcal{O}(-1)$ called tautological line bundle because in C3.4 course each (closed) point of P^n is a 1-dim vector subspace $V \subseteq k^{n+1}$ ($P^n = k^{n+1} / \sim$ k^n -rescaling)
so get obvious line bundle: over the point $[V] \in P^n$ have the line V .

Hwk 4 $\text{Pic}(P^n) \cong \mathbb{Z}$ generated by the $\mathcal{O}(m)$
 $\Gamma(P^n, \mathcal{O}(m)) = \begin{cases} k[x_0, \dots, x_n]_m & \text{if } m \geq 0 \\ 0 & \text{if } m < 0 \end{cases}$ $\leftarrow$ so homogeneous polys of deg $= m$ so on A_i get polys of deg $\leq m$ in the variables $\frac{x_0}{x_i}, \dots, \frac{x_n}{x_i}$

8.8 Product on Čech cohomology

(Non-examinable section) $(X, \mathcal{O}_X)$ any ringed space
 $\check{H}^p_{\{U_i\}}(X, F) \times \check{H}^q_{\{U_i\}}(X, G) \rightarrow \check{H}^{p+q}_{\{U_i\}}(X, F \otimes_{\mathcal{O}_X} G)$
 $((s_1), (t_1)) \mapsto (s_1 \otimes t_1)$

Rmk In 8.6 where we took constant coefficients $F = G = \mathbb{Z}$ we recover the cup product on singular cohomology (respectively on de Rham cohomology)
 $\leftarrow$ note $\mathbb{Z} \otimes_{\mathcal{O}_X} \mathbb{Z} \cong \mathbb{Z}$
 $\leftarrow$ using $F = G = \mathbb{R}$
 $\mathcal{O}_X = \text{smooth real functions}$
 $\leftarrow$ so $\mathbb{R} \otimes_{\mathcal{O}_X} \mathbb{R} \cong \mathbb{R}$

10. Qcoh(P^n), GRADED MODULES, PROJ(R) (Non-examinable chapter)

10.1 Graded modules and QCoh(P^n)

Def graded ring means a ring R s.t.

$$R = R_0 \oplus R_1 \oplus R_2 \oplus \dots \text{ as abelian groups (so a graded abelian gp graded by } \mathbb{N})$$

$$R_i \cdot R_j \subseteq R_{i+j} \leftarrow \text{Rmk } R_0 \subseteq R \text{ subring since } R_0 \cdot R_0 \subseteq R_0$$

The elements of R_n are called homogeneous elements of degree n

Graded module means R -mod M s.t.

$$M = \dots \oplus M_{-2} \oplus M_{-1} \oplus M_0 \oplus M_1 \oplus M_2 \oplus \dots \text{ as abelian groups (so graded by } \mathbb{Z})$$

$R_i \cdot M_j \subseteq M_{i+j}$

$\leftarrow$ (often write M_0 to emphasize $\exists$ grading $\circ$)

A morphism of graded R -mods is R -mod hom $M \xrightarrow{\varphi} N$, with $\varphi(M_n) \subseteq N_n \quad \forall n$

From now on: $R = k[x_0, \dots, x_n]$ R_n = homogeneous polys of $\deg = n$ (so $R_0 = k$)

$$X = \mathbb{P}_k^n = A_0 \vee A_1 \vee \dots \vee A_n \text{ for}$$

$$A_i = \text{Spec } k \left[\frac{x_0}{x_i}, \dots, \frac{x_n}{x_i} \right] = \text{Spec} \left(k[x_0, \dots, x_n]_{(x_i)} \right)$$

$\leftarrow$ means take 0-th graded part
so $p(x_0, x_1, \dots, x_n) \mapsto \text{poly}$
 $x_i \cdot \deg(p)$

$$A_i \cap A_j = \text{Spec } k \left[\frac{x_0}{x_i}, \dots, \frac{x_n}{x_i}, \frac{x_0}{x_j}, \dots, \frac{x_n}{x_j} \right] = \text{Spec} \left(k[x_0, \dots, x_n]_{(x_i, x_j)} \right)$$

Claim $\exists$ exact, full & faithful functor

$$\{ \text{graded } R\text{-mods} \} \longrightarrow \text{QCoh}(\mathbb{P}^n)$$

$$M \longmapsto \tilde{M}$$

Pf Let $M_i = (M_{x_i})_{x_i=0}$ 0-th graded piece and $M_{ij} = (M_{x_i x_j})_{x_i x_j=0}$
Define $\tilde{M}|_{A_i} = \tilde{M}_i$ these glue since $\tilde{M}_i|_{A_i \cap A_j} \cong \tilde{M}_{ij} \cong \tilde{M}_j|_{A_i \cap A_j}$
Exactness is a local condition, so it holds since it holds in affine case.

$$\text{Full \& faithful: } \text{Hom}(\tilde{M}|_{A_i}, \tilde{N}|_{A_i}) = \text{Hom}(\tilde{M}_i, \tilde{N}_i) = \text{Hom}_{(R_{x_i})\text{-mods}}((M_{x_i})_0, (N_{x_i})_0)$$

This reduces the problem to an exercise in graded R -mods. (omitted here) $\square$

Warning Not an equivalence of categories because:

$$\text{HwK 4 if } M_n = N_n \text{ for } n \geq N \text{ then } \tilde{M} \cong \tilde{N}$$

$\leftarrow$ unlike case from 7.6:
 $R\text{-Mods} \cong \text{QCoh}(\text{Spec } R)$
 $M \mapsto \tilde{M}$
 $F(M) \hookrightarrow F$

Fact If work with graded R -mods "modulo" identifying those which would give rise to "same"

$\tilde{M}$, then get equivalence of categories. So work with $\{R\text{-mods } M\} / \{R\text{-mods } M : \tilde{M} = 0\}$.

For $X = \mathbb{P}^n$, $\tilde{M} = 0 \Leftrightarrow M$ is locally nilpotent, i.e. $\forall m \in M, \exists d$ s.t. $x_i^d \cdot m = 0 \quad \forall i$.

In reverse direction: $\{ \text{graded } R\text{-mods} \} \longleftarrow \text{QCoh}(\mathbb{P}^n)$

$$\left(\text{stands for } \left(\text{grading } \geq 0 \right) \right) \Gamma_+(F) := \bigoplus_{d \geq 0} \Gamma(\mathbb{P}^n, F(d)) \longleftarrow F \text{ where } F(d) = F \otimes_{\mathcal{O}_{\mathbb{P}^n}} \mathcal{O}(d) \leftarrow \text{called twisting}$$

$$\text{Fact } F \cong \Gamma_+(\tilde{F})$$

When we mod out by the M with $\tilde{M} = 0$ as in $\otimes$, this functor together with the functor of claim define an equivalence of cats.

$\text{Coh}(\mathbb{P}^n)$ corresponds to the f.g. graded modules under the equivalence.

Rmk The preferred representative of M in the quotient $\otimes$ is the saturation $\Gamma_+(\tilde{M})$ of M .

call M a saturated module if $M \cong \Gamma_+(\tilde{M})$. $\leftarrow$ (think of this like a sheafification)

Def $M[d]$ new graded R -mod with $M[d]_i = M_{d+i}$

Example $\mathcal{L} := \tilde{R}[d]$ on $\mathbb{P}^n \leftarrow$ (so $k[x_0, \dots, x_n][d]$)

$$\mathcal{L}(A_i) = (R[d]_{x_i}) = x_i^d \cdot k \left[\frac{x_0}{x_i}, \dots, \frac{x_n}{x_i} \right] = x_i^d \cdot (R_{x_i})_0$$

$\leftarrow$ exercise

$\leftarrow$ line bundle with $\sigma_{ij} = (x_i/x_j)^d$. Hence $\mathcal{L} = \mathcal{O}(d)$.

$\leftarrow$ $(\mathcal{O}_i)_{x_i} \cong \mathcal{L}|_{A_i} = \mathcal{L}|_{A_j} \cong \mathcal{O}_j|_{A_i} \cong \mathcal{O}_j \otimes_{\mathcal{O}_i} \mathcal{O}_i \cong \mathcal{O}_j$ $f \mapsto x_i^d \cdot f \mapsto x_j^{-d} \cdot x_i^d \cdot f$

Exercise $\tilde{M}[d] \cong \tilde{M}(d) (= \tilde{M} \otimes_{\mathcal{O}_{\mathbb{P}^n}} \mathcal{O}(d)) \leftarrow$ (e.g. $\tilde{R}[d] = \tilde{R}(d) = \mathcal{O}_{\mathbb{P}^n} \otimes (d) = \mathcal{O}(d)$)

Rmk $k[x_0, \dots, x_n] = \bigoplus_{d \geq 0} \Gamma(\mathbb{P}^n, \mathcal{O}(d))$ (but this does not generalise due to above issue about cats)

The construction of $\tilde{M}$ is so similar to the Spec R case of $\tilde{M}$, because $\exists$ analogue of Spec R : Proj R

10.2 Proj(R) and QCoh(Proj R)

$$\text{Proj}(R) = \{ \text{graded prime ideals } I \subseteq R \text{ not containing the irrelevant ideal} \}$$

$\leftarrow$ R any graded ring

$\leftarrow$ means $I = \bigoplus_{n \geq 0} (I_n R_n)$

$\leftarrow$ (generated by homogeneous elts)

$\leftarrow$ in $\mathbb{P}^n$ we remove the max ideal $(x_0, \dots, x_n)$ (irrelevant ideal) because don't allow the closed point $\{0 : \dots : 0\}$

$V(I) = \{ p \in \text{Proj } R : p \supseteq I \}$ define closed sets of Zariski topology

f homogeneous of degree $> 0 \Rightarrow D_f = \text{Proj } R \setminus V(f) = \{ p \in \text{Proj } R : f \notin p \}$ basis of open sets

Warning $\text{Proj } R = \bigcup D_i \Leftrightarrow R \subseteq \sqrt{\text{call } f_i} \leftarrow$ (example: $\mathbb{P}^n = D_{x_0} \cup \dots \cup D_{x_n}$ and $(x_0, \dots, x_n) = k[x_0, \dots, x_n] +$

Fact $D_f \cong \text{Spec}((R_f)_0)$ as topological spaces

$$p \mapsto p R_f \cap (R_f)_0 \quad (\text{inverse map: } p_0 \mapsto \bigoplus_{k \geq 0} \{ a_k \in R_k : \frac{a_k f^k}{f^{k+1}} \in p_0 \})$$

Sheaf $\mathcal{O} := \mathcal{O}_{\text{Proj}(R)} :$

$$\mathcal{O}|_{D_f} = \mathcal{O}_{\text{Spec}((R_f)_0)} \text{ then glue.} \leftarrow \text{(on } D_{f_1} = D_f \cap D_{g_1} \text{ get } \mathcal{O}_{\text{Spec}((R_{f_1 g_1})_0)})$$

Warning Proj is not functorial like Spec

If $\varphi: R \rightarrow S$ graded hom of rings, $\varphi(R_+) \not\subseteq S_+$ then get morph $\varphi^\#: \text{Proj } S \rightarrow \text{Proj } R$ but not all morphs arise in this way.

$I \mapsto \varphi^{-1}(I)$

Examples

1) $S = R[x_0, \dots, x_n]$ with usual grading $\Rightarrow \text{Proj } R = \mathbb{P}^n_R$ (or $\mathbb{P}^n_{\text{Spec } R}$)

2) $R^{(d)} := \bigoplus_{n \geq 0} R_{d+n}$ then the inclusion $R^{(d)} \hookrightarrow R$ induces an iso $\text{Proj } R \cong \text{Proj } R^{(d)}$

(recall $R_0 \hookrightarrow R$ subring)

3) S graded ring generated as an S_0 -algebra by $n+1$ elements $s_0, \dots, s_n \in S_1$

$\Rightarrow S_0[x_0, \dots, x_n] \xrightarrow{\varphi} S \Rightarrow S \cong S_0[x_0, \dots, x_n] \Rightarrow \text{Proj } S \cong \mathbb{V}(I) \subseteq \mathbb{P}^n_{S_0}$
closed subscheme

Example $k[x, y]^{(2)} = k[x^2, xy, y^2]$ $\xleftarrow{\text{Ker } \varphi}$ call this I

$k[X, Y, Z] \rightarrow k[x^2, xy, y^2], X \mapsto x^2, Y \mapsto xy, Z \mapsto y^2$

$\Rightarrow \mathbb{P}^1 = \text{Proj } k[x, y] \cong \text{Proj } k[x^2, xy, y^2] \cong \text{Proj } k[X, Y, Z]/(XZ - Y^2)$ closed subscheme of $\mathbb{P}^2$

is the Veronese embedding $\nu_2: \mathbb{P}^1 \hookrightarrow \mathbb{P}^2$. Similarly get $\nu_d: \mathbb{P}^n \hookrightarrow \mathbb{P}^N$ $N = \# \text{degree } d \text{ monomials in } x_0, \dots, x_n$

4) every closed subscheme of $\text{Proj } R$ arises as $\text{Proj}(R/I)$ some graded ideal I . so $N = \binom{n+d}{d}$

Fact $R = \bigoplus_{n \geq 0} R_n$ graded ring $\Rightarrow$ get line bundles $\mathcal{O}(d) = \widetilde{R}(d)$ on $\text{Proj } R$, and

$\exists$ exact, full & faithful functor

$$\begin{array}{ccc} \{\text{graded } R\text{-mods}\} & \rightarrow & \text{QCoh}(\text{Proj } R) \\ M & \mapsto & \widetilde{M} \\ \Gamma_*(F) & \longleftarrow & F \end{array}$$

Note: this tells us $\text{QCoh}(\cdot) \forall \text{Proj variety!}$

$\widetilde{M}$ built by gluing as in 10.1 namely

$\widetilde{M}(D_f) = M_f$ is homogeneous localization at f
(so localize at f and take 0-th graded part)
stalk $\widetilde{M}_I = M_{(I)}$ is homogeneous localization at the homogeneous prime ideal I
 $= 0$ -th graded part of M_I

where $\Gamma_d(F) := \Gamma(\text{Proj } R, F(d)) \leftarrow (F(d) = F \otimes_{\mathcal{O}_X} \mathcal{O}(d) \text{ and } \mathcal{O}_X = \widetilde{R} \text{ on } X = \text{Proj } R)$

again, not an equivalence of cats, but $\widetilde{\Gamma_*(F)} \cong F$ and the two functors define an equivalence of cats if we work with saturated graded R -mods $(M_* \cong \Gamma_*(\widetilde{M}))$

Fact If R_0 Noetherian, R generated as R_0 -algebra by finitely many elts $\in R_1 \Rightarrow$ Example: $R = k[x_0, \dots, x_n]/I$ then $x_0, \dots, x_n \in R_1$ generate.

then $\{f.g. R\text{-mods}\} / \{f.g. \text{"torsion"} R\text{-mods}\} \rightarrow \text{Coh}(\text{Proj } R)$ is equiv. of cats.

$M \mapsto \widetilde{M}$ and quasi-inverse $\Gamma_*(F) \leftarrow F$

Here "torsion" means $\forall m \in M, \exists n \in \mathbb{N}: (R_+)^n \cdot m = 0$. For M f.g. A -mod: this holds $\Leftrightarrow M_R = 0$ for large k

So $\otimes$ same as working with f.g. R -mods modulo identifying those that "agree" in large degrees.

Exercise M "torsion" $\Rightarrow M_f = 0 \forall$ homogeneous $f \in R_+ \Rightarrow \widetilde{M}(D_f) = M_{(f)} = 0 \Rightarrow \widetilde{M} = 0$. (homogeneous localization at f)

Now assume only R Noether graded ring.

Exercise Show R_0 Noeth, and R generated as R_0 -alg. by finitely many $f_1, \dots, f_n \in R_1$.

Let $d := \text{lcm}(\deg f_i)$. Call homogeneous $m \in M$ irrelevant if $(R_+ \cdot m)_{n \cdot d} = 0$ for all large N . M called irrelevant if all $\curvearrowright$ are irrelevant. Fact $\otimes$ holds if replace "torsion" by "irrelevant".