

Numerical Solution of Partial Differential Equations

Endre Süli

Mathematical Institute
University of Oxford
2022

Lecture 9

Stability of finite difference schemes

To replicate the stability property of the heat equation in the L_2 norm at the discrete level, we need a suitable notion of stability.

Stability of finite difference schemes

To replicate the stability property of the heat equation in the L_2 norm at the discrete level, we need a suitable notion of stability.

We shall say that a finite difference scheme for the unsteady heat equation is **(practically) stable in the ℓ_2 norm**, if

$$\|U^m\|_{\ell_2} \leq \|U^0\|_{\ell_2}, \quad m = 1, \dots, M,$$

where

$$\|U^m\|_{\ell_2} = \left(\Delta x \sum_{j=-\infty}^{\infty} |U_j^m|^2 \right)^{1/2}.$$

Stability of finite difference schemes

To replicate the stability property of the heat equation in the L_2 norm at the discrete level, we need a suitable notion of stability.

We shall say that a finite difference scheme for the unsteady heat equation is **(practically) stable in the ℓ_2 norm**, if

$$\|U^m\|_{\ell_2} \leq \|U^0\|_{\ell_2}, \quad m = 1, \dots, M,$$

where

$$\|U^m\|_{\ell_2} = \left(\Delta x \sum_{j=-\infty}^{\infty} |U_j^m|^2 \right)^{1/2}.$$

We shall use the semidiscrete Fourier transform to explore the stability of finite difference schemes.

Definition

The semidiscrete Fourier transform of a function U defined on the infinite mesh $x_j = j\Delta x$, $j = 0, \pm 1, \pm 2, \dots$, is:

$$\hat{U}(k) = \Delta x \sum_{j=-\infty}^{\infty} U_j e^{-ikx_j}, \quad k \in [-\pi/\Delta x, \pi/\Delta x].$$

Definition

The semidiscrete Fourier transform of a function U defined on the infinite mesh $x_j = j\Delta x$, $j = 0, \pm 1, \pm 2, \dots$, is:

$$\hat{U}(k) = \Delta x \sum_{j=-\infty}^{\infty} U_j e^{-ikx_j}, \quad k \in [-\pi/\Delta x, \pi/\Delta x].$$

We shall also need the inverse semidiscrete Fourier transform, as well the discrete counterpart of Parseval's identity that connect these transforms, similarly as in the case of the Fourier transform and its inverse considered earlier.

Definition

Let \hat{U} be defined on the interval $[-\pi/\Delta x, \pi/\Delta x]$. The inverse semidiscrete Fourier transform of \hat{U} is defined by

$$U_j := \frac{1}{2\pi} \int_{-\pi/\Delta x}^{\pi/\Delta x} \hat{U}(k) e^{ikj\Delta x} dk.$$

Definition

Let \hat{U} be defined on the interval $[-\pi/\Delta x, \pi/\Delta x]$. The inverse semidiscrete Fourier transform of \hat{U} is defined by

$$U_j := \frac{1}{2\pi} \int_{-\pi/\Delta x}^{\pi/\Delta x} \hat{U}(k) e^{ikj\Delta x} dk.$$

We then have the following result.

Lemma (Discrete Parseval's identity)

Let

$$\|U\|_{\ell_2} = \left(\Delta x \sum_{j=-\infty}^{\infty} |U_j|^2 \right)^{1/2} \quad \text{and} \quad \|\hat{U}\|_{L_2} = \left(\int_{-\pi/\Delta x}^{\pi/\Delta x} |\hat{U}(k)|^2 dk \right)^{1/2}.$$

If $\|U\|_{\ell_2}$ is finite, then also $\|\hat{U}\|_{L_2}$ is finite, and

$$\|U\|_{\ell_2} = \frac{1}{\sqrt{2\pi}} \|\hat{U}\|_{L_2}.$$

Lemma (Discrete Parseval's identity)

Let

$$\|U\|_{\ell_2} = \left(\Delta x \sum_{j=-\infty}^{\infty} |U_j|^2 \right)^{1/2} \quad \text{and} \quad \|\hat{U}\|_{L_2} = \left(\int_{-\pi/\Delta x}^{\pi/\Delta x} |\hat{U}(k)|^2 dk \right)^{1/2}.$$

If $\|U\|_{\ell_2}$ is finite, then also $\|\hat{U}\|_{L_2}$ is finite, and

$$\|U\|_{\ell_2} = \frac{1}{\sqrt{2\pi}} \|\hat{U}\|_{L_2}.$$

The proof of this is similar to that of Parseval's identity discussed earlier, and we shall therefore leave its proof as an exercise.

Stability analysis of the explicit Euler scheme

By inserting

$$U_j^m = \frac{1}{2\pi} \int_{-\pi/\Delta x}^{\pi/\Delta x} e^{ikj\Delta x} \hat{U}^m(k) dk$$

into the Euler scheme we deduce that

$$\begin{aligned} & \frac{1}{2\pi} \int_{-\pi/\Delta x}^{\pi/\Delta x} e^{ikj\Delta x} \frac{\hat{U}^{m+1}(k) - \hat{U}^m(k)}{\Delta t} dk \\ &= \frac{1}{2\pi} \int_{-\pi/\Delta x}^{\pi/\Delta x} \frac{e^{ik(j+1)\Delta x} - 2e^{ikj\Delta x} + e^{ik(j-1)\Delta x}}{(\Delta x)^2} \hat{U}^m(k) dk. \end{aligned}$$

Stability analysis of the explicit Euler scheme

By inserting

$$U_j^m = \frac{1}{2\pi} \int_{-\pi/\Delta x}^{\pi/\Delta x} e^{ikj\Delta x} \hat{U}^m(k) dk$$

into the Euler scheme we deduce that

$$\begin{aligned} & \frac{1}{2\pi} \int_{-\pi/\Delta x}^{\pi/\Delta x} e^{ikj\Delta x} \frac{\hat{U}^{m+1}(k) - \hat{U}^m(k)}{\Delta t} dk \\ &= \frac{1}{2\pi} \int_{-\pi/\Delta x}^{\pi/\Delta x} \frac{e^{ik(j+1)\Delta x} - 2e^{ikj\Delta x} + e^{ik(j-1)\Delta x}}{(\Delta x)^2} \hat{U}^m(k) dk. \end{aligned}$$

Therefore, we have that

$$\begin{aligned} & \frac{1}{2\pi} \int_{-\pi/\Delta x}^{\pi/\Delta x} e^{ikj\Delta x} \frac{\hat{U}^{m+1}(k) - \hat{U}^m(k)}{\Delta t} dk \\ &= \frac{1}{2\pi} \int_{-\pi/\Delta x}^{\pi/\Delta x} e^{ikj\Delta x} \frac{e^{ik\Delta x} - 2 + e^{-ik\Delta x}}{(\Delta x)^2} \hat{U}^m(k) dk. \end{aligned}$$

By comparing the left-hand side with the right-hand side we get

$$\hat{U}^{m+1}(k) = \hat{U}^m(k) + \mu(e^{ik\Delta x} - 2 + e^{-ik\Delta x})\hat{U}^m(k)$$

for all **wave numbers** $k \in [-\pi/\Delta x, \pi/\Delta x]$.

¹After: Richard Courant, Kurt Friedrichs, and Hans Lewy (*Über die partiellen Differenzgleichungen der mathematischen Physik*. *Mathematische Annalen*, 100:32–74, 1928).

By comparing the left-hand side with the right-hand side we get

$$\hat{U}^{m+1}(k) = \hat{U}^m(k) + \mu(e^{ik\Delta x} - 2 + e^{-ik\Delta x})\hat{U}^m(k)$$

for all **wave numbers** $k \in [-\pi/\Delta x, \pi/\Delta x]$. Thus we have

$$\hat{U}^{m+1}(k) = \lambda(k)\hat{U}^m(k),$$

where

$$\lambda(k) = 1 + \mu(e^{ik\Delta x} - 2 + e^{-ik\Delta x})$$

is the **amplification factor** and

$$\mu := \frac{\Delta t}{(\Delta x)^2}$$

is called the **CFL number**¹.

¹After: Richard Courant, Kurt Friedrichs, and Hans Lewy (*Über die partiellen Differenzgleichungen der mathematischen Physik*. *Mathematische Annalen*, 100:32–74, 1928).

By the discrete Parseval identity stated in Lemma 3 we have that

$$\begin{aligned}\|U^{m+1}\|_{\ell_2} &= \frac{1}{\sqrt{2\pi}} \|\hat{U}^{m+1}\|_{L_2} \\ &= \frac{1}{\sqrt{2\pi}} \|\lambda \hat{U}^m\|_{L_2} \\ &\leq \frac{1}{\sqrt{2\pi}} \max_k |\lambda(k)| \|\hat{U}^m\|_{L_2} \\ &= \max_k |\lambda(k)| \|U^m\|_{\ell_2}.\end{aligned}$$

By the discrete Parseval identity stated in Lemma 3 we have that

$$\begin{aligned}\|U^{m+1}\|_{\ell_2} &= \frac{1}{\sqrt{2\pi}} \|\hat{U}^{m+1}\|_{L_2} \\ &= \frac{1}{\sqrt{2\pi}} \|\lambda \hat{U}^m\|_{L_2} \\ &\leq \frac{1}{\sqrt{2\pi}} \max_k |\lambda(k)| \|\hat{U}^m\|_{L_2} \\ &= \max_k |\lambda(k)| \|U^m\|_{\ell_2}.\end{aligned}$$

In order to mimic the L_2 norm bound, we would like to ensure that

$$\|U^{m+1}\|_{\ell_2} \leq \|U^m\|_{\ell_2}, \quad m = 0, 1, \dots, M-1.$$

By the discrete Parseval identity stated in Lemma 3 we have that

$$\begin{aligned}\|U^{m+1}\|_{\ell_2} &= \frac{1}{\sqrt{2\pi}} \|\hat{U}^{m+1}\|_{L_2} \\ &= \frac{1}{\sqrt{2\pi}} \|\lambda \hat{U}^m\|_{L_2} \\ &\leq \frac{1}{\sqrt{2\pi}} \max_k |\lambda(k)| \|\hat{U}^m\|_{L_2} \\ &= \max_k |\lambda(k)| \|U^m\|_{\ell_2}.\end{aligned}$$

In order to mimic the L_2 norm bound, we would like to ensure that

$$\|U^{m+1}\|_{\ell_2} \leq \|U^m\|_{\ell_2}, \quad m = 0, 1, \dots, M-1.$$

Thus we demand that

$$\max_k |\lambda(k)| \leq 1,$$

i.e., that

$$\max_k |1 + \mu(e^{ik\Delta x} - 2 + e^{-ik\Delta x})| \leq 1.$$

Using Euler's formula

$$e^{i\varphi} = \cos \varphi + i \sin \varphi$$

and the trigonometric identity

$$1 - \cos \varphi = 2 \sin^2 \frac{\varphi}{2}$$

we can restate this as follows:

$$\max_k \left| 1 - 4\mu \sin^2 \left(\frac{k\Delta x}{2} \right) \right| \leq 1.$$

Using Euler's formula

$$e^{i\varphi} = \cos \varphi + i \sin \varphi$$

and the trigonometric identity

$$1 - \cos \varphi = 2 \sin^2 \frac{\varphi}{2}$$

we can restate this as follows:

$$\max_k \left| 1 - 4\mu \sin^2 \left(\frac{k\Delta x}{2} \right) \right| \leq 1.$$

Equivalently, we need to ensure that

$$-1 \leq 1 - 4\mu \sin^2 \left(\frac{k\Delta x}{2} \right) \leq 1 \quad \forall k \in [-\pi/\Delta x, \pi/\Delta x].$$

Using Euler's formula

$$e^{i\varphi} = \cos \varphi + i \sin \varphi$$

and the trigonometric identity

$$1 - \cos \varphi = 2 \sin^2 \frac{\varphi}{2}$$

we can restate this as follows:

$$\max_k \left| 1 - 4\mu \sin^2 \left(\frac{k\Delta x}{2} \right) \right| \leq 1.$$

Equivalently, we need to ensure that

$$-1 \leq 1 - 4\mu \sin^2 \left(\frac{k\Delta x}{2} \right) \leq 1 \quad \forall k \in [-\pi/\Delta x, \pi/\Delta x].$$

This holds if, and only if, $\mu = \frac{\Delta t}{(\Delta x)^2} \leq \frac{1}{2}$.

Thus we have shown the following result.

Thus we have shown the following result.

Theorem

Suppose that U_j^m is the solution of the explicit Euler scheme

$$\frac{U_j^{m+1} - U_j^m}{\Delta t} = \frac{U_{j+1}^m - 2U_j^m + U_{j-1}^m}{(\Delta x)^2}, \quad j = 0, \pm 1, \pm 2, \dots,$$

$$U_j^0 = u_0(x_j), \quad j = 0, \pm 1, \pm 2, \dots,$$

and $\mu = \frac{\Delta t}{(\Delta x)^2} \leq \frac{1}{2}$. Then,

$$\|U^m\|_{\ell_2} \leq \|U^0\|_{\ell_2}, \quad m = 1, 2, \dots, M. \quad (1)$$

Thus we have shown the following result.

Theorem

Suppose that U_j^m is the solution of the explicit Euler scheme

$$\frac{U_j^{m+1} - U_j^m}{\Delta t} = \frac{U_{j+1}^m - 2U_j^m + U_{j-1}^m}{(\Delta x)^2}, \quad j = 0, \pm 1, \pm 2, \dots,$$

$$U_j^0 = u_0(x_j), \quad j = 0, \pm 1, \pm 2, \dots,$$

and $\mu = \frac{\Delta t}{(\Delta x)^2} \leq \frac{1}{2}$. Then,

$$\|U^m\|_{\ell_2} \leq \|U^0\|_{\ell_2}, \quad m = 1, 2, \dots, M. \quad (1)$$

Hence, the explicit Euler scheme is **conditionally practically stable**, the condition for stability being that $\mu = \Delta t / \Delta x^2 \leq 1/2$. One can also show that if $\mu > 1/2$, then (1) will fail.

Stability analysis of the implicit Euler scheme

We shall now perform a similar analysis for the **implicit Euler scheme** for the heat equation:

$$\frac{U_j^{m+1} - U_j^m}{\Delta t} = \frac{U_{j+1}^{m+1} - 2U_j^{m+1} + U_{j-1}^{m+1}}{(\Delta x)^2}, \quad j = 0, \pm 1, \pm 2, \dots$$

$$U_j^0 = u_0(x_j), \quad j = 0, \pm 1, \pm 2, \dots$$

Stability analysis of the implicit Euler scheme

We shall now perform a similar analysis for the **implicit Euler scheme** for the heat equation:

$$\frac{U_j^{m+1} - U_j^m}{\Delta t} = \frac{U_{j+1}^{m+1} - 2U_j^{m+1} + U_{j-1}^{m+1}}{(\Delta x)^2}, \quad j = 0, \pm 1, \pm 2, \dots$$

$$U_j^0 = u_0(x_j), \quad j = 0, \pm 1, \pm 2, \dots$$

Equivalently,

$$U_j^{m+1} - \mu(U_{j+1}^{m+1} - 2U_j^{m+1} + U_{j-1}^{m+1}) = U_j^m$$

$$U_j^0 = u_0(x_j), \quad j = 0, \pm 1, \pm 2, \dots,$$

where, again,

$$\mu = \frac{\Delta t}{(\Delta x)^2}.$$

Using an identical argument as for the explicit Euler scheme, we find that the amplification factor is now

$$\lambda(k) = \frac{1}{1 + 4\mu \sin^2\left(\frac{k\Delta x}{2}\right)}.$$

Using an identical argument as for the explicit Euler scheme, we find that the amplification factor is now

$$\lambda(k) = \frac{1}{1 + 4\mu \sin^2\left(\frac{k\Delta x}{2}\right)}.$$

Clearly,

$$\max_k |\lambda(k)| \leq 1$$

for all values of

$$\mu = \frac{\Delta t}{(\Delta x)^2}.$$

Using an identical argument as for the explicit Euler scheme, we find that the amplification factor is now

$$\lambda(k) = \frac{1}{1 + 4\mu \sin^2\left(\frac{k\Delta x}{2}\right)}.$$

Clearly,

$$\max_k |\lambda(k)| \leq 1$$

for all values of

$$\mu = \frac{\Delta t}{(\Delta x)^2}.$$

Thus we have the following result.

Theorem

Suppose that U_j^m is the solution of the implicit Euler scheme

$$\frac{U_j^{m+1} - U_j^m}{\Delta t} = \frac{U_{j+1}^{m+1} - 2U_j^{m+1} + U_{j-1}^{m+1}}{(\Delta x)^2}, \quad j = 0, \pm 1, \pm 2, \dots,$$

$$U_j^0 = u_0(x_j), \quad j = 0, \pm 1, \pm 2, \dots$$

Then, for all $\Delta t > 0$ and $\Delta x > 0$,

$$\|U^m\|_{\ell_2} \leq \|U^0\|_{\ell_2}, \quad m = 1, 2, \dots, M. \quad (2)$$

Theorem

Suppose that U_j^m is the solution of the implicit Euler scheme

$$\frac{U_j^{m+1} - U_j^m}{\Delta t} = \frac{U_{j+1}^{m+1} - 2U_j^{m+1} + U_{j-1}^{m+1}}{(\Delta x)^2}, \quad j = 0, \pm 1, \pm 2, \dots,$$

$$U_j^0 = u_0(x_j), \quad j = 0, \pm 1, \pm 2, \dots$$

Then, for all $\Delta t > 0$ and $\Delta x > 0$,

$$\|U^m\|_{\ell_2} \leq \|U^0\|_{\ell_2}, \quad m = 1, 2, \dots, M. \quad (2)$$

Thus, the implicit Euler scheme is **unconditionally practically stable**, meaning that the bound (2) holds without any restrictions on Δx and Δt .

Stability analysis of the θ -scheme

Consider the θ -scheme:

$$\frac{U_j^{m+1} - U_j^m}{\Delta t} = (1 - \theta) \frac{U_{j+1}^m - 2U_j^m + U_{j-1}^m}{(\Delta x)^2} + \theta \frac{U_{j+1}^{m+1} - 2U_j^{m+1} + U_{j-1}^{m+1}}{(\Delta x)^2},$$

$$U_j^0 = u_0(x_j), \quad j = 0, \pm 1, \pm 2, \dots,$$

where $\theta \in [0, 1]$ is a parameter.

Stability analysis of the θ -scheme

Consider the θ -scheme:

$$\frac{U_j^{m+1} - U_j^m}{\Delta t} = (1 - \theta) \frac{U_{j+1}^m - 2U_j^m + U_{j-1}^m}{(\Delta x)^2} + \theta \frac{U_{j+1}^{m+1} - 2U_j^{m+1} + U_{j-1}^{m+1}}{(\Delta x)^2},$$

$$U_j^0 = u_0(x_j), \quad j = 0, \pm 1, \pm 2, \dots,$$

where $\theta \in [0, 1]$ is a parameter.

For $\theta = 0$ it is the explicit Euler scheme, for $\theta = 1$ it is the implicit Euler scheme, and for $\theta = 1/2$ it is the arithmetic average of the two Euler schemes, and is called the **Crank–Nicolson scheme**.

Using an identical argument as in the case of the two Euler methods, we find that

$$\lambda(k) - 1 = -4(1 - \theta) \mu \sin^2 \left(\frac{k\Delta x}{2} \right) - 4\theta \mu \lambda(k) \sin^2 \left(\frac{k\Delta x}{2} \right).$$

Using an identical argument as in the case of the two Euler methods, we find that

$$\lambda(k) - 1 = -4(1 - \theta)\mu \sin^2\left(\frac{k\Delta x}{2}\right) - 4\theta\mu \lambda(k) \sin^2\left(\frac{k\Delta x}{2}\right).$$

Therefore,

$$\lambda(k) = \frac{1 - 4(1 - \theta)\mu \sin^2\left(\frac{k\Delta x}{2}\right)}{1 + 4\theta\mu \sin^2\left(\frac{k\Delta x}{2}\right)}.$$

For practical stability, we demand that

$$|\lambda(k)| \leq 1 \quad \forall k \in [-\pi/\Delta x, \pi/\Delta x],$$

which holds if, and only if,

$$2(1 - 2\theta)\mu \leq 1.$$

For practical stability, we demand that

$$|\lambda(k)| \leq 1 \quad \forall k \in [-\pi/\Delta x, \pi/\Delta x],$$

which holds if, and only if,

$$2(1 - 2\theta)\mu \leq 1.$$

Thus we have shown that:

For practical stability, we demand that

$$|\lambda(k)| \leq 1 \quad \forall k \in [-\pi/\Delta x, \pi/\Delta x],$$

which holds if, and only if,

$$2(1 - 2\theta)\mu \leq 1.$$

Thus we have shown that:

- For $\theta \in [1/2, 1]$ the θ -scheme is **unconditionally practically stable**;
- For $\theta \in [0, 1/2)$ the θ -scheme is **conditionally practically stable**, the stability condition being that

$$\mu \leq \frac{1}{2(1 - 2\theta)}.$$