Problem Sheet 4

Distributions

- 1. <u>Scalar property of delta function</u>. Show from its interpretation as an integral that $\delta(ax) = \frac{1}{|a|}\delta(x)$ for any constant a.
- 2. Convergence of distributions. Show that the following function sequences converge to the δ -distribution as $n \to \infty$.

(a)
$$f_n(x) = \begin{cases} n/2 & \text{for } -1/n < x < 1/n, \\ 0 & \text{else} \end{cases}$$

(b)
$$f_n(x) = \frac{e^{-nx^2/4}}{\sqrt{4\pi/n}}$$
 (using $\int_{-\infty}^{\infty} e^{-x^2} dx = \sqrt{\pi}$ without proof)

You have to show that $\lim_{n\to\infty} \langle f_n, \phi \rangle = \langle \delta, \phi \rangle$ holds for all $\phi \in C_0^{\infty}(\mathbb{R})$, i.e. that $\lim_{n\to\infty} \int_{-\infty}^{\infty} f_n(x)\phi(x)dx = \phi(0)$

Hints: In (a), split the integral as

$$\int_{-\infty}^{\infty} f_n(x)\phi(x)dx = \int_{-\infty}^{\infty} f_n(x) \left[\phi(x) - \phi(0)\right] dx + \phi(0) \int_{-\infty}^{\infty} f_n(x)dx.$$

The second integral can be evaluated explicitly. What can you say about the first integral? A similar approach for (b) though a bit more work. In my model solution I split the integral into 4 parts, including a term

$$\int_{-\rho}^{\rho} f_n(x) \left[\phi(x) - \phi(0)\right] dx$$

with ρ chosen appropriately.

- 3. <u>Derivatives of distributions.</u>
 - (a) For $n \ge 0$, $a \in \mathbb{R}$, let $D_n : C_0^{\infty} \to \mathbb{R}$ be defined by

$$\langle D_n, \phi \rangle = \left. \frac{d^n \phi}{dx^n} \right|_{x=a} = \phi^{(n)}(a) \quad \text{for all } \phi \in C_0^\infty(\mathbb{R}).$$

Show that D_n is a distribution, and then, by induction, that

$$D_n = (-1)^n \delta^{(n)}(x-a),$$

where $\delta^{(n)}$ is the n-th distributional derivative of the δ -distribution.

(b) Let $T : C_0^{\infty} \to \mathbb{R}$ be a distribution (not necessarily one that is induced by a continuous function). Using the definition of distributional derivative, translation of distributions and convergence of distributions, show that

$$\lim_{\alpha \to 0} \frac{T(x+\alpha) - T(x)}{\alpha} = T'(x)$$

You may use (without proof) that for any $\phi \in C_0^{\infty}(\mathbb{R})$,

$$\rho_{\alpha}(x) \equiv \frac{\phi(x) - \phi(x - \alpha)}{\alpha}$$

converges to $\phi'(x)$ uniformly in x as $\alpha \to 0$.