

Infinite Groups

Cornelia Druțu

University of Oxford

Part C course MT 2023, Oxford

Inspirational quotation about Mathematics

William Thurston: “As one reads mathematics, one needs to have an active mind, asking questions, forming mental connections between the current topic and other ideas from other contexts, **so as to develop a sense of the structure, not just familiarity with a particular tour through the structure.**”

Burnside Theorem and applications

Theorem (Burnside's Theorem)

If $A \subset \text{End}(V)$ is a subalgebra which acts absolutely irreducibly on a finite-dimensional vector space V , then $A = \text{End}(V)$. In particular, if $G \leq \text{End}(V)$ is a subsemigroup acting irreducibly, then G spans $\text{End}(V)$ as a vector space, i.e. $\mathbb{K}[G] = \text{End}(V)$.

Theorem

Suppose that $G \leq \text{GL}_n(\mathbb{K})$ is irreducible and that

$$|\{\text{tr}(g) \mid g \in G\}| = q < \infty.$$

Then $|G| \leq q^{n^2}$.

Second application to Burnside. Nilpotent and unipotent.

Corollary

Suppose that $G \leq GL_n(\mathbb{K})$ is completely reducible and that $g^k = 1, \forall g \in G$. Then $|G| \leq k^{n^3}$.

Proof. See Ex. Sheet 4. □

$h \in \text{End}(V)$ is **nilpotent** if $h^k = 0$ for some $k > 0$

\Leftrightarrow in some basis, h can be written as an upper triangular matrix with zeroes on the diagonal.

\Leftrightarrow the only eigenvalue of h is 0.

$g \in \text{End}(V)$ is **unipotent** if $g = \text{id} + h$, where h is nilpotent

\Leftrightarrow the only eigenvalue of g is 1.

$g \in \text{End}(V)$ is **quasi-unipotent** if g^k is unipotent for some $k > 0$

\Leftrightarrow all the eigenvalues of g are roots of unity.

A subgroup $G < GL(V)$ is **unipotent** (respectively **quasi-unipotent**) if every element of G is unipotent (respectively, quasi-unipotent).

Kolchin's theorem

Theorem (Kolchin's theorem)

Suppose that $\mathbb{K} = \bar{\mathbb{K}}$ and $G < GL(V)$ is a unipotent subgroup. Then, for an appropriate choice of basis, G is isomorphic to a subgroup of the group of invertible upper-triangular matrices $\mathcal{T}_n(\mathbb{K})$. In particular G is nilpotent.

Proof. The conclusion is equivalent to the statement that G preserves a full flag

$$0 \subset V_1 \subset \dots \subset V_{n-1} \subset V,$$

where $i = \dim(V_i)$ for each i .

The proof is by induction on the dimension n of V .

Clear for $n = 1$.

We assume that $n > 1$ and that the statement is true for dimensions $< n$.

Proof of Kolchin's theorem continued

Suppose first that the action of G on V is reducible: G preserves a proper subspace $V' \subset V$.

We obtain two induced actions of G on V' (by restriction) and on $V'' = V/V'$ (by projection).

Both actions preserve full flags in V' , V'' (induction assumption), and the combination of these flags yields a full G -invariant flag in V .

Assume now that the action of G on V is irreducible.

For $g \in G$ arbitrary the endomorphism $g' = g - I$ is nilpotent, hence it has zero trace.

Therefore, for every $x \in G$, we have

$$\operatorname{tr}(g'x) = \operatorname{tr}(gx - x) = \operatorname{tr}(I) - \operatorname{tr}(I) = 0.$$

By Burnside's theorem, G spans $\operatorname{End}(V) \Rightarrow$ for each $x \in \operatorname{End}(V)$ and each $g \in G$, $\operatorname{tr}(g'x) = 0$.

$\tau : \operatorname{End}(V) \times \operatorname{End}(V) \rightarrow \mathbb{K}$ is nondegenerate $\Rightarrow g' = 0$ for all $g \in G$, i.e. $G = \{1\}$. □

Variation of Kolchin's theorem

Theorem (Variation of Kolchin's theorem)

Suppose $\mathbb{K} = \bar{\mathbb{K}}$, $G < GL(V)$ *quasiunipotent* and, moreover, there exists an upper bound α on the orders of eigenvalues of elements $g \in G$.

Then G contains a finite index subgroup isomorphic to a subgroup of the group $\mathcal{U}_n(\mathbb{K})$ of upper triangular matrices with 1 on the diagonal. *The index depends only on V and on α . In particular G is virtually nilpotent.*

Proof. By induction on the dimension of V .

As before, it suffices to consider the case when G acts irreducibly on V .

The orders of the eigenvalues of elements of G are uniformly bounded \Rightarrow the set of traces of elements of G is a certain finite set $C \subset \mathbb{K}$ of cardinality $q = q(\alpha)$.

Our first application to Burnside $\Rightarrow G$ is finite, of cardinality at most q^{n^2} , where $n = \dim V$. □

Solvable linear groups

Theorem (Lie-Kolchin-Mal'cev Theorem)

Let $G \leq GL(V)$ be solvable linear, with $n = \dim V$ (G solvable linear of degree n). Then G has a triangularizable normal subgroup K of finite index at most $\mu(n)$, a number that depends only on n .

Definition

Let \mathcal{X} and \mathcal{Y} be two classes of groups.

A group G is \mathcal{X} -by- \mathcal{Y} if there exists a short exact sequence

$$\{1\} \longrightarrow N \xrightarrow{i} G \xrightarrow{\pi} Q \longrightarrow \{1\},$$

such that $N \in \mathcal{X}$ and $Q \in \mathcal{Y}$.

Solvable linear groups versus nilpotent and polycyclic

Corollary

Let G be a solvable linear group of degree n .

- (i) G is virtually unipotent-by-abelian;
- (ii) (the *Zassenhaus Theorem*) the derived length of G is at most $\beta(n) := n + \log_2 \mu(n)$.

This can be combined with the following general result.

Theorem

Every nilpotent subgroup of $GL(n, \mathbb{Z})$ is finitely generated.

The two previous results imply the following.

Corollary

Every finitely generated solvable group linear over \mathbb{Z} is polycyclic.

Solvable groups linear over \mathbb{Z}

Proof. Such a group G has a finite index subgroup that is unipotent-by-abelian, hence a finite index normal subgroup N that is polycyclic.

The quotient G/N is solvable and finite, hence noetherian, hence polycyclic. Hence G is polycyclic. □

The converse is also true.

Theorem (Auslander's Theorem)

Every polycyclic group is linear over \mathbb{Z} .

Corollary

Every polycyclic group is virtually (finitely generated nilpotent)-by-(f.g. abelian).

Another comparison between solvable and polycyclic

We have thus obtained the following way of distinguishing polycyclic groups among solvable groups.

Theorem

Given a f.g. solvable group G , the following are equivalent:

- *G is polycyclic;*
- *G is linear over \mathbb{Z} .*