

Geometric Group Theory

Cornelia Druțu

University of Oxford

Part C course HT 2024

HNN extensions

Definition

Suppose we have $A \subseteq G$ and $\theta : A \rightarrow G$ an **injective homomorphism**. The **HNN extension of G on A with respect to θ** is

$$\begin{aligned} G *_A &:= \langle G, t \mid tat^{-1} = \theta(a), \forall a \in A \rangle \\ &= G * \langle t \rangle / \langle\langle \{tat^{-1}\theta(a)^{-1} : a \in A\} \rangle\rangle \end{aligned}$$

t is called the **stable letter** of the HNN extension.

The name comes from **Graham Higman, Bernhard Neumann and Hanna Neumann**.

A definition with a **Universal Property** can be formulated using

$$G *_A = \langle G, t \mid tat^{-1} = \theta(a), \forall a \in A \rangle.$$

HNN extensions

Definition

Let $A \subseteq G$ and $\theta : A \rightarrow G$ be **injective homomorphism**. The **HNN extension** of G on A with respect to θ is

$$G *_A := \langle G, t \mid tat^{-1} = \theta(a), \forall a \in A \rangle$$

Remark

If $G = \langle S \mid R \rangle$ then $G *_A = \langle S \cup \{t\} \mid R \cup \{tat^{-1} = \theta(a) : a \in A\} \rangle$.

Examples

- The **Baumslag-Solitar groups** $BS(m, n) = \langle a, t \mid ta^m t^{-1} = a^n \rangle$, where $m, n \in \mathbb{Z}$.
- When $m = n = 1$, we have \mathbb{Z}^2 (fundamental group of the **torus**.)
- When $m = 1, n = -1$, the fundamental group of the **Klein bottle**.
- When $m = 1$ (or $n = 1$) the group is **solvable**.

Reduced words of HNN extensions

Suppose A_1 is a set of right coset representatives for A and A_2 is a set of right coset representatives for $\theta(A)$ such that $1 \in A_1 \cap A_2$.

A **reduced word** of $G*_A$ is a sequence $(g_0, t^{\epsilon_1}, g_1, t^{\epsilon_2}, g_2, \dots, t^{\epsilon_n}, g_n)$ such that

- $\epsilon_i = \pm 1$
- $g_0 \in G$
- $g_i \in A_1$ if $\epsilon_i = 1$, $g_i \in A_2$ if $\epsilon_i = -1$
- $g_i \neq 1$ if $\epsilon_{i+1} = -\epsilon_i$

A **reduced element** of $G*_A$ is an element of the form $g_0 t^{\epsilon_1} g_1 \dots t^{\epsilon_n} g_n$.

Theorem

*Each $g \in G*_A$ is represented by a unique reduced word.*

Reduced words of HNN extensions

Theorem

*Each $g \in G *_A$ is represented by a unique reduced word.*

Proof:

Existence of a representation: We induct on the length of g as a reduced word in $G \cup \{t, t^{-1}\}$. The length 1 case is obvious.

Assume true for n . Length $n + 1$ means either $g = ut^{\pm 1}$, $\text{length}(u) \leq n$, or

$$g \in \{wth, wt^{-1}h\}$$

where $\text{length}(w) \leq n - 1$ and $h \in G$. If $g = ut^{\pm 1}$, apply induction. If

$$g = wth = wtah_1 = wtat^{-1}th_1 = w\theta(a)th_1$$

then $\text{length}(w\theta(a)) \leq n$ so we can apply the inductive assumption. The case $g = wt^{-1}h$ is similar.

Reduced words of HNN extensions

Uniqueness of representation: Let X be the set of reduced words. $G *_A$ acts on it (i.e. there exists a group homomorphism $G *_A \rightarrow \text{Bij}(X)$) as follows:

$$\phi(g)(g_0, t^{\epsilon_1}, g_1, \dots, t^{\epsilon_n}, g_n) = (gg_0, t^{\epsilon_1}, g_1, \dots, t^{\epsilon_n}, g_n)$$

and $\phi(t)(g_0, t^{\epsilon_1}, g_1, \dots, t^{\epsilon_n}, g_n)$ equals

$$\begin{cases} (\theta(g_0), t, 1, t^{\epsilon_1}, \dots, t^{\epsilon_n}, g_n) & \text{if } g_0 \in A \text{ and } \epsilon_1 = 1 \\ (\theta(g_0)g_1, t^{\epsilon_2}, \dots, t^{\epsilon_n}, g_n) & \text{if } g_0 \in A \text{ and } \epsilon_1 = -1 \\ (\theta(a), t, g'_0, t^{\epsilon_1}, \dots, t^{\epsilon_n}, g_n) & \text{if } g_0 = ag'_0 \text{ and } g'_0 \in A_1 \setminus \{1\} \end{cases}$$

Exercise: Prove that $\phi(t)$ is a bijection. For instance, prove that $\phi(t^{-1})$ is its inverse.

We thus have a homomorphism $\phi : G * \langle t \rangle \rightarrow \text{Bij}(X)$.

Exercise: prove that $\phi(tat^{-1}) = \phi(\theta(a))$, $\forall a \in A$.

Hence ϕ defines $\bar{\phi} : G *_A \rightarrow \text{Bij}(X)$. And if $g = g_0 t^{\epsilon_1} g_1 \dots t^{\epsilon_n} g_n$ then $\phi(g)(1) = (g_0, t^{\epsilon_1}, g_1, \dots, t^{\epsilon_n}, g_n)$. □

HNN extensions

Theorem

*Each $g \in G*_A$ is represented by a unique reduced word.*

Corollary

*The group G embeds into $G*_A$.*

Corollary (Britton's lemma)

If $g_0 t^{\epsilon_1} g_1 \dots t^{\epsilon_n} g_n$ is such that $g_i \in G \setminus A$ when $(\epsilon_i, \epsilon_{i+1}) = (1, -1)$ and $g_i \in G \setminus \theta(A)$ when $(\epsilon_i, \epsilon_{i+1}) = (-1, 1)$ then it is non-trivial.

Graphs of groups

Definition

If $G = A *_H B$ or $G = A *_H$ then we say that G **splits over H** .

Definition

Let Y be an **oriented graph** such that the corresponding unoriented graph is **connected** and each of its edges appears with both orientations in Y .

A **graph of groups** is a pair (G, Y) , where G is a **map** that assigns a group G_v to each vertex $v \in V(Y)$ and a group G_e to each edge $e \in E(Y)$ such that

- 1 $G_e = G_{\bar{e}}$
- 2 for all edges e , there exists an injective homomorphism $\alpha_e : G_e \rightarrow G_{t(e)}$

where $t(e)$ is the terminus of the edge $e = [o(e), t(e)]$.

Graphs of groups

Graphs of groups appear naturally when G acts on a graph X without inversions.

When this happens, we define the **quotient graph** $Y = X/G$ and the **projection** $p : X \rightarrow Y$ as follows:

- **Vertices** are orbits Gv , $v \in X$
- Gv , Gw are **joined** if there exists an edge $[v_1, w_1]$ such that $v_1 \in Gv$, $w_1 \in Gw$.

We define $p : X \rightarrow X/G$ by $p(v) = Gv$, $p(e) = \{Go(e), Gt(e)\}$.

In this case,

- $\forall v \in Y$, define $G_v = \text{Stab}(\hat{v})$ where \hat{v} is some element of $p^{-1}(v)$
- $\forall e \in Y$, define $G_e = \text{Stab}(\hat{e})$ where \hat{e} is some element of $p^{-1}(e)$

taking care that, whenever we can, \hat{v} is an endpoint of \hat{e} such that $G_e \subseteq G_v$.

For some edges, we might have to define α_e not as an inclusion, but as an inclusion composed with a conjugation.

Graphs of groups

Definition

The **path group** of the graph of groups (G, Y) is

$$F(G, Y) = \langle \bigcup_{v \in V} G_v \cup E(Y) \mid \bar{e} = e^{-1}, e\alpha_e(g)e^{-1} = \alpha_{\bar{e}}(g), \forall e \in E(Y), g \in G_e \rangle.$$

If $G_v = \langle S_v \mid R_v \rangle$ then

$$F(G, Y) = \langle \bigcup_{v \in V} S_v \cup E(Y) \mid \bigcup_{v \in V(Y)} R_v, \bar{e} = e^{-1}, e\alpha_e(g)e^{-1} = \alpha_{\bar{e}}(g) \rangle.$$

Graphs of groups

Remarks

- 1 *If all $G_v = \{1\}$ then $F(G, Y) = F(E^+(Y))$.*
- 2 *There exists an epimorphism $F(G, Y) \rightarrow F(E^+(Y))$ defined by sending each G_v to $\{1\}$.*
- 3 *If all $G_e = 1$ then*

$$F(G, Y) = *_{v \in V(Y)} G_v * F(E^+(Y)).$$

Graphs of groups

Definition

A **path** in (G, Y) is a sequence

$$c = (g_0, e_1, g_1, e_2, \dots, g_{n-1}, e_n, g_n)$$

such that $t(e_i) = o(e_{i+1})$ and $g_i \in G_{t(e_i)} = G_{o(e_{i+1})}$. If $v_0 = o(e_1)$, $v_n = t(e_n)$ then we call this **a path from v_0 to v_n** . We call

$$v_0, v_1 = t(e_1) = o(e_2), \dots, v_i = t(e_i) = o(e_{i+1}), \dots, v_n$$

its **sequence of vertices**. We define $|c|$ to be the **element of the path group** $g_0 e_1 g_1 \dots e_n g_n$. If $a_0, a_1 \in V(Y)$ then we define

$$\pi[a_0, a_1] = \{|c| : c \text{ a path from } a_0 \text{ to } a_1\}$$

Graphs of groups

Remark

If $a_0, a_1, a_2 \in V(Y)$ and $\gamma \in \pi[a_0, a_1]$, $\delta \in \pi[a_1, a_2]$ then $\gamma\delta \in \pi[a_0, a_2]$.

Proposition

Let (G, Y) be a graph of groups and suppose $a_0 \in V(Y)$. The set $\pi[a_0, a_0]$ is a subgroup of $F(G, Y)$.

Proof.

If

$$c = (g_0, e_1, g_1, e_2, \dots, g_{n-1}, e_n, g_n)$$

is a path from a_0 to a_0 then

$$|c|^{-1} = g_n^{-1} \bar{e}_n g_{n-1}^{-1} \dots \bar{e}_1 g_0^{-1} \in \pi[a_0, a_0]$$



Graphs of groups with basepoint

Proposition

Let (G, Y) be a graph of groups and suppose $a_0 \in V(Y)$. *The set $\pi[a_0, a_0]$ is a subgroup of $F(G, Y)$.*

We call this subgroup *the fundamental group of the graph of groups (G, Y) with basepoint a_0* and denote it $\pi_1(G, Y, a_0)$.