

B2.2 Commutative Algebra

Sheet 2 — HT25

Sections 1-8

Section A

1. Consider the ideals $\mathfrak{p}_1 = (x, y)$, $\mathfrak{p}_2 = (x, z)$ and $\mathfrak{m} = (x, y, z)$ of $K[x, y, z]$, where K is a field. Show that $\mathfrak{p}_1 \cap \mathfrak{p}_2 \cap \mathfrak{m}^2$ is a minimal primary decomposition of $\mathfrak{p}_1 \cdot \mathfrak{p}_2$. Determine the isolated and the embedded prime ideals of $\mathfrak{p}_1 \cdot \mathfrak{p}_2$.

Solution: For future reference, note that we have

$$\mathfrak{m}^2 = ((x) + (y) + (z))^2 = (x^2, y^2, z^2, xy, xz, yz)$$

and

$$\mathfrak{p}_1 \cdot \mathfrak{p}_2 = ((x) + (y))((x) + (z)) = (x^2, xz, yx, yz).$$

We have $\mathfrak{p}_1 \cdot \mathfrak{p}_2 \subseteq \mathfrak{p}_1 \cap \mathfrak{p}_2$ and we also clearly have $\mathfrak{p}_1 \cdot \mathfrak{p}_2 \subseteq \mathfrak{m}^2$ since $\mathfrak{p}_1, \mathfrak{p}_2 \subseteq \mathfrak{m}$. Thus we have $\mathfrak{p}_1 \cdot \mathfrak{p}_2 \subseteq \mathfrak{p}_1 \cap \mathfrak{p}_2 \cap \mathfrak{m}^2$. Note that \mathfrak{p}_1 and \mathfrak{p}_2 are prime since the rings $K[x, y, z]/\mathfrak{p}_1 \simeq K[z]$ and $K[x, y, z]/\mathfrak{p}_2 \simeq K[y]$ are domains. Note also that \mathfrak{m} is a maximal ideal, since $K[x, y, z]/\mathfrak{m} \simeq K$ is a field. Thus \mathfrak{p}_1 , \mathfrak{p}_2 and \mathfrak{m}^2 are primary (see after Lemma 6.4 for the latter). The radicals of the ideals \mathfrak{p}_1 , \mathfrak{p}_2 and \mathfrak{m}^2 are \mathfrak{p}_1 , \mathfrak{p}_2 and \mathfrak{m} (see again Lemma 6.4 for the latter). These three ideals are distinct. Finally, we have $\mathfrak{p}_1 \not\supseteq \mathfrak{p}_2 \cap \mathfrak{m}^2$ (because $z^2 \notin \mathfrak{p}_1$ but $z^2 \in \mathfrak{p}_2 \cap \mathfrak{m}^2$), $\mathfrak{p}_2 \not\supseteq \mathfrak{p}_1 \cap \mathfrak{m}^2$ (because $y^2 \notin \mathfrak{p}_2$ but $y^2 \in \mathfrak{p}_1 \cap \mathfrak{m}^2$) and $\mathfrak{m}^2 \not\supseteq \mathfrak{p}_1 \cap \mathfrak{p}_2$ (because $x \notin \mathfrak{m}^2$ but $x \in \mathfrak{p}_2 \cap \mathfrak{p}_1$). Hence if $\mathfrak{p}_1 \cdot \mathfrak{p}_2 = \mathfrak{p}_1 \cap \mathfrak{p}_2 \cap \mathfrak{m}^2$ then this decomposition is indeed primary and minimal. Thus we only have to show that $\mathfrak{p}_1 \cdot \mathfrak{p}_2 \supseteq \mathfrak{p}_1 \cap \mathfrak{p}_2 \cap \mathfrak{m}^2$.

From the above, we have to show that

$$(x, y) \cap (x, z) \cap (x^2, y^2, z^2, xy, xz, yz) \subseteq (x^2, xz, yx, yz).$$

This is immediate, since all the ideals we are considering have the property that a polynomial lies in such an ideal if and only if all of the monomial summands of the polynomial lie in the ideal.

Section B

2. Let K be a field. Show that the ideal $(x^2, xy, y^2) \subseteq K[x, y]$ is a primary ideal, which is not irreducible.

Solution: We first show that (x^2, xy, y^2) is primary. This simply follows from the fact that (x, y) is maximal ideal and from the fact that $(x^2, xy, y^2) = (x, y)^2$ (see after Lemma 6.4).

Now note that $(x^2, xy, y^2) = (x^2, y) \cap (x, y^2)$. This is again immediate, since all three ideals have the same property as in Question 1.

3. Let R be a noetherian ring and let T be a finitely generated R -algebra. Let G be a finite subgroup of the group of automorphisms of T as a R -algebra. Let T^G be the fixed point set of G (ie the subset of T , which is fixed by all the elements of G).
- (a) Show that T is integral over T^G .
- (b) Show that T^G is a subring of T , which contains the image of R and that T^G is finitely generated over R .

Solution: It is clear from the definitions that T^G is a subring which contains the image of R . Let $t \in T$. Then t satisfies the polynomial equation

$$\prod_{g \in G} (t - g(t)) = 0$$

The polynomial $M_t(x) = \prod_{g \in G} (x - g(t))$ has coefficients in T^G , because the coefficients are symmetric polynomials in the variables $g(t)$, and therefore are invariant under G . Hence t is integral over T^G . Since t was arbitrary, T is integral over T^G . Since T is also finitely generated as a T^G -algebra (because it is finitely generated as an R -algebra), we thus see that T is finite over T^G (see after Lemma 8.3). Hence T^G is finitely generated over R by the Artin–Tate Theorem.

4. Show that \mathbb{Z} is integrally closed and that the integral closure of \mathbb{Z} in $\mathbb{Q}(i)$ is $\mathbb{Z}[i]$.

Solution: We first prove that \mathbb{Z} is integrally closed. Let $p/q \in \mathbb{Q}$, where p and q are coprime integers, and let $P(x) = x^n + a_{n-1}x^{n-1} + \dots + a_0 \in \mathbb{Z}[x]$ be a monic polynomial. Suppose that $P(p/q) = 0$. Then we have

$$q^n P(p/q) = p^n + a_{n-1}p^{n-1}q + a_{n-2}p^{n-2}q^2 + \dots + a_0q^n = 0.$$

Since $a_{n-1}p^{n-1}q + a_{n-2}p^{n-2}q^2 + \dots + a_0q^n$ is divisible by q and p^n is coprime to q , this implies that $q = \pm 1$, so $p/q \in \mathbb{Z}$.

To prove that the integral closure of \mathbb{Z} in $\mathbb{Q}(i)$ is $\mathbb{Z}[i]$, note first that i is a root of the monic polynomial $x^2 + 1$ over \mathbb{Z} .

So we only have to prove that $\mathbb{Z}[i]$ is integrally closed in $\mathbb{Q}(i)$ (see Lemma 8.6). Note furthermore that $\mathbb{Q}(i)$ is the fraction field of $\mathbb{Z}[i]$. To see this, write let $r + it \in \mathbb{Q}(i)$, where $r, t \in \mathbb{Q}$ (any element of $\mathbb{Q}(i)$ can be written in this form because $\mathbb{Q}(i) \simeq \mathbb{Q}[x]/(x^2 + 1)$). Let $r = p/q$ and $t = u/v$. We then have $r + it = (vp + uqi)/(vq)$, which is a fraction of elements of $\mathbb{Z}[i]$, proving our claim. Finally, recall that we know from Rings and Modules that $\mathbb{Z}[i]$ is a Euclidean domain, where the Euclidean function is given by the norm (the norm of $c + id$ is $c^2 + d^2$ if $c + id \in \mathbb{Z}[i]$). In particular, $\mathbb{Z}[i]$ is a PID and every ideal in $\mathbb{Z}[i]$ is generated by an element of smallest norm.

To prove that $\mathbb{Z}[i]$ is integrally closed in $\mathbb{Q}(i)$, we may now proceed as for \mathbb{Z} . Let

$$P(x) = x^n + a_{n-1}x^{n-1} + \dots + a_0 \in \mathbb{Z}[i][x]$$

and let $r + it = B/A$, be its root where $A, B \in \mathbb{Z}[i]$. Since $\mathbb{Z}[i]$ is a UFD, we may assume that A and B do not share any irreducible factors. We can now write as before

$$A^n P(B/A) = B^n + a_{n-1}B^{n-1}A + a_{n-2}B^{n-2}A^2 + \dots + a_0A^n = 0.$$

Since $a_{n-1}B^{n-1}A + a_{n-2}B^{n-2}A^2 + \dots + a_0A^n$ is divisible by A and B^n is coprime to A , this implies that A is a unit, so $B/A \in \mathbb{Z}[i]$.

Note that the proof above actually shows that any UFD (Unique Factorisation Domain) is integrally closed.

5. Let S be a ring and let $R \subseteq S$ be a subring of S . Suppose that R is integrally closed in S . Let $P(x) \in R[x]$ and suppose that $P(x) = Q(x)J(x)$, where $Q(x), J(x) \in S[x]$ and $Q(x)$ and $J(x)$ are monic. Show that $Q(x), J(x) \in R[x]$. Use this to give a new proof of the fact that if $T(x) \in \mathbb{Z}[x]$ and $T(x) = T_1(x)T_2(x)$, where $T_1(x), T_2(x) \in \mathbb{Q}[x]$ are monic polynomials, then $T_1(x), T_2(x) \in \mathbb{Z}[x]$.

Solution: We will need the following.

Lemma. Let A be a ring and let $U(x) \in A[x]$ be a non zero monic polynomial. Then there exists a ring B containing A , which is integral over A and such that

$$U(x) = \prod_{i=1}^{\deg(U)} (x - b_i)$$

for some $b_i \in B$, where we set $\prod_{i=1}^{\deg(U)} (x - b_i) = 1$ if $\deg(U) = 0$.

Proof of the lemma. By induction on the degree $d = \deg(U)$ of $U(x)$. If $d \in \{0, 1\}$, there is nothing to prove. So suppose that $d > 1$ and that the result holds for any smaller value of d . The ring $C = A[y]/(U(y))$ is integral over A by Proposition 8.2. The element y of C satisfies the equation $U(y) = 0$ by construction. By Euclidean division (see Preamble), we thus have $U(x) = (x - y)Z(x)$ for some $Z(x) \in C[x]$. Since $Z(x)$ has degree less than d , we may apply the inductive hypothesis and we obtain a ring B , which contains C and where $Z(x)$ splits. The polynomial $U(x)$ also splits in B , so we are done. \square

We now apply the lemma to $Q(x)$ and $J(x)$ successively and we obtain a ring B , which contains S , such that B is integral over S and such that

$$Q(x) = \prod_{i=1}^{\deg(Q)} (x - b_i)$$

and

$$J(x) = \prod_{i=1}^{\deg(J)} (x - c_i)$$

where $b_i, c_i \in B$. Now we have $P(b_i) = P(c_i) = 0$ by construction, so the b_i and c_i are actually integral over R . Since the integral closure of R in B is a subring, we conclude that the coefficients of $Q(x)$ and $J(x)$ are integral over R (and in S , by assumption). But since R is integrally closed in S , this means that these coefficients lie in R .

Note that we did not actually use the fact that B was integral over S in the proof.

6. Let R be a subring of a ring T and suppose that T is integral over R . Let \mathfrak{p} be prime ideal of R and let \mathfrak{q} be a prime ideal of T . Suppose that $\mathfrak{q} \cap R = \mathfrak{p}$. Let $\mathfrak{p}_1 \subseteq \mathfrak{p}_2 \subseteq \dots \subseteq \mathfrak{p}_k$ be primes ideal of R and suppose that $\mathfrak{p}_1 = \mathfrak{p}$. Show that there are prime ideals $\mathfrak{q}_1 \subseteq \mathfrak{q}_2 \subseteq \dots \subseteq \mathfrak{q}_k$ of T such that $\mathfrak{q}_1 = \mathfrak{q}$ and such that $\mathfrak{q}_i \cap R = \mathfrak{p}_i$ for all $i \in \{1, \dots, k\}$.

Solution: By induction on k , we only need to treat the case $k = 2$. Consider the extension of rings $R/\mathfrak{p} \subseteq T/\mathfrak{q}$. This is also an integral extension. Furthermore, there is a unique prime ideal \mathfrak{p}'_2 in R/\mathfrak{p} , which corresponds to \mathfrak{p}_2 via the quotient map. By the Going-up Theorem, there is a prime ideal \mathfrak{q}'_2 in T/\mathfrak{q} , which is such that $\mathfrak{q}'_2 \cap R/\mathfrak{p} = \mathfrak{p}'_2$. The prime ideal \mathfrak{q}_2 corresponding to \mathfrak{q}'_2 via the quotient map has the required properties.

7. Let R be a ring. Let \mathcal{S} be the set of ideals in R that are not finitely generated; assume that $\mathcal{S} \neq \emptyset$.
- (a) Show that \mathcal{S} has at least one maximal element.
 - (b) Let I be maximal element of \mathcal{S} (with respect to the relation of inclusion). Show that I is prime.
 - (c) Suppose that all the prime ideals of R are finitely generated. Prove that R is noetherian.

[Hint: exploit the fact that R/I is noetherian.]

Solution:

- (a) Zorn's lemma.
- (b) Let $x, y \notin I$ and suppose for contradiction that $xy \in I$. Let $I_x = (x) + I$ and $I_y = (y) + I$. Write $J = I_x \cdot I_y$. By assumption I_x, I_y and hence J are finitely generated, and we have $J \subset I$. Consider the image $I \bmod J$ of I in the R/I_y -module I_x/J . Note that I_x/J is finitely generated as an R/I_y -module since I_x is finitely generated as an R -module. Note also that the ring R/I_y is noetherian, since every ideal of R/I_y is the image of either the zero ideal or of an ideal of R strictly containing I . Hence $I \bmod J$ is also finitely generated as an R/I_y -module, and therefore as an R -module. So is J , and the extension of finitely generated modules is finitely generated, yielding finite generation of I as an R -module. This is exactly the same as being finitely generated as an ideal; contradiction.
- (c) This is immediate now – if \mathcal{S} is non-empty, then it has a prime member.

Section C

8. Let R be a ring. Let \mathcal{S} be the set of non-principal ideals in R ; assume that $\mathcal{S} \neq \emptyset$. Prove that \mathcal{S} admits maximal elements, and that every such element is a prime ideal.

Solution: The existence of maximal elements follows from Zorn's lemma. Let I be one such. Let $x, y \notin I$ and suppose for contradiction that $xy \in I$. Let $I_x = (x) + I$. By assumption, we have $I_x = (g_x)$ for some $g_x \in R$. Let $\phi: R \rightarrow I_x$ be the surjection of R -modules given by the formula $\phi(r) = rg_x$. We then have $I \subseteq \phi^{-1}(I)$.

Suppose first that $I = \phi^{-1}(I)$. In other words, for all $r \in R$, we have $rg_x \in I$ if and only if $r \in I$. This contradicts the fact that $yg_x \in I$. So we conclude that $I \subset \phi^{-1}(I)$. From the definition of I , we then see that $\phi^{-1}(I)$ is a principal ideal of R , and hence so is $I = \phi(\phi^{-1}(I))$. This is a contradiction.