

## Geometric Group Theory

### Problem Sheet 0-Solutions

1. Show that a subgroup of index 2 is normal.

*Solution.* If  $H$  is a subgroup of index 2 of  $G$  then for any  $g \notin H$  we have  $G = H \cup gH = H \cup Hg$  so  $gH = Hg$  and  $H$  is normal.

2. Let  $A, B$  be finite index subgroups of  $G$ . Show that  $A \cap B$  is a finite index subgroup of  $G$ .

*Solution.*

We show first that if  $K < H < G$  then  $|G : K| = |G : H| \cdot |H : K|$ : Say  $G = \bigcup Ha_i, H = \bigcup Kb_j$  then

$$G = \bigcup Kb_j a_i. \text{ (disjoint union)}$$

Indeed assume  $Kb_j a_i = Kb_k a_l$  then  $a_l a_i^{-1} \in H$  so  $a_l = a_i$  and  $b_k = b_j$ .

We remark now that  $|A : A \cap B|$  is finite since  $|G : B|$  is finite and the map  $a(A \cap B) \rightarrow aB$  is 1-1.

So  $|G : A \cap B| = |G : A| \cdot |A : A \cap B| < \infty$ .

3. Let  $G$  be a finitely generated group and let  $H$  be a subgroup of  $G$  of finite index. Show that  $H$  is finitely generated.

*Solution.* Let  $A = \{a_1, \dots, a_n\}$  be generators of  $G$  and let  $X = \{x_1 = 1, \dots, x_k\}$  be right coset representatives for  $H$  in  $G$ . Consider the set

$$S = \{x_i a_j x_k^{-1} : x_i, x_k \in X, a_j \in A, \text{ such that } x_i a_j x_k^{-1} \in H\}$$

If  $g \in H$  then  $g = g_1 \dots g_r, (g_i \in A)$ . Clearly there are  $y_1, \dots, y_{r-1} \in X$  such that all

$$g_1 y_1^{-1}, y_1 g_2 y_2^{-1}, \dots, y_{r-2} g_{r-1} y_{r-1}^{-1}$$

lie in  $S$ . Note that

$$(g_1 y_1^{-1}) \cdot (y_1 g_2 y_2^{-1}) \cdot \dots \cdot (y_{r-2} g_{r-1} y_{r-1}^{-1}) \cdot (y_{r-1} g_r) = g_1 \dots g_r \in H$$

It follows that  $y_{r-1} g_r \in S$ , so  $S$  is a finite set of generators of  $H$ .

4. Show that if  $G$  is a finitely generated group such that every element of  $G$  has order 2 then  $G$  is finite.

*Solution.* Let  $a, b \in G$ . Then  $abab = 1$  so  $ab = ba$  and  $G$  is abelian. Since  $G$  is finitely generated  $G \cong \mathbb{Z}_2^n$  for some  $n \geq 0$ .

5. Let  $H$  be a finite index subgroup of  $G$ . Show that there is a normal finite index subgroup of  $G$ ,  $N$  such that  $N \subset H$ .

*Solution.* Consider the action of  $G$  on the left cosets  $aH$  where  $g(aH) = (ga)H$ . If  $|G : H| = n$  this action induces a homomorphism from  $G$  to the

finite symmetric group  $S_n$  so its kernel is a finite index normal subgroup  $N$  which is clearly contained in  $H$ . Since  $|S_n| = n!$ ,  $|G : N| \leq n!$ .

**6.** Let  $G$  be a finitely generated group. Show that  $G$  has finitely many subgroups of index  $n$ . (*hint*: use the previous exercise).

*Solution* By the previous exercise any subgroup  $H$  of index  $n$  contains a normal subgroup  $N$  of index bounded by  $n!$ . So there is a homomorphism  $f : G \rightarrow G/N$ . where  $|G/N| \leq n!$  and  $\ker f \subseteq H$ .

Consider the set of homomorphisms from  $G$  to groups of order at most  $n!$ . Since  $G$  is finitely generated and a homomorphism is given by assigning values to generators there are finitely many such homomorphisms say  $f_1, \dots, f_k$ . If  $H_1, H_2$  are subgroups of index  $n$  then for some  $f_i$ ,  $f_i(H_1) \neq f_i(H_2)$ . Indeed take  $f_i$  such that  $N = \ker f_i \subseteq H_1$ . If  $h_2 \in H_2 - H_1$  and  $f_i(h_2) = f_i(h_1)$  with  $h_1 \in H_1$  then  $h_2 h_1^{-1} \in N$  so  $h_2 \in N h_1 \subset H_1$ , a contradiction.

In other words the map  $H \rightarrow (f_1(H), \dots, f_k(H))$  is 1-1.