
BO1.1. History of Mathematics Lecture V Newton's *Principia*

MT24 Week 3

Summary

- ▶ Isaac Newton (1642–1727)
- Kepler's laws,
 Descartes' theory,
 Hooke's conjecture
- ► The Principia
- Editions and influence of the *Principia*



Newton summarised

Alexander Pope, 1730:

Nature and Nature's Laws lay hid in Night. God said, Let Newton be! and All was Light.

Woolsthorpe Manor




Newton born at Woolsthorpe Manor, 25th December 1642

Woolsthorpe Manor




Grantham Grammar School



Now The King's School, Grantham

John Bate: The mysteries of nature and art (1634)



John Bate: The mysteries of nature and art (1634)

The first Booke

16



little from it on the face Hawing thus prepared the barrell, fit a good thick board unto it, fo that it may lip eafily up and down from the top of the barrell unto the bottom, nayle a lether about the edges offit, and another upon the top of it? on the underfide of it let there he fathered a good diffe.

but flexible fpring of fleele, which may thrust the board from the bottom to the top of the barrell: let the foot of this spring reft upon a barre fashed acros the bottom of the barrell; let this board also have tied at the middle a little tope of slength sifficient. When you used it, bore a little the lein the table that you fer it on, to put the rope thorow, and pull the rope down, which will contract the spring, and with it draw down the board: then poure in water at the hafin until the vessel let first a you let slack the rope, the water will spring the the water will run into the vessel as you pull it straight, the water will run into the vessel as you pull it straight, the water will run into the vessel as you go the spring the straight should be suffered to the pipe, and which the water may break.

of Water-workes.

17

Another manner of forcing water, whereby the water of any firing may be forced unto the tob of a hill.

Let there be two hollow posts, with a succur at the bottom of each, also a succur night the top of each: let there be sastned unto both these posts a strong pecce of




timber, having, as it were, a beame or scale pinned in it, and having two handles, at each end one. In the tops of

Another

John Bate: The mysteries of nature and art (1634)



Trinity College, Cambridge



Isaac Newton (1642–1727)

Newton's major interests:

1650s: model-building

1660s: optics; (pure) mathematics

1670s: alchemy, theology

1684+: mathematics

1696-1727: Warden of the Mint


Johannes Kepler (1571–1630)

Engaged to sift through the astronomical data gathered by the Danish astronomer Tycho Brahe (1546–1601)

Major works: Astronomia nova (1609) Harmonices mundi (1619)



Kepler: Astronomia nova (1609)



PHYSICA COELESTIS,

DE MOTIBUS STELLÆ

MARTIS,

Ex observationibus G. V.

TTCHONIS BRAHE:

Jussu & sumptibus

RVDOLPHI II

Plurium annorum pertinaci studio elaborata Praga ,

JOANNE KEPLERO,

Comejusdem Co. M. in privilegio speciali

Anno ata Dionysiana elo lo e 1x.



184 DE MOTIB STELLÆ MARTIS

milli methodo; traffing determin timina terpeladerseps. A bias distinct and traffic and tr

prodiderit diffantiam a sjudam e him index peritandebarterranium pilma a a herium, quatrodom efficon in e libena quod verifilment, pilma a a herium quatrodom efficon in e de libena quod verifilment, pilma a salerum estam si a qua focel o si e s. Effe ignor a forma a salerum peritandomen juli a diffantia, film 8c looprodinen 8c s. s. a anomalis ver e cocquara. Manifestim eff autemno est sa se a compara somo estam si a quatrodomi pilma estam si a quatrodomi peritandomen juli a diffantia, film 8c looprodinen 8c s. s. a anomalis ver e cocquara. Manifestim eff autemno el 1 sa setto fiere o le lineam, looc disperiori folicer in r. ixaque anguli s. a 8c s a o differant quantities a s. s.

Argumentatio mea talis furt, qualis cap. XIXXI. & VVI. Circu lus cap. XIIII. peccat excellu, elliptis capitis XIV peccat defectu. E fiint excellus ille & hic defectus #quales. Inter circulum vero & elliptis excellus ille & hic defectus #quales.


Kepler's laws

Kepler's laws of planetary motion (1609, 1619):

- 1. Planets move in elliptical orbits with the sun as focus
- 2. Planets sweep out equal areas in equal times
- 3. T^2 is proportional to R^3 (where T is time of one revolution, R is mean distance to sun)

All from empirical evidence


Descartes' theory



Descartes' views of planetary motion in *Principia philosophiae* (1644):

- the sun is one star among many
- asserted that planets are carried round their suns by vortices of the surrounding 'ether'
- claimed that theory could also explain magnetism and static electricity

An inverse square law?




An inverse square law?

Speculations and calculations on an inverse square law of gravity:

- 1645 Ismaël Bullialdus refutes a claim of Kepler that 'gravity' drops off linearly with distance, instead suggesting an inverse square law
- c. 1679 Hooke corresponds with Newton and suggests that an inverse square law might lead to elliptical orbits
 - 1684 Halley visits Newton and asks whether this might be the case; Newton sends a short treatise on motion to Halley
 - 1687 Publication of Newton's *Principia* at Halley's expense

Isaac Newton: The mathematical principles of natural philosophy (London, 1687)



Contents of the Principia

- Eight definitions of matter, motion, innate force, impressed force, acceleration, ...
- Three axioms or Laws of Motion (as taught in school), together with six corollaries
- Book I: The motion of bodies
- ▶ Book II: The motion of bodies in resisting media
- ▶ Book III: The system of the world

The laws of motion

T 12 7

AXIOMATA SIVE LEGES MOTUS

Lex. I.

Corpus omne perfeverare in statu suo quiescendi vel movendi uniformiter in directum, nisi quatenus a viribus impressis cogitur statum illum mutare.

Projectifia perfeverant in motibus finis nill quatents a refiftentra acris teardantur & vi gravitatis impellantur deorfum. Trochus, cujus partes colarendo perpetuo retralunte fed a motibus refilincis, non ceffir rotari nili quatentus ab acre retardatur. Majora attene Plasetarum & Cometarum corpora motus futos & progreffivos & circulares in spatis minus resistentibus fados conferenta dintius.

Lex. II.

Mutationem motus proportionalem esse vi motrici impresse, & sieri secundum lineam restam qua vis illa imprimitur.

Si via aliqua motum quemvis generes, dupla duplum, tripla triplum generalis, five fimul & femel, five gradatim & fincedire imperfia fiterir. Et hie motus quoniam jin eandem femper plagam quin vigerenarise determinatur, fi corpus anten movebatur, moturi pass vle onifijanti additur, vel contrano fiabilacitur, vel obliquo oblique adjicitur, & cum co fecundum utriufiq determinatiorem componitur. [13]

Astioni contrariam semper & aqualem esse reactionem: siwe corporum duorum actiones in se mutuo semper esse aquales & in partes contrarias dirigi.

Quicquid premit vel traiti alterum, tantundemale co premiture Velusius el traitium. Siquii alpidem digito premit; premiture Schuius digitus a lapides. Si equus lapidem fumi alle gatum traitir, retraheture eriam Se equus sequalitere in lapidem; ruam fumis tutrina, differents codem relixandis Feotoma traitis. Se forqua slapidem, as elapidem verfüs equum, tantunaçi impediet progreffum unitus quantum promovet progreffum alterius. Si corqua silauden incerpus aliud impingems, mostum ejus vi fina quomodocune; muttaverit, i-dem quoque viculifim in motur proprio candem unutationem in partem contrariam vi alterius (ob aqualitatem prefiloris mutus) fibibite. His alcionibus aquales funt mutationes non velocitatum fed motuum, (feilice in corporibus non altunde impeditis:) Mutationes enim velocitatum, in contrarias intelm partes false, quia motus aqualiter mutantur, fiunt corporibus reciproce proportionales.

Corol. L

Corpus viribus conjunctis diagonalem parallelogrammi eodem tempore defcribere, quo latera feparatis.

Si corpus dato tempore, vi íola M, ferretur ab A ad B, & vi íola N, ab A ad C, compleatur parallelogrammum ABDC, & vi utraq; feretur id eodem tempore ab A ad D. Nam quoniam vis N agit fecundum lineam



AC ipf BD parallelam, hac vis nihil mutabit velocitatem accedendi ad lineam illam B D a vi altera genitam. Accedet igitum corpus codem tempore ad lineam BD five vis N imprimatur, five non, atq; adeo in fine illius temporis reperietur alicubi in linea

Book I: Motion of bodies

Book I, Section I: On the method of first and last ratios

Lemma I: Quantities, and ratios of quantities, which [...] approach nearer to each other than by any given difference, become ultimately equal.

For suppose they are ultimately unequal, and their ultimate difference is D. Then they cannot approach nearer to equality than by that difference.

Book I, Lemma II

[27] Lemma II.

Si in figura quavis AacE rechis Aa, AE, & curva AcE comprehenfa, inferibantur parallelogramma quoteunq; Ab, Bc, Cd, &c. fub bafibus AB, BC, CD, &c.

equalibus, & latevibus Bb, Cc, D d, &cc, figure luteri Aa parallelis contenta; & compleantur parallelogramma aKbl, bl.cm, cMdn, &cc, Dein burum parallelogrammorum laitudo mimuatur, & munevus augeatur in infuitum: dico quod ultime rationes, quas bahen ad fe invicem figura inferipta AKbl.cMdD, cireumferipta Aalbmend oE, & curvalinea AabedE, funt rationes ornalitatis.



Nam figura inferipte & circumferipte differentia eft funma parallelogrammorum Kl+Lm+Mn+Do, hos eft (ob æquales omnium bales) rechangulum fub unius balt Kb & alritudinum fumma As, id eft rechangulum RbLa. Sed hoe rechangulum, co quod hatriudocjus AB in infinitum minuitur, fit minus quovis dato. Ergo, per Lemma I, figura inferipta & circumferipta & multo magis figura curvilinea intermedia fiunt ultimo equales, O, E, D.

Lemma III.

Exdem rationes ultime funt etiem equalitatis, ubi parallelogramomrum latitudines AB, BC, CD, &c. funt inequales, & omnes minuumtur in infinitism.

Sit enim AF aqualis latitudini maxima, & compleatur parallelogrammum FA of the erit majus quam differentia figure inferipta & fi une circumferipra, at latitudine fua AF

Lemma II: Ultimate equality of inscribed figure, circumscribed figure, and curved area


Motion under centripetal forces

E 37] S E C T. II. De Inventione Virium Centripetarum. Prop. I. Theorema. I. Areas quas corpora in gyros acta radiis ad immobile centrum virium ductis defenium, & im planis immobilius confifere, & elle tendentical defenium, o im planis immobilius confifere, & elle tendentical defenium.

poribus proportionales.

Dividatur tempus in partes æquales, & prima temporis parte describat corpus vi instra restam AB. Idem secunda temporis partes, si in li impedires, resta pergeret ad c, sper Leg. 1) describens lineam Bc æqualem ipsi AB, adeo ut radiis AS, BS, cS ad

centrum adis, confedar forent aquales area A SB, B Se. Verum ubi corpus venit ad B, agat viscentripetaim-pulfu unico fed magno, faciatq; corpus a refla Be deflectere & pergere in refla BC. Ipfi B S parallela agature C cocurrens BC in



C, & completa fecunda temporis parte, corpus ("per Legum Coriol. 1) reperietur in C, in codem plano cum triangulo A SB. Junge SC, & triangulum SBC, do parallelas SB, Cc, aquale erit triangulo SBC, act, adeo eriam triangulo SAB. Simili argumento fi Book I, Section II: Motion under centripetal forces.

Proposition I: Bodies constrained by a central force to orbit a fixed point move in a plane and sweep out equal areas in equal times.

(Kepler's second law)

NB. independent of the 'law of force' involved.

Book I, Section II: Circular motion

[41]

Prop. IV. Theor. IV.

Corporum que diversos circulos sequabili motu describum, vires centripeas ad centra contudem circulorum tendere, & esse in arcuna simul descriptorum quadrata applicata ad circulorum rados.

Corpora B, b in circumferentiis circulorum BD, bd gyrantia, finul deferibant arcus BD, bd. Quoniam fola vi infita deferiberent tangentes BC, bc his arcubus æquales, manifeftum eft quad vires centripetæ funt quæ

etrquod vires centriperes unit que perpeturo retrabunt corpora de tangentibus ad circumierentias circulorum, atq, adeo he fun ad invicem in racione prima fpatiorum nafeentum CD, cd: tendunt vero ad centra circulorum per Theor. II, propterea quod areæ radiis deferipte ponuntur temporibus proportionales. Fat figura Lb figura D CB fimilis, & per Lemma V, lincola CD critad Bnoolam ktut



deoqs. (ob equales rationes $\frac{b}{S}\frac{t}{b} \& \frac{B}{S}\frac{D}{B}$) ut $\frac{B}{S}\frac{D}{B}$ quad. ad $\frac{b}{S}\frac{d}{b}$

Corol. 1. Hine vires centripetæ funt ut velocitatum quadrata applicata ad radios circulorum.

Corol, 2. Et reciproce ut quadrata temporum periodicorum appli-

Book I, Sect. II, Prop. IV: Motion under centripetal forces: motion in a circle.

Corollary 1: For motion in a circle centripetal force is proportional to $\frac{v^2}{r}$.

Corollary 6: For motion in a circle Kepler's third law implies an inverse square law of force.

Book I, Section III: orbits that are conic sections

[50]

SECT. III.


De motu Corporum in Conicis Sectionibus excentricis.

Prop. XI. Prob. VI.

Revolvatur corpus in Ellipsi: Requiritur lex vis centripetæ tendentis ad umbilicum Ellipseos.

Esto Ellipseos superioris umbilicus S. Agatur SP secans Ellipseos tum diametrum DK in E, tum ordinatim applicatam Q = 0 in X, S compleatur parallelogrammum $Q \times PR$. Patet EP as

qualem effe temiaxi majori AC, oquod aĉa ab altero quod aĉa ab altero Ellipfeos umbilico Hlinca HIIpfi EC parallela, (ob aquales CS, CH) acquentur ES, EL, adeo ut EP femiliumma fit ipfarum PS, PI, id eft (ob parallelas HI, PR & angulos æquales IP R, HPZ) ipforum PS, PI, qua



conjunction axem torum 2AC adequant. Ad SP demittatur perpendicularis QT, & Ellipfeos latere recto principali (feu 2BC quad.) dicto L, crit $L \times QR$ ad $L \times P = v$ ut QR ad P = v;


AC
id eft ut PE (feu AC) ad PC: & LxPvad GvP ut LadGv;

Proposition XI: Motion under centripetal forces: Kepler's First Law (orbit is an ellipse with sun at focus) implies an inverse square law of force.

Proposition XII: Motion under centripetal forces: hyperbolic orbit implies an inverse square law of force.

Proposition XIII: Motion under centripetal forces: parabolic orbit implies an inverse square law of force.

Book I, Sections II and III summarised



Book I, later sections

More mechanics of motion:

- converses: an inverse square law of force implies that orbits are conic sections;
- trajectories;
- much more besides.

All treated geometrically

Books II and III

Book II: Motion of bodies in resisting media:

Conclusion: "... it is manifest that the planets are not carried round in corporeal vortices ..." (Scholium to Proposition LIII)

Book III: The system of the world:

- Reconciliation of observation and theory
- Shape of the earth (correct?)
- Motion of the moon (wrong)
- Prediction of tides
- Comets


Influence of the Principia

Principia showed how mathematical methods could be used to study physical, especially but not exclusively, cosmological phenomena.

New ways of thinking: for example the method of ultimate ratios, though expressed geometrically, came close to a modern concept of limits.


Predictions could be verified by observation and experiment — verified (after some controversy) in the case of the shape of the earth, contradicted in the case of the motion of the moon.


For more on the *Principia*...




(Colin Pask, Magnificent Principia, Prometheus Books, 2013)

Three (very different) books among many...







And a lecture given at Gresham College: https://www.gresham.ac.uk/lectures-and-events/isaac-newtons-world