


Infinite Groups

Cornelia Druţu

University of Oxford

Part C course MT 2024, Oxford

Mikhael Leonidovich Gromov



“This common and unfortunate fact of the lack of adequate presentation of basic ideas and motivations of almost any mathematical theory is probably due to the binary nature of mathematical perception. Either you have no inkling of an idea, or, once you have understood it, the very idea appears so embarrassingly obvious that you feel reluctant to say it aloud.”

“But anything that can be called “rigor” is lost exactly where the things become interesting and non trivial.”

The Baumslag-Solitar group

An example of solvable (even metabelian) **finitely presented** group that is not polycyclic is the **Baumslag-Solitar group**.

$$G = BS(1, p) = \langle a, b \mid aba^{-1} = b^p \rangle \text{ for } |p| \geq 2.$$

The matrices

$$a = \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix} \text{ and } b = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$$

generate a subgroup of $SL(2, \mathbb{R})$ isomorphic to $BS(1, p)$.

Baumslag-Solitar group continued

- The **derived subgroup** G' of G is (isomorphic to)

$$G' = \left\{ \begin{pmatrix} 1 & mp^k \\ 0 & 1 \end{pmatrix} ; m, k \in \mathbb{Z} \right\}.$$

- Therefore $G = BS(1, p)$ is **metabelian**.
- The derived subgroup G' is **not finitely generated**.

Hence G is **not polycyclic**.

Nilpotency class and derived length

Every nilpotent group is solvable.

Question: find a relationship between nilpotency class and derived length.

Proposition

① For every group G and every $i \geq 0$,

$$G^{(i)} \leq C^{2^i} G.$$

② If G is k -step nilpotent then its derived length is at most

$$[\log_2 k] + 1.$$

Proof (1) by induction on $i \geq 0$.

The statement is obviously true for $i = 0$. Assume that it is true for i .

Then

$$G^{(i+1)} = [G^{(i)}, G^{(i)}] \leq [C^{2^i} G, C^{2^i} G] \leq C^{2^{i+1}} G.$$

(2) follows immediately from (1). □

Remark

*The derived length can be much smaller than the nilpotency class:
the dihedral group D_{2n} with $n = 2^k$ is k -step nilpotent and metabelian.
In particular we do not have $\ell_{\text{der}}(G) \geq f(k)$, with $\lim_{k \rightarrow \infty} f(k) = \infty$.*

Linear groups

In what follows \mathbb{K} is an algebraically closed field (e.g. \mathbb{C}), V is a finite-dimensional vector space over \mathbb{K} .

$\text{End}(V)$ is the algebra of (linear) endomorphisms of V .

$GL(V)$ is the group of invertible endomorphisms of V .

A linear action of a group G on V is called **representation** of G on V .

It amounts to the existence of a **group homomorphism** $\rho : G \rightarrow GL(V)$.

The representation may not be **faithful** (i.e. ρ might have a **kernel**.)

A group G that is **isomorphic** to a subgroup of $GL(V)$, for some V , is called a **matrix group** or a **linear group**.

The subalgebra of $\text{End}(V)$ generated by a linear group G will be denoted by $\mathbb{K}[G]$; **this is just the linear span of G over \mathbb{K} .**

Trace, $GL(V)$ and $End(V)$

Lemma

The map $\tau : End(V) \times End(V) \rightarrow \mathbb{K}$, $\tau(A, B) = \text{trace}(AB)$ is a non-degenerate bi-linear form.

Fixing a basis for V determines:

- an isomorphism of groups $GL(V) \simeq GL_n(\mathbb{K})$, where $GL_n(\mathbb{K})$ is the group of invertible $n \times n$ matrices over \mathbb{K} ;
- an isomorphism of algebras $End(V) \simeq M_n(\mathbb{K})$, where the latter is the algebra of all $n \times n$ matrices over \mathbb{K} .

Irreducible, reducible and triangularizable actions

If V is a vector space and $A \leq \text{End}(V)$ is a subgroup, then A is said to act **irreducibly** on V if V contains no proper subspace $\{0\} \subsetneq V' \subsetneq V$ such that $aV' \subset V'$ for all $a \in A$.

We say that the action of A on V is **completely reducible** if V decomposes as a direct sum of irreducible subspaces.

A linear group $G \leq GL(V)$ is called **triangularizable** if there exists a basis of V with respect to which G is represented by upper-triangular matrices.

Actions of abelian groups

Lemma

If A is an abelian group acting irreducibly on V then V has dimension 1.

Proof. \mathbb{K} algebraically closed \Rightarrow every $a \in A$ has at least an eigenvalue. A abelian \Rightarrow the corresponding space of eigenvectors is b -invariant for every $b \in A$, hence it must coincide with V .

Thus, every $a \in A$ is a multiple of the identity map on V , hence by irreducibility V must have dimension 1. □

Proposition

If A is an abelian group acting on V then there exists a basis of V with respect to which A becomes upper triangular.

Actions of abelian groups, continued

Proof. By induction on the dimension of V . Obvious in dimension 1. Assume true for dimension $< n$, take V of dimension n .

If A acts irreducibly apply previous Lemma.

Assume A acts reducibly, and preserves a proper subspace $V' < V$.

We obtain two induced actions of A : on V' (by **restriction**) and on $V'' = V/V'$ (by **projection**).

Both actions become actions by triangular matrices with the right choice of basis.

The combination of the two bases yields a basis in V with respect to which A becomes upper triangular. □

Our goal in this lecture is to generalize this last result to **solvable groups**.

Burnside Theorem and applications

Theorem (Burnside's Theorem)

If $A \subset \text{End}(V)$ is a subalgebra which acts absolutely irreducibly on a finite-dimensional vector space V , then $A = \text{End}(V)$. In particular, if $G \leq \text{End}(V)$ is a subsemigroup acting irreducibly, then G spans $\text{End}(V)$ as a vector space, i.e. $\mathbb{K}[G] = \text{End}(V)$.

Theorem

Suppose that $G \leq \text{GL}_n(\mathbb{K})$ is irreducible and that

$$|\{\text{tr}(g) \mid g \in G\}| = q < \infty.$$

Then $|G| \leq q^{n^2}$.

Proof of first application to Burnside

Proof. By Burnside's Theorem, G contains $m = n^2$ linearly independent matrices $w(1), \dots, w(m)$.

For $\underline{\mu} \in k^m$ let

$$G(\underline{\mu}) = \{g \in G \mid \operatorname{tr}(w(s)g) = \mu_s \ (s = 1, \dots, m)\}.$$

Observe that $g = (g_{ij}) \in G(\underline{\mu})$ if and only if it satisfies the equations

$$\sum_{i=1}^n \sum_{l=1}^n w(s)_{il} g_{li} = \mu_s \ (s = 1, \dots, m).$$

This is a system of $m = n^2$ linearly independent equations, so it has at most one solution (g_{ij}) . The result follows as there are just q^{n^2} possibilities for $\underline{\mu}$. \square

Second application to Burnside. Nilpotent and unipotent.

Corollary

Suppose that $G \leq \mathrm{GL}_n(\mathbb{K})$ is completely reducible and that $g^k = 1$, $\forall g \in G$. Then $|G| \leq k^{n^3}$.

Proof. See Ex. Sheet 4.

