

# Numerical Solution of Partial Differential Equations

Endre Süli

Mathematical Institute  
University of Oxford  
2025

Lecture 3

## Finite difference approximation of a two-point b.v.p.

We illustrate the method of finite difference approximation on a simple two-point boundary-value problem for a second-order linear (ordinary) differential equation:

$$\begin{aligned} -u'' + c(x)u &= f(x), \quad x \in (0, 1), \\ u(0) &= 0, \quad u(1) = 0, \end{aligned} \tag{1}$$

where  $f$  and  $c$  are real-valued functions, which are defined and continuous on the interval  $[0, 1]$  and  $c(x) \geq 0$  for all  $x \in [0, 1]$ .

## The first step

The first step in the construction of a finite difference scheme for this boundary-value problem is to define the mesh.

Let  $N$  be an integer,  $N \geq 2$ , and let  $h = 1/N$  be the mesh-size; the mesh-points are  $x_i = ih$ ,  $i = 0, \dots, N$ .

We define the set of interior mesh-points:

$$\Omega_h := \{x_i : i = 1, \dots, N-1\}$$

the set of boundary mesh-points:

$$\Gamma_h := \{x_0, x_N\},$$

and the set of all mesh-points:

$$\overline{\Omega}_h := \Omega_h \cup \Gamma_h.$$

## The second step

Suppose that  $u$  is sufficiently smooth (e.g.  $u \in C^4([0, 1])$ ). Then, by Taylor series expansion,

$$\begin{aligned}u(x_{i\pm 1}) &= u(x_i \pm h) \\&= u(x_i) \pm hu'(x_i) + \frac{h^2}{2}u''(x_i) \pm \frac{h^3}{6}u'''(x_i) + \mathcal{O}(h^4),\end{aligned}$$

so that

$$D_x^+ u(x_i) := \frac{u(x_{i+1}) - u(x_i)}{h} = u'(x_i) + \mathcal{O}(h),$$

$$D_x^- u(x_i) := \frac{u(x_i) - u(x_{i-1})}{h} = u'(x_i) + \mathcal{O}(h),$$

and

$$\begin{aligned}D_x^+ D_x^- u(x_i) &= D_x^- D_x^+ u(x_i) \\&= \frac{u(x_{i+1}) - 2u(x_i) + u(x_{i-1}))}{h^2} \\&= u''(x_i) + \mathcal{O}(h^2).\end{aligned}$$

$D_x^+$  and  $D_x^-$  are called the **forward** and **backward first divided difference** operator, respectively, and  $D_x^+ D_x^-$  ( $= D_x^- D_x^+$ ) is called the (symmetric) **second divided difference** operator.

Thus we replace the second derivative  $u''$  in the differential equation by the second divided difference  $D_x^+ D_x^- u(x_i)$ ; hence,

$$\begin{aligned} -D_x^+ D_x^- u(x_i) + c(x_i)u(x_i) &\approx f(x_i), \quad i = 1, \dots, N-1, \\ u(x_0) &= 0, \quad u(x_N) = 0. \end{aligned} \tag{2}$$

Now (2) motivates us to seek the approximate solution  $U$  as the solution of the system of difference equations:

$$\begin{aligned} -D_x^+ D_x^- U_i + c(x_i)U_i &= f(x_i), \quad i = 1, \dots, N-1, \\ U_0 &= 0, \quad U_N = 0. \end{aligned} \tag{3}$$

This is a system of  $N - 1$  linear algebraic equations for the  $N - 1$  unknowns,  $U_i$ ,  $i = 1, \dots, N - 1$ . Using matrix notation,

$$AU = F,$$

where  $A$  is the  $(N - 1) \times (N - 1)$  matrix

$$A = \begin{bmatrix} \frac{2}{h^2} + c(x_1) & -\frac{1}{h^2} & & & & 0 \\ -\frac{1}{h^2} & \frac{2}{h^2} + c(x_2) & -\frac{1}{h^2} & & & \\ & \ddots & \ddots & \ddots & & \\ & & -\frac{1}{h^2} & \frac{2}{h^2} + c(x_{N-2}) & -\frac{1}{h^2} & \\ 0 & & & -\frac{1}{h^2} & \frac{2}{h^2} + c(x_{N-1}) & \end{bmatrix}$$

$$U = (U_1, U_2, \dots, U_{N-2}, U_{N-1})^T$$

and

$$F = (f(x_1), f(x_2), \dots, f(x_{N-2}), f(x_{N-1}))^T.$$

## Existence and uniqueness of a solution

We begin the analysis of the finite difference scheme (3) by showing that it has a unique solution. It suffices to show that the matrix  $A$  is non-singular (i.e.  $\det A \neq 0$ ), and therefore invertible.

We shall develop a technique which we shall, in subsequent sections, extend to the finite difference approximation of PDEs.

For this purpose, we introduce, for two functions  $V$  and  $W$  defined at the interior mesh-points  $x_i$ ,  $i = 1, \dots, N - 1$ , the inner product

$$(V, W)_h = \sum_{i=1}^{N-1} h V_i W_i,$$

which resembles the  $L_2((0, 1))$ -inner product

$$(v, w) = \int_0^1 v(x) w(x) dx.$$

The argument is based on mimicking, at the discrete level, the following procedure based on integration-by-parts, noting that the solution of the boundary-value problem (1) satisfies the homogeneous boundary conditions  $u(0) = 0$  and  $u(1) = 0$ :

$$\begin{aligned} \int_0^1 (-u''(x) + c(x)u(x)) u(x) dx &= \int_0^1 |u'(x)|^2 + c(x)|u(x)|^2 dx \\ &\geq \int_0^1 |u'(x)|^2 dx, \end{aligned} \tag{4}$$

because  $c(x) \geq 0$  for all  $x \in [0, 1]$ . Thus if, for example,  $f \equiv 0$  on  $[0, 1]$ , then  $-u'' + c(x)u \equiv 0$  on  $[0, 1]$ , and therefore by (4) also  $u' \equiv 0$  on  $[0, 1]$ . Consequently,  $u$  is a constant function on  $[0, 1]$ , but because  $u(0) = 0$  and  $u(1) = 0$ , necessarily  $u \equiv 0$  on  $[0, 1]$ . Hence, the only solution to the homogeneous boundary-value problem is the function  $u(x) \equiv 0$ ,  $x \in [0, 1]$ .



For the finite difference approximation of the boundary-value problem, if we can show by an analogous argument that the homogeneous system of linear algebraic equations corresponding to  $f(x_i) = 0$ ,  $i = 1, \dots, N - 1$ , has the trivial solution  $U_i = 0$ ,  $i = 0, \dots, N$ , as its unique solution, then the desired invertibility of the matrix  $A$  will directly follow.

Our key tool is a summation-by-parts identity, which is the discrete counterpart of the integration-by-parts identity

$$(-u'', u) = (u', u') = \|u'\|_{L_2((0,1))}^2 = \int_0^1 |u'(x)|^2 dx$$

satisfied by the function  $u$ , obeying the homogeneous boundary conditions  $u(0) = 0$ ,  $u(1) = 0$ , used in (4) above.

## Summation by parts identity

### Lemma

*Suppose that  $V$  is a function defined at the mesh-points  $x_i$ ,  $i = 0, \dots, N$ , and let  $V_0 = V_N = 0$ ; then,*

$$(-D_x^+ D_x^- V, V)_h = \sum_{i=1}^N h \left| D_x^- V_i \right|^2. \quad (5)$$

## PROOF.

By the definitions of  $(\cdot, \cdot)_h$  and  $D_x^+ D_x^- V_i$  we have that

$$\begin{aligned}(-D_x^+ D_x^- V, V)_h &= - \sum_{i=1}^{N-1} h (D_x^+ D_x^- V_i) V_i \\&= - \sum_{i=1}^{N-1} \frac{V_{i+1} - V_i}{h} V_i + \sum_{i=1}^{N-1} \frac{V_i - V_{i-1}}{h} V_i \\&= - \sum_{i=2}^N \frac{V_i - V_{i-1}}{h} V_{i-1} + \sum_{i=1}^{N-1} \frac{V_i - V_{i-1}}{h} V_i \\&= - \sum_{i=1}^N \frac{V_i - V_{i-1}}{h} V_{i-1} + \sum_{i=1}^N \frac{V_i - V_{i-1}}{h} V_i \\&= \sum_{i=1}^N \frac{V_i - V_{i-1}}{h} (V_i - V_{i-1}) = \sum_{i=1}^N h \left| D_x^- V_i \right|^2.\end{aligned}$$

In the transition to the 3rd line we shifted the index in the first sum; in the transition to the 4th line used that  $V_0 = V_N = 0$ .  $\square$

Returning to the finite difference scheme (3), let  $V$  be as in the above lemma and note that as, by hypothesis,  $c(x) \geq 0$  for all  $x \in [0, 1]$ , we have

$$\begin{aligned}(AV, V)_h &= (-D_x^+ D_x^- V + cV, V)_h \\ &= (-D_x^+ D_x^- V, V)_h + (cV, V)_h \\ &\geq \sum_{i=1}^N h \left| D_x^- V_i \right|^2.\end{aligned}\tag{6}$$

Thus, if  $AV = 0$  for some  $V$ , then  $D_x^- V_i = 0$ ,  $i = 1, \dots, N$ . Because  $V_0 = V_N = 0$ , this implies that  $V_i = 0$ ,  $i = 0, \dots, N$ . Hence  $AV = 0$  if and only if  $V = 0$ .

It therefore follows that  $A$  is a non-singular matrix, and thereby (3) has a unique solution,  $U = A^{-1}F$ .

We record this result in the next theorem.

### Theorem

*Suppose that  $c$  and  $f$  are continuous real-valued functions defined on the interval  $[0, 1]$ , and  $c(x) \geq 0$  for all  $x \in [0, 1]$ ; then, the finite difference scheme (3) possesses a unique solution  $U$ .*

## Stability, consistency, and convergence

Next, we investigate the approximation properties of the finite difference scheme (3). A key ingredient in our analysis is that the scheme (3) is stable (or discretely well-posed) in the sense that “small” perturbations in the data result in “small” perturbations in the corresponding finite difference solution.

To prove this, we define the *discrete  $L_2$ -norm*

$$\|U\|_h := (U, U)_h^{1/2} = \left( \sum_{i=1}^{N-1} h |U_i|^2 \right)^{1/2},$$

and the **discrete Sobolev norm**

$$\|U\|_{1,h} := (\|U\|_h^2 + \|D_x^- U\|_h^2)^{1/2},$$

where

$$\|V\|_h^2 := \sum_{i=1}^N h |V_i|^2.$$

Using this notation, the inequality (6) can be rewritten as follows:

$$(AV, V)_h \geq \|D_x^- V\|_h^2. \quad (7)$$

In fact, by employing a discrete version of the Poincaré–Friedrichs inequality, stated in the next lemma, we shall be able to prove that

$$(AV, V)_h \geq c_0 \|V\|_{1,h}^2,$$

where  $c_0$  is a positive constant, independent of  $h$ .

### Lemma (Discrete Poincaré–Friedrichs inequality)

*Let  $V$  be a function defined on the mesh  $\{x_i, i = 0, \dots, N\}$ , and such that  $V_0 = V_N = 0$ ; then, there exists a positive constant  $c_\star$ , independent of  $V$  and  $h$ , such that*

$$\|V\|_h^2 \leq c_\star \|D_x^- V\|_h^2 \quad (8)$$

*for all such  $V$ .*

PROOF. Thanks to the definition of  $D_x^- V_i$  and by use of the Cauchy–Schwarz inequality,

$$|V_i|^2 = \left| \sum_{j=1}^i h(D_x^- V_j) \right|^2 \leq \left( \sum_{j=1}^i h \right) \sum_{j=1}^i h |D_x^- V_j|^2 = ih \sum_{j=1}^i h |D_x^- V_j|^2.$$

Thus, because  $\sum_{i=1}^{N-1} i = \frac{1}{2}(N-1)N$  and  $Nh = 1$ , we have that

$$\begin{aligned} \|V\|_h^2 &= \sum_{i=1}^{N-1} h |V_i|^2 \leq \sum_{i=1}^{N-1} ih^2 \sum_{j=1}^i h |D_x^- V_j|^2 \\ &\leq \frac{1}{2}(N-1)Nh^2 \sum_{j=1}^N h |D_x^- V_j|^2 \\ &\leq \frac{1}{2} \|D_x^- V\|_h^2. \end{aligned}$$

We note that the constant  $c_\star = 1/2$  in the inequality (8).  $\square$



Using the inequality (8) to bound the right-hand side of the inequality (7) from below we obtain

$$(AV, V)_h \geq \frac{1}{c_\star} \|V\|_h^2. \quad (9)$$

Adding the inequality (7) to the inequality (9) we arrive at the inequality

$$(AV, V)_h \geq (1 + c_\star)^{-1} \left( \|V\|_h^2 + \|D_x^- V\|_h^2 \right).$$

Letting  $c_0 = (1 + c_\star)^{-1}$  it follows that

$$(AV, V)_h \geq c_0 \|V\|_{1,h}^2. \quad (10)$$

Now the stability of the finite difference scheme (3) easily follows.

### Theorem

*The scheme (3) is stable in the sense that*

$$\|U\|_{1,h} \leq \frac{1}{c_0} \|f\|_h. \quad (11)$$

PROOF. From (10) and (3) we have that

$$\begin{aligned} c_0 \|U\|_{1,h}^2 &\leq (AU, U)_h = (f, U)_h \leq |(f, U)_h| \\ &\leq \|f\|_h \|U\|_h \leq \|f\|_h \|U\|_{1,h}, \end{aligned}$$

and hence (11).  $\square$

Using this stability result it is easy to derive an estimate of the error between the exact solution  $u$ , and its finite difference approximation,  $U$ . We define the **global error**,  $e$ , by

$$e_i := u(x_i) - U_i, \quad i = 0, \dots, N.$$

Obviously  $e_0 = 0$ ,  $e_N = 0$ , and

$$\begin{aligned} Ae_i &= Au(x_i) - AU_i = Au(x_i) - f(x_i) \\ &= -D_x^+ D_x^- u(x_i) + c(x_i)u(x_i) - f(x_i) \\ &= u''(x_i) - D_x^+ D_x^- u(x_i), \quad i = 1, \dots, N-1. \end{aligned}$$

Thus,

$$\begin{aligned} Ae_i &= \varphi_i, & i &= 1, \dots, N-1, \\ e_0 &= 0, & e_N &= 0, \end{aligned} \tag{12}$$

where  $\varphi_i := u''(x_i) - D_x^+ D_x^- u(x_i)$  is the **consistency error** (sometimes also called the **truncation error**).

By applying ineq. (11) to the finite difference scheme (12):

$$\|u - U\|_{1,h} = \|e\|_{1,h} \leq \frac{1}{c_0} \|\varphi\|_h. \quad (13)$$

It remains to bound  $\|\varphi\|_h$ . We showed that, if  $u \in C^4([0, 1])$ , then

$$\varphi_i = u''(x_i) - D_x^+ D_x^- u(x_i) = \mathcal{O}(h^2),$$

i.e. there exists a positive constant  $C$ , independent of  $h$ , such that

$$|\varphi_i| \leq Ch^2, \quad i = 1, \dots, N-1.$$

Consequently, we have proved **consistency**:

$$\|\varphi\|_h = \left( \sum_{i=1}^{N-1} h |\varphi_i|^2 \right)^{1/2} \leq Ch^2. \quad (14)$$

Combining the inequalities (13) and (14), it follows that

$$\|u - U\|_{1,h} \leq \frac{C}{c_0} h^2. \quad (15)$$

In fact, a more careful treatment of the remainder term in the Taylor series expansion reveals that

$$\varphi_i = u''(x_i) - D_x^+ D_x^- u(x_i) = -\frac{h^2}{12} u^{IV}(\xi_i), \quad \xi_i \in [x_{i-1}, x_{i+1}].$$

Thus

$$|\varphi_i| \leq h^2 \frac{1}{12} \max_{x \in [0,1]} |u^{IV}(x)|, \quad i = 1, \dots, N-1,$$

and hence

$$C = \frac{1}{12} \max_{x \in [0,1]} |u^{IV}(x)|$$

in inequality (14). Recalling that  $c_0 = (1 + c_\star)^{-1}$  and  $c_\star = 1/2$ , we deduce that  $c_0 = 2/3$ . Substituting the values of the constants  $C$  and  $c_0$  into inequality (15) it follows that

$$\|u - U\|_{1,h} \leq \frac{1}{8} h^2 \|u^{IV}\|_{C([0,1])}.$$

Thus we have proved the following result.

### Theorem

*Let  $f \in C([0, 1])$ ,  $c \in C([0, 1])$ , with  $c(x) \geq 0$  for all  $x \in [0, 1]$ , and suppose that the corresponding (weak) solution of the boundary-value problem (1) belongs to  $C^4([0, 1])$ ; then*

$$\|u - U\|_{1,h} \leq \frac{1}{8} h^2 \|u^{IV}\|_{C([0,1])}. \quad (16)$$

In other words,

*stability + consistency  $\Rightarrow$  convergence.*