

C2.6 Introduction to Schemes

Prof. Alexander F. Ritter
University of Oxford
HT 2020 - HT 2022
ritter@maths.ox.ac.uk

Feedback and corrections are welcome!

References

2018-2019 Course Lecture Notes by Prof. Damian Rössler ← on course page
Ravi Vakil, The Rising Sea, Foundations of Algebraic Geometry ← online
http://stacks.math.columbia.edu ← Search defs, theorems/proofs in algebra & alg-geomtry
Qing Liu, Algebraic Geometry and Arithmetic Curves, OUP 2002 ← modern book, seems rather nice
Eisenbud & Harris, The Geometry of Schemes, Springer GTM 197 ← classic
George R. Kempf, Algebraic Varieties, LMS Lecture notes 172
Classic books by: Mumford (Red Book of Varieties & Schemes)
Hartshorne (Algebraic Geometry)

Shafarevich (Basic Algebraic Geometry 2) ← or my website

My C3.4 Algebraic geometry notes (see C2.6 course webpage) try to fill the gap between classical algebraic geometry (C3.4) and C2.6
For the brave, you can look at the original works by the masters in French: Grothendieck, "Éléments de géométrie algébrique" series on www.numdam.org
Serre, "Faisceaux Algébriques Cohérents", Annals of Math. 1955.

Prerequisites

Commutative algebra (e.g. Atiyah - MacDonald, Introduction to Comm. Alg.)
Category theory - or willingness to read things up as necessary
Homological algebra - or willingness to read things up as necessary

Expectations

That you read the notes regularly after each class.
(This is a 16-lecture course, 2 lectures/week across 8 weeks.)
Not everything can be covered in detail in class, so you need to be willing to look things up as necessary.

Conventions

Diagrams commute unless we say otherwise
Ring means commutative ring with unit 1
Ring homomorphisms are by definition unital i.e. 1 maps to 1

Arrows:
← means injective
→ means surjective

CONTENTS

0. INTRODUCTION

- 0.1 Classical Algebraic Geometry: Affine varieties
- 0.2 Why schemes?
- 0.3 What is a point? (reducible, irreducible)

1. DEFINITION OF SCHEMES

- 1.1 Examples of affine schemes (Spec R , $V(I)$) generic/closed point, covering trick, quasi-compact (ringed space, locally ringed space, affine scheme, scheme)
- 1.2 Definition of a scheme (pre-sheaf, morph of presheaves, sub-presheaf)
- 1.3 Pre-sheaves (sheaf, local-to-global condition, skyscraper sheaf, $\mathcal{A}_k(X)$)
- 1.4 Sheaves (stalk, direct limits, checking inj/surj at stalk level)
- 1.5 Stalks (sheafification $F^\#$, universal property of $F^\#$)
- 1.6 Sheafification (abelian categories, additive categories, additive functor)
- 1.7 Kernels, cokernels, images (cochain complex/cohomology in abelian cats, left/right exact)
- 1.8 Exactness (sheaf image $(F_\#, F, F^{-1}, F|_U, \Gamma(F, U))$, adjointness of $F_\#$ & F^{-1})
- 1.9 Push-forward (direct image) and inverse image (max ideals in local rings \leftrightarrow points)
- 1.10 Morphisms of ringed spaces (A sheaf defined on a topological basis (B-sheaf, inverse limits, extending morphs defined on basis) (Using $B = \{D_+^i\}$ for Spec R , structure sheaf \mathcal{O}_X classical alg. geom.) (Spec: Rings \leftrightarrow equivalence Aff. faithfully Locally ringed spaces)
- 1.11 A sheaf defined on a topological basis (max ideals in local rings \leftrightarrow points)
- 1.12 Construction of $\mathcal{O}_{\text{Spec } R}$ (Using $B = \{D_+^i\}$ for Spec R , structure sheaf \mathcal{O}_X classical alg. geom.)
- 1.13 Morphisms between Specs (Spec: Rings \leftrightarrow equivalence Aff. faithfully Locally ringed spaces)
- 1.14 Closed affine subschemes (ideal sheaf for $I \subseteq R$ on Spec R , quasi-coherence)
- 1.15 Closed subschemes (sheaf of ideals on a scheme, quasi-coherence, support of a sheaf)

2. GLOBAL SECTIONS AND THE FUNCTOR OF POINTS

- 2.0 Points of Spec R (not necessarily closed) (max ideals in local rings \leftrightarrow points)
- 2.1 Global sections and basic open sets for locally ringed spaces (X canonical, Spec $\Gamma(X, \mathcal{O}_X)$, $D_+(f)$)
- 2.2 What it means to be affine (Yoneda lemma/embedding, $\text{Mor}(X, \text{Spec } R) \cong \text{Hom}(R, \Gamma(X, \mathcal{O}_X))$)
- 2.3 Functor of points h_Y

3. PROPERTIES OF SCHEMES

- 3.0 Useful facts from commutative algebra: localisation (localisation of modules exactness)
- 3.1 Noetherian (locally Noetherian schemes, useful trick: basis \subseteq overlap of affines)
- 3.2 Properties that are affine-local (locality of finite type, reduced, Noetherian)
- 3.3 Reduced schemes (stalk-local property, extending morphisms onto closures)
- 3.4 Irreducible schemes (nilradical as generic point, connectedness, irred. components, primary decomp.)
- 3.5 Integral schemes (integral \leftrightarrow reduced & irreducible, injectivity of restrictions, function field $K(X)$)
- 3.6 Properties of morphisms (affine quasi-compact, locally finite type, finite type, closed/open immersion, closed/open subschemes, flat, flatness & deformations, closeness in Spec R)

4. GLUING THEOREMS

- 4.1 Gluing sheaves (gluing data, compatibility conditions, morphisms defined by local data)
- 4.2 Gluing schemes (gluing conditions, gluing lemma, functor of points is a sheaf of sets)
- 4.3 Affine n -space by gluing (see Homework for projective space) (\mathbb{A}^n and \mathbb{P}^n as representable functors)

5. PRODUCTS

- 5.0 Products in category theory (product, coproduct, category \mathcal{C}/B , fiber product, pushout)
- 5.1 Fiber products exist in Schemes/ B (A -algebras, tensor products, fiber products in Aff & Sch)
- 5.2 Fibers and preimages (Mumford's picture, underlying topological space of products)
- 5.3 Base change (separated, universally closed, proper, projective morphism)
- 5.4 More properties of schemes (abstract varieties, complete, affine and quasi-projective vars)
- 5.5 Varieties (induced scheme structure, locally closed subsets)
- 5.6 Scheme structure on subsets (induced scheme structure, locally closed subsets)

Conventions

Diagrams commute unless we say otherwise
Ring means commutative ring with unit 1
Ring homomorphisms are by definition unital i.e. 1 maps to 1

0.1 Classical Algebraic Geometry : Affine varieties

$R = k[x_1, \dots, x_n]$ polynomial ring over algebraically closed field k

$I \subseteq R$ ideal

$X = V(I) = \{a \in k^n : f(a) = 0 \forall f \in I\}$ affine variety

The topological space

Affine space: $\mathbb{A}^n = k^n$ with Zariski topology: $\left\{ \begin{array}{l} \text{closed sets: } V(I) \\ \text{open sets: } U_I = \mathbb{A}^n \setminus V(I) \end{array} \right.$

$X \subseteq \mathbb{A}^n$ subspace topology: $X \cap U_I$

basis of open sets: $D_f = \{a \in k^n : f(a) \neq 0\}, f \in R$

The functions on it

$R \cong \text{Hom}(\mathbb{A}^n, \mathbb{A}^1), f \mapsto (a \mapsto f(a))$

$\mathbb{I}(X) = \{f \in R : f(X) = 0\}$

Remark $V(\mathbb{I}(X)) = X$ for affine varieties X

Coordinate ring: $k[X] = R/\mathbb{I}(X)$

Key facts: 1) Hilbert's basis theorem: R Noetherian, so $k[X]$ Noetherian

2) Hilbert's weak nullstellensatz: Maximal ideals of R (and of $k[X]$) are $m_a = \mathbb{I}(\{a\}) = \langle x_1 - a_1, \dots, x_n - a_n \rangle$, so correspond to points: $\{a\} = V(m_a)$

3) Hilbert's Nullstellensatz: $\mathbb{I}(V(I)) = \sqrt{I}$ (radical of I)
 Hence: $\mathbb{I}(V(\mathbb{I}(X))) = \sqrt{\mathbb{I}(X)} = \mathbb{I}(X)$

Lemma There are enough functions to separate points

Pf $a \neq b \in X \subseteq \mathbb{A}^n \Rightarrow$ some coordinate $a_i \neq b_i \Rightarrow x_i \in k[X]$ separates a, b

Morphisms between affine varieties

$\text{Hom}(\mathbb{A}^n, \mathbb{A}^m) \cong R^m \leftarrow$ polynomial maps $a \mapsto (f_1(a), \dots, f_m(a))$

$\text{Hom}(X, Y) = \{ \text{restriction of a polynomial map } \mathbb{A}^n \rightarrow \mathbb{A}^m \text{ s.t. } X \rightarrow Y \}$

Facts: 1) $k[X] \cong \text{Hom}(X, \mathbb{A}^1) \leftarrow$ "values of functions are enough to determine the abstract function"

2) $\text{Hom}(X, Y) \cong \text{Hom}_{k\text{-alg}}(k[X], k[Y])$

$(F: X \rightarrow Y) \mapsto (F^*: \text{Hom}(Y, \mathbb{A}^1) \rightarrow \text{Hom}(X, \mathbb{A}^1)) \leftarrow$ "pullback"

$f \mapsto F^*f = f \circ F$

Equivalence of categories

$\{\text{affine varieties}\} \longleftrightarrow \{\text{finitely generated reduced } k\text{-algebras } \Delta \text{ homs of } k\text{-algs.}\}$

$X \longmapsto k[X]$ (no nilpotents)

$(F: X \rightarrow Y) \longmapsto F^*$ (f nilpotent $\iff f^* = 0$ some N)

Remark The "same" (up to isomorphism) X can be embedded in various \mathbb{A}^n .

E.g. cuspidal cubic $V(y^2 - x^3) = \mathbb{A}^2_{x,y} \subseteq \mathbb{A}^3_{x,y,z}$ is $\cong V(y^2 - x^3, z - x) \subseteq \mathbb{A}^3_{x,y,z}$

6. SHEAVES OF MODULES

- 6.1 \mathcal{O}_X -modules
- 6.2 Modules generated by sections
- 6.3 Vector bundles and coherent modules (locally free, invertible sheaf, coherent, loc. finitely presented)
- 6.4 \mathcal{O}_X -module \mathcal{F} on $X = \text{Spec } R$, for R -mod $M \leftarrow R\text{-Mods} \rightarrow \text{Spec } R\text{-Mods}$ fully faithful exact
- 6.5 Direct image and inverse image $(\xi_* F, f^{-1} F)$
- 6.6 Operations on \mathcal{O}_X -mods $(\text{Hom}_{\mathcal{O}_X}(\mathcal{F}, \mathcal{G}), \otimes \mathcal{F}, F \otimes_{\mathcal{O}_X} \mathcal{G})$
- 6.7 Pullback $(f^* F, \text{adjointness of } f_*$ and $f^*)$
- 6.8 \mathcal{F} on any scheme $(f^* \mathcal{F}$ vs. changing rings)
- 6.9 Classification of \mathcal{O}_X -homs $\mathcal{F} \rightarrow \mathcal{F} \leftarrow \text{Hom}_{\mathcal{O}_X}(\mathcal{F}, \mathcal{F}) = \text{Hom}_{\mathcal{O}_X}(M, \Gamma(X, F))$ on $X = \text{Spec } R$
- 6.10 Flatness $(f: X \rightarrow Y \text{ flat} \iff f^*: \mathcal{O}_Y\text{-mod} \rightarrow \mathcal{O}_X\text{-mod}$ exact, flat resolutions)

7. (QUASI-) COHERENT SHEAVES

- 7.1 $\mathcal{Q}\text{Coh}(X)$ (locally finitely presented vs. coherence, coherent modules)
- 7.2 Overview of general properties of $\mathcal{Q}\text{Coh}(X)$ and $\text{Coh}(X)$ for X scheme
- 7.3 Pull-back preserves quasi-coherence
- 7.4 Push-forwards for X Noetherian
- 7.5 Gluing modules
- 7.6 $\mathcal{Q}\text{Coh}(X), \text{Coh}(X), \text{Vect}(X)$ for $X = \text{Spec } R \leftarrow R\text{-Mods} \cong \mathcal{Q}\text{Coh}(\text{Spec } R), \text{Coh } R\text{-Mods} = \text{Coh}(\text{Spec } R)$ (cocycle condition, gluing lemma)

8. ČECH COHOMOLOGY

- 8.1 Čech complex $(\check{C}^i(U); \check{C}$ Čech differential, $\check{H}^n(X, F)$, chain map, chain homotopy (Serre's trick))
- 8.2 Čech complex with ordering
- 8.3 Affines have no cohomology except $H^0 \leftarrow \check{H}^n(\text{Spec } R, F) = 0 \forall n \geq 1$ for $F \in \mathcal{Q}\text{Coh}(X)$
- 8.4 Independence of cover $(X \text{ separated } \& \text{ quasi-compact} \Rightarrow \check{H}^i(U; \cdot)$ indep. of cover for $\mathcal{Q}\text{Coh}$)
- 8.5 Induced LES on H $(\Gamma(U, \cdot)$ exact on $\mathcal{Q}\text{Coh}$ for affine U)
- 8.6 Dealing with infinite covers (refinements of covers, H^* vs. singular cohomology)
- 8.7 Application: line bundles and $\check{H}^1(X, \mathcal{O}_X^*)$ (trivialization, vector bundles, sheaf \mathcal{O}_X^* of invertible \mathcal{O}_X)
- 8.8 Divisors (Picard group, $\text{Pic}(\mathbb{P}^1), \text{Pic}(\mathbb{P}^n)$)
- 8.9 Čech cohomology computations on \mathbb{P}^n (Cartier divisor vs line bundle, Weil divisors)
- 8.10 Product on Čech cohomology $(H^*(\mathbb{P}^n, \mathcal{O}(d)), \mathcal{O}(d))$ for $d \in \mathbb{Z}$

9. SHEAF COHOMOLOGY

- 9.1 Resolutions (injective/projective, left/right-derived functors, "enough injectives")
- 9.2 Acyclic resolutions
- 9.3 Čech cohomology vs Sheaf cohomology (characterization of \check{H}^i (separated quasi-compact schemes) for $\mathcal{Q}\text{Coh}$, separated Noeth. $\Rightarrow \check{H}^i = H^i$ on $\mathcal{Q}\text{Coh}$, Serre's Theorem)
- 9.4 Product on sheaf cohomology

10. $\mathcal{Q}\text{Coh}(\mathbb{P}^n)$, GRADED MODULES, $\text{PROJ}(R)$

- 10.1 Graded modules and $\mathcal{Q}\text{Coh}(\mathbb{P}^n) \leftarrow$ (graded rings/mods, Graded $k[x_0, \dots, x_n]$ Mods $\xrightarrow{\text{full \& faithful}} \mathcal{Q}\text{Coh}(\mathbb{P}^n)$ (line bundles via graded mods)
- 10.2 $\text{Proj}(R)$ and $\mathcal{Q}\text{Coh}(\text{Proj } R) \leftarrow$ ($\text{Proj } R$, irrelevant ideal, $V(\text{graded ideal}), \mathcal{O}_{\text{Proj}(R)}$, $(\mathbb{P}^n = \text{Proj } k[x_0, \dots, x_n], \text{Graded } R\text{-Mods} \xrightarrow{\text{exact full \& faithful}} \mathcal{Q}\text{Coh}(\text{Proj } R))$)

0.2 Why schemes?

Some reasons:

- 1) Why always have spaces embedded in \mathbb{A}^n ? (extrinsic)
Can you make sense of X without reference to \mathbb{A}^n ? (intrinsic)
- 2) Why not let R be any ring?
- 3) When you deform varieties, nilpotents arise naturally and should not be ignored:

$f = (x-a) \cdot (x-b)$
 $X = \mathbb{V}(f) = \{a, b\} \subseteq \mathbb{A}^1 \leftarrow$ two points
 $k[X] \cong k[x]/(x-a) \oplus k[x]/(x-b) \cong k^2 \leftarrow$ a value at each point
 Deform: a, b become 0 :
 $f = (x-0) \cdot (x-0) = x^2$
 $X = \mathbb{V}(f) = \{0\} \subseteq \mathbb{A}^1$
 $k[X] \cong k[x]/\sqrt{(x^2)} = k[x]/(x) \cong k$
 notice $k[X]$ is the reduced ring, not $k[X]/(x^2)$

We lost information: classically you cannot tell $x=0$ apart from $x^2=0$. In the theory of schemes, the key role is not played by the topological space. The key role is played by the ring of functions, or rather, the sheaf of functions \mathcal{O} : on each open set $U \subseteq X$ get a ring of functions $\mathcal{O}(U)$.

Example above: $\mathcal{O}(X) = k[x]/(x^2)$ \leftarrow we do not reduce the ring of functions
 At what cost? Values of functions need not determine the abstract function:
 $\mathcal{O}(X) \ni \alpha + \beta x \mapsto (\alpha + \beta x : X = \{0\} \rightarrow \mathbb{A}^1) \in \text{Hom}(X, \mathbb{A}^1)$
 $0 \mapsto \alpha$ do not recover β .
 Idea: the abstract " β " remembers that X arose from the collision of two points, so β records tangential information: $\frac{\partial}{\partial x} \Big|_{x=0} (\alpha + \beta x) = \beta$.

0.3 What is a point?

X topological space is reducible if $X = X_1 \cup X_2$ for proper closed $X_i \subseteq X$.
 Euclidean world (more generally if X Hausdorff): $Y \subseteq X$ irreducible $\Leftrightarrow Y = \text{point}$ or $Y = \emptyset$
 Classical Alg. Geom. \leftarrow point $a \in X \Leftrightarrow$ max ideal $m_a \subseteq k[X]$
 closed $\emptyset \neq Y \subseteq X$ irreducible $\Leftrightarrow \Pi(Y) \subseteq k[X]$ prime ideal

R ring \Rightarrow "points" of R are $\text{Spec}(R) = \{\text{prime ideals of } R\}$ not just max ideals
 Categorically a good choice since functorial:
 $\varphi: R \rightarrow S$ hom of rings $\Rightarrow \varphi^{-1}(\text{prime ideal}) = \text{prime ideal}$
 $\Rightarrow \text{Spec } S \xrightarrow{\varphi^{-1}} \text{Spec } R$
 fails for max ideals e.g. $\mathbb{Z} \xrightarrow{\varphi} \mathbb{Q}, \varphi^{-1}(0) = 0$
 We were just lucky that homs $k[X] \rightarrow k[X]$ send max ideal \rightarrow max ideal.

1. DEFINITION OF SCHEMES

1.1 Examples of affine schemes

$\text{Spec}(R)$ some ring R (always: comm. ring with 1)
 As a set: $\text{Spec}(R) = \{\text{prime ideals } P \subseteq R\} \leftarrow$ (prime) Spectrum
 Zariski topology: e.g. $V(R) = \emptyset$
 $V(0) = \text{Spec } R$

closed sets: $\mathbb{V}(I) = \{\text{prime ideals containing } I\} \subseteq \text{Spec } R$
 which we construct later.
 sheaf $\mathcal{O}_{\text{Spec } R}$

The global functions are: $\mathcal{O}_{\text{Spec } R}(\text{Spec } R) = R$.
 so spaces of frns can recover the top space!
 Key exercise \Rightarrow axioms for a topology
 $V(I) \cup V(J) = V(I \cdot J) = V(I \cap J)$
 $V(I) \cap V(J) = V(\sum I_i)$
 so $\sqrt{I \cdot J} = \sqrt{I \cap J}$ but $I \cdot J$ and $I \cap J$ may be \neq

Key: $V(I) = \emptyset \Leftrightarrow I = R \Leftrightarrow 1 \in I$, since any proper ideal \subseteq some maximal ideal
 Topological consequences:
 open sets: $U_I = \text{Spec } R \setminus \mathbb{V}(I) = \bigcup_{f \in I} D_f$
 basis of open sets: $D_f = \{P \in \text{Spec } R : f \notin P\}$
 $= \{P \in \text{Spec } R : f(P) \neq 0\}$

"value of $f \in R$ at P ": $R \rightarrow R/P \xrightarrow{f} k(P) = \text{Frac}(R/P) \xrightarrow{f} f(P)$
 localisation of R at P
 Remarks: P prime, R/P is integral domain

Remark: $f(P) = 0 \Leftrightarrow f \in P$
 Examples 1) $R = k[X] \leftarrow$ affine variety $X \subseteq \mathbb{A}^n$
 $\text{Spec } R \xrightarrow{\text{bijection}} \text{irreducible subvarieties } Y \subseteq X$
 $\text{Spec } m \leftarrow$ (max. ideals) $X \leftarrow$ and Zariski: topologies agree

Value of $f \in R$ at m_a : $m_a \rightarrow R/m_a \cong k \xrightarrow{f} f(a)$
 $(m_a = \langle x_1 - a_1, \dots, x_n - a_n \rangle)$
 in this case the target field does not depend on the point
 2) $\text{Spec } \mathbb{Z} = \{0\} \cup \{P : P \in \mathbb{N} \text{ prime}\}$
 value of $f \in \mathbb{Z}$ at (0) : $\mathbb{Z} \rightarrow \text{Frac}(\mathbb{Z}/0) = \mathbb{Q} \xrightarrow{f} f$
 so lost no information.

$\mathbb{V}(0) = \{\text{prime ideals containing } (0)\} = \text{Spec } \mathbb{Z}$ so the point (0) is dense!
 $\mathbb{V}(P) = \{P\}$ are "closed points". Value of $f \in \mathbb{Z} : f(P) = (f \in \mathbb{Z}/P) = (f \text{ mod } P)$
 In general Prime ideals P with $\mathbb{V}(P) = \text{Spec } R$ are called generic points
 Prime ideals P with $\mathbb{V}(P) = \{P\}$ are called closed points
 Exercise: $\{\text{closed points}\} = \{\text{max ideals of } R\}$

Motivation: M $n \times n$ matrix over \mathbb{C}
 Then $\mathbb{C}[x] \rightarrow \mathbb{C}[M], x \mapsto M$ has $\text{Ker} = \langle m \rangle$
 so $\mathbb{C}[M] \cong \mathbb{C}[x]/\langle m \rangle \cong \mathbb{C}[\lambda]/(x-\lambda)^n$
 $\text{Spec } \mathbb{C}[M] = \{(\lambda, \lambda) : \lambda \text{ eigenvalue of } A\}$

Exercises • a prime ideal \Rightarrow a radical $(a = \sqrt{a})$

For a, b radical, $a \subseteq b \Leftrightarrow V(a) \supseteq V(b)$ ← order reversing!

Cor $V(I) \subseteq V(J) \Leftrightarrow \sqrt{I} \supseteq \sqrt{J}$

Pf $V(I) = V(\sqrt{I})$, so: $V(\sqrt{I}) \subseteq V(\sqrt{J}) \Leftrightarrow \sqrt{I} \supseteq \sqrt{J}$ by exercise. \square

Cor $V(a) = V(b) \Leftrightarrow \sqrt{a} = \sqrt{b}$

\Rightarrow [closed sets of $\text{Spec } R$] $\xleftrightarrow{\text{order-reversing correspondence}}$ [radical ideals of R]

Proposition $f \in R$ vanishes at all $p \in \text{Spec } R \Leftrightarrow f$ nilpotent

Covering Trick $\text{Spec } R = \cup D_{f_i} \Leftrightarrow 1 \in \langle \text{all } f_i \rangle \Leftrightarrow \langle \text{all } f_i \rangle = R$

Pf $\text{Spec } R \setminus \cup D_{f_i} = \cap V(f_i) = V(\langle \text{all } f_i \rangle)$, now use previous key. \square

Theorem $\text{Spec } R$ is quasi-compact \leftarrow (quasi-compact = compact = open covers have finite subcovers)

Pf $\text{Spec } R = \cup_i U_i$. As $U_i = \cup_j D_{f_{ij}}$, wlog $U_i = D_{f_i}$.

Trick $\Rightarrow 1 = \sum_{\text{finite}} r_i f_i \leftarrow$ so finitely many f_i generate R , so those D_{f_i} cover. \square

Basic Exercises

1) $\varphi: R \rightarrow S$ ring hom $\Rightarrow \alpha: \text{Spec } S \rightarrow \text{Spec } R, p \mapsto \varphi^{-1}(p)$ is continuous

indeed $\alpha^{-1}(D_f) = D_{\varphi(f)} \leftarrow$ (Hint: $f \notin p \Leftrightarrow \varphi(f) \notin \varphi(p)$ has $\varphi(p) \in \alpha^{-1}(p)$)

2) Show that $\text{Spec}(R/I)$ "is" the subspace $V(I) \subseteq \text{Spec } R$ and the quotient

map $\pi: R \rightarrow R/I$ induces via (1) the inclusion map on Specs.

Example $\text{Spec}(R/(f)) = \{ \text{prime ideals of } R \text{ containing } f \}$
 $= \text{the points of } \text{Spec } R \text{ where } f \text{ vanishes}$
 $= V(f)$

3) Show that $\text{Spec}(S^{-1}R)$ "is" a subspace of $\text{Spec } R$, where $S^{-1}R$ is localisation of R at a multiplicative set $S \subseteq R$, and $R \rightarrow S^{-1}R, r \mapsto \frac{r}{1}$ induces via (1) the inclusion

Example $S = \{1, f, f^2, \dots\}$, so $S^{-1}R = R_f$, then:

$\text{Spec } R_f = \{ \text{prime ideals of } R \text{ not containing } f \}$
 $= \text{the points of } \text{Spec } R \text{ where } f \text{ does not vanish}$
 $= D_f$

4) $D_f \cap D_g = D_{fg}$, so $\text{Spec } R_f \cap \text{Spec } R_g = \text{Spec } R_{fg}$

5) $D_f \subseteq D_g \Leftrightarrow V(f) \supseteq V(g) \Leftrightarrow \forall f \in \sqrt{g} \Leftrightarrow f \in (g)$ some $n \Leftrightarrow g \in R_f$ invertible

6) $p \subseteq R$ prime ideal $\Rightarrow R_p = S^{-1}R$ for $S = R \setminus p$, then $\exists!$ closed point $m_p = p \in R_p \subseteq \text{Spec } R_p$
 so local ring: $\exists!$ max ideal $m(\Leftrightarrow$ elts outside m are invertible)

Also: $m_p \in U \subseteq \text{Spec } R_p$ open $\Rightarrow U = \text{Spec } R_p$.

1.2 Definition of a scheme

Def A ringed space is

- a topological space X
 - with a sheaf of rings \mathcal{O}_X on X
- Locally ringed space if also:
- all stalks $\mathcal{O}_{X,x}$ are local rings
 - (so \exists unique maximal ideal $\mathfrak{m}_{X,x} \subseteq \mathcal{O}_{X,x}$ and \exists residue field at $x: k(x) = \mathcal{O}_{X,x} / \mathfrak{m}_{X,x}$)

Def An affine scheme is a locally ringed space for some ring R . isomorphic to $(\text{Spec } R, \mathcal{O}_{\text{Spec } R})$

Def A scheme is a locally ringed space which is locally isomorphic to an affine scheme.

means:

$\forall x \in X \exists$ some open neighbourhood $x \in U \subseteq X$ s.t. $(U, \mathcal{O}_X|_U) \cong (\text{Spec } R, \mathcal{O}_{\text{Spec } R})$

1.3 Pre-sheaves

Ab = category of abelian groups and group homs

X = any topological space

$\text{Top } X$ = category with objects: open sets $U \subseteq X$ morphs: inclusion maps

Def A presheaf (of abelian groups) on X is a contravariant functor $F: \text{Top } X \rightarrow \text{Ab}$

So: \forall open $U \subseteq X$ have an abelian group $F(U)$ ← elements called sections (over U)

\forall inclusion $U \rightarrow V$ have a "restriction" group hom $F(V) \rightarrow F(U)$

$F(\text{id}: U \rightarrow U): F(U) \xrightarrow{\text{id}} F(U)$ so $s|_U = s$ for $s \in F(U)$.

$U \subseteq V \subseteq W \Rightarrow F(W) \rightarrow F(V) \rightarrow F(U)$ so: $(s|_V)|_U = s|_U$ for $s \in F(W)$.

Example X topological space, $F(U) = \{ \text{continuous functions } U \rightarrow \mathbb{R} \}$ with obvious restrictions Morphism of pre-sheaves = natural transformation of such functors: $\varphi: F \rightarrow G$

So: \forall open $U \subseteq X$ have $\varphi_U: F(U) \rightarrow G(U)$ group hom

\forall inclusion $U \rightarrow V$ have $F(U) \xrightarrow{\varphi_U} G(U) \xrightarrow{\varphi_V} G(V)$ ← restriction homs

Sub pre-sheaf $F \subseteq G$ means $F(U) \subseteq G(U)$ subgt, compatibly with restrictions

RED: WORDS TO BE DEFINED LATER

IDEA

- ← the points
- ← the functors
- ← the germs of functions near point x
- ← the "value" of a function at x lives here

if use category \mathcal{C} get (pre)sheaves with values in \mathcal{C} e.g. $\mathcal{C} = \text{Rings}$ get presheaf of rings

$(\text{Mor}(U, V) = \{ \emptyset \text{ if } U \not\subseteq V \})$ (incl. if $U \subseteq V$)

$F(V) \rightarrow F(U)$
 $s \mapsto s|_U$

so the homs // are compatible with restrictions

i.e. this diagram with $\varphi_U = \text{inclusion}$


Fact $Ab(X)$ is an **abelian category**
 idea: it "behaves like" category of abelian grps

Fact In additive cat, $\text{mono} \Leftrightarrow H \rightarrow F \rightarrow G$ then $H \rightarrow F$ epi $\Leftrightarrow F \rightarrow G \rightarrow H$ then $G \rightarrow H$ categorical Ker & Coker, see below

Def **abelian category** = **additive category** such that morphisms have **Ker, Coker** and i) $\varphi: F \rightarrow G$ monomorph is the Ker of its Coker ii) $\varphi: F \rightarrow G$ epimorph is Coker

Def **additive category** means $\text{Mor}(A, B)$ abelian gr (so often write $\text{Hom}(A, B)$) s.t.
 • Composition of morphisms distributes over addition
 • \exists products $A \times B$ ($\forall \text{ obj. } X, (\exists! \text{ morph } X \rightarrow 0)$)
 • \exists zero object 0 (an object that is both **initial** & **terminal**)

Functor F of additive/abelian cats is additive if $\text{Hom}(A, B) \rightarrow \text{Hom}(FA, FB)$ is gp. hom.



Example For abelian grps, (i) says: $\text{Ker } \pi = \frac{A}{\langle \text{Im } \pi \rangle} \rightarrow B/A$ as expected!

I will now stop underlining Ker, Coker, Im.
RMK These categorical definitions can be cumbersome to work with. It turns out: \forall small abelian category \mathcal{A} , \exists a possibly non-commutative ring R with 1 and full faithful exact functor $\mathcal{A} \rightarrow \{\text{left } R\text{-modules}\}$ (in particular preserves $(\text{Obj}(\mathcal{A}) \text{ and } \text{Homs})$ \Rightarrow can "pretend" you work with modules. (example you just apply the theorem to the small abelian subcategory involved in your diagram/sequence of maps - don't need to use the whole category))

1-8 Exactness
 A (cochain) complex $F^\bullet = (\dots \rightarrow F^{i-1} \xrightarrow{d^{i-1}} F^i \xrightarrow{d^i} F^{i+1} \rightarrow \dots)$ in an abelian cat means composite of two consecutive morphs is zero: $d^{i+1} \circ d^i = 0 \quad \forall i$

(Co)homology $H^i(F^\bullet) = \text{Ker } d^{i+1} / \text{Im } d^i$
 F^\bullet exact means $\text{Im } d^i = \text{Ker } d^{i+1}$ (\Leftrightarrow complex with zero homology $H^i = 0$)
 (\exists mono $\text{Im } d^i \hookrightarrow \text{Ker } d^{i+1}$ (and H^i is its Coker))

Proposition complex F^\bullet in $Ab(X)$ exact $\Leftrightarrow F^\bullet$ is exact sequence of abelian grps $\forall X \in X$
 (mediate by Facts on previous page)

RMK For SES (short exact sequences) $0 \rightarrow F \xrightarrow{\alpha} G \xrightarrow{\beta} H \rightarrow 0$ of sheaves you usually check exactness at level of stalks, but can equivalently check:

- i) $0 \rightarrow F(U) \rightarrow G(U) \rightarrow H(U)$ exact \forall open U
- ii) H is smallest subsheaf containing $\text{pre-Im } \beta$, meaning every section of H can be obtained by gluing local sections of type $\beta(\frac{\text{local section}}{\text{open}})$

A functor of abelian cats is **left exact** if: $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ exact $\Rightarrow 0 \rightarrow FA \rightarrow FB \rightarrow FC$ exact
right exact if $\Rightarrow FA \rightarrow FB \rightarrow FC \rightarrow 0$ exact

Example $\text{Hom}_R(M, \cdot)$ is left exact, $\otimes_R M$ is right exact, as functors on $R\text{-mods}$ (any $R\text{-mod } M$)

1.9 Push-forward (direct image) and inverse image

$f: X \rightarrow Y$ continuous
 \Rightarrow additive functor $f_*: Ab X \rightarrow Ab Y$
Def $F \in Ab(X)$ gives $f_* F \in Ab(Y)$:
 $(f_* F)(V) = F(f^{-1}(V))$
Exercise $(g \circ f)_* F = g_*(f_* F)$ for $X \xrightarrow{f} Y \xrightarrow{g} Z$.

\Rightarrow additive functor $f^{-1}: Ab Y \rightarrow Ab X$
Def $F \in Ab(Y)$ gives $f^{-1}F \in Ab(X)$ is $(\text{pre-}f^{-1}F)^+$ where
 $(\text{pre-}f^{-1}F)(U) = \varinjlim_{V \supseteq f(U)} F(V)$
Exercise $(f^{-1}F)_x = F_{f(x)}$ and $(g \circ f)^{-1} \cong f^{-1} \circ g^{-1}$ canonical



Examples 1) $i: S \rightarrow X$ inclusion of an open subset:
 $F \in Ab(S) \quad i_* F: V \mapsto F(V \cap S)$
 $F \in Ab(X) \quad i^{-1}F: U \mapsto F(U)$ \leftarrow denoted $F|_S$ called **restriction of F**
 2) $\lambda_x: \text{point} \rightarrow X, \quad i_x(\text{point}) = x$
 $F \in Ab(X) \quad i_x^{-1}F = F_x$
 more precisely: $(i_x^{-1}F)(U) = \begin{cases} F_x & \text{if } U = \{\text{point}\} \\ 0 & \text{if } U = \emptyset \end{cases}$
 will not make such remarks again.

Proposition 1) f_* is left exact \leftarrow in particular $\Gamma(X, \cdot)$ is left exact
 2) f^{-1} is exact

For f_* : exercise
Proof for f^{-1} : $0 \rightarrow (f^{-1}A)_x \rightarrow (f^{-1}B)_x \rightarrow (f^{-1}C)_x \rightarrow 0$
 $0 \rightarrow A_{f(x)} \rightarrow B_{f(x)} \rightarrow C_{f(x)} \rightarrow 0$ which by assumption is exact

RMK f_* left exact } would follow by category theory from next proposition
 f^{-1} right exact

F exact $\Rightarrow F$ both left & right exact



also follows by uniqueness up to unique iso of adjoint functors see next page.

Proposition f^{-1} is the left adjoint functor of f_* , meaning \exists natural iso

$\text{Mor}(f^{-1}F, G) \cong \text{Mor}(F, f_*G)$ which is natural in F and G

Sketch pt
 $\text{In} \rightarrow \text{direction:}$ $F(V) \xrightarrow{\text{since } W \subseteq V \text{ is allowed}} \varinjlim_{W \supseteq fU} F(W) \xrightarrow{\text{given}} G(U)$
 $\parallel \leftarrow \text{pick } U = f^{-1}V$
 $G(f^{-1}V) = f_*G(V)$

$\text{In} \leftarrow \text{direction:}$ $F(V) \xrightarrow{\text{given}} G(f^{-1}V) \xrightarrow{\text{assume } V \supseteq fU} \varinjlim_{V \supseteq fU} G(f^{-1}V) \xrightarrow{\text{restriction}} G(U)$

Now check these two are natural transformations, inverse to each other, and natural in F, G, \square

Rmk Another example of adjoint functors, for R -modules, are $\text{Hom}(M, -)$ and $\otimes M$:
 $\text{Hom}(F \otimes M, G) \cong \text{Hom}(F, \text{Hom}(M, G))$ for R -mods F, G .

1.10 Morphisms of ringed spaces

Def $(f, \varphi) : (X, \mathcal{O}_X) \rightarrow (Y, \mathcal{O}_Y)$ morph of ringed spaces means
 $X \xrightarrow{f} Y$ continuous map of topological spaces
 $f_* \mathcal{O}_X \xleftarrow{\varphi} \mathcal{O}_Y$ morph of sheaves of rings (on Y)
 (So: $\mathcal{O}_X(f^{-1}V) \xleftarrow{\varphi_V} \mathcal{O}_Y(V)$ for $V \subseteq Y$, compatibly with restrictions)

often write $\varphi = f^\#$
 For a morphism of locally ringed spaces want in addition:
 $\mathcal{O}_{X,x} \xleftarrow{\varphi_x} \mathcal{O}_{Y,f(x)}$ is local ring hom
 (Explanation: $\varphi_V(s) \in \mathcal{O}_X(f^{-1}V)$ is a representative for $\varphi_x(s_{f(x)})$)

Can compose: $(X, \mathcal{O}_X) \xrightarrow{f_1} (Y, \mathcal{O}_Y) \xrightarrow{g_2} (Z, \mathcal{O}_Z)$
 $(g \circ f)_* \mathcal{O}_X = g_* f_* \mathcal{O}_X \xleftarrow{g_*(f^\#)} g_* \mathcal{O}_Y \xleftarrow{g^\#} \mathcal{O}_Z$

Notice in the definition we cannot just talk about a morphism $\mathcal{O}_X \leftarrow \mathcal{O}_Y$ because the sheaves are not defined over the same topological space.

\Rightarrow either need a morph $f_* \mathcal{O}_X \leftarrow \mathcal{O}_Y$ of sheaves on Y or a morph $\mathcal{O}_X \leftarrow f^{-1} \mathcal{O}_Y$ of sheaves on X

By the proposition, this is the same information since $\text{Mor}(f^{-1} \mathcal{O}_Y, \mathcal{O}_X) \cong \text{Mor}(\mathcal{O}_Y, f_* \mathcal{O}_X)$

(Notice also the map on stalks $\mathcal{O}_{X,x} \leftarrow (f^{-1} \mathcal{O}_Y)_x = \mathcal{O}_{Y,f(x)}$ is the φ_x above)

Rmk φ local \Rightarrow also get hom on residue fields: $\varphi_x : k(f(x)) = \mathcal{O}_{Y,f(x)} / \mathfrak{m}_{Y,f(x)} \rightarrow \mathcal{O}_{X,x} / \mathfrak{m}_{X,x} = k(x)$
 \Rightarrow field extension $\varphi_x : k(f(x)) \hookrightarrow k(x)$ (in classical algebraic geometry: k alg. closed and x closed point get id: $k \rightarrow k, p(f(x)) \mapsto (f \circ p)(x)$ where $\{x \in X\}$)

1.11 A sheaf defined on a topological basis

X top space with a basis B of open subsets \leftarrow means: basic sets cover X , and:
 $(\forall \text{ basic } B, B_2, x \in B, B_2, \exists \text{ basic } B' \text{ with } x \in B' \subseteq B, B_2)$

Def B -sheaf F means
 $F(U) \in \text{Ab}, \forall \text{ basic } U$ with horns $F(U) \rightarrow F(V), s \mapsto s|_V \forall \text{ basic } V \subseteq U$
 and as usual: $F(U) \xrightarrow{\text{id}} F(U)$ and $F(U) \rightarrow F(V) \rightarrow F(W) \rightarrow F(U)$ for $W \subseteq V \subseteq U$

\bullet local-to-global condition:
 $\forall \text{ basic } U$ with $U = \cup U_i$
 $\forall s_i \in F(U_i)$ "agreeing locally on overlaps":
 $\forall x \in U_i \cap U_j \exists \text{ basic } x \in U_k \subseteq U_i \cap U_j$ with $s_i|_{U_k} = s_j|_{U_k} \in F(U_k)$

$\Rightarrow \exists$ unique $s \in F(U)$ with $s|_{U_i} = s_i$.

Rmk stalk $F_x = \varinjlim_{x \in \text{basic } U} F(U)$.

Theorem 1) B -sheaf F extends uniquely (up to unique iso) to a sheaf \tilde{F} on X .
 (Hence also the stalk is \tilde{F}_x up to canonical iso.)
 (So $F(\text{basic } U)$ and restrictions for basic sets are same up canonical isomorphisms.)


2) B -sheaves F, G then morph $F \rightarrow G$ on the extended sheaves is uniquely defined by data:
 \bullet horns $F(U) \rightarrow G(U)$ for basic U , commuting with restrictions (for basic opens)

Proof (1):
 Uniqueness: Such an extension \tilde{F} is unique (if it exists) because we can canonically identify $\tilde{F}(U)$ for any open U in terms of the B -sheaf data:
 $\tilde{F}(U) \xrightarrow{\text{bijection}} \{s_V \in F(V) \text{ for } (\text{basic } V) \subseteq U : s_V|_W = s_W, s_V|_{W'} = s_{W'} \text{ for basic } W \subseteq V \subseteq W'\}$
 $s \mapsto (s_V := s|_V \in \tilde{F}(V) = F(V))$

Explanation: given s , notice that this holds: $s_V|_W = (s|_V)|_W = s|_W = (s|_{W'})|_W = s_{W'}|_W$.
 Conversely, given such $s_V \in F(V) = \tilde{F}(V)$, then $s_V|_{V \cap V'} \in \tilde{F}(V \cap V')$ and $s_V|_{V \cap V'} \in \tilde{F}(V \cap V')$ must equal because their restrictions to a covering of $V \cap V'$ by basic W agree ($= s_W$). (and then use sheaf property of \tilde{F})

Existence
 \leftarrow inverse limit over restrictions for basics
 \leftarrow "compatible families of local sections on basic open sets"
 \leftarrow basic sections on basic open sets: $s_V|_W = s_W \forall W \subseteq V \subseteq U$

With obvious restriction maps (for $U' \subseteq U$ a subset of the basic $V \subseteq U$ are $\subseteq U'$)



Notice: $F(\text{basic } U)$ has not changed up to canonical identification:

$$F(U) \cong \lim_{(\text{basic } V) \subseteq U} F(V) \xrightarrow{s} (S|_U) \text{ which includes } s|_U = s.$$

and for stalks:

$$\lim_{x \in U} F(V) \cong \lim_{x \in U} \lim_{(\text{basic } V) \subseteq U} F(V) \cong \lim_{(\text{basic } V) \subseteq U} G(V).$$

easy check: if sections agree on $x \in W$ then agree on $x \in V \subseteq W$ some basic V .

Proof (2): by functoriality of \lim_{\leftarrow} :

Rmk Equivalently, it is enough to remember germs around each point:

$$F(U) = \left(\lim_{(\text{basic } V) \subseteq U} F(V) \right) \cong \left\{ s: U \rightarrow \coprod_{x \in X} F_x : s(x) \in F_x \text{ which are "locally compatible": } \forall x \in U, \exists x \in (\text{basic } V) \subseteq U \text{ with } \exists t \in F(V) \text{ } \exists \text{ open } x \in W \subseteq V \text{ } t|_y = s(y) \forall y \in W \right\}$$

with obvious restriction maps for these (just restrict the map $U \rightarrow \coprod F_x$).

Rmk Can simplify. WLOG W also basic (just pick $x \in (\text{basic } V) \subseteq U$ so $\exists t \in F(V)$ with $t|_y = s(y) \forall y \in V$). WLOG replace V by W , so $V = W$ basic. $\exists t \in F(V)$ with $t|_y = s(y) \forall y \in V$ so \otimes holds so can extend to unique global section.

1.12 Construction of $\mathcal{O}_{\text{Spec } R}$

$X = \text{Spec } R$, we define \mathcal{O}_X first on basic open sets: $\{g : g \text{ does not vanish on } D_f\}$

$$\mathcal{O}_X(D_f) = R \xrightarrow{\cong} R_f \xrightarrow{\text{natural}} \mathcal{O}_X(D_g) \xrightarrow{\cong} R_f$$

For $D_f \subseteq D_g$ define natural restriction homs: (which are compatible under composition)

$$\mathcal{O}_X(D_g) \xrightarrow{\cong} \mathcal{O}_X(D_f) \xrightarrow{\cong} R_f \xrightarrow{\cong} R_f$$

Lemma 1 This is a B -sheaf on X for $B = \{\text{basic open sets } D_f, f \in R\}$

Pf Uniqueness: $\alpha, \beta \in R_f = \mathcal{O}_X(D_f)$ and $D_f = \cup D_{f_i}$ if $\alpha|_{D_{f_i}} = \beta|_{D_{f_i}} \forall i$ then $\alpha = \beta$

Proof By redefining X, R by D_f, R_f we can assume $f=1, R_f=R, D_f=X$.

$\alpha - \beta = 0 \in R_f \Rightarrow f_i^n \cdot (\alpha - \beta) = 0$ some $N \in \mathbb{N} \leftarrow N \text{ may depend on } i, \text{ but WLOG finite subcover } D_{f_i}$

$\Rightarrow \langle \text{all } f_i^n \rangle = (\alpha - \beta) = 0$ (quasi-compactness) \rightarrow so pick maximal N "Covering Trick" $\rightarrow R$ since $X = D_{f_1} \cup \dots \cup D_{f_n} \leftarrow (\text{recall } D_f = D_{f^n})$

$\Rightarrow 1 \cdot (\alpha - \beta) = 0$ so $\alpha = \beta$ \square

Existence in \otimes : as before WLOG $U = D_f, R_f$ become X, R . Uniqueness \Rightarrow in \otimes can assume sections $s_i \in \mathcal{O}_X(D_{f_i})$ agree on overlaps $D_{f_i} \cap D_{f_j} = D_{f_i f_j}$

(applies Uniqueness to $D_{f_i f_j}$) $s_i|_{D_{f_i f_j}} = s_j|_{D_{f_i f_j}} \in R_{f_i f_j}$

WLOG $X = D_{f_1} \cup \dots \cup D_{f_n}$ finite cover, $s_i = \frac{g_i}{f_i^{n_i}}$ since $D_{f_i} = D_{f_i^{n_i}}$, WLOG $n_i=1$, so $s_i = \frac{g_i}{f_i}$

$s_i = s_j$ on $D_{f_i f_j} \Rightarrow (f_i f_j)^N (f_j g_i - f_i g_j) = 0 \in R \leftarrow N \text{ depends on } i, j \text{ but can pick largest } N \text{ over finitely many } i, j \text{ so } N \text{ works } \forall i, j$

rewrite: $(f_j^{N+1}) \cdot \underbrace{(f_i g_i)}_{a_i} - \underbrace{(f_i^{N+1})}_{b_i} \cdot \underbrace{(f_j g_j)}_{c_j} = 0$ notice $s_i = \frac{a_i}{b_i}, D_{f_i} = D_{b_i}$ so WLOG $N=0!$ "Covering Trick": $X = D_{f_1} \cup \dots \cup D_{f_n}$ so $1 = \sum r_i f_i \leftarrow$ ("partition of unity" trick)

$$1 \cdot g_j = \left(\sum_i r_i f_i \right) g_j = \sum_i r_i (f_i g_j) = \sum_i r_i (f_j g_i) = f_j \left(\sum_i r_i g_i \right)$$

$\Rightarrow s_j = \frac{g_j}{f_j} = \frac{\sum_i r_i g_i}{1} \in R_f \forall j$ so we globalised the $s_i \in \mathcal{O}_X(D_{f_i})$ to $\sum_i r_i g_i \in \mathcal{O}_X(X) = R$ \square

Corollary \mathcal{O}_X extends uniquely to a sheaf on $X = \text{Spec } R$ called structure sheaf (or sheaf of regular functions)

Messy unpacking of definitions:

we identify $\frac{f_m}{f_n} \in R_f \cong \mathcal{O}_X(D_f)$ and $\frac{s}{g} \in R_g \cong \mathcal{O}_X(D_g)$ iff $\frac{f_m}{f_n} = \frac{s}{g} \in R_h$ some $h \in R$ with $p \in D_h \subseteq D_f \cap D_g$ (iff $R \setminus (R g^n - s f^m) = \emptyset$ for some n) $D_{f g}$

Lemma 2

$$\mathcal{O}_{X, p} \cong R_p \xrightarrow{\text{rest.}} \mathcal{O}_X(X) \cong R$$

straightforward algebra exercise \leftarrow Recall in R_p you invert all elements $f \notin p$

$$\text{Pf } \lim_{D_f \ni p} \mathcal{O}_X(D_f) \cong \lim_{f \notin p} R_f \cong R_p \cdot \square$$

1.13 Morphisms between Specs

$$\varphi: R \rightarrow S \text{ hom of rings} \Rightarrow \text{Spec}(\varphi): \text{Spec} S \rightarrow \text{Spec} R$$

Example $\varphi: R \rightarrow R_f, r \mapsto \frac{r}{f}$ localisation

$\text{Spec} R \leftarrow \text{Spec} R_f, r \mapsto \frac{r}{f}$ is an "inclusion" with image = D_f .

$\alpha = \text{Spec}(\varphi): Y \rightarrow X, p \mapsto \varphi^{-1}(p)$

Lemma $\alpha^{-1}(D_f) = D_{\varphi(f)}$ automatically true!

$$\text{Pf } \alpha^{-1}\{q \in X: f \notin q\} = \{p \in Y: \varphi^{-1}(p) = q \text{ some } q \in X, f \notin \varphi^{-1}(p)\} = \{p \in Y: \varphi(f) \notin p\} \quad \square$$

Claim $\exists \varphi^\#: \theta_X \rightarrow \alpha_* \theta_Y$ such that $\varphi^\#: \theta_X(X) = R \xrightarrow{\varphi} S = \alpha_* \theta_Y(X)$

Pf Enough to build $\varphi^\#$ on basic opens, compatibly with restrictions

$$\varphi^\#: \theta_X(D_f) \xrightarrow{\cong} \alpha_* \theta_Y(D_f) = \theta_Y(\alpha^{-1}D_f) = \theta_Y(D_{\varphi(f)}) \xrightarrow{\cong} S_{\varphi(f)}$$

(By Theorem on B-sheaves)

Easy check: compatible with restriction maps for $D_g \subseteq D_f \cdot \square$

Claim $\theta_{X,p}$ is local and $\varphi^\#$ is local

Pf Lemma 2: $\theta_{X,p} \cong R_p$ so local with max ideal $m_p = p \cdot R_p$.

For $p \in Y, \varphi^\# : \theta_{X,p} \rightarrow \theta_{Y,p}$ is direct limit of maps hence: natural map: $\frac{r}{f} \mapsto \frac{\varphi(r)}{\varphi(f)}$
 Hint: $\varphi(r) \notin p \Rightarrow r \notin \varphi^{-1}(p)$ so $\varphi(f) \notin p$

Theorem (ring R) \rightarrow locally ringed space $(\text{Spec} R, \theta_{\text{Spec} R})$
 (ring hom $R \xrightarrow{\varphi} S$) \rightarrow $(\text{Spec} \varphi, \varphi^\#): (\text{Spec} S, \theta_{\text{Spec} S}) \rightarrow (\text{Spec} R, \theta_{\text{Spec} R})$ (easy to check)

contravariant functor $\text{Spec}: \text{Rings} \rightarrow \text{Locally Ringed Spaces}$

Claim The functor is fully faithful \leftarrow i.e. surj & inj. (so iso) on morphism spaces

Pf Given a hom of loc. ringed spaces $(f, f^\#): (Y, \theta_Y) \rightarrow (X, \theta_X)$ $X = \text{Spec} R, Y = \text{Spec} S$

Let $\varphi := f^\#: R \cong \theta_X(X) \xrightarrow{f^\#} \theta_Y(X) = \theta_Y(Y) \cong S$ ring hom.

$\downarrow \varphi$ localisation maps (Lemma 2) for $\theta_{X_i}, \theta_{Y_i}$

$$R \cong \theta_{X,fp} \xrightarrow{f^\#} \theta_{Y,p} \cong S_p \supseteq m_p = p \cdot S_p$$

$$\Rightarrow \varphi^{-1}(p) = \varphi^{-1}(l_p^{-1}(m_p)) = l_{f^\#}^{-1}(f^\#^{-1}(m_p)) = f(p)$$

diagram m_{fp} since $f^\#$ local ring hom

$\Rightarrow \theta_X(U) = \{s: U \rightarrow \coprod_{p \in X} R_p : s(p) \in R_p \text{ which are locally compatible}\}$

with the obvious restriction maps. $\forall p \in U, \exists$ open nbhd $p \in D \subseteq U$ with $s(x) = t_x$ $\forall x \in D_f$ $\frac{f_m}{f_n} \in \theta_{X,p}$ some $f \in R$

Remark could assume $t = \frac{f}{f}$ since can replace D_f with $D_{f_m} (= D_f)$.

could just ask $s(x) = t_x$ on a smaller open $p \in V \subseteq D_f$.

Comparison with classical algebraic geometry

- X affine variety, $p \in U \subseteq X$ open nbhd $f: U \rightarrow k$ is regular at p if \exists open nbhd $p \in W \subseteq U$ with $h(w) \neq 0 \forall w \in W$ $f = \frac{g}{h}$ on $W, g, h \in k[X], h(w) \neq 0 \forall w \in W$

Remark In fact can assume $W = D_h$ basic open (if $f = \frac{g}{h}$, replace D_h by $D_{h^n} = D_h$)

$\theta_X(U) = k$ -algebra of functions $U \rightarrow k$ regular at all $p \in U$

$\theta_{X,p} = k$ -algebra of germs of functions near p , regular at p

(so pairs (U, f) with $p \in U \subseteq X$ open, $f: U \rightarrow k$ regular at p (and identify $(U, f) \sim (V, g) \Leftrightarrow f|_W = g|_W$ on some open $p \in W \subseteq U \cap V$)

Theorem $\theta_X(X) \cong k[X] \leftarrow$ (Remark This theorem is not obvious in C3.4 course. $X = \text{Spec} k[X]$ so by Lemma 1 get $\theta_X(X) = k[X]$)

$\theta_X(X) = k[X] \cong k[X]$ so by Lemma 1 get $\theta_X(X) = k[X]$

$\theta_X(X) = k[X] \cong k[X]$ so by Lemma 1 get $\theta_X(X) = k[X]$

$\theta_X(X) = k[X] \cong k[X]$ so by Lemma 1 get $\theta_X(X) = k[X]$

$\theta_X(X) = k[X] \cong k[X]$ so by Lemma 1 get $\theta_X(X) = k[X]$

$\theta_X(X) = k[X] \cong k[X]$ so by Lemma 1 get $\theta_X(X) = k[X]$

$\theta_X(X) = k[X] \cong k[X]$ so by Lemma 1 get $\theta_X(X) = k[X]$

$\theta_X(X) = k[X] \cong k[X]$ so by Lemma 1 get $\theta_X(X) = k[X]$

$\theta_X(X) = k[X] \cong k[X]$ so by Lemma 1 get $\theta_X(X) = k[X]$

$\theta_X(X) = k[X] \cong k[X]$ so by Lemma 1 get $\theta_X(X) = k[X]$

$\theta_X(X) = k[X] \cong k[X]$ so by Lemma 1 get $\theta_X(X) = k[X]$

$\theta_X(X) = k[X] \cong k[X]$ so by Lemma 1 get $\theta_X(X) = k[X]$

$\theta_X(X) = k[X] \cong k[X]$ so by Lemma 1 get $\theta_X(X) = k[X]$

$\theta_X(X) = k[X] \cong k[X]$ so by Lemma 1 get $\theta_X(X) = k[X]$

$\theta_X(X) = k[X] \cong k[X]$ so by Lemma 1 get $\theta_X(X) = k[X]$

$\theta_X(X) = k[X] \cong k[X]$ so by Lemma 1 get $\theta_X(X) = k[X]$

2. GLOBAL SECTIONS AND THE FUNCTOR OF POINTS

2.0 Points of Spec R (not necessarily closed)

$$R \xrightarrow{\text{quotient}} R_p = R_p / \mathfrak{m}_p \Rightarrow \text{Spec } K(p) \hookrightarrow \text{Spec } R_p \hookrightarrow \text{Spec } R$$

$$\text{Loc}^{-1}(\mathfrak{m}_p) = \mathfrak{p} \leftarrow \text{p.R.} = \mathfrak{m}_p \leftarrow (0) \quad \text{via } \{0\} \quad \text{via } \mathfrak{p} \quad \text{via } \mathfrak{m}_p \quad \text{via } \mathfrak{p} \quad \text{via } \mathfrak{m}_p \quad \text{via } \mathfrak{p}$$

So points of Spec R correspond to the max ideals in the local rings.

2.1 Global sections and basic open sets for locally ringed spaces

(X, \mathcal{O}_X) locally ringed space $\Gamma(\cdot, \mathcal{O}_X) : \text{Top}(X)^{\text{op}} \rightarrow \text{Rings}$, $U \xrightarrow{\Gamma} \mathcal{O}_X(U)$
 sections functor \downarrow restrict $\Gamma \downarrow \mathcal{O}_X(V)$

Global sections functor: Locally ringed spaces $\text{op} \rightarrow \text{Rings}$, $(X, \mathcal{O}_X) \mapsto \Gamma(X, \mathcal{O}_X) = \mathcal{O}_X(X)$
 \exists canonical map $X \rightarrow \text{Spec } \mathcal{O}_X(X)$, $x \mapsto \text{res}_x^{-1}(\mathfrak{m}_{x,x})$ where $\text{res}_x: \mathcal{O}_X(X) \rightarrow \mathcal{O}_{x,x}$ restricts.

Trick $f \in \mathcal{O}_X(X)$ then $f_x \in \mathcal{O}_{x,x}$ invertible $\Leftrightarrow f(x) \neq 0 \in K(x) = \mathcal{O}_{x,x} / \mathfrak{m}_x$
 $\text{Pf } f_x \in \mathcal{O}_{x,x} \setminus \mathfrak{m}_x = \{\text{invertibles of } \mathcal{O}_{x,x}\} \Leftrightarrow f_x \notin \mathfrak{m}_x \square$
 image of f via $\mathcal{O}_X(X) \rightarrow \mathcal{O}_{x,x} \rightarrow \mathcal{O}_{x,x} / \mathfrak{m}_x \rightarrow K(x)$
 $f \mapsto f_x \mapsto f_x \mapsto f(x)$

Lemma $f \in \mathcal{O}_X(X) \Rightarrow D_f = \{x \in X : f(x) \neq 0 \in K(x)\}$ is open in X .
 $\Leftrightarrow f \notin \mathfrak{m}_x \Leftrightarrow \{f_x \in \mathcal{O}_{x,x} \text{ invertible}\}$

Pf Trick $\Rightarrow \exists g \in \mathcal{O}_{x,x} : f \cdot g = 1$ so \exists open $U \subseteq X$ s.t. $f, g \in \mathcal{O}_X(U)$, $f \cdot g = 1 \in \mathcal{O}_X(U)$
 $\Rightarrow x \in U \subseteq D_f$ since $\forall y \in U, f_y \cdot g_y = (f \cdot g)_y = 1_y \in \mathcal{O}_{y,y}$ so $f_y \in \{\text{invertibles of } \mathcal{O}_{y,y}\}$ so $f(y) \neq 0$, so $y \in D_f \square$

Lemma $f|_{D_f} \in \mathcal{O}_X(D_f)$ is invertible
Pf Lemma $\Rightarrow f$ is locally invertible. If $f \cdot h = 1$ on U then $h = g$ on $U \cap V$. So can globalize. \square
 uniqueness of inverses ($h = g, 1 = h \cdot g = 1 \cdot g = g$)

2.2 What it means to be affine
 \hookrightarrow locally ringed space

(X, \mathcal{O}_X) affine $\Leftrightarrow \exists$ ring $R : \exists X \xrightarrow{\alpha} Y = \text{Spec } R$ homeomorph, and $\exists \mathcal{O}_Y \xrightarrow{\cong} \alpha_* \mathcal{O}_X$
 But $\mathcal{O}_Y(Y) = R$ so $R \xrightarrow{\cong} \mathcal{O}_X(X)$ so $\text{Spec } \mathcal{O}_X(X) \xrightarrow{\cong} Y$.
 via $\varphi^{-1}(\cdot)$

$\varphi_x \text{ local} \Rightarrow \mathcal{O}_Y(Y) = R \xrightarrow{\cong} \mathcal{O}_X(X) \xrightarrow{\cong} \mathcal{O}_x(x) \xrightarrow{\cong} \text{res}_x^{-1}(\mathfrak{m}_x) \subseteq \mathcal{O}_X(X)$
 $\mathcal{O}_Y(Y) = R \xrightarrow{\alpha(x)} \mathcal{O}_x(x) \xrightarrow{\alpha(x)} \mathcal{O}_x(x) \xrightarrow{\alpha(x)} \text{res}_x^{-1}(\mathfrak{m}_x) \rightarrow \mathcal{O}_x(x)$
 $\alpha(x) : R \rightarrow \mathcal{O}_x(x)$ so $X \xrightarrow{\text{canonical}} \text{Spec } \mathcal{O}_X(X) \cong Y$
 $x \mapsto \text{res}_x^{-1}(\mathfrak{m}_x) \mapsto \alpha(x)$

So a locally ringed space (X, \mathcal{O}_X) is affine precisely if:

- the canonical map $X \rightarrow \text{Spec } \Gamma(X, \mathcal{O}_X)$ is homeomorph
- $\mathcal{O}_X(D_f) \cong \Gamma(X, \mathcal{O}_X)_f$, $\forall f \in \Gamma(X, \mathcal{O}_X)$ and restrictions are localizations \leftarrow (by Sec. 1.12)

2.3 Functor of points h_Y

MOTIVATION Y set, you recover set Y from $\text{Mor}(\text{point}, Y)$
 Y group, " " set " " $\text{Mor}(\mathbb{Z}, Y)$

$\Rightarrow f(p) = \varphi^{-1}(p)$ so $f = \text{Spec}(\varphi)$ is the map on Specs induced by $\varphi: R \rightarrow S$.

Upshot: have two morphs of sheaves $f^\#, \varphi^\# : \mathcal{O}_X \rightarrow \text{Spec}(\varphi)_* \mathcal{O}_Y$
 and $f^\# = \varphi^\#$ since equal on stalks (by the diagram have $f^\# = \varphi^\#$) \square

Def Aff = category of affine schemes (and morphs of locally ringed spaces)
 (locally ringed spaces $\cong (\text{Spec } R, \mathcal{O}_{\text{Spec } R})$ some ring R)
 $\varphi(x) = \varphi(\frac{1}{s} \cdot \frac{t}{s}) = \varphi(\frac{t}{s}) \cdot \varphi(\frac{1}{s})$
 because: $\frac{t}{s} \mapsto \frac{\varphi(t)}{\varphi(s)}$

$\Rightarrow \text{Spec} : \text{Rings}^{\text{op}} \rightarrow \text{Aff}$ is an equivalence of categories.
 (op = opposite category = reverse arrows so artificially make Spec covariant)

1.14 Closed affine subschemes full, faithful, essentially surjective functor
 $X = \text{Spec } R$, $I \subseteq R$ ideal (link same as specifying a subs) each object in target category is iso to an object in image
 $Y = V(I) \cong \text{Spec}(R/I)$ are called closed (affine) subschemes of X
 (as top space, $V(I) = V(I)$ but sheaf remembers $I : \mathcal{O}_Y(Y) = R/I$)

Example $I = \mathfrak{m}$ max ideal \Rightarrow get a closed point $\{\mathfrak{m}\} = \text{Spec } R/\mathfrak{m} \hookrightarrow X$.
Link $\text{Spec}(R/I)$ is closed subscheme of $\text{Spec}(R/I)$ means $J \supseteq I \Rightarrow V(J) \subseteq V(I)$
Def $\text{Spec } R/I \cap \text{Spec } R/J = \text{Spec}(R/(I+J))$, $\text{Spec } R/I \cup \text{Spec } R/J = \text{Spec } R_{I \cap J}$
 $\sqrt{J} \supseteq \sqrt{I}$

Def sheaf of ideals $\mathcal{J} = \mathcal{J}_x \times Y$ on X :
 (also: ideal sheaf) $\mathcal{J}(D_f) = I \cdot R_f \subseteq R_f = \mathcal{O}_X(D_f)$ ideal
 Notice $\mathcal{O}_Y(D_f) = (R/I)_f \cong R_f/I \cdot R_f = \mathcal{O}_X(D_f) / \mathcal{J}(D_f)$

$\Rightarrow \mathcal{J} = \text{Ker}(\mathcal{O}_X \rightarrow j_* \mathcal{O}_Y)$
 $\mathcal{O}_Y = \mathcal{O}_X / \mathcal{J}$ where $j : Y \rightarrow X$ inclusion.
 more precisely this is $j_* \mathcal{O}_Y$

1.15 Closed subschemes (later in course: sheaves of R -modules and quasi-coherence)
 (X, \mathcal{O}_X) scheme, sheaf of ideals \mathcal{J} means $\mathcal{J}(U) \subseteq \mathcal{O}_X(U)$ ideal compatibly with restrictions.
Def A sheaf of ideals on $X = \text{Spec } R$ is quasi-coherent if it arises as \mathcal{J} as above, some ideal $I \subseteq R$ on $X = \text{scheme}$ " if \forall affine open U , $\mathcal{J}|_U$ is quasi-coherent.
 (later revisit these in Sec. 3.6) \leftarrow see 1.14

closed subscheme means $Y \subseteq X$ closed topological space
 $\mathcal{O}_Y = \mathcal{O}_X / \mathcal{J}$ some quasi-coherent sheaf of ideals \mathcal{J} on X ,
 s.t. $Y \cap (\text{affine open } U) \subseteq U$ is closed affine subscheme for the ideal $\mathcal{J}(U) \subseteq \mathcal{O}_X(U)$.

Rmk $\exists !: 1$ correspondence $\{\text{closed subschemes of } X\} \leftrightarrow \{\text{quasi-coh. sheaves of ideals on } X\}$
 Can recover $Y \subseteq X$ from \mathcal{J} from the support of $\mathcal{O}_X / \mathcal{J} : \leftarrow$ if $I \subseteq R$ then $\mathbb{A}^1 \cap \text{Spec } R/I = \text{Spec } R/I$ since $I \cap \mathfrak{p} \in \mathfrak{p}$
 $Y = \text{Supp } \mathcal{O}_X / \mathcal{J} = \{x \in X : (\mathcal{O}_X / \mathcal{J})_x \neq 0\} = \{x \in X : \mathcal{J}_x \neq \mathcal{O}_{X,x}\}$

Example closed point $p \in X$ (so $\{p\} = \{p\}$) \Rightarrow pick affine $p \in \text{Spec } R \hookrightarrow X$ then $p \in \text{Spec } R \subseteq \text{Spec } R$
 \Rightarrow sheaf \mathcal{J} on $\text{Spec } R \Rightarrow$ extend \mathcal{J} to X by $\mathcal{J}(V) = \mathcal{O}_X(V)$ if $p \notin V$ (so $\mathcal{O}_Y(V) = 0$)

Functor of points $h_Y : \text{Sch}^{\text{op}} \rightarrow \text{Sets}$, $h_Y(X) = \text{Mor}(X, Y)$
 $X \xrightarrow{f} Z \rightarrow$ on morphs: $h_Y(X \xrightarrow{f} Z) = (\text{Mor}(X, Y) \xrightarrow{\text{of}} \text{Mor}(Z, Y))$

MOTIVATION

$Y = \text{Spec } \mathbb{Z}[x]/(x^2+1)$. \mathbb{C} -valued points of Y ?
 $\mathbb{Z}[x]/(x^2+1) \rightarrow \mathbb{C}, x \mapsto i \Rightarrow \text{morph } X = \text{Spec } \mathbb{C} \rightarrow Y$ so $i \in h_Y(X) \leftarrow$ (often write $Y(\mathbb{C})$)

Yoneda Lemma Nat $(h_Y, F) \cong F(Y)$
 Take image of $\text{id}_Y \in \text{Mor}(Y, Y) = h_Y(Y)$ given $F(Y)$
 Conversely given $\alpha \in F(Y)$, $\varphi \in h_Y(X)$ get $F(\varphi)(\alpha) = F(X)$

Yoneda embedding $h_Y : \text{Sch} \rightarrow \text{Sets}^{\text{Sch}^{\text{op}}}$
 h_Y is fully faithful (iso on morphisms: $\text{Nat}(h_Y, h_W) \cong \text{Mor}(Y, W)$)

UPSHOT $h_Y \cong h_W \iff Y \cong W$

Can now ask which functors $\text{Sch}^{\text{op}} \rightarrow \text{Sets}$ are represented by a scheme Y .
 Sets $\text{Sch}^{\text{op}} = \text{category: } \{\text{Obj are functors } \text{Sch}^{\text{op}} \rightarrow \text{Sets} \text{ Morph are natural transformations}\}$

Example Will show that $\mathbb{A}^1 = \text{Spec } \mathbb{Z}[x_1, \dots, x_n]$ represents \mathbb{A}^1 (tell me who your friends are and I will tell you you are \mathbb{A}^1)
 $\text{Sch}^{\text{op}} \rightarrow \text{Sets}, X \mapsto \{\text{morphs } \mathbb{A}^1 \rightarrow X \text{ which are } \theta_X\text{-linear}\}$

Example 1 $h_{\text{Spec } R} \Rightarrow \text{Mor}(X, \text{Spec } R) \rightarrow \text{Hom}(R, \Gamma(X, \theta_X))$
 $\text{Mor}(X, \text{Spec } R) = \text{Mor}_{\text{Spec } R}(\text{Spec } R, X)$

KEY EXAMPLE $Y = \mathbb{A}^1 = \text{Spec } \mathbb{Z}[x]$
 $\text{Mor}(X, \mathbb{A}^1) \cong \text{Mor}(X, \mathbb{A}^1) \cong \text{Mor}(X, \mathbb{A}^1)$
 $\text{Mor}(X, \mathbb{A}^1) \cong \text{Mor}(X, \mathbb{A}^1)$

Cor 1 (X, θ_X) scheme \Rightarrow canonical morph $X \rightarrow \text{Spec } \Gamma(X, \theta_X)$
 Explicitly: on sets $x \mapsto \text{res}^{-1}(m_{X,x}) \in \theta_X(x)$
 on sheaves over $D_f \subseteq X: \theta_X(X)_f \xrightarrow{\text{res}} \theta_X(D_f)$ now localise at f using that f invertible

Cor 2 $x \in X \Rightarrow \exists$ canonical morph $\text{Spec } \theta_{X,x} \rightarrow X$
 Any $\text{Spec } R \rightarrow X$ factors as $\text{Spec } R \rightarrow \text{Spec } \theta_{X,x} \rightarrow X$ some $x \in X$
 Any $f: X \rightarrow Y$ of schemes get $\text{Spec } \theta_{X,x} \rightarrow X \xrightarrow{f} Y$
 $\text{Spec } \theta_{X,x} \rightarrow Y$ induced by f_x

Example 2 $h_Y(\text{Spec } \mathbb{K}) \leftarrow$ (also written $Y(\mathbb{K})$)
 Thus: $\{f \in \text{Mor}(\text{Spec } \mathbb{K}, Y) \mid f(\text{pt}) = y\} \xrightarrow{1:1} \text{Hom}(\mathbb{K}(y), \mathbb{K})$ and any $\text{Spec } \mathbb{K} \rightarrow Y$ factors:
 $\text{Spec } \mathbb{K} \rightarrow \text{Spec } \mathbb{K}(y) \rightarrow Y$

UPSHOT: Morphs from local rings or fields don't give more information than already know from $\text{Spec } \theta_{X,x} \rightarrow X$ and $\text{Spec } \mathbb{K}(x) \rightarrow X$.
 \mathbb{K} -valued point if $\mathbb{K}(y) \cong \mathbb{K}$, then $\text{id}_{\mathbb{K}}$ defines a morph $\text{Spec } \mathbb{K} \rightarrow Y$
 (or \mathbb{K} -point or \mathbb{K} -rational point)

Non-examinable: If Y comes with a morph $Y \rightarrow \text{Spec } \mathbb{K}$ (hence $\theta_{Y,U}$ are \mathbb{K} -algebras) and above require morphs to commute with π , then get $\text{Hom}_{\mathbb{K}}(\mathbb{K}(y), \mathbb{K})$, and if $\mathbb{K}(y) \cong \mathbb{K}$ then $\text{Hom}_{\mathbb{K}}(\mathbb{K}, \mathbb{K}) = \{\text{id}_{\mathbb{K}}\}$. E.g. $\text{Spec } (\mathbb{C})$ has many \mathbb{C} -points: one for each automorphism of \mathbb{C} (e.g. $\sigma \rightarrow \sigma, \tau \rightarrow \bar{\tau}$) but if work over \mathbb{C} get only one \mathbb{C} -point.

Example 1 $\text{Spec } \mathbb{K} \rightarrow \text{Spec } \mathbb{K}(y)$
 A local hom $R \xrightarrow{\varphi} \mathbb{K} = \text{field}$ factors $R \xrightarrow{\text{quot}} \mathbb{K} \rightarrow \mathbb{K}$
 (since $\ker \varphi = \varphi^{-1}(0) = \mathfrak{m}$) (since local hom)

Example 2 $\text{Spec } \mathbb{K} \rightarrow \text{Spec } \mathbb{K}(y)$
 Thus: $\{f \in \text{Mor}(\text{Spec } \mathbb{K}, Y) \mid f(\text{pt}) = y\} \xrightarrow{1:1} \text{Hom}(\mathbb{K}(y), \mathbb{K})$ and any $\text{Spec } \mathbb{K} \rightarrow Y$ factors:
 $\text{Spec } \mathbb{K} \rightarrow \text{Spec } \mathbb{K}(y) \rightarrow Y$

UPSHOT: Morphs from local rings or fields don't give more information than already know from $\text{Spec } \theta_{X,x} \rightarrow X$ and $\text{Spec } \mathbb{K}(x) \rightarrow X$.
 \mathbb{K} -valued point if $\mathbb{K}(y) \cong \mathbb{K}$, then $\text{id}_{\mathbb{K}}$ defines a morph $\text{Spec } \mathbb{K} \rightarrow Y$
 (or \mathbb{K} -point or \mathbb{K} -rational point)

Non-examinable: If Y comes with a morph $Y \rightarrow \text{Spec } \mathbb{K}$ (hence $\theta_{Y,U}$ are \mathbb{K} -algebras) and above require morphs to commute with π , then get $\text{Hom}_{\mathbb{K}}(\mathbb{K}(y), \mathbb{K})$, and if $\mathbb{K}(y) \cong \mathbb{K}$ then $\text{Hom}_{\mathbb{K}}(\mathbb{K}, \mathbb{K}) = \{\text{id}_{\mathbb{K}}\}$. E.g. $\text{Spec } (\mathbb{C})$ has many \mathbb{C} -points: one for each automorphism of \mathbb{C} (e.g. $\sigma \rightarrow \sigma, \tau \rightarrow \bar{\tau}$) but if work over \mathbb{C} get only one \mathbb{C} -point.

Example 1 $\text{Spec } \mathbb{K} \rightarrow \text{Spec } \mathbb{K}(y)$
 A local hom $R \xrightarrow{\varphi} \mathbb{K} = \text{field}$ factors $R \xrightarrow{\text{quot}} \mathbb{K} \rightarrow \mathbb{K}$
 (since $\ker \varphi = \varphi^{-1}(0) = \mathfrak{m}$) (since local hom)

Example 2 $\text{Spec } \mathbb{K} \rightarrow \text{Spec } \mathbb{K}(y)$
 Thus: $\{f \in \text{Mor}(\text{Spec } \mathbb{K}, Y) \mid f(\text{pt}) = y\} \xrightarrow{1:1} \text{Hom}(\mathbb{K}(y), \mathbb{K})$ and any $\text{Spec } \mathbb{K} \rightarrow Y$ factors:
 $\text{Spec } \mathbb{K} \rightarrow \text{Spec } \mathbb{K}(y) \rightarrow Y$

UPSHOT: Morphs from local rings or fields don't give more information than already know from $\text{Spec } \theta_{X,x} \rightarrow X$ and $\text{Spec } \mathbb{K}(x) \rightarrow X$.
 \mathbb{K} -valued point if $\mathbb{K}(y) \cong \mathbb{K}$, then $\text{id}_{\mathbb{K}}$ defines a morph $\text{Spec } \mathbb{K} \rightarrow Y$
 (or \mathbb{K} -point or \mathbb{K} -rational point)

Non-examinable: If Y comes with a morph $Y \rightarrow \text{Spec } \mathbb{K}$ (hence $\theta_{Y,U}$ are \mathbb{K} -algebras) and above require morphs to commute with π , then get $\text{Hom}_{\mathbb{K}}(\mathbb{K}(y), \mathbb{K})$, and if $\mathbb{K}(y) \cong \mathbb{K}$ then $\text{Hom}_{\mathbb{K}}(\mathbb{K}, \mathbb{K}) = \{\text{id}_{\mathbb{K}}\}$. E.g. $\text{Spec } (\mathbb{C})$ has many \mathbb{C} -points: one for each automorphism of \mathbb{C} (e.g. $\sigma \rightarrow \sigma, \tau \rightarrow \bar{\tau}$) but if work over \mathbb{C} get only one \mathbb{C} -point.

Example 2 $X = \text{Spec } R \Rightarrow \{f \in \text{Mor}(\text{Spec } R, Y) \mid f(\text{pt}) = y\} \xrightarrow{1:1} \text{Hom}_{\text{local hom of rings}}(\theta_{Y,y}, R)$
 $m \subseteq R = \text{local ring}$ with $f(m) = \mathfrak{m}$

Pf $\text{Spec } R \xrightarrow{f} Y$
 $\text{Spec } R \xrightarrow{f} Y$
 $m \xrightarrow{f} \mathfrak{m}$

Affine case $Y = \text{Spec } S$
 $\varphi: S_y \rightarrow R \Rightarrow \text{Spec } R \rightarrow \text{Spec } S$
 $\varphi^{-1}(m) = y \subseteq S_y$
 $m \xrightarrow{\varphi} \mathfrak{m}$

General case
 $y \in U \subseteq Y$ open affine, then $\theta_{U,y} = \theta_{Y,y} \circ \varphi$ gives $\text{Spec } R \rightarrow U \subseteq Y$
 uniqueness: Suppose $f: \text{Spec } R \rightarrow Y$ gives same φ

pick $y \in V \subseteq Y$ affine open $\Rightarrow f^{-1}(V)$ open $\ni m = (\text{unique closed point of } \text{Spec } R) \Rightarrow f^{-1}(m) = \text{Spec } R$
 so $f: \text{Spec } R \rightarrow V \subseteq Y$ so reduce to affine case. \square

Cor 2 $x \in X \Rightarrow \exists$ canonical morph $\text{Spec } \theta_{X,x} \rightarrow X$
 Any $\text{Spec } R \rightarrow X$ factors as $\text{Spec } R \rightarrow \text{Spec } \theta_{X,x} \rightarrow X$ some $x \in X$
 Any $f: X \rightarrow Y$ of schemes get $\text{Spec } \theta_{X,x} \rightarrow X \xrightarrow{f} Y$
 $\text{Spec } \theta_{X,x} \rightarrow Y$ induced by f_x

Example Case $X = \text{Spec } \mathbb{K}$ for field \mathbb{K} .
 R local \Rightarrow residue field $\mathbb{K} = R/\mathfrak{m}$
 A local hom $R \xrightarrow{\varphi} \mathbb{K} = \text{field}$ factors $R \xrightarrow{\text{quot}} \mathbb{K} \rightarrow \mathbb{K}$
 (since $\ker \varphi = \varphi^{-1}(0) = \mathfrak{m}$) (since local hom)

Example 1 $\text{Spec } \mathbb{K} \rightarrow \text{Spec } \mathbb{K}(y)$
 Thus: $\{f \in \text{Mor}(\text{Spec } \mathbb{K}, Y) \mid f(\text{pt}) = y\} \xrightarrow{1:1} \text{Hom}(\mathbb{K}(y), \mathbb{K})$ and any $\text{Spec } \mathbb{K} \rightarrow Y$ factors:
 $\text{Spec } \mathbb{K} \rightarrow \text{Spec } \mathbb{K}(y) \rightarrow Y$

UPSHOT: Morphs from local rings or fields don't give more information than already know from $\text{Spec } \theta_{X,x} \rightarrow X$ and $\text{Spec } \mathbb{K}(x) \rightarrow X$.
 \mathbb{K} -valued point if $\mathbb{K}(y) \cong \mathbb{K}$, then $\text{id}_{\mathbb{K}}$ defines a morph $\text{Spec } \mathbb{K} \rightarrow Y$
 (or \mathbb{K} -point or \mathbb{K} -rational point)

Non-examinable: If Y comes with a morph $Y \rightarrow \text{Spec } \mathbb{K}$ (hence $\theta_{Y,U}$ are \mathbb{K} -algebras) and above require morphs to commute with π , then get $\text{Hom}_{\mathbb{K}}(\mathbb{K}(y), \mathbb{K})$, and if $\mathbb{K}(y) \cong \mathbb{K}$ then $\text{Hom}_{\mathbb{K}}(\mathbb{K}, \mathbb{K}) = \{\text{id}_{\mathbb{K}}\}$. E.g. $\text{Spec } (\mathbb{C})$ has many \mathbb{C} -points: one for each automorphism of \mathbb{C} (e.g. $\sigma \rightarrow \sigma, \tau \rightarrow \bar{\tau}$) but if work over \mathbb{C} get only one \mathbb{C} -point.

Example 1 $\text{Spec } \mathbb{K} \rightarrow \text{Spec } \mathbb{K}(y)$
 A local hom $R \xrightarrow{\varphi} \mathbb{K} = \text{field}$ factors $R \xrightarrow{\text{quot}} \mathbb{K} \rightarrow \mathbb{K}$
 (since $\ker \varphi = \varphi^{-1}(0) = \mathfrak{m}$) (since local hom)

Example 2 $\text{Spec } \mathbb{K} \rightarrow \text{Spec } \mathbb{K}(y)$
 Thus: $\{f \in \text{Mor}(\text{Spec } \mathbb{K}, Y) \mid f(\text{pt}) = y\} \xrightarrow{1:1} \text{Hom}(\mathbb{K}(y), \mathbb{K})$ and any $\text{Spec } \mathbb{K} \rightarrow Y$ factors:
 $\text{Spec } \mathbb{K} \rightarrow \text{Spec } \mathbb{K}(y) \rightarrow Y$

UPSHOT: Morphs from local rings or fields don't give more information than already know from $\text{Spec } \theta_{X,x} \rightarrow X$ and $\text{Spec } \mathbb{K}(x) \rightarrow X$.
 \mathbb{K} -valued point if $\mathbb{K}(y) \cong \mathbb{K}$, then $\text{id}_{\mathbb{K}}$ defines a morph $\text{Spec } \mathbb{K} \rightarrow Y$
 (or \mathbb{K} -point or \mathbb{K} -rational point)

Non-examinable: If Y comes with a morph $Y \rightarrow \text{Spec } \mathbb{K}$ (hence $\theta_{Y,U}$ are \mathbb{K} -algebras) and above require morphs to commute with π , then get $\text{Hom}_{\mathbb{K}}(\mathbb{K}(y), \mathbb{K})$, and if $\mathbb{K}(y) \cong \mathbb{K}$ then $\text{Hom}_{\mathbb{K}}(\mathbb{K}, \mathbb{K}) = \{\text{id}_{\mathbb{K}}\}$. E.g. $\text{Spec } (\mathbb{C})$ has many \mathbb{C} -points: one for each automorphism of \mathbb{C} (e.g. $\sigma \rightarrow \sigma, \tau \rightarrow \bar{\tau}$) but if work over \mathbb{C} get only one \mathbb{C} -point.

Example 1 $\text{Spec } \mathbb{K} \rightarrow \text{Spec } \mathbb{K}(y)$
 A local hom $R \xrightarrow{\varphi} \mathbb{K} = \text{field}$ factors $R \xrightarrow{\text{quot}} \mathbb{K} \rightarrow \mathbb{K}$
 (since $\ker \varphi = \varphi^{-1}(0) = \mathfrak{m}$) (since local hom)

Example 2 $\text{Spec } \mathbb{K} \rightarrow \text{Spec } \mathbb{K}(y)$
 Thus: $\{f \in \text{Mor}(\text{Spec } \mathbb{K}, Y) \mid f(\text{pt}) = y\} \xrightarrow{1:1} \text{Hom}(\mathbb{K}(y), \mathbb{K})$ and any $\text{Spec } \mathbb{K} \rightarrow Y$ factors:
 $\text{Spec } \mathbb{K} \rightarrow \text{Spec } \mathbb{K}(y) \rightarrow Y$

UPSHOT: Morphs from local rings or fields don't give more information than already know from $\text{Spec } \theta_{X,x} \rightarrow X$ and $\text{Spec } \mathbb{K}(x) \rightarrow X$.
 \mathbb{K} -valued point if $\mathbb{K}(y) \cong \mathbb{K}$, then $\text{id}_{\mathbb{K}}$ defines a morph $\text{Spec } \mathbb{K} \rightarrow Y$
 (or \mathbb{K} -point or \mathbb{K} -rational point)

Non-examinable: If Y comes with a morph $Y \rightarrow \text{Spec } \mathbb{K}$ (hence $\theta_{Y,U}$ are \mathbb{K} -algebras) and above require morphs to commute with π , then get $\text{Hom}_{\mathbb{K}}(\mathbb{K}(y), \mathbb{K})$, and if $\mathbb{K}(y) \cong \mathbb{K}$ then $\text{Hom}_{\mathbb{K}}(\mathbb{K}, \mathbb{K}) = \{\text{id}_{\mathbb{K}}\}$. E.g. $\text{Spec } (\mathbb{C})$ has many \mathbb{C} -points: one for each automorphism of \mathbb{C} (e.g. $\sigma \rightarrow \sigma, \tau \rightarrow \bar{\tau}$) but if work over \mathbb{C} get only one \mathbb{C} -point.

Example 1 $\text{Spec } \mathbb{K} \rightarrow \text{Spec } \mathbb{K}(y)$
 A local hom $R \xrightarrow{\varphi} \mathbb{K} = \text{field}$ factors $R \xrightarrow{\text{quot}} \mathbb{K} \rightarrow \mathbb{K}$
 (since $\ker \varphi = \varphi^{-1}(0) = \mathfrak{m}$) (since local hom)

Example 2 $\text{Spec } \mathbb{K} \rightarrow \text{Spec } \mathbb{K}(y)$
 Thus: $\{f \in \text{Mor}(\text{Spec } \mathbb{K}, Y) \mid f(\text{pt}) = y\} \xrightarrow{1:1} \text{Hom}(\mathbb{K}(y), \mathbb{K})$ and any $\text{Spec } \mathbb{K} \rightarrow Y$ factors:
 $\text{Spec } \mathbb{K} \rightarrow \text{Spec } \mathbb{K}(y) \rightarrow Y$

UPSHOT: Morphs from local rings or fields don't give more information than already know from $\text{Spec } \theta_{X,x} \rightarrow X$ and $\text{Spec } \mathbb{K}(x) \rightarrow X$.
 \mathbb{K} -valued point if $\mathbb{K}(y) \cong \mathbb{K}$, then $\text{id}_{\mathbb{K}}$ defines a morph $\text{Spec } \mathbb{K} \rightarrow Y$
 (or \mathbb{K} -point or \mathbb{K} -rational point)

Non-examinable: If Y comes with a morph $Y \rightarrow \text{Spec } \mathbb{K}$ (hence $\theta_{Y,U}$ are \mathbb{K} -algebras) and above require morphs to commute with π , then get $\text{Hom}_{\mathbb{K}}(\mathbb{K}(y), \mathbb{K})$, and if $\mathbb{K}(y) \cong \mathbb{K}$ then $\text{Hom}_{\mathbb{K}}(\mathbb{K}, \mathbb{K}) = \{\text{id}_{\mathbb{K}}\}$. E.g. $\text{Spec } (\mathbb{C})$ has many \mathbb{C} -points: one for each automorphism of \mathbb{C} (e.g. $\sigma \rightarrow \sigma, \tau \rightarrow \bar{\tau}$) but if work over \mathbb{C} get only one \mathbb{C} -point.

Non-examinable (see C3.4 Notes on Lasker-Noether theorem)

To recover the scheme $\text{Spec}(R) = \bigcup \mathbb{V}(q_i)$, $\mathbb{V}(q_i) \not\subseteq \bigcup_{j \neq i} \mathbb{V}(q_j)$ need primary decomposition \leftarrow (like "unique factorisation" but for ideals)

$\{0\} = q_1 \cap q_2 \cap \dots \cap q_n \cap \dots \cap q_m$ where q_i are **primary ideals** s.t. $q_i \not\subseteq q_j$ for $i \neq j$

$q \in R$ primary ideal if zero divisors of R/q are nilpotent (Equivalently: $ab \in q \Rightarrow a \in q$ or $b \in q$ some n (\Rightarrow if $a, b \notin q$ then $a, b \in \sqrt{q}$))

Example p^2 is primary if p prime ideal, e.g. $(3^2) \subseteq \mathbb{Z}$

Example $(18) = (2 \cdot 3^2) = (2) \cap (3^2) \subseteq \mathbb{Z}$ is primary decomposition.

The q_i are not unique, but the $\sqrt{q_i}$ are unique (up to reordering) (the p_i are precisely the prime ideals arising as radicals of annihilators of e.l.s of R)

The $\mathbb{V}(q_i)$ are called **primary components**: not unique as schemes, but are unique topologically.

WLOG $p_1 = \sqrt{q_1}, \dots, p_n = \sqrt{q_n}$ are as in previous exercise: the **minimal prime ideals** give the **isolated components** $\mathbb{V}(q_i)$ (as top subspace $= \mathbb{V}(p_i)$ irreducible comp.). These q_1, \dots, q_n are unique.

The other q_{n+1}, \dots, q_m give rise to the **embedded components** $\mathbb{V}(q_j)$, $j > n+1$ (not unique).

(Note $p_j \supseteq p_i$ so $\mathbb{V}(p_j) \subseteq \mathbb{V}(p_i) \subseteq \mathbb{V}(q_i)$ are closed subschemes, but $\mathbb{V}(q_j) \not\subseteq \mathbb{V}(p_i)$ as scheme)

RMK Can apply above to R/I to get $\sqrt{I} = p_1 \cap \dots \cap p_n$, $I = q_1 \cap \dots \cap q_n \cap \dots \cap q_m$, etc.

Example $I = (y^2, xy) \subseteq k[x, y] = R$, $X = \text{Spec}(R/I) = \mathbb{A}^1$ (as top space) $\mathbb{A}^1 = \mathbb{A}^1 \cup \{0\}$ for $q_1 = (y)$, $p_1 = (y)$ min prime, $\mathbb{V}(q_1)$ is isolated, irreducible

Think: functions vanishing on $q_2 = (x, y)^2$ embedded prim $\mathbb{V}(q_2) = \{0\}$ "fattened origin" is embedded

(max length of chain of ideals \mathbb{A}^1 is 2, $2 = \text{max length}$)

order 2, $2 = \text{max length}$ (max length of chain of ideals \mathbb{A}^1 is 2, $2 = \text{max length}$)

order 2, $2 = \text{max length}$ (max length of chain of ideals \mathbb{A}^1 is 2, $2 = \text{max length}$)

order 2, $2 = \text{max length}$ (max length of chain of ideals \mathbb{A}^1 is 2, $2 = \text{max length}$)

order 2, $2 = \text{max length}$ (max length of chain of ideals \mathbb{A}^1 is 2, $2 = \text{max length}$)

order 2, $2 = \text{max length}$ (max length of chain of ideals \mathbb{A}^1 is 2, $2 = \text{max length}$)

order 2, $2 = \text{max length}$ (max length of chain of ideals \mathbb{A}^1 is 2, $2 = \text{max length}$)

order 2, $2 = \text{max length}$ (max length of chain of ideals \mathbb{A}^1 is 2, $2 = \text{max length}$)

order 2, $2 = \text{max length}$ (max length of chain of ideals \mathbb{A}^1 is 2, $2 = \text{max length}$)

Claim $X = \bigcup \text{Spec } R_i$ each has $\star \Rightarrow$ every open affine in X has \star \leftarrow "if holds for a cover, it holds for affine open"

Pf $\text{Spec } R \rightarrow X \Rightarrow \text{Spec } R = \bigcup_{\text{finite } i} D_{f_i} \Rightarrow \text{Spec } R_i \Rightarrow D_{f_i} \Rightarrow \text{Spec } R \Rightarrow \text{Spec } R_i$ (use useful trick in 3.1)

Examples of \star : "ring is reduced", "ring is Noetherian", "ring is f.g. B-algebra" (some fixed ring B "base")

"locally of finite type over B " \leftarrow "locally of finite type over B "

so \exists surj: hom of B -alg $B[x_1, \dots, x_n] \rightarrow \text{ring}$ e.g. fixed k : Affine vars $X \subseteq \mathbb{A}^n$ loc. finite type/ k .

3.3 Reduced schemes (X, \mathcal{O}_X) reduced if all $\mathcal{O}_X(U)$ reduced rings (= no nilpotents $\neq 0$)

Hwk 1 reduced \Leftrightarrow stalks $\mathcal{O}_{X, x}$ are reduced $\Leftrightarrow \forall p \in X$ has an open affine neighbourhood for a reduced ring

RMK By 3.2: $\text{Spec } R$ reduced $\Leftrightarrow R$ reduced

Lemma X reduced, $f, g \in \mathcal{O}_X(U)$ take same values $f(x) = g(x) \in X(x) = \mathcal{O}_X(x)/\mathfrak{m}_x \Rightarrow f = g$

Pf. Take $f - g$, WLOG $g = 0$. On affine, $K(p) \subseteq \text{Frac}(R_p)$ so $f \in \cap p = \text{Nilradical}(R) = \{0\}$ (Don't confuse this with general fact \forall scheme: $f_x = g_x \in \mathcal{O}_{X, x} \forall x \in U \Rightarrow f = g \in \mathcal{O}_X(U)$)

Claim (not that strong a condition e.g. $f, g: \mathbb{C} \rightarrow \mathbb{C}, f(z) = z, g(z) = \bar{z}$ different, but $f(0) = g(0)$, $\text{Spec } \mathbb{C} = \{0\}$)

X reduced, $f, g: X \rightarrow Y, f = g$ as topological maps, $f = g$ on open dense set $\Rightarrow f = g$. \mathbb{A}^1 (Spec R) enough show $f = g$ locally by sheaf property. WLOG $Y = \text{Spec } R, X = \text{Spec } S \subseteq \mathbb{A}^1$ (Spec R)

$\varphi := f^* - g^*: R \rightarrow S$: to show φ vanishes it is enough to show $S \cap \ker \varphi = \{0\}$ is zero \leftarrow (if $\varphi = \varphi(\text{nil})$)

$\{p \in \text{Spec } S: S(p) = 0 \in K(p)\} = \mathbb{V}(S)$ closed & contains an open dense set, hence $S = 0$ by Lemma. \square

3.4 Irreducible schemes Def Topological space X is **irreducible** if X is not a union of 2 proper closed sets: $X = C_1 \cup C_2 \Rightarrow X = C_1$ or $X = C_2$ (where C_i closed)

Easy exercise If X irreducible: Any non-empty open $U \subseteq X$ is dense and irreducible. Any two " " U_1, U_2 have $U_1 \cap U_2 \neq \emptyset$ (open, dense, irred)

Recall: $\text{Nil}(R) = \text{nilradical}(R) = \{ \text{nilpotent elements} \} = \sqrt{(0)} = \bigcap \{ p \in \text{Spec } R \}$ (R ring)

Hwk 2 (X, \mathcal{O}_X) irreducible \Leftrightarrow nilpotent elements are irreducible \Rightarrow all affine opens are irreducible

Hwk 1 $\text{Spec } R$ irreducible $\Leftrightarrow \text{Nil}(R)$ prime ideal $\Rightarrow R/\text{Nil}(R)$ integral domain

Recall $p \in X$ generic point if closure $\bar{p} = X$ (p is dense) $\Leftrightarrow \exists!$ generic point, namely $\text{Nil}(R)$

Claim (X, \mathcal{O}_X) irreducible $\Rightarrow \exists!$ generic point y , and $y \in$ every affine open $\neq \emptyset$

Pf affine open $\emptyset \neq U \subseteq X \Rightarrow U$ irred. $\Rightarrow \exists!$ generic point $x \in U \Rightarrow \bar{x} = U = X$ (\bar{x} in X closed and $2U$)

Suppose $y \in X$ generic \Rightarrow if $y \in X \cap U$ then $\bar{y} \subseteq X \cap U$ not dense, so $y \in U$, so $y = x$. \square

Hwk 2 irreducible \Leftrightarrow connected. Fact $\text{Spec } R$ connected \Leftrightarrow no idempotents $\neq 0, 1$ ($\neq x \cup U, U_2$ for disjoint open $U_1 \neq \emptyset$)

\leftarrow classifies connected components of $\text{Spec } R$ in terms of idempotents $\leftarrow r \in R$ with $r^2 = r$

Exercise R Noetherian $\Rightarrow \exists!$ sequence of prime ideals p_1, \dots, p_n (up to reordering): $\bigcap p_i = \text{Nil}(R)$ (Same pt. as in C3.4)

$\Rightarrow \exists!$ sequence of irred. closed subsets $C_i = \mathbb{V}(p_i)$ (up to reordering): $\text{Spec } R = \bigcup C_i, C_i \not\subseteq \bigcup_{j \neq i} C_j$

(which as top. subspaces are the irreducible components) as topological spaces

Warning: $q = (x^2) \subseteq k[x] = R \Rightarrow p = \text{Nil}(R_q) = \{0\} = \text{Spec}(R_q) = \{0\} = \text{Spec}(R/p) = \{0\} = \text{Spec}(R/p)$ as top. spaces, not as schemes

RMK $p = \sqrt{q}$ is prime ideal ("associated prime ideal") and is smallest prime ideal containing q . So: $a \in q, a \notin q \Rightarrow b \in p$ ($\mathbb{V}(q_i) = \mathbb{V}(p_i)$) (as closed sets)

so "irredundant!" \leftarrow can't omit q_i

non-examinable fact if X is locally Noetherian \leftarrow X integral $\Leftrightarrow X$ locally Noetherian & connected

X integral $\Leftrightarrow X = \text{USpec } R$ \leftarrow integral $\Leftrightarrow R$ integral

Spec R integral $\Leftrightarrow R$ integral domain \leftarrow Example All irreducible affine varieties $X \subseteq \mathbb{A}^n$ ($\text{Spec } k[X]$)

(X, \mathcal{O}_X) integral \Rightarrow restrictions $\mathcal{O}_X(U) \rightarrow \mathcal{O}_X(V)$ are injective for $V \neq \emptyset$

\Rightarrow all sections can be compared in $\mathcal{O}_{X, y} \leftarrow \mathcal{O}_X(U) \rightarrow \mathcal{O}_{X, y}$ generic point

$\cdot K(y) \cong \mathcal{O}_{X, y} \cong \text{Frac } \mathcal{O}_X(U)$ via restriction (any $U \neq \emptyset$)

Pf $\mathcal{O}_X(U) \rightarrow \mathcal{O}_X(V) \rightarrow \mathcal{O}_{X, y}$ so enough show $s_x = 0$ all $x \in U$ so $s = 0 \in \mathcal{O}_X(U)$.

If show $s = 0$ on every open affine $\subseteq U$ then $s_x = 0$ all $x \in U$ so $s = 0 \in \mathcal{O}_X(U)$.

\Rightarrow WLOG $U = \text{Spec } R, y = \text{Nil}(R) = \{0\}$ (since R is ID). Thus $s_y = 0 \Rightarrow s = 0$.

$R \hookrightarrow R_{(0)} = \text{Frac } R, r \mapsto \frac{r}{1}$ inj. since R is ID. Thus $s_y = 0 \Rightarrow s = 0$.

Classical Alg. Geometry $X \subseteq \mathbb{A}^n$ irred. affine var $\Rightarrow \mathcal{O}_X(X) \rightarrow \mathcal{O}_X(Df) \rightarrow \mathcal{O}_{X, p}$ \leftarrow (so $\text{Spec } k[X] \subseteq k[X]_p \subseteq \text{Frac } k[X]$)

non-examinable fact if X is locally Noetherian & connected \leftarrow X integral $\Leftrightarrow X = \text{USpec } R$ \leftarrow integral $\Leftrightarrow R$ integral

$k[x, y]/(xy) \cong k[x] \oplus k[y]$ reducible: union of two axes

2 Key Non-Examples \leftarrow "fat lines" \leftarrow $k[x, y]/(x^2)$ not reduced

Hwk 2 X integral \Leftrightarrow reduced and irreducible

Fact Localisation Direct limits \lim preserve ID property

Cor X integral $\Rightarrow \mathcal{O}_{X, x}$ ID (but not \leftarrow)

Hwk 2 X integral \Leftrightarrow reduced and irreducible

Spec R integral $\Leftrightarrow R$ integral domain \leftarrow Example All irreducible affine varieties $X \subseteq \mathbb{A}^n$ ($\text{Spec } k[X]$)

(X, \mathcal{O}_X) integral if all $\mathcal{O}_X(U)$ ID \leftarrow (integral domain = no zero divisors $\neq 0$)

Hwk 2 X affine open U \leftarrow preserve ID property

Fact X integral $\Rightarrow \mathcal{O}_{X, x}$ ID (but not \leftarrow)

Hwk 2 X integral \Leftrightarrow reduced and irreducible

Spec R integral $\Leftrightarrow R$ integral domain \leftarrow Example All irreducible affine varieties $X \subseteq \mathbb{A}^n$ ($\text{Spec } k[X]$)

(X, \mathcal{O}_X) integral if all $\mathcal{O}_X(U)$ ID \leftarrow (integral domain = no zero divisors $\neq 0$)

Hwk 2 X affine open U \leftarrow preserve ID property

Fact X integral $\Rightarrow \mathcal{O}_{X, x}$ ID (but not \leftarrow)

Hwk 2 X integral \Leftrightarrow reduced and irreducible

Spec R integral $\Leftrightarrow R$ integral domain \leftarrow Example All irreducible affine varieties $X \subseteq \mathbb{A}^n$ ($\text{Spec } k[X]$)

(X, \mathcal{O}_X) integral if all $\mathcal{O}_X(U)$ ID \leftarrow (integral domain = no zero divisors $\neq 0$)

Hwk 2 X affine open U \leftarrow preserve ID property

Fact X integral $\Rightarrow \mathcal{O}_{X, x}$ ID (but not \leftarrow)

Hwk 2 X integral \Leftrightarrow reduced and irreducible

Spec R integral $\Leftrightarrow R$ integral domain \leftarrow Example All irreducible affine varieties $X \subseteq \mathbb{A}^n$ ($\text{Spec } k[X]$)

(X, \mathcal{O}_X) integral if all $\mathcal{O}_X(U)$ ID \leftarrow (integral domain = no zero divisors $\neq 0$)

Hwk 2 X affine open U \leftarrow preserve ID property

Fact X integral $\Rightarrow \mathcal{O}_{X, x}$ ID (but not \leftarrow)

Hwk 2 X integral \Leftrightarrow reduced and irreducible

Spec R integral $\Leftrightarrow R$ integral domain \leftarrow Example All irreducible affine varieties $X \subseteq \mathbb{A}^n$ ($\text{Spec } k[X]$)

(X, \mathcal{O}_X) integral if all $\mathcal{O}_X(U)$ ID \leftarrow (integral domain = no zero divisors $\neq 0$)

Hwk 2 X affine open U \leftarrow preserve ID property

Fact X integral $\Rightarrow \mathcal{O}_{X, x}$ ID (but not \leftarrow)

3.6 Properties of morphisms ← all properties we list are preserved when compose such morphs

A morph of schemes $f: X \rightarrow Y$ is: (will suppress f^*, θ_x, θ_y from notation)

- affine: Equivalent conditions:
 - f^{-1} (affine open) is **affine**
 - \exists affine open cover V_i of Y , $f^{-1}(V_i)$ **affine**
 - \forall affine open cover V_i of Y , $f^{-1}(V_i)$ **affine**
- quasi-compact: replace **affine** by **quasi-compact**
- locally of finite type: \forall affine opens $U \subseteq X, V \subseteq Y$ with $f(U) \subseteq V$, $f^*: \mathcal{O}_Y(V) \rightarrow \mathcal{O}_X(U)$ finite type
 (Rings: $A \rightarrow B$ finite type means B f.g. as A -alg., i.e. \exists sgs $A[x_1, \dots, x_n] \rightarrow B$ of A -mods.)
 (meaning: $\theta_y(V) \xrightarrow{f^*} \theta_x(f^{-1}V) \xrightarrow{\text{rest}} \mathcal{O}_X(U)$)
- finite type: \exists open affine covers $Y = \cup V_i$, $f^{-1}(V_i) = \cup U_{ij}$ finite type
 $f^*: \theta_y(V_i) \rightarrow \theta_x(U_{ij})$ finite type
- closed immersion: iso onto a closed subscheme.

Explicitly: $f: X \xrightarrow{\text{homeo}} f(X) \subseteq Y$
 $f^*: \theta_y \rightarrow f_* \theta_x$ surjective (so ideal sheaf $J = \ker f^*$)
 \forall aff. open $U = \text{Spec } R \subseteq Y \exists$ ideal $I \subseteq R$ s.t. $f^{-1}(U) \cong \text{Spec}(R/I)$
 \exists aff. cover $Y = \cup \text{Spec } R_i$, ideals $I_i \subseteq R_i$, $f^{-1}(\text{Spec } R_i) = \text{Spec}(R_i/I_i)$

Example $X = Y_{\text{red}} \subseteq Y$ closed subscheme: $X = Y$ as topological space and sheaf of ideals $J(U) = \{s \in \mathcal{O}_Y(U) : s(p) = 0 \in \mathcal{K}(p), \forall p \in U\}$ (so $\theta_x = \theta_y/J$)

Note locally: on $U = \text{Spec } R, J(U) = \{s \in R : s \in \mathcal{P} = \text{Nil}(R) = \{nilpotents\}\}$, so locally J agrees with $\text{Nil}(\theta_y)$, indeed J is the sheafification of $\text{Nil}(\theta_y)$ ← need not be sheaf e.g. $Y = \mathbb{A}^1, X_n = \text{Spec}(\mathbb{Z}/p^n)$ $z \in \theta_y(Y), z \notin \text{Nil}(\theta_y(Y))$ but $z \in \text{Nil}(\theta_y(Y_n)), z \in J(X)$

- open immersion: iso onto an open subscheme $\leftarrow U \subseteq Y, U = \theta_y|_U$
 (idea: functions on X are the same as θ_y locally)
- flat: all $\theta_{y,fx} \rightarrow \theta_{x,x}$ are flat ring homs

Not intuitively clear, but ensures that fibers of f vary in a controlled way: Many invariants of fibers like dimension, do not change unless you "expect" it! It is weaker than saying the fibers are locally iso e.g. it allows two points to collide as vary fiber

Algebra: R -mod M is flat if $M \otimes_R \cdot$ is exact functor on R -mods

$\varphi: R \rightarrow S$ flat ring hom means S flat R -mod (using $r \cdot s = \varphi(r) \cdot s$)

Basic facts

- $M \otimes_R \cdot$ always right exact, so M flat R -mod $\Leftrightarrow M \hookrightarrow N_1 \hookrightarrow M \otimes_R N_2$
 Fact Enough to check $M \otimes_R I \hookrightarrow M \otimes_R R \forall$ f.g. ideal $I \subseteq R$.
 - M free $\Rightarrow M$ flat (pf. $M \cong \bigoplus_{i \in I} R \Rightarrow M \otimes N \cong \bigoplus_{i \in I} N$)
- Example $\prod_{\text{infinite}} \mathbb{Z}$ is not free \mathbb{Z} -mod, but it is flat. An abelian gp is flat \mathbb{Z} -mod \Leftrightarrow torsion free

Non-example \mathbb{Z}/n is not flat \mathbb{Z} -mod $\leftarrow \mathbb{Z} \xrightarrow{\cdot n} \mathbb{Z}$ then $\cdot \otimes \mathbb{Z}/n$ get \mathbb{Z}/n not inj.

Fact (Lazard) R -mod M is flat $\Leftrightarrow M = \varinjlim M_i$ some f.g. free R -mods M_i

3) R local, M finite R -mod (so $M = \sum_{\text{finite}} R m_i$): M flat $\Leftrightarrow M$ free $\leftarrow \theta_{y,fx}$ local but $\theta_{y,x}$ is rarely finite over C

4) $A \rightarrow B$ flat, $B \rightarrow C$ flat $\Rightarrow A \rightarrow C$ flat

pf $N_1 \hookrightarrow N_2$ A -mods $\Rightarrow B \otimes_A N_1 \hookrightarrow B \otimes_A N_2$ B -mods $\Rightarrow C \otimes_B B \otimes_A N_1 \hookrightarrow C \otimes_B B \otimes_A N_2$ \square

5) $A \rightarrow B$ flat $\Rightarrow A_p \rightarrow B_p = B \otimes_A A_p$ flat $\forall p \in \text{Spec } A$

pf $N_1 \hookrightarrow N_2$ A_p -mods $\Rightarrow N_{1,p} \hookrightarrow N_{2,p}$ A_p -mods (via $A \rightarrow A_p$) $\Rightarrow B \otimes_{A_p} N_{1,p} \hookrightarrow B \otimes_{A_p} N_{2,p}$ \square

6) Ring hom $\varphi: A \rightarrow B$, multiplicative sets $S \subseteq A, T \subseteq B$ with $\varphi(S) \subseteq T$, then $\psi: S^{-1}B = S^{-1}A \otimes_A B \rightarrow T^{-1}B, \frac{s}{t} \otimes b \mapsto \frac{\varphi(s)b}{\varphi(t)}$ factorizes as $S^{-1}B \xrightarrow{\varphi(S)^{-1}B} T^{-1}B \xrightarrow{\varphi(S)}$

Since isos of rings and localisation are exact functors, get ψ flat.

Example: $P \subseteq B$ prime ideal, $q = \varphi^{-1}(P)$ prime ideal, $S = A \setminus q, T = B \setminus P \Rightarrow B_q = B \otimes_A A_q \rightarrow B_p$ flat


Theorem $\varphi: A \rightarrow B$ flat ring hom $\Leftrightarrow \varphi^*: \text{Spec } B \rightarrow \text{Spec } A$ flat

pf \Rightarrow $A \rightarrow B$ flat $\Rightarrow A_q \rightarrow B_q$ flat for $q = \varphi^{-1}(p)$ by (5), $B_q \rightarrow B_p$ flat by (4) $\Rightarrow A_q \rightarrow B_p$ flat.

\Leftarrow Recall $\ker(B \otimes_A N_1 \rightarrow B \otimes_A N_2) \neq 0 \Leftrightarrow \ker \psi_p \neq 0 \forall p \in \text{Spec } B$.

$\ker(N_1 \rightarrow N_2) = 0 \Rightarrow \ker(A_q \otimes_A N_1 \rightarrow A_q \otimes_A N_2) = 0 \Rightarrow \ker(B_p \otimes_{A_q} A_q \otimes_A N_1 \rightarrow B_p \otimes_{A_q} A_q \otimes_A N_2) = 0$ flatness

Motivation: Deformations (see Homework 2 ex. 6) "Flatness \Rightarrow 1-parameter families of schemes have limits."



Fact Another nice properties of flat morphs $f: X \rightarrow B$, for B, X locally Noeth.: $\dim_x f^{-1}(b) = \dim_x X - \dim_b B$ where $b = f(x)$

So dimensions of fibers don't "jump" unexpectedly.

Geometrical motivation (very loosely) $X_f = V(y_j - t) \subseteq \mathbb{A}^2, X_0 = V(x_j)$



how many times does a line in \mathbb{A}^2 intersect fiber? $X = V(y_j - t) \subseteq \mathbb{A}^2, X_0 = \text{Spec } k[t, x_j]$

$\mathbb{A}^1 = \text{Spec } k[t]$

if have a family for which intersection number is constant, it may be easy to calculate for a degenerate fiber

example: \mathbb{A}^2 has $\dim = 2$
 $\{p\} \subseteq \text{line} \subseteq \text{plane}$
 $\mathbb{Z}_0 \subseteq \mathbb{Z}_1 \subseteq \mathbb{Z}_2$

in such theorems you will almost always see the flatness assumption

defined rigorously later in S.1, for now $X_B = \pi^{-1}(b) = \text{Spec } k(b)[x] \otimes_{k(b)} X$
 $= \text{Spec } (k(b) \otimes_{k(b)} R)$ if $X = \text{Spec } R$
 \leftarrow fiber X_0 is "limit" $\lim_{b \rightarrow 0} X_b$
 \leftarrow fiber over 0 of closure of $X^* = \pi^{-1}(b^*)$
 \leftarrow so $\Leftrightarrow X^* = X$ (see S.1: $(B^* \times B) \times X$)

Remarks about calculating closures of sets in $X = \text{Spec } R$

1) $p \in \text{Spec } R \Rightarrow \overline{p} = V(p)$

Pf $p \in V(p) \Rightarrow \overline{p} \subseteq V(p)$ (since $V(p)$ closed)
 Converse: $p \in \overline{p} \subseteq V(I) \Rightarrow I \subseteq p \Rightarrow q \in V(I) \cap p$
 $q \in V(p) \Rightarrow p \subseteq q$

Example $X^* = V(p_1, p_2, \dots, p_k) \subseteq \mathbb{A}_{\mathbb{B}}^n$, $p_j \in R[x_1, \dots, x_n, t^*]$ prime ideals
 $= V_{\mathbb{B}}(p_1) \cup \dots \cup V_{\mathbb{B}}(p_k)$ where $V_{\mathbb{B}}(\cdot)$ is $V(\cdot)$ calculated in $\mathbb{A}_{\mathbb{B}}^n$
 $\Rightarrow \overline{X^*} = V(p_1) \cup \dots \cup V(p_k) \subseteq \mathbb{A}_{\mathbb{B}}^n$ since $p_i \in X^* \subseteq X^*$
 $= V(p_1, p_2, \dots, p_k)$ and $p_i \in V_{\mathbb{B}}(p_i) \subseteq V(p_i) = \overline{p_i}$

2) For $\varphi: R \rightarrow S$ ring hom, $\alpha: \text{Spec } S \rightarrow \text{Spec } R$, $\alpha(p) = \varphi^{-1}p$:

Given $C = V(J) \subseteq \text{Spec } S$, $\overline{\alpha(C)} = V(\varphi^{-1}J)$

Pf $J = \sqrt{J} \Rightarrow \varphi^{-1}J = \bigcap_{\substack{J \subseteq P \\ P \in \text{Spec } S}} P$ since $\alpha(C) \subseteq \overline{\alpha(C)} = V(I)$, $I \subseteq \varphi^{-1}P$
 $\Rightarrow \varphi^{-1}P \in \text{Spec } S$ so $\varphi^{-1}P \in V(\varphi^{-1}J)$
 $\alpha(C) \subseteq V(\varphi^{-1}J)$

Example $S = R_f$ localisation, $f \in R$, if $\varphi: R \hookrightarrow R_f$ injection then $\varphi^{-1}J = R \cap J$
 e.g. $X^* = V(J) \subseteq \mathbb{A}_{\mathbb{B}}^n$ for $\mathbb{B} = \text{Spec } R[t]$, $\mathbb{B}^* = \text{Spec } R[t, t^{-1}]$
 so $\mathbb{A}_{\mathbb{B}}^n = \text{Spec } R[x_1, \dots, x_n, t]$, $\mathbb{A}_{\mathbb{B}^*}^n = R[x_1, \dots, x_n, t, t^{-1}]$
 $\Rightarrow \overline{X^*} = V(R[x_1, \dots, x_n, t] \cap J) \subseteq \mathbb{A}_{\mathbb{B}}^n$ is the closure

RMK Also know inverse images of closed sets: $\alpha^{-1}(V(I)) = V(\langle \varphi I \rangle)$

Pf $I = \langle f_i \rangle$, $\text{Spec } R \setminus V(I) = U D_{f_i}$,
 $U D_{\varphi f_i} = \alpha^{-1}(U D_{f_i}) = \alpha^{-1}(\text{Spec } R \setminus V(I)) = \text{Spec } S \setminus \alpha^{-1}V(I)$
 $\Rightarrow \alpha^{-1}V(I) = \text{Spec } S \setminus U D_{\varphi f_i} = V(\langle \varphi f_i \rangle)$
 (recall $\alpha^{-1}D_f = D_{\varphi f}$)

4- GLUING THEOREMS

4-1 Gluing sheaves


$X = \cup U_i$ open cover, abbreviate $U_{ij} = U_i \cap U_j$, $U_{ijk} = U_i \cap U_j \cap U_k$

F_i sheaf on U_i

$\varphi_{ij}: F_i|_{U_{ij}} \xrightarrow{\sim} F_j|_{U_{ij}}$

Compatibility conditions

- $\varphi_{ii} = \text{id}$
- $\varphi_{ji} = \varphi_{ij}^{-1}$
- $\varphi_{ik}|_{U_{ijk}} = \varphi_{jk} \circ \varphi_{ij}|_{U_{ijk}}$



Example F sheaf on X , $F_i := F|_{U_i}$ (so $F_i(V) = F(U_i \cap V) = F(U_i \cap V)$, $\forall \text{ open } V \subseteq U_i$)

φ_{ij} = isos induced by double restrictions (iso of functors $\cdot|_{U_i} \cdot|_{U_j} \cong \cdot|_{U_{ij}}$)


Theorem \exists , up to unique iso, a sheaf F on X with isos

$\psi_i: F|_{U_i} \xrightarrow{\sim} F_i$

s.t. $\psi_j^{-1} \circ \varphi_{ij} \circ \psi_i|_{U_{ij}} \cong F|_{U_{ij}} \cong F_j|_{U_{ij}}$ is the natural iso $F|_{U_i} \cong F|_{U_j}|_{U_{ij}}$

Pf Let $E = \bigsqcup_i (F_i)_x / \sim$ equivalence relation $(F_i)_x \sim (F_j)_x$ for $x \in U_{ij}$


$F(U) = \{s: U \rightarrow E : s \text{ is locally a section of some } F_i\}$ (using conditions)
 ($\forall x \in U, \exists i, \exists \text{ open } \mathcal{U}_x \subseteq U_i, \exists t \in F_i(\mathcal{U}_x), s|_{\mathcal{U}_x} = t$)



Theorem Given sheaves F, G constructed as above from local data F_i, φ_{ij} on U_i

a morph $f: F \rightarrow G$ can be uniquely defined from data:

- morphs $f_i: F_i \rightarrow G_i$
- compatibility condition: $\psi_j \circ f_i|_{U_{ij}} = f_j \circ \varphi_{ij}|_{U_{ij}}$



s.t. via identifications $F|_{U_i} \cong F_i, G|_{U_i} \cong G_i$ recover $f|_{U_i} = f_i$

4-2 Gluing schemes


U_i schemes, $U_{ij} \subseteq U_i$ open subschemes ($U_{ii} = U_i$)

$\varphi_{ij}: U_{ij} \xrightarrow{\cong} U_{ji}$ isos \leftarrow (think "go from U_i to U_j ")


gluing conditions 1) $\varphi_{ii} = \text{id}$

2) $\varphi_{ij}(U_{ij} \cap U_{ik}) \subseteq U_{ji} \cap U_{jk}$

3) $\varphi_{ik} = \varphi_{jk} \circ \varphi_{ij}$ when restricted as maps $U_{ij} \cap U_{ik} \rightarrow U_{jk}$



Example if $U_i \subseteq X$ open subschemes, can take $U_{ij} = U_i \cap U_j \subseteq X$ with $\varphi_{ij} = \text{id}$
Claim (exercise) \exists unique (up to iso) scheme X with open cover $X = \cup X_i$



Gluing Lemma Suppose we built X as above

$\Rightarrow f: X \rightarrow Y$ morph can be uniquely defined from morphs $f_i: X_i \rightarrow Y$ st.
 compatibility condition:
 $X_i \cap X_j \xrightarrow{\text{id}} X_i \xrightarrow{f_i} Y$
 $X_i \cap X_j \xrightarrow{\text{id}} X_j \xrightarrow{f_j} Y$ (Compatibility)

Pf Continuous map: $f: X \rightarrow Y$ defined by $f|_{X_i} = f_i$ (Compatibility)

on sheaves need $f^{-1}\theta_Y \rightarrow \theta_X \leftarrow$ (recall get $\theta_Y \rightarrow f_*\theta_X$ by adjunction)
 $(f^{-1}\theta_Y)|_{X_i} = f_i^{-1}\theta_Y = f_i^{-1}\theta_Y \leftarrow (X_i \xrightarrow{\varphi_i} X \text{ inclusion, then } \varphi_i^{-1}f_i^{-1}\theta_Y = (f \circ \varphi_i)^{-1}\theta_Y$
 $f_i^{\#} \in \text{Mor}(\theta_Y|_{X_i} \otimes \theta_{X_i}) \cong \text{Mor}(f_i^{-1}\theta_Y, \theta_{X_i})$ and $\theta_{X_i} = \theta_X|_{X_i}$ since open subs.
 Finally we can glue the $f_i^{\#}: f_i^{-1}\theta_Y \rightarrow \theta_X|_{X_i}$ by \oplus to get $f^{-1}\theta_Y \rightarrow \theta_X$. \square

Consequence $h_Y|_{\text{Top}(X)^{\text{op}}} : \text{Top}(X)^{\text{op}} \rightarrow \text{Sets}$
 $U \mapsto h_Y(U) = \text{Mor}(U, Y)$ is a sheaf of sets.

4.3 Affine space by gluing (see Homework for projective space)

Affine n -space over $\text{Spec } R: \mathbb{A}_R^n := \text{Spec } R[x_1, \dots, x_n] (= \mathbb{A}_{\text{Spec } R}^n)$ (comes with morph $\mathbb{A}_R^n \rightarrow \text{Spec } R$ via constant polys $R \rightarrow R[x_1, \dots, x_n]$)

Rmk $R \rightarrow S$ ring hom \Rightarrow hom on polys (so: $R[x_1, \dots, x_n] \rightarrow S[x_1, \dots, x_n] \cong \mathbb{A}_S^n \rightarrow \mathbb{A}_R^n$)

Example $R \rightarrow R_f \Rightarrow \mathbb{A}_R^n \rightarrow \mathbb{A}_R^n$ is the basic open set of \mathbb{A}_R^n for $f \in R \setminus \{0\}$

If $U \subseteq \text{Spec } R$ open $\Rightarrow U = \cup D_f$ (some $f_i \in R$)

X scheme, affine n -space over $X: \mathbb{A}_X^n := \cup \mathbb{A}_X^n$ where $X = \cup U_i$ affine open cover

(notice $\mathbb{A}_X^n = \cup \mathbb{A}_{X_i}^n$, then identify these copies) glued along $\mathbb{A}_{X_i \cap X_j}^n$ open in affine X_i

Claim $\mathbb{A}^n = \text{Spec } \mathbb{Z}[x_1, \dots, x_n]$ represents functor $F: \text{Sch}^{\text{op}} \rightarrow \text{Sets}, X \mapsto \{ \text{Morps } \mathbb{A}^n \times_{\mathbb{Z}} X \rightarrow X \text{ s.t. } \forall U, \mathbb{A}^n \times_{\mathbb{Z}} U \rightarrow U \text{ is hom of } \theta_U \text{ (sheaf)} \}$

Pf $F|_{\text{Top}(X)^{\text{op}}}$ is a sheaf of sets (easy to check: can glue morphs since θ_X sheaf)

$h_{\mathbb{A}^n}|_{\text{Top}(X)^{\text{op}}}$ by consequence above. Thus if the two functors agree on affines then

by sheaf property they agree everywhere. For affine $X = \text{Spec } R$ just need compare global sections

$F(\text{Spec } R) = \text{Hom}_R(R^n, R) \leftarrow$ (here: R -mod homs!) in both cases just $\{e_i = (0, \dots, 1, \dots, 0)\} \rightarrow R$


$h_{\mathbb{A}^n}(\text{Spec } R) = \text{Mor}(\text{Spec } R, \mathbb{A}^n) \cong \text{Hom}(\mathbb{Z}[x_1, \dots, x_n], R)$ where generators go $(x_i \mapsto r_i)$

5. PRODUCTS


5.0 Products in category theory

Category theory: \mathcal{C} Cat., $C_i \in \mathcal{C}$

Product $C_1 \times \dots \times C_n$ (if exists) is an object with morphs π_i to C_i , s.t.



coproduct $C_1 \sqcup \dots \sqcup C_n$:



Examples Sets / Top spaces: $X = \text{product}$, $\pi_i = \text{projections}$, $\sqcup = \text{disjoint union}$, π_i are inclusions.
 Vector spaces/abeliangps/modules: $\sqcup = \text{direct sum}$, π_i are inclusions.
 Rings: $\sqcup = \text{tensor product}$, $\pi_i(r) = 1 \otimes \dots \otimes r \otimes \dots \otimes 1$

Fix $B \in \mathcal{C}$ ("base")

Category of B -objects: \mathcal{C}/B

obj: morphs $C \rightarrow B$, morphs: $C \rightarrow D$ in \mathcal{C}

(think of B as a parameter space and C as a family parametrised by B)

fiber product $C \times_B D$ is the product in \mathcal{C}/B of $C \xrightarrow{f} B, D \xrightarrow{g} B$ (if exists)

(or pullback, or Cartesian square)



Similarly get $C_1 \times \dots \times C_n$

Example for sets or Top spaces: $C \times_B D = \{ (c, d) \in C \times D : f(c) = g(d) \in B \}$

for example if f, g are inclusions of subsets (subspaces) then $C \times_B D = C \cap D$

Pushout The opposite diagram (reverse arrows)

Example: for Rings the pushout of $B \rightarrow C, B \rightarrow D$ is the tensor product $C \otimes_B D$

Exercise: (co)product, fiber product, pushout are Unique up to Unique iso if they exist.

(Hint: compose unique maps between them (s.t. diagram commutes) then composites=id by uniqueness of self-map)

Examples of fiber products in cat. of Sets or Top spaces: $C \times_B D = \{ (c, d) : f(c) = g(d) \} \subseteq C \times D$

$B = \text{point} \Rightarrow C \times_B D = C \times D$

$C \xrightarrow{f} B, D \xrightarrow{g} B \Rightarrow C \times_B D \cong C \cap D$

$D \xrightarrow{g} B \Rightarrow C \times_B D \cong f^{-1}(D) \subseteq C$ for example $D = \text{point} = b \in B$ get fiber $f^{-1}(b)$

$C = D \Rightarrow C \times_B D = \{ (x, y) : f(x) = g(y) \} \subseteq C \times D$ ("equaliser")

IMPORTANT EXAMPLES:

All schemes X have canonical $X \rightarrow \text{Spec } \mathbb{Z}$ by giving canonical maps on affines:

$\text{Spec } R \rightarrow \text{Spec } \mathbb{Z}$ from $\mathbb{Z} \rightarrow R, 1 \mapsto 1$

Schemes over field k means have $X \rightarrow \text{Spec } k$, same as saying all $\theta_X(U)$ are k -algebras and restrictions are k -alghoms

Functor of points interpretation:
 $\text{Hom}(\mathbb{Z}, C \times_B D) \cong \text{Hom}(\mathbb{Z}, C) \times_{\text{Hom}(\mathbb{Z}, B)} \text{Hom}(\mathbb{Z}, D)$
 So we are asking whether $h_c \times_{h_b} h_d$ is representable

sec. 4.2

$C \sqcup B$ is the gluing!

Uniqueness of Unique iso if they exist.

then composites=id by uniqueness of self-map

5.1 Fiber products exist in Schemes/B

Fix scheme B, consider category Schemes/B
 Theorem fiber products $X_1 \times_B \dots \times_B X_n$ exist

Inductively suffices to do case $n=2$. First need some algebraic preliminaries

An A-algebra R is a ring R together with a ring hom $A \rightarrow R$
 (A ring) $(\Rightarrow R$ is A-mod via $a \cdot r = \psi(a)r$)


R, S A-algebras $\Rightarrow (R \otimes_A S) =$ free R-alg. on R x S relations

relations: i) \otimes is bilinear
 ii) $a \cdot (r \otimes s) = (\psi(a) \cdot r) \otimes s = r \otimes (\psi(a) \cdot s)$.
 In particular $A \rightarrow R \otimes_A S$ is $a \mapsto a \cdot (1 \otimes 1) = \psi(a) \otimes 1 = 1 \otimes \psi(a)$
 The product on generators: $(r \otimes s) \cdot (r' \otimes s') = rr' \otimes ss'$.

Rank R, S rings $\Rightarrow R \otimes S = R \otimes_{\mathbb{Z}} S$


- Facts
- $R \otimes_{\mathbb{Z}} S \cong S$ (via $\sum r_i \otimes s_i \mapsto \sum r_i s_i$)
 - $R[x_1, \dots, x_n] \otimes_{\mathbb{Z}} R[y_1, \dots, y_m] \cong R[x_1, \dots, x_n, y_1, \dots, y_m]$
 - $(S/I) \otimes_{\mathbb{Z}} T \cong (S \otimes_{\mathbb{Z}} T) / (I \otimes 1) \cdot (S \otimes T)$ where S, T are R-algebras
 - k field, A k-alg, for A-algs R, S get: $R \otimes_A S \cong (R \otimes_k S) / \langle \psi_R(a) \otimes 1 - 1 \otimes \psi_S(a) : a \in A \rangle$

Affine case: $\text{Spec } R \times_{\text{Spec } A} \text{Spec } S = \text{Spec}(R \otimes_A S)$ exists in Aff/Spec A:



have pushout: $R \otimes_A S \leftarrow S$
 (in category of A-algs) $\uparrow \psi_S$
 $R \xleftarrow{\psi_R} A$

Claim: this is fiber product also in Sch/Spec A: let $X = \text{Spec } R$
 $Y = \text{Spec } S$
 $B = \text{Spec } A$
 $F = \text{Spec}(R \otimes_A S)$



Recall fiber products are unique up to unique iso if they exist.
 By construction (as U_i affine) $\exists!$ U_i $\rightarrow F$ making diagram commute

Rank $B = \text{Spec } \mathbb{Z}$ gives $X \times_B Y = X \times Y$

If can show these agree on overlaps $U_{ij} = U_i \cap U_j$, then glue to unique $Z \rightarrow F$.
 If U_{ij} were affine, this would have been immediate.

$U_{ij} \subseteq$ affine U_i , so running same argument with Z replaced by U_{ij} ,

we can cover U_{ij} by basic open affines $D_{f_k} \subseteq U_i$ and now $D_{f_k} \cap D_{f_l} = D_{f_k f_l}$ affine.
 \Rightarrow glue uniquely to give $U_{ij} \rightarrow F$

Recall trick that can pick open cover of U_{ij} that are basic opens simultaneously for U_i, U_j
 $\Rightarrow U_{ij} \rightarrow F$ and $U_{ji} \rightarrow F$ agree.

General case build schemes/morphs by 3 gluing procedures (tedious!)

- case $U_i \times_B Y$ with B, Y affine, $X = U_i$: affine open cover $\Rightarrow \exists X \times$ affine
- case $X \times_B V_j$ with B affine, $Y = U_j$: " " $\Rightarrow \exists X \times$ affine
- case $X \times_{W_k} Y$ with $B = U_{W_k}$: " " $\Rightarrow \exists X \times$ affine

Gluing work because agreement on overlaps is ensured by uniqueness up to iso of fiber products. Sketch:
 (preimage of open set viewed as open subscheme of U_i)

1) If know $U_i \times_B Y$ exist, then $\Pi_1^{-1}(U_{ij})$ is fiber product
 $U_{ij} \times_B Y$ so by uniqueness \exists iso $\Pi_1^{-1}(U_{ij}) \rightarrow \Pi_1^{-1}(U_{ij})$, so glue & get $X \times_B Y$
 (indeed a natural identification since $U_{ij} = U_i$ with sheaf $\mathcal{O}_{U_{ij}}$)


2) as in 1, swapping roles X, Y . \leftarrow again: open subschemes since preimages of opens

3) let $X_k = f^{-1}(W_k)$, $Y_k = g^{-1}(W_k) \Rightarrow X_k \times_{W_k} Y_k$ exists by 2) $(X_k, Y_k$ general)
 Key trick: notice $X_k \times_{W_k} Y_k = X_k \times_B Y_k$
 "because images are trapped in W_k, Y_k anyway"
 Then use argument in 1) to glue the $X_k \times_B Y_k$. \square

Rank Proof shows that $X \times_B Y$ has affine open cover by $U(U_i \times_{W_k} V_j)$
 where $X = U_i, Y = U_j$ are " " "
 $B = U_{W_k}$ with $U_i \rightarrow W_k \subseteq B, V_j \rightarrow W_k \subseteq B$

- Examples
- $\mathbb{A}^n \times_{\text{Spec } \mathbb{Z}} \mathbb{A}^m = \text{Spec } R[x_1, \dots, x_n, y_1, \dots, y_m] = \mathbb{A}^{n+m}$
 - $\text{Spec } \mathbb{Z}_2 \times_{\text{Spec } \mathbb{Z}} \text{Spec } \mathbb{Z}_3 = \text{Spec}(\mathbb{Z}_2 \otimes_{\mathbb{Z}} \mathbb{Z}_3) = \text{Spec}(0) = \emptyset$
- Exercise $X \times_B Y \cong X, X \times_B Y \cong Y \times_B X, (X \times_B Y) \times_B Z \cong X \times_B (Y \times_B Z), X \times_B Y \times_B Z \cong X \times_B Y$.

5.2 Fibers and preimages




$f: X \rightarrow B$ morph of schemes

fiber over point $b \in B$: $f^{-1}(b) = \text{Spec } \kappa(b) \times_B X$

preimage of closed subscheme $Y \subseteq B$: $f^{-1}(Y) = Y \times_B X$


Examples



$k =$ algebraically closed field \leftarrow (so classical alg. geometry)
 $f: A_k^1 \rightarrow A_k^1$ induced by $f^\#: k[x] \rightarrow k[y], x \mapsto y^2$
 fiber over $0: (\text{view point } 0 \text{ as } \text{Spec } k \rightarrow \text{Spec } k[x] \rightarrow A_k^1 \text{ so } k \cong k[x])$
 fiber = $\text{Spec } k[x] \times_{\text{Spec } k[x]} \text{Spec } k[x] = \text{Spec } (k[x] \otimes_k k[x]) = \text{Spec } (k[x] \otimes_k k[x])$ where $f(x) = y^2$
 $= \text{Spec } (k[x^2] \otimes_{(y^2)} k[x^2]) = \text{Spec } (k[x^2] \otimes_{(y^2)} k[x^2])$ where $f(x) = y^2$
 (e.g. use facts about \otimes from 5.1)

Link Notice how a product of affine varieties gave a scheme that was not an affine variety.

4) Mumford's picture of $\text{Spec } \mathbb{Z}[x]$:



$X = \text{Spec } \mathbb{Z}[x] = \dots$
 $B = \text{Spec } \mathbb{Z} = \dots$
 π is induced by inclusion $\mathbb{Z} \rightarrow \mathbb{Z}[x]$
 $\Rightarrow \pi^{-1}((p)) = V((p)) = \{(p), (p, f(x)) : f(x) \text{ mod } p \text{ is irreducible in } \mathbb{F}_p[x]\}$
 (so (p) is a dense point in $\pi^{-1}((p))$) if $p \in \mathbb{I}$ then $\mathbb{Z}[x]/\mathbb{I} \cong \mathbb{F}_p[x]/\mathbb{I}$ where $\mathbb{F}_p = \mathbb{Z}/p$
 PID, so (f) prime $\Leftrightarrow f$ irred or 0
Rmk curve $V(x^2+1)$ passes through $(p, x+j)$ iff x^2+1 vanishes at that point, so iff $x^2+1=0$ in $\mathbb{F}_p[x]/(x+j) \cong \mathbb{F}_p, x \mapsto -j$, so iff $j^2 = -1$.
 Classical number theory says a square root of -1 exists in $\mathbb{F}_p \Leftrightarrow (p \equiv 1 \pmod 4)$ (or $p=2$)
 fiber over $(p) : K(p) = \mathbb{Z}(p)/p \cdot \mathbb{Z}(p) = (\mathbb{Z}/p)(p) = \mathbb{F}_p = \mathbb{Z}/p$
 $\Rightarrow \pi^{-1}(p) = \text{Spec } (k(p) \otimes_{\mathbb{Z}} \mathbb{Z}[x]) = \text{Spec } \mathbb{F}_p[x] = \{(0), (f(x))\}$ irred in $\mathbb{F}_p[x]$ nonconstant
 fiber over $(0) : K(0) = \mathbb{Z}(0) = \mathbb{Q}$
 $\Rightarrow \pi^{-1}(0) = \text{Spec } (K(0) \otimes_{\mathbb{Z}} \mathbb{Z}[x]) = \text{Spec } \mathbb{Q}[x] = \{(0), (f(x))\}$ irred in $\mathbb{Q}[x]$ nonconstant
 [Gauss's Lemma: For $f \in \mathbb{Z}[x]$ primitive (gcd(coeffs)=1) irred in $\mathbb{Q}[x] \Leftrightarrow f$ irred in $\mathbb{Z}[x]$ nonconstant
 f irred in $\mathbb{Z}[x] \Leftrightarrow f$ irred in $\mathbb{Q}[x]$ nonconstant]
Consequence $\text{Spec } \mathbb{Z}[x] = \{(0), (p), (f), (p, f)\}$ $f \in \mathbb{Z}[x]$ irred, mod p nonconstant
 $\leftarrow p \in \mathbb{Z}$ prime $f \in \mathbb{Z}[x]$ irred, nonconstant

Forgetful functor $|\cdot|: \text{Sch} \rightarrow \text{Top Spaces}, X \mapsto |X| =$ underlying topological space.
 morph \mapsto underlying continuous map

Claim $f: X \rightarrow B$ morph schemes $\Rightarrow |f^{-1}(b)| = |f|^{-1}(b)$ \leftarrow fiber is homeomorphic to topological fiber

Pf WLOG B affine = $\text{Spec } S$ and b is prime ideal $p \subseteq S$
 $f^{-1}(B) = \cup \text{Spec } R_i$ given by $\varphi_i: S \rightarrow R_i$
 WLOG just consider one affine, so $R = R_i$, so WLOG $X = \text{Spec } R$
 $\Rightarrow \text{Spec } k(b) \times_B X = \text{Spec } (k(b) \otimes_S R)$

$k(b) = (S/p)_p \Rightarrow k(b) \otimes_S R = (S/p)_p \otimes_S R = S_p \otimes_S R = R_p / (\varphi(p)R_p)$
 $\Rightarrow \text{Spec } (k(b) \otimes_S R) \xrightarrow{|\cdot|} \{q \subseteq R \text{ prime ideal containing } \varphi(p) \text{ but not intersecting } \varphi(S \setminus p)\}$
 $q \cdot R_p \xrightarrow{|\cdot|} q \quad (= \text{preimage of } qR_p \text{ via localisation } R \rightarrow R_p = S_p \otimes_S R) \xrightarrow{f^{-1}} p$
 $q \subseteq R \setminus \varphi(S \setminus p) \Rightarrow \varphi^{-1}q \subseteq S \setminus (S \setminus p) = p$ so get $\{q \in \text{Spec } R : \varphi^{-1}q = p\} \cdot \square$
 $q \supseteq \varphi(p) \Rightarrow \varphi^{-1}q \supseteq p$ (apply 1.1 to diagram defining X, Y then by universal property in category of topological spaces get unique map \otimes)

Cor Given $f: X \rightarrow B, g: Y \rightarrow B$, fiber of $|X \times_B Y| \rightarrow |X| \times |B| \times |Y|$ over (x, y) is $|\text{Spec } (k(x) \otimes_{k(b)} k(y))|$ where $f(x) = g(y) = b$

Pf fiber of $X \times_B Y \rightarrow X$ over $x : \text{Spec } k(x) \times_X (X \times_B Y) = \text{Spec } k(x) \times_B Y$
 fiber of $\text{Spec } k(x) \times_B Y \rightarrow Y$ over $y : \text{Spec } k(x) \times_B Y \times_Y \text{Spec } k(y) = \text{Spec } k(x) \times_B \text{Spec } k(y)$
 fiber of $\text{Spec } k(x) \times_B \text{Spec } k(y) \rightarrow B$ over $b : \text{Spec } k(x) \times_{\text{Spec } k(b)} \text{Spec } k(y) = \text{Spec } (k(x) \otimes_{k(b)} k(y))$

by Claim can work with fiber in Sch before applying 1.1.
 at algebra level: if A_1, A_2 are modules over $S = R_p/R_p$ then $S \otimes (A_1 \otimes_R A_2) \cong A_1 \otimes_S A_2$ namely: $\frac{R_p \otimes (R_p/R_p) \otimes R}{\mathbb{I}} \otimes_{\mathbb{I}} (a_1 \otimes a_2) \rightarrow \frac{R_p \otimes (a_1 \otimes a_2)}{\mathbb{I}}$
 or at category level, with abuse of notation: hence $\rightarrow \exists!$ $\frac{z \cdot x \cdot y}{z \cdot x \cdot y} \rightarrow \frac{z \cdot x \cdot y}{z \cdot x \cdot y}$

Examples $|\text{Spec } \mathbb{Z}_2 \times_{\text{Spec } \mathbb{Z}} \text{Spec } \mathbb{Z}_3| = |\text{Spec } \mathbb{Z}_2| \times_{|\text{Spec } \mathbb{Z}|} |\text{Spec } \mathbb{Z}_3| = \emptyset$ since 1^{st} factor $\mapsto (2)$ and 2^{nd} factor $\mapsto (3) \in \text{Spec } \mathbb{Z}$

$\cdot A_k^2 = A_k^1 \times_{\text{Spec } k} A_k^1 = \text{Spec } k[x, y]$ then $(x+y) \mapsto (0)$ via both projections to A_k^1 but $(x+y) \neq (0)$ (field k) so $|A_k^2| \neq |A_k^1| \times |A_k^1|$: the fiber over $(0), (0)$ is complicated.
 note $\text{Spec } k = \text{point} = \{(0)\}$ so often omit "Spec k " from notation.

Rmk If x, y closed points of scheme X, Y finite type over k, k algebraically closed, then fiber over (x, y) of $X \times_{\text{Spec } k} Y$ is $\text{Spec } (k(x) \otimes_k k(y)) = \text{Spec } (k \otimes_k k) = \text{Spec } k = (0)$ so over closed points you get the product of sets. \leftarrow (so classical alg. geom.)

Warning $A_k^2 = A_k^1 \times A_k^1$ does not have the product topology, e.g. consider $\forall (x, y)$

Non-Examinable Rmk Working over an algebraically closed field k , the stalk of $X \times_{\text{Spec } k} Y$ at (x, y) is $\mathcal{O}_{X, x} \otimes_k \mathcal{O}_{Y, y}$ localised at max ideal $\mathfrak{m}_{X, x} \otimes \mathfrak{m}_{Y, y} + \mathcal{O}_{X, x} \otimes \mathfrak{m}_{Y, y}$

5.6 Scheme structure on subsets

Motivation: classically, a projective variety is a closed subset of \mathbb{P}^n . A quasi-proj. var. is an open \subseteq proj. var., so \cong locally closed subset of \mathbb{P}^n .

Claim Any closed subset $C \subseteq X$ of a scheme $\Rightarrow \exists!$ closed reduced subscheme $(C, \mathcal{O}_C) \rightarrow X$
Pf $\mathcal{J}(U) := \{s \in \mathcal{O}_X(U) : s(p) = 0 \in k(p) \forall p \in C \cap U\}$ is sheaf of ideals
 Locally: $U = \text{Spec } R, C \cap U = V(I)$ for unique radical ideal $I \Rightarrow \mathcal{J}(\text{Spec } R) = I$

Then $s(p) = 0 \in k(p) = (R/p) \Leftrightarrow s \in \bigcap_{p \in V(I)} p = \sqrt{I} = I$
 Same trick shows $\mathcal{J}(D_f) = I_f$, so \mathcal{J} is the quasi-coherent ideal sheaf corresponding to I .
 Note: $C = \text{supp}(\mathcal{O}_X/\mathcal{J})$ and $C \cap U = \text{Spec } R/I$, and we define $\mathcal{O}_C = \mathcal{O}_X/\mathcal{J}$. \square

Def call this the induced reduced scheme structure on C .
Example When we consider an irreducible component $Z \subseteq X$, we use this scheme structure or: $\text{Mod}_{\mathcal{O}_X}$
Exercise For $C = X \subseteq X$ get the reduced scheme X_{red} (see 3.6)

Def $Z \subseteq X$ locally closed means $\forall z \in Z, \exists$ open $U \ni z$ s.t. $Z \cap U$ is closed in U .
Lemma Z locally closed $\Leftrightarrow Z$ open in $\bar{Z} \leftarrow$ (i.e. $Z = \bar{Z} \cap U$ some open $U \subseteq X$) by Lemma, $C = \bar{Z} \cap U$ works
Pf \Leftarrow : $Z = \bar{Z} \cap U$ for open $U \subseteq X \Rightarrow \bar{Z} \cap U = Z = \bar{Z} \cap U$
 \Rightarrow : $Z \cap U$ closed in U so equals its closure in U which is: $\bar{C}_U(Z \cap U) = \bar{Z} \cap U$
 $\Rightarrow z \in \bar{Z} \cap U = \bar{Z} \cap U \subseteq \bar{Z}$ so Z contains an open neighbourhood of z in \bar{Z}
 $\Leftrightarrow \forall$ open $U \subseteq U \Rightarrow \bar{Z} \cap U = Z$
Rmk $Z \subseteq X$ closed, so $\exists!$ induced reduced scheme structure $\mathcal{O}_{\bar{Z}}$ on \bar{Z}
 $Z \subseteq \bar{Z}$ is open so get " " $\mathcal{O}_Z = \mathcal{O}_{\bar{Z}}|_Z$ (so $\mathcal{O}_Z(V) = \mathcal{O}_{\bar{Z}}(V)$)

The local description is the same as above: $Z \cap U = \bar{Z} \cap U = \text{Spec}(R/I), \mathcal{O}_Z|_U \cong \mathcal{O}_{\text{Spec}(R/I)}$
Rmk If Z irreducible ($\Rightarrow \bar{Z}$ irreducible) then $I = p \in \text{Spec } R$ where p is a generic point for both Z, \bar{Z}
Hwk 3 \bar{Z} irred. locally closed \subseteq Variety $(X, \mathcal{O}_X) \Rightarrow (\bar{Z}, \mathcal{O}_{\bar{Z}})$ variety

Hwk 3 (X, \mathcal{O}_X) variety, $Z \subseteq X$ irreducible subspace \leftarrow (the irreducibility is not so important if allow varieties to be reducible)
Define sheaf \mathcal{O}_Z on Z : for open $V \subseteq Z$,
 $\mathcal{O}_Z(V) = \left\{ s: V \rightarrow \coprod_{z \in V} k(x) : \forall x \in V \exists$ open $U \subseteq X, t \in \Gamma(U, \mathcal{O}_X) \right\}$
 such that $s(x) = t(x) \in k(x), \forall x \in V \cap U$

Prove that:
 (Z, \mathcal{O}_Z) variety $\Rightarrow Z$ locally closed and \mathcal{O}_Z is the induced reduced scheme structure
 (universal property for the above sheaf)

Lemma With that definition, if Y reduced scheme, $f: Y \rightarrow X$ morph of sch. if $f(Y) \subseteq Z$ (as topological spaces) then f factorizes $f: Y \rightarrow Z \rightarrow X$
Pf Need check sheaves: $s \in \mathcal{O}_Z(U \cap Z)$ for $U \subseteq X$ open then \exists open cover $U \cap Z = \cup U_i \cap Z$ and $s_i \in \mathcal{O}_X(U_i)$, $s(x) = s_i(x) \in k(x), \forall x \in U_i \cap Z$
 $\Rightarrow f^\#(s_i) \in \mathcal{O}_Y(f^{-1}U_i), f^\#(s_i)(y) = f^\#(s_i)(y) \in k(y), \forall y \in f^{-1}(U_i \cap Z)$
 \Rightarrow by Sec. 3.3 since Y reduced: $f^\#(s_i)_y = f^\#(s)_y \in \mathcal{O}_{Y,y}, \forall y \in f^{-1}(U_i \cap Z)$
 $\Rightarrow f^\#(s)$ glue to a unique section $r \in \mathcal{O}_Y(f^{-1}U)$. Define $\mathcal{O}_Z(U) \rightarrow \mathcal{O}_Y(f^{-1}U), s \mapsto r$ and note $\mathcal{O}_X(U_i) \rightarrow \mathcal{O}_Z(U_i \cap Z) \rightarrow \mathcal{O}_Y(f^{-1}U_i), s_i \mapsto s|_{U_i \cap Z} \mapsto r|_{f^{-1}U_i}$. \square

Rmk Applying the Lemma to the case $Y =$ locally closed $Z \subseteq X$ with induced reduced sheaf, implies $\mathcal{O}_Z \cong \mathcal{O}_Z$.

6. SHEAVES OF MODULES

6.1 \mathcal{O}_X -modules

Def \mathcal{O}_X -module is: • sheaf $F \in \text{Ab}(X)$ (often abbreviate $\mathcal{O}_X := \mathcal{O}_X|_U$)
 (or sheaf of/in \mathcal{O}_X -mods) • restrictions are compatible with module structure

Morphism $F \rightarrow G$ of \mathcal{O}_X -module is: • morph $F \xrightarrow{\psi} G$ of sheaves (if monomorph, i.e. ψ_U injective, F is \mathcal{O}_X -submod of G) • $F(U) \xrightarrow{\psi_U} G(U)$ is hom of $\mathcal{O}_X(U)$ -mods
Rmk stalk F_x is $\mathcal{O}_{X,x}$ -mod, and for morph $F \rightarrow G$ get $F_x \rightarrow G_x$ is $\mathcal{O}_{X,x}$ -mod hom.

Example A sheaf of ideals is an \mathcal{O}_X -submod of $\mathcal{O}_X \leftarrow$ (just like R -submods of R are ideals)
Fact \mathcal{O}_X -Mods = (category of \mathcal{O}_X -mods on X) is an abelian cat \leftarrow (proof similar to $\text{Ab}(X)$)
 indeed notions of submod, quotient mod, ker, coker, Im agree with what get in $\text{Ab}(X)$
 e.g. $F \rightarrow G \rightarrow H$ exact \Leftrightarrow exact in $\text{Ab}(X) \Leftrightarrow$ exact on stalks

Will write $\text{Hom}_{\mathcal{O}_X}$ for morphisms in this category.

6.2 Modules generated by sections

$\text{Hom}_{\mathcal{O}_X}(\mathcal{O}_X, F) \xrightarrow{\cong} F(X) \quad \forall F \in \mathcal{O}_X\text{-Mods} \leftarrow$ analogue of $\text{Hom}_R(R, M) \cong M$
 $(\varphi: \mathcal{O}_X \rightarrow F) \leftrightarrow s = \varphi(1)$ since $\varphi_U(r) = \varphi_U(r \cdot 1) = r \cdot s|_U \quad \forall r \in \mathcal{O}_X(U)$
 Similarly $\text{Hom}_{\mathcal{O}_X}(\mathcal{O}_X^n, F) \xrightarrow{\cong} F(X)^{\oplus n}$ defined by n global sections $s_1, \dots, s_n \in F(X)$
Def F is generated by global sections if \exists surjection $\bigoplus_{i \in I} \mathcal{O}_X \rightarrow F$ of \mathcal{O}_X -mods ($\Leftrightarrow s_i|_Z$ generate $\mathcal{O}_{X,x}$ -mod $F_x \quad \forall x \in X$)
 same as picking sections $s_i \in F(X)$ (as \mathcal{O}_X -module, $\bigoplus \mathcal{O}_X \rightarrow F|_U$)
Rmk Can produce \mathcal{O}_X -submods from given local sections $s_i \in F(U_i) \leftarrow$ possible $\mathcal{O}_X(U_i)$ -linear combos of $\{s_i|_{U_i} : U_i \subseteq U\}$
Def A sheaf has finite type if locally generated by finitely many sections. \leftarrow sheafify $U \mapsto$ possible combos of $\{s_i|_{U_i} : U_i \subseteq U\}$
 (see Hwk 4) \leftarrow (equivalent definitions) so $\mathcal{O}_U^n \rightarrow F|_U$ some open $U \subseteq U$ (not fixed)

6.3 Vector bundles and coherent modules
Def \mathcal{O}_X -mod F is locally free \mathcal{O}_X -mod of finite rank (or vector bundle) if $\forall x \in X \exists$ open $U \ni x$ s.t. $F|_U \cong \mathcal{O}_U^n$ (as \mathcal{O}_U -module, $\mathcal{O}_U^n \rightarrow F|_U$)
 locally $\mathcal{O}_U^n \cong \mathcal{O}_U^n \rightarrow \mathcal{O}_U^n = F|_U$
 locally $\mathcal{O}_U^n \cong \mathcal{O}_U^n \rightarrow \mathcal{O}_U^n = F|_U$ if $n=1$ (fixed) \leftarrow generated by one section $s \in F(U)$
Def X invertible sheaf ("or" line bundle) if $n=1$ (fixed) \leftarrow generated by one section $s \in F(U)$
Question Is it enough to ask $F_x \cong \mathcal{O}_{X,x} \quad \forall x$ some $n \in \mathbb{N}$ depending on x ? (\Leftarrow : can fail)
Lemma F finite type, $\mathcal{O}_X^n \xrightarrow{\psi} F_x \text{ surj} \Rightarrow \exists x \in U \subseteq X$ with surj $\mathcal{O}_U^n \xrightarrow{\psi} F|_U, \varphi|_U = \varphi_x$.
Pf finite type $\Rightarrow \exists$ surj $\mathcal{O}_U^m \xrightarrow{\psi} F|_U$. Let $s_i = \varphi_x(e_i) \in F_x = \mathcal{O}_{X,x}$ -s. Now $s_i \in F(U_i)$ some U_i . Replace U by $U \cap U_1 \cap \dots \cap U_n$ so wlog $s_i \in F(U)$. Let $f_i = 1 \in U_i$ -th copy of $\mathcal{O}_U \xrightarrow{\psi} F|_U$ is $\Sigma f_i \cdot s_i$ some $s_i \in \mathcal{O}_X$. So $\psi(f_i) \in \Sigma \mathcal{O}_U \cdot s_i$ surj, some $v_j \in U$, again wlog $v_j = U$ (replace U by $U \cap V_1 \cap \dots \cap V_m$)
 $\Rightarrow \psi(f_i) \in \text{Im } \varphi$ for $\varphi: \mathcal{O}_U^n \rightarrow F|_U$ with $\varphi(e_i) = s_i$ on U . So φ hits \mathcal{O}_U -mod generators $\psi(f_i)$.
 Continuing above question: We know φ_x is inj at x , but we don't know if the same φ works also for y close to x , so we do not know whether φ_y inj $\Leftrightarrow \varphi_y$ inj at all stalks at $y \in U$.

5.6 Scheme structure on subsets

Claim Any closed subset $C \subseteq X$ of a scheme $\Rightarrow \exists!$ closed reduced subscheme $(C, \mathcal{O}_C) \rightarrow X$
Pf $\mathcal{J}(U) := \{s \in \mathcal{O}_X(U) : s(p) = 0 \in k(p) \forall p \in C \cap U\}$ is sheaf of ideals
 Locally: $U = \text{Spec } R, C \cap U = V(I)$ for unique radical ideal $I \Rightarrow \mathcal{J}(\text{Spec } R) = I$

Then $s(p) = 0 \in k(p) = (R/p) \Leftrightarrow s \in \bigcap_{p \in V(I)} p = \sqrt{I} = I$
 Same trick shows $\mathcal{J}(D_f) = I_f$, so \mathcal{J} is the quasi-coherent ideal sheaf corresponding to I .
 Note: $C = \text{supp}(\mathcal{O}_X/\mathcal{J})$ and $C \cap U = \text{Spec } R/I$, and we define $\mathcal{O}_C = \mathcal{O}_X/\mathcal{J}$. \square

Def call this the induced reduced scheme structure on C .
Example When we consider an irreducible component $Z \subseteq X$, we use this scheme structure or: $\text{Mod}_{\mathcal{O}_X}$
Exercise For $C = X \subseteq X$ get the reduced scheme X_{red} (see 3.6)

Def $Z \subseteq X$ locally closed means $\forall z \in Z, \exists$ open $U \ni z$ s.t. $Z \cap U$ is closed in U .
Lemma Z locally closed $\Leftrightarrow Z$ open in $\bar{Z} \leftarrow$ (i.e. $Z = \bar{Z} \cap U$ some open $U \subseteq X$) by Lemma, $C = \bar{Z} \cap U$ works
Pf \Leftarrow : $Z = \bar{Z} \cap U$ for open $U \subseteq X \Rightarrow \bar{Z} \cap U = Z = \bar{Z} \cap U$
 \Rightarrow : $Z \cap U$ closed in U so equals its closure in U which is: $\bar{C}_U(Z \cap U) = \bar{Z} \cap U$
 $\Rightarrow z \in \bar{Z} \cap U = \bar{Z} \cap U \subseteq \bar{Z}$ so Z contains an open neighbourhood of z in \bar{Z}
 $\Leftrightarrow \forall$ open $U \subseteq U \Rightarrow \bar{Z} \cap U = Z$
Rmk $Z \subseteq X$ closed, so $\exists!$ induced reduced scheme structure $\mathcal{O}_{\bar{Z}}$ on \bar{Z}
 $Z \subseteq \bar{Z}$ is open so get " " $\mathcal{O}_Z = \mathcal{O}_{\bar{Z}}|_Z$ (so $\mathcal{O}_Z(V) = \mathcal{O}_{\bar{Z}}(V)$)

The local description is the same as above: $Z \cap U = \bar{Z} \cap U = \text{Spec}(R/I), \mathcal{O}_Z|_U \cong \mathcal{O}_{\text{Spec}(R/I)}$
Rmk If Z irreducible ($\Rightarrow \bar{Z}$ irreducible) then $I = p \in \text{Spec } R$ where p is a generic point for both Z, \bar{Z}
Hwk 3 \bar{Z} irred. locally closed \subseteq Variety $(X, \mathcal{O}_X) \Rightarrow (\bar{Z}, \mathcal{O}_{\bar{Z}})$ variety

Hwk 3 (X, \mathcal{O}_X) variety, $Z \subseteq X$ irreducible subspace \leftarrow (the irreducibility is not so important if allow varieties to be reducible)
Define sheaf \mathcal{O}_Z on Z : for open $V \subseteq Z$,
 $\mathcal{O}_Z(V) = \left\{ s: V \rightarrow \coprod_{z \in V} k(x) : \forall x \in V \exists$ open $U \subseteq X, t \in \Gamma(U, \mathcal{O}_X) \right\}$
 such that $s(x) = t(x) \in k(x), \forall x \in V \cap U$

Prove that:
 (Z, \mathcal{O}_Z) variety $\Rightarrow Z$ locally closed and \mathcal{O}_Z is the induced reduced scheme structure
 (universal property for the above sheaf)

Lemma With that definition, if Y reduced scheme, $f: Y \rightarrow X$ morph of sch. if $f(Y) \subseteq Z$ (as topological spaces) then f factorizes $f: Y \rightarrow Z \rightarrow X$
Pf Need check sheaves: $s \in \mathcal{O}_Z(U \cap Z)$ for $U \subseteq X$ open then \exists open cover $U \cap Z = \cup U_i \cap Z$ and $s_i \in \mathcal{O}_X(U_i)$, $s(x) = s_i(x) \in k(x), \forall x \in U_i \cap Z$
 $\Rightarrow f^\#(s_i) \in \mathcal{O}_Y(f^{-1}U_i), f^\#(s_i)(y) = f^\#(s_i)(y) \in k(y), \forall y \in f^{-1}(U_i \cap Z)$
 \Rightarrow by Sec. 3.3 since Y reduced: $f^\#(s_i)_y = f^\#(s)_y \in \mathcal{O}_{Y,y}, \forall y \in f^{-1}(U_i \cap Z)$
 $\Rightarrow f^\#(s)$ glue to a unique section $r \in \mathcal{O}_Y(f^{-1}U)$. Define $\mathcal{O}_Z(U) \rightarrow \mathcal{O}_Y(f^{-1}U), s \mapsto r$ and note $\mathcal{O}_X(U_i) \rightarrow \mathcal{O}_Z(U_i \cap Z) \rightarrow \mathcal{O}_Y(f^{-1}U_i), s_i \mapsto s|_{U_i \cap Z} \mapsto r|_{f^{-1}U_i}$. \square

Rmk Applying the Lemma to the case $Y =$ locally closed $Z \subseteq X$ with induced reduced sheaf, implies $\mathcal{O}_Z \cong \mathcal{O}_Z$.

Cor

$F \in \text{QCoh}(X) \iff \forall x \in X \exists \text{ affine open } U \ni x \in U \ni \text{Spec } R \subseteq U \cong \tilde{M}$ some R -mod M
 $F \in \text{Coh}(X) \iff$ in addition require M is coherent R -mod

$\left\{ \begin{array}{l} M \text{ finitely generated} \\ \ker(R^n \rightarrow M) \text{ is f.g., any n} \in \mathbb{N} \end{array} \right.$
 (Pick U so that lemma applies.)
 Idea: want $\forall f, g$ submod of M to have finite presentation, indeed get exact sequence $R^m \rightarrow R^n \rightarrow \text{Im } \varphi \rightarrow 0$ $\xrightarrow{\text{map to gens of ker } \varphi}$ $\text{Im } \varphi \rightarrow 0$

Rmk If R Noeth., coherent = f.g. (since R^n f.g., so its submod as f.g. as R Noeth.)
Example X loc. Noeth. scheme $\implies \mathcal{O}_X$ is coherent \implies ideal sheaf of any closed subsc. is coherent.

Rmk \forall scheme: $F \in \text{QCoh}(X) \iff \exists$ affine open cover $X = \cup U_i$ s.t. $F|_{U_i} \cong \tilde{M}_i$ for R_i -mod M_i $U_i = \text{Spec } R_i$ (immediate from Cor)
 $(\text{WLOG: } R_i = \mathcal{O}_X(U_i), M_i = F(U_i))$

Rmk restriction to open $V \subseteq X$: $\text{QCoh}(X) \rightarrow \text{QCoh}(V)$, $\text{Coh}(X) \rightarrow \text{Coh}(V)$
Pf $x \in V \cap U = \cup D_{f_i}$ for $f_i \in R$ then $F|_U \cong \tilde{M}_i$ $\cong \tilde{M}_i|_{D_{f_i}}$ (and use fact that localization preserves coherent properties)
 so again locally module. \square

Why is quasi-coherence a good notion?
 Rings $\rightarrow \text{Aff}$, $R \rightarrow \text{Spec}(R)$, $\mathcal{O}_{\text{Spec}(R)}$ equivalence of cats
 R -Mods $\rightarrow \mathcal{O}_{\text{Spec}(R)}$ -Mods, $M \mapsto \tilde{M}$ not equivalence of cats \leftarrow notice $F \in \mathcal{O}_X$ -Mods if $\mathcal{O}_X \in \mathcal{O}_X$
Example $X = \text{Spec } k[x] = \mathbb{A}^1_k$, skyscraper sheaf at 0 : $F(U) = \begin{cases} k[x] & \text{if } 0 \in U \\ 0 & \text{else} \end{cases}$
 \implies if the above were an equivalence of cats, then $F \cong \tilde{M}$ some $k[x]$ -mod M
 so $k[x] = F(X) \cong \tilde{M}(X) = M$. But $k[x] = \mathcal{O}_X$ is not isomorphic to F !
Solution restrict which \mathcal{O}_X -mods you allow: want them locally to look like \tilde{M} , just like when we studied sheaves of ideals that locally look like \tilde{I}

Will show later: $\boxed{\text{For } X = \text{Spec } R: R\text{-Mods} \rightarrow \text{QCoh}(X) \text{ equivalence of categories } M \mapsto \tilde{M} \leftarrow F(X) \leftarrow F}$

7.2 Overview of general properties of QCoh(X) and Coh(X) for X scheme

- $\text{Coh}(X)$ abelian category, and $\text{Coh}(X) \xrightarrow{\text{incl}} \mathcal{O}_X\text{-Mod}$
 $\text{QCoh}(X) \xrightarrow{\text{incl}} \text{QCoh}(X)$ are exact functors
 In particular can take Ker, Coker, Image in both (not in $\text{Vect}(X)$) \leftarrow Easy for QCoh since locally hom of mods $M_i \rightarrow M_j$ so take Ker, Coker, Im
- $0 \rightarrow F_1 \rightarrow F_2 \rightarrow F_3 \rightarrow 0$ exact in \mathcal{O}_X -Mods.
 Two of the $F_i \in \text{QCoh}(X) \implies$ all three are. Same holds for $\text{Coh}(X)$ (not for $\text{Vect}(X)$)
Trick $0 \rightarrow F_1 \rightarrow F_2 \rightarrow F_3 \rightarrow 0$ exact, and F_2, F_3 are, then F_1 is. (Pf. $F_1 \cong \text{Ker}(F_2 \rightarrow F_3)$, use (1.10))
- Can take finite \oplus , $\cdot \mathcal{O}_X$, $\text{Hom}_{\mathcal{O}_X}(\cdot, \cdot)$ in $\text{QCoh}(X)$, $\text{Coh}(X)$ and $\text{Vect}(X)$
 for $\text{QCoh, Hom}_{\mathcal{O}_X}(F, G)$ need assume F loc. finitely presented
- Gabriel-Rosenberg thm
 X quasi-compact & separated (e.g. variety) $\implies X$ is determined up to iso by $\text{QCoh } X$!
- X loc. Noeth. scheme, $Z \hookrightarrow X$ closed subsc. $\implies 0 \rightarrow \mathcal{I}_Z \rightarrow \mathcal{O}_X \rightarrow \mathcal{O}_Z \rightarrow 0$ exact in $\text{Coh } X$
 finite type subsheaf $F \subseteq G$, $G \in \text{Coh}(X) \implies F \in \text{Coh}(X)$ \leftarrow combine to prove kernels exist in $\text{Coh } X$
 $\varphi: F \rightarrow G$, $G \in \text{Coh } X$, F finite type $\implies \text{Ker } \varphi$ finite type
 $\varphi: F \rightarrow G$, $G \in \text{Coh } X$, F finite type, $\varphi_x: F_x \rightarrow G_x$ injective $\implies \varphi|_U: F|_U \rightarrow G|_U$ inj. some U
Hwk 4: Picard group $\text{Pic}(X) = \{ \text{isomorphism classes of invertible sheaves} \}$ \leftarrow we proved it in case $F = \mathcal{O}_X$ in Pf. claim
 group operation is $\cdot \mathcal{O}_X$. (abelian group as $F \otimes_{\mathcal{O}_X} G \cong G \otimes_{\mathcal{O}_X} F$) in Sec. 6.3

7.3 Pullback preserves quasi-coherence

$f: X \rightarrow Y$ morph ringed spaces
Claim $f^*: \text{QCoh}(Y) \rightarrow \text{QCoh}(X)$. If X loc. Noeth. scheme $\implies f^*: \text{Coh } Y \rightarrow \text{Coh } X$.
 Without this can fail e.g. $f^* \mathcal{O}_Y = \mathcal{O}_X$ so if \mathcal{O}_Y coh, \mathcal{O}_X not coh, then fails

Pf if $\bigoplus_{i=1}^n \mathcal{O}_Y|_U \rightarrow \bigoplus_{j=1}^m \mathcal{O}_Y|_U \rightarrow 0$ exact $(f_x \in U \subseteq Y$ open)
 apply g^* where $g = f|_{f^{-1}U}: f^{-1}U \rightarrow U$, using g^* right exact & commutes with \oplus :
 $\bigoplus_{i=1}^n \mathcal{O}_X|_{f^{-1}U} \rightarrow \bigoplus_{j=1}^m \mathcal{O}_X|_{f^{-1}U} \rightarrow 0$ exact, and $x \in f^{-1}U$ open. \leftarrow using X loc. Noeth.

Rmk $F \in \text{Coh}(Y) \implies F$ locally finitely presented $\implies f^*F$ loc. finitely presented $\implies f^*F \in \text{Coh}(X)$
 issue is f^* affine need not be affine. For affine morphs you get result by Sec. 6.5

7.4 Push-forwards for X Noetherian
Claim $f: X \rightarrow Y$ morph of schemes, X Noetherian $\implies f_*: \text{QCoh } X \rightarrow \text{QCoh } Y$
Pf $0 \rightarrow F \rightarrow \Pi F|_{U_i} \rightarrow \Pi F|_{U_j} \rightarrow 0$ exact by sheaf property, where $X = \cup U_i$ affine open cover

Rmk X quasi-compact & commutes with limits e.g. with $\Pi \implies 0 \rightarrow f_*F \rightarrow \Pi f_*F|_{U_i} \rightarrow \Pi f_*F|_{U_j} \rightarrow 0$ exact
 WLOG Y open affine = $\text{Spec } R$ (replace X by $f^{-1}(\text{Spec } R)$), $\text{WLOG } F|_{U_i} = \tilde{F}(U_i)$, so $f_*F|_{U_i} = \tilde{F}(U_i)$
 similarly for U_j . If show $\Pi f_*F|_{U_i} = \tilde{F}(U_i)$, then $f_*F \in \text{QCoh}(Y)$ \leftarrow Trick (2) in Sec. 6.5

Rmk X quasi-compact, separated $\implies f_*: \text{QCoh } X \rightarrow \text{QCoh } Y \leftarrow$ proof above but easier \leftarrow take differences of sections on overlaps (Sec. 1.4) $U_i, U_j = U_{i,j}$ " " \leftarrow issue is f^* affine need not be affine. For affine morphs you get result by Sec. 6.5

Non-examinable fact f proper, X, Y loc. Noeth. $\implies f_*: \text{Coh } X \rightarrow \text{Coh } Y$
 otherwise in general f_* can ruin (quasi)-coherence
 (e.g. $\mathbb{A}^1 \xrightarrow{\pi} \mathbb{A}^1$ obvious morph, $F = \Pi k[x]$, $f_*F = \Pi k[x]$ if assume $\mathcal{O}_{\text{Coh}}(X)$ but notice $(\frac{1}{x}) \in F|_{D_x} = (f_*F)|_{D_x}$ but $\notin (\Pi k[x])|_{D_x} = (\Pi k[x])|_{D_x}$) $\neq \Pi k[x]|_{D_x}$ \leftarrow by 7.6 \leftarrow e.g. $\mathbb{A}^1 \xrightarrow{\pi} \mathbb{A}^1$ Spec $k[x] \rightarrow \text{Spec } k[x]$ not finite k -mod

7.5 Gluing modules
 Similar to Sec. 4.1: R ring $\ni f_1, \dots, f_n$ s.t. $1 \in \langle f_i \rangle$
 data: $M_i: R_{f_i}$ -mod \leftarrow (so have M_i on $D_{f_i} \subseteq \text{Spec } R$) cocycle $(M_i)_{f_i, f_j} \xrightarrow{\psi_{ij}} (M_j)_{f_i, f_j}$
 $\psi_{ij}: (M_i)_{f_j} \rightarrow (M_j)_{f_i}$ iso of R_{f_i, f_j} -mods \leftarrow condition

Define $M := \text{Ker} \left(\bigoplus_i M_i \xrightarrow{\varphi} \bigoplus_i (M_i)_{f_i} \right)$
 $(m_i) \mapsto (\frac{m_i}{f_j} - \psi_{ji}(\frac{m_j}{f_i}))$
 Call $\Pi_i: M \rightarrow M_i$ the projections.

Gluing Lemma Π_i induces isos $M_{f_i} \rightarrow M_i$ and $\psi_{ij} \circ \Pi_i = \Pi_j \circ \psi_{ji}$
 see Sec. 3.0 \leftarrow Idea: local data which agrees on overlaps

Pf Enough to show Π_i iso after localising at every prime $\mathfrak{q} \in \text{Spec } R$
 $\implies \mathfrak{q} = P_{f_i}$ with $f_i \notin P \in \text{Spec } R$. By exactness of localisation
 $(M_{f_i})_{\mathfrak{q}} = M_P = \text{Ker} \left(\bigoplus_i (M_i)_P \xrightarrow{\varphi_P} \bigoplus_i ((M_i)_P)_{f_i} \right)$

$f_i \in R_P$ is unit so WLOG replace $R \rightarrow R_P$, $M \rightarrow M_P$, $M_i \rightarrow (M_i)_P$, $f_i \rightarrow 1$.
 Abbreviate $N = M_{\mathfrak{q}}$ so: $\Pi_i: N = \text{Ker} \varphi_P \subseteq (N \oplus \bigoplus_{i \neq j} M_i) \rightarrow N$
 $\psi_{ji}: N \xrightarrow{\cong} (M_i)_P = M_i$

\leftarrow R_P -mods \leftarrow WLOG in some k mod localising at \mathfrak{q} is like localising at 1 since f_i is a unit in R_P

WLOG $M_i = N_{f_i}$ (identifies via ψ_{f_i}), so cocycle cond. becomes:

$$\Rightarrow 0 \rightarrow N \xrightarrow{\text{natural}} \bigoplus_i N_{f_i} \xrightarrow{\psi} \bigoplus_{i,j} N_{f_i f_j}$$

$$(N \rightarrow N \oplus \bigoplus_{i \neq j} N_{f_i} \xrightarrow{n \rightarrow n \oplus \frac{n}{1}} \bigoplus_{i \neq j} N_{f_i} \xrightarrow{(x_i) \mapsto (x_i - \frac{x_i}{1})} \bigoplus_{i,j} N_{f_i f_j})$$

Sub-claim: This is exact ($\Rightarrow N = \text{Ker } \psi = M$, \mathbb{T}_e iso, $\psi_{f_k} = \text{id}$ under identifications via π maps)
 Pf: Enough to prove after localising at each max ideal \mathfrak{m} ← See 3.0
 By \otimes not all $f_i \in \mathfrak{m}$ otherwise $1 \in \langle \text{all } f_i \rangle \subseteq \mathfrak{m} \subseteq \mathbb{Z}$
 Say $f_k \notin \mathfrak{m}$, so WLOG replace $N \rightarrow N_{f_k}, R \rightarrow R_{f_k} \xrightarrow{1} \dots$

$$\Rightarrow 0 \rightarrow N \rightarrow N \oplus \bigoplus_{i \neq k} N_{f_i} \xrightarrow{\psi} \bigoplus_{i,j} N_{f_i f_j}$$

clearly injective
 $n \oplus \bigoplus_{i \neq k} n_i \in \text{Ker then } \frac{n}{1} = \frac{n_i}{1} \in N_{f_i f_k} = N_{f_i} \quad \forall i$
 hence $= n \oplus \bigoplus_{i \neq k} n_i$ so image of n via previous map

7.6 Qcoh(X), Coh(X), Vect(X) for $X = \text{Spec } R$

Theorem
 For $X = \text{Spec } R$, \exists equivalence of categories

$$R\text{-Mods} \xrightarrow{\sim} \text{QCoh}(X)$$

$$M \xrightarrow{\sim} \tilde{M}$$

$$F(X) = \Gamma(X, F) \xleftarrow{\sim} \tilde{F}$$

Pf. Easy direction: $M \mapsto F = \tilde{M} \mapsto F(X) = \tilde{M}(X) = M$. Converse: given F want $F \cong \tilde{F}(X)$.
 \Rightarrow locally $\forall p \in X, \exists p \in D_f$ s.t. $F|_{D_f} \cong \tilde{F}|_{D_f}$ some $R_f\text{-mod } N$
 cover X by finitely many such, say $N_i \text{ on } D_{f_i}, i=1, \dots, n$, so $1 \in \langle \text{all } f_i \rangle$ and $\text{Spec } R$ quasi-compact
 \Rightarrow On overlaps: $\psi_{ij} : (N_i)_{f_j} \xrightarrow{\cong} (N_j)_{f_i}$ satisfy cocycle condition \leftarrow since $(N_i)_{f_j f_k}$ and other two are identified with $F|_{D_{f_i f_j f_k}}$
 \Rightarrow by gluing thm $\exists M$ with $M_{f_i} = N_i$ compatibly with the ψ_{ij}
 But then \tilde{M}, F have isomorphic local gluing data for cover $X = D_{f_1} \cup \dots \cup D_{f_n}$ so $\tilde{M} \cong F$.
 (Explicitly: $m \in M \mapsto m_i = \frac{m}{1} \in M_{f_i} = N_i \xrightarrow{\psi_i} s_i \in F(D_{f_i})$ and $s_i|_{D_{f_i f_j}} = s_j|_{D_{f_i f_j}}$ so globalises to unique $s \in F(X)$. Recall $M \rightarrow F(X)$ determines $\tilde{M} \rightarrow F$ by Sec. 6.9)

Cor $X = \text{Spec } R: F \in \text{Coh } X \Leftrightarrow F = \tilde{M}$ for coherent module M \leftarrow and if R Noeth., get:

$$F(X) = \tilde{F}(X) \Leftrightarrow F(X) \text{ f.g. } R\text{-mod}$$
 and conversely if M coherent get \tilde{M} coherent since \sim is exact & fully faithful. \square

Fact $X = \text{Spec } R: F \in \text{Vect } X \Leftrightarrow (F = \tilde{M} \text{ for finitely presented}) \Leftrightarrow \text{f.g. projective } R\text{-mod}$
 (see thm 4)
 means in $R\text{-mods}$
 $\text{Hom}(M, \cdot)$ exact
 $(\Leftrightarrow M$ is a direct summand of some free $R\text{-mod})$

8. Čech Cohomology
8.1 Čech complex

$$U_I = U_{i_0} \cap \dots \cap U_{i_n}$$

$$C^n = \check{C}_c^n\{U_i\} = \prod_{|I|=n} \Gamma(U_I, F)$$

$$d = d^n: C^n \rightarrow C^{n+1}$$

Motivation for cohomology: assign group or ring of n -invariants to a space i.e. iso. spaces give iso. of $e.g. H^*(X) \cong H^*(Y)$ then $X \cong Y$ are not iso. of spaces.
 top. space, $X = \cup U_i$ open cover
 $U_{ij} = U_i \cap U_j$
 $U_{ijk} = U_i \cap U_j \cap U_k$
 multi-index, abbreviate $|I|=n$
 ordered, allow repetitions
 see is actually $n+1$
 $S_I \in F(U_I)$
 called cochain
 where $I_j = (i_0, \dots, i_j, \dots, i_{n+1})$
 later also use notation I_{j_1, \dots, j_r} if omit i_j, i_{k_1}, \dots
 if you took C3.1 Algebraic Top. notice similar to simplicial differential
 $I_0 = (i_0, i_0)$
 $I_0 = (i_1) = j$
 $i_0 = i, i_1 = j$
 since i, j, k missing in I_k

Example $C^0 = \prod_i \Gamma(U_i) \xrightarrow{d} \prod_{i,j} \Gamma(U_{ij}) = C^1$
 $(s_i) \mapsto (s_j|_{U_{ij}} - s_i|_{U_{ij}})$
 $C^1 = \prod_{i,j} \Gamma(U_{ij}) \xrightarrow{d} \prod_{i,j,k} \Gamma(U_{ijk}) = C^2$
 $(s_{ij}) \mapsto (s_{jk}|_{U_{ijk}} - s_{ik}|_{U_{ijk}} + s_{ij}|_{U_{ijk}})$
 $U_{j,k} = U_j \cap U_k$
 $U_j \cap U_k = \sum_{j < k} (-1)^{k+j} s_{j,k} \Big|_{U_j \cap U_k}$
 \leftarrow anti-symmetry if swap j, k (notice full sum is over all $j \neq k$)
 since i, j, k missing in I_k
 called coboundaries
 cycles
 omitted
 sometimes
 Emphasizes doing
 Cohomology

Claim $d^2 = 0$, so (C^*, d) is a complex
Pf

$$(dds)_j = \sum_{k=0}^{n+2} (-1)^k (ds)_{j,k} = \sum_{k=0}^{n+2} \sum_{l=0}^{n+2} (-1)^{k+l} s_{j,k,l}$$

$$= 0. \square$$

$$H^n(X, F) = \check{H}^n\{U_i, F\} = \text{Ker } d^n / \text{Im } d^{n-1}$$

Lemma $H^0(X, F) = \Gamma(X, F)$
Pf $s_j|_{U_{ij}} = s_i|_{U_{ij}}$ says s glues to global section. \square

Terminology 1) hom of complexes $f: C^n \rightarrow C^n$ is chain map if $f \circ d = d \circ f$
 2) $f: C^n \rightarrow C^{n-1}$ is chain homotopy between chain maps f, g if $f - g = d \circ h + h \circ d$
Consequences: 1) $f: H^n \rightarrow H^n \leftarrow (dc = 0 \Rightarrow [f - g] = [fc] = [d \circ h] + h \circ d)$ well-defined
 2) $f = g: H^n \rightarrow H^n \leftarrow (dc = 0 \Rightarrow [f - g] = [d \circ h] + h \circ d)$
Key trick To show $H^* = 0$ can find chain homotopy between $id, 0$.
 i.e. C^* is exact, also called acyclic

Rmk If a homomorphism $d_n: C_n \rightarrow C_{n-1}$ decreases the degree by 1, and $d_{n-1} \circ d_n = 0$ then $H_n = \text{Ker } d_n / \text{Im } d_{n+1}$ is called the homology of (C_*, d_*) . In this case a chain homotopy is degree increasing: $f: C_n \rightarrow C_{n+1}$ with $f_n - g_n = d_{n+1} \circ f_n - f_{n-1} \circ d_n$.

Rmk \mathcal{L} line bundle with transition maps α_{ij}

$\Rightarrow \mathcal{L}^{-1}$ " " " " α_{ij}^{-1}

and $\mathcal{L} \otimes \mathcal{L}^{-1} \cong \mathcal{O}_X$ = trivial line bundle

FACT line bundles on A^n are always trivial

indeed vector bundles on A^n are always trivial

(Serre's conjecture 1955 (Quillen-Suslin Theorem 1976))

EXAMPLE $\text{Pic}(\mathbb{P}^1) \cong \mathbb{Z} \cong \mathbb{Z} \oplus \mathbb{Z} \cong \mathbb{Z} \oplus \mathbb{Z}$

$\mathbb{P}^1 = A_0 \cup A_1$

$\text{Spec } k[t] \cong \mathbb{A}^1$ $\Rightarrow \mathcal{L}|_{A_0} \cong \mathcal{L}|_{A_1}$ trivial since $A_i \cong \mathbb{A}^1$

Have homogeneous coordinates $[x_0 : x_1]$ and A_0 corresponds to $\{[t : 1] : t \in A^1\}$ where $t = x_1/x_0$

$\alpha_{10} : \mathcal{L}|_{A_1} \rightarrow \mathcal{L}|_{A_0} \in k[t, t^{-1}]^*$

$\beta_0 \in k[t]^* = k^*$, $\beta_1 \in k[t^{-1}]^* = k^*$

$\Rightarrow \text{Pic}(\mathbb{P}^1) \cong \mathbb{H}^1(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}^*) \cong \mathbb{Z} \oplus \mathbb{Z}$

$\mathcal{O}(i) \leftarrow (\alpha_{10} = t^i) \leftarrow i$

so define $\mathcal{O}(i)$ by using $\alpha_{10} = t^i$, $\alpha_{01} = t^{-i}$

Rmk $\mathcal{O}(0) = \mathcal{O}_{\mathbb{P}^1}$ trivial line bundle.

Hwk 4 Ideal sheaf of a closed point in \mathbb{P}^1 is $\mathcal{O}(-1)$ for disjoint union of n closed pts get $\mathcal{O}(-n)$

for order n point $(t^i) \in k[t]$ (i.e. closed subscheme $\text{Spec } k[t]/(t^i) \subseteq A_0 \subseteq \mathbb{P}^1$) get $\mathcal{O}(-n)$.

Non-examinable Rmk (for differential geometers): i determines the Chern class $c_1(\mathcal{L}) : i = \int_{\mathbb{P}^1} c_1(\mathcal{L})$

$T\mathbb{P}^1$ is $\mathcal{O}(2)$ since $2 = \chi(\mathbb{P}^1) = \chi(S^2)$ and $c_1(T\mathbb{P}^1) = \text{Euler class of } \mathbb{P}^1$, and $T^*\mathbb{P}^1 = \mathcal{O}(-2)$.

$\mathcal{O}(-1) \rightarrow \mathbb{P}^1$ is blow-up of \mathbb{P}^2 at 0 : the lines through 0 in \mathbb{P}^2 are the fibres.

Theorem

Cultural Rmk symmetry is "Serre duality" for \mathbb{P}^1 : $\mathbb{H}^1(\mathcal{O}(i)) \cong \mathbb{H}^0(\mathcal{O}(-i-2)) = \mathcal{O}(-i-2)$

1) $\mathbb{H}^0(\mathbb{P}^1, \mathcal{O}(i)) = \begin{cases} 0 & i < 0 \\ \{f \in k[t] : \deg f \leq i\} \cong k[x_0, x_1] & i \geq 0 \end{cases}$

2) $\mathbb{H}^1(\mathbb{P}^1, \mathcal{O}(i)) = \begin{cases} 0 & i > -1 \\ k[t^{-1}] / k + t \cdot k[t^{-1}] \cong k[x_0, x_1] & i < -1 \end{cases}$

3) $\mathbb{H}^n(\mathbb{P}^1, \mathcal{O}(i)) = 0$ for $n \geq 2$

Pf By 8.6, since \mathbb{P}^1 separated & quasi-compact enough to calculate $\mathbb{H}_{\{A_0, A_1\}}^*(\mathbb{P}^1, \mathcal{O}(i))$

3) no triple order overlaps or higher $\Rightarrow g \in \mathcal{O}_{A_1} \otimes_{\mathcal{O}_{A_0}} \mathcal{O}_{A_1} \otimes_{\mathcal{O}_{A_0}} \mathcal{O}_{A_1}$ where α_{10} is defined on $A_0 \cap A_1$.

1) $\mathbb{H}^0 = \Gamma = g(t^{-1}) \in k[t^{-1}]$ on A_1 , $f(t) \in k[t]$ on A_0 , on overlap: $t \cdot g(t^{-1}) = f(t) \in k[t, t^{-1}]$

$\Rightarrow \deg f \leq i$ and g is determined by f from equation

2) $\mathcal{L} = \mathcal{O}(i) \oplus \Gamma(A_0, \mathcal{L}) \oplus \Gamma(A_1, \mathcal{L}) \xrightarrow{d} \Gamma(A_0 \cap A_1, \mathcal{L}) \xrightarrow{d} 0$

(strictly speaking $\mathcal{L}(A_0) \cong \mathcal{O}_{A_0}(i)$, $\mathcal{L}(A_1) \cong \mathcal{O}_{A_1}(i)$)

$\mathcal{L}(A_0) \cong \mathcal{O}_{A_0}(i) \cong k[t]$, $\mathcal{L}(A_1) \cong \mathcal{O}_{A_1}(i) \cong k[t^{-1}]$

$(f, g) \mapsto t \cdot g(t^{-1}) - f(t)$

$\mathbb{H}^1 = k[t, t^{-1}] / k[t] + t \cdot k[t^{-1}]$

is all of $k[t, t^{-1}]$ if $i \geq -1$

does not contain $t^{-1}, t^{-2}, \dots, t^{-i-1}$ if $i < -1$

need to transition from $g(t^{-1}) \in \mathcal{O}_{A_1}(i)$ to $\mathcal{O}_{A_0}(i)$ via $\alpha_{10} : \mathcal{O}_{A_1} \cong \mathcal{L}|_{A_1} \cong \mathcal{L}|_{A_0} \cong \mathcal{O}_{A_0}(i)$

EXAMPLE: \mathbb{P}^n

$X = \mathbb{P}^n = A_0 \cup A_1 \cup \dots \cup A_n$

$A_i = \text{Spec } k[\frac{x_0}{x_i}, \dots, \frac{x_n}{x_i}]$ (omit $\frac{x_i}{x_i}$)

called hyperplane bundle or Serre's twisting sheaf

$\mathcal{O}(1) = \text{line bundle with } \alpha_{ij} = (\frac{x_i}{x_j}) : k[\frac{x_0}{x_i}, \dots, \frac{x_n}{x_i}] \rightarrow k[\frac{x_0}{x_j}, \dots, \frac{x_n}{x_j}]$

$\mathcal{O}(m) = \mathcal{O}(1)^{\otimes m}$ so $\alpha_{ij} = (\frac{x_i}{x_j})^m$

both equal to $\Gamma(A_i \cap A_j, \mathcal{O}_X)$ is multiplication by $\frac{x_0}{x_i} = t^{-1}$ ✓

Rmk $\mathcal{O}(-1)$ called tantological line bundle because in C3.4 course each (closed) point of \mathbb{P}^n is a 1-dim vector subspace $V \subseteq k^{n+1}$ ($\mathbb{P}^n = k^{n+1} / k^*$ -rescaling)

so get obvious line bundle: over the point $[V] \in \mathbb{P}^n$ have the line V .

Hwk 4 $\text{Pic}(\mathbb{P}^n) \cong \mathbb{Z}$ generated by the $\mathcal{O}(1)$

$\Gamma(\mathbb{P}^n, \mathcal{O}(m)) = \begin{cases} k[x_0, \dots, x_n]_m & \text{if } m \geq 0 \\ 0 & \text{if } m < 0 \end{cases}$

So homogeneous polys of deg $\leq m$. So on A_i get polys of deg $\leq m$ in the variables x_0, \dots, x_n

8.8 Divisors

Let $(X, \mathcal{O}_X = \mathcal{O})$ be an integral scheme (i.e. reduced & irreducible) ← see Sec. 3.5

Recall from Sec. 3.5 that \forall open $\emptyset \neq U \subseteq X$ can view $\mathcal{O}(U) \cong K(U)$ inside $K(X) =$ function field

Abbreviate: $K = K(X)$, $K^* = K \setminus \{0\}$ (non-zero rational functions are invertible)

$\mathcal{O}^* \subseteq \mathcal{O}$ subsheaf of sections of \mathcal{O} admitting inverses in \mathcal{O} (so can view $\mathcal{O}^* \subseteq K^*$)

$X = \cup U_i$ open cover

$f_i \in K^*$ s.t. $f_i |_{U_i \cap U_j} = \frac{f_j}{f_i} |_{U_i \cap U_j}$ (see below)

\Rightarrow get line bundle $\mathcal{L} \subseteq K$ via $\mathcal{L}(U) := \mathcal{O}(U) \cdot \frac{f_i}{f_j} \subseteq K$

Exercise

1) Obvious trivializations $\varphi_i : \mathcal{L}(U_i) \rightarrow \mathcal{O}(U_i)$, $g \cdot \frac{f_i}{f_j} \mapsto g$

Yields transition maps $\alpha_{ij} = \varphi_j \circ \varphi_i^{-1} |_{U_i \cap U_j} = \frac{f_j}{f_i}$ (from U_i to U_j)

2) If $D_1 = (U_1, f_1), D_2 = (U_2, g_2)$ are two Cartier divisors on X yielding line bundles $\mathcal{L}_1, \mathcal{L}_2$ then $D_1 + D_2 = (U_1 \cap U_2, f_1 g_2)$ is a Cartier divisor yielding the line bundle $\mathcal{L}_1 \otimes \mathcal{L}_2$ [in particular $-D_1 - D_2 = (U_1 \cap U_2, \frac{1}{f_1 g_2})$ yields $\mathcal{L}_1^{-1} \otimes \mathcal{L}_2^{-1}$]

Key Example Recall $\mathbb{P}^n = \cup U_i$ for $U_i = \text{Spec } Z[\frac{x_0}{x_i}, \dots, \frac{x_n}{x_i}]$ ($\frac{x_i}{x_i} = 1$)

Let $m \in \mathbb{Z}$, $f_0 = 1, f_i = (\frac{x_0}{x_i})^m \in K(\mathbb{P}^n) = \mathbb{Q}(x_0, \dots, x_n)$, $\forall i \in \{0, \dots, n\}$

$\mathcal{L}(U_0) = \mathcal{O}_{\mathbb{P}^n}(U_0) \cdot 1 \subseteq K(\mathbb{P}^n)$ (side remark: $K(\mathbb{P}^n) \cong k(U) \cong k(A^n) \cong \mathbb{Q}(x_1, \dots, x_n)$)

$\mathcal{L}(U_i) = \mathcal{O}_{\mathbb{P}^n}(U_i) \cdot (\frac{x_0}{x_i})^m \subseteq K(\mathbb{P}^n)$ transition $\alpha_{ij} = (\frac{x_0}{x_j} \frac{x_i}{x_0})^m = (\frac{x_i}{x_j})^m$ (from U_i to U_j) so $\mathcal{L} = \mathcal{O}_{\mathbb{P}^n}(m)$

Rmk This does not look very "symmetric" in the x_i . One can define an $\mathcal{O}_{\mathbb{P}^n}$ -module F by $F(U_i) = \mathcal{O}_{\mathbb{P}^n}(U_i) \cdot x_i^m$ which is a line bundle with the same transitions $\alpha_{ij} = (\frac{x_i}{x_j})^m$

So $F \cong \mathcal{L}$ above, but we cannot pick $f_i = x_i^m$ for the Cartier divisor since $x_i \notin K(\mathbb{P}^n)$.

Actually want to identify Cartier divisors related by refining the cover, so if $X = \cup U_i = \cup V_j$ and $V_j \subseteq U_i$ compare Sec. 8.6 then identify (U_i, f_i) and (V_j, f_j) .

(Also identify (U_i, f_i) with $(U_i, f_i g)$ if $g \in \mathcal{O}^*(U_i)$ ← i.e. rescaling f_i by invertible regular-fun)

Viewing K, K^* as constant sheaves, have an exact sequence

$$0 \rightarrow \mathcal{O}^* \rightarrow K^* \rightarrow K^*/\mathcal{O}^* \rightarrow 0$$

Because of \otimes , a Cartier divisor is just a global section of K^*/\mathcal{O}^* so $\check{H}^0(X, K^*/\mathcal{O}^*)$

Take LES: $0 \rightarrow \check{H}^0(X, \mathcal{O}^*) \rightarrow \check{H}^0(X, K^*) \rightarrow \check{H}^0(X, K^*/\mathcal{O}^*) \rightarrow \check{H}^1(X, \mathcal{O}^*) \rightarrow \check{H}^1(X, K^*) \rightarrow \dots$

A Cartier divisor in image of \check{H}^0 is called principal (i.e. use cover X and one $f \in K^*$)
 Two Cartier divisors D, D' are linearly equivalent if $D - D'$ is principal. Write $D \sim D'$.

Get abelian group $\text{CaCl}(X)$ of Cartier divisors modulo linear equivalence.
 $\Rightarrow \text{CaCl}(X) \cong \text{Pic}(X)$ by the LES in particular (bundles $\mathcal{L}(D) \cong \mathcal{L}(D')$) $\Leftrightarrow D \sim D'$.

Cultural Rmks (Non-examinable) There is another notion of divisor: Weil divisor.
 This means a formal sum $\sum_{i=1}^n \mathbb{Z} \cdot Z_i$ of integral closed subschemes Z_i of codim=1 (think hypersurfaces)

Example rational function $f \in K(X) \Rightarrow \exists$ an "order of vanishing" $\text{ord}_Z(f)$ of f along such subschemes Z .

\Rightarrow Weil divisor $\text{div}(f) := \sum \text{ord}_Z(f) \cdot Z$ called principal Weil divisor

Example Cartier divisor (U, f) yields Weil divisor $W = \sum_{i=1}^m \mathbb{Z} \cdot Z_i$ (notice compute the order of $f|_{Z_i} \in K(U_i)$ along Z_i)

On \mathbb{P}^1 : Cartier divisor $(U_0, 1)$, $(U_1, \frac{z_0}{z_1})$ yields $W = + \text{point } [0:1] - \text{point } [0:0]$ but ignore pole $z_1=0$

Cartier divisor $(U_0, 1)$, $(U_1, \frac{z_0}{z_1})$ yields $W = m \cdot p$ where $m \in \mathbb{Z}$, $p = [0:1]$ since $[0:0] \notin U_1$

On \mathbb{P}^n : $(U_i, 1)$, $(U_j, \frac{z_0}{z_j})$ yields $W = H$ where $H \in \mathbb{P}^{n-1}$ is the hyperplane (case $m=0$ is when f has a pole at $p = [0:0]$)

The lack of "symmetry" mentioned in Rmk above is because it involves a choice of Weil divisor H . We could have picked any hyperplane to get $\mathcal{L} \cong \mathcal{O}(H)$. More complicated choices are possible

e.g. Cartier divisor D on \mathbb{P}^1 with $W = \sum n_i \cdot p_i$ any points p_i and $n_i \in \mathbb{Z}$, yields $\mathcal{L}(D) \cong \mathcal{O}(\sum n_i)$ (compare Hw 4)

Weil divisors $\text{Div}(X)$ modulo principal Weil divisors define the class group $\text{Cl}(X)$ (abelian group).

Weil divisor D defines an \mathcal{O}_X -module $\mathcal{O}_X(D)$ by $\Gamma(U, \mathcal{O}_X(D)) = \{f \in K \mid \text{div}(f) + D \geq 0 \text{ on } U\}$

But $\mathcal{O}_X(D)$ need not be a line bundle (i.e. invertible sheaf). When it is a line bundle the Weil divisor is Cartier since on some cover $X = \cup U_i$ have trivializations $\mathcal{O}(U_i) \cong \Gamma(\mathcal{O}_X(D)|_{U_i})$

\Rightarrow Cartier divisor (U_i, f_i) and $\mathcal{L}(U_i) = \mathcal{O}(U_i) \cdot f_i = \Gamma(U_i, \mathcal{O}_X(D))$

Weil divisor is Cartier if locally principal: so locally looks like $\text{div}(\text{div}(f))$ (e.g. $D = \text{div}(f)$ gives $\mathcal{O}_X(D) \cong \mathcal{O}$ via $g \mapsto gf$)

(also need mild condition: X is "normal")

X non-singular variety $\Rightarrow \text{CaCl}(X) \cong \text{Cl}(X) \cong \text{Pic}(X)$ e.g. get \mathcal{L} for \mathbb{P}^n

For X singular it can fail: $X = \text{Spec } k[x, y, z]/(xy - z^2) \leq \mathbb{A}_k^3$ has $\text{CaCl}(X) = 0$ but $\text{Cl}(X) = \mathbb{Z}/2$ generated by the hypersurface $Z = (y = z = 0)$. (At $\mathcal{O} \in \mathbb{Z}$ we really need 2 equations to cut out \mathbb{Z} , one is not enough, so not locally principal.)

Cultural Remark: Riemann-Roch Theorem (non-examinable)
 C projective non-singular algebraic curve/adg. closed field k
 $F = \mathcal{O}_C(D)$ for divisor D of degree d $\leftarrow \dim(\text{global sections})$ often written $\ell(D)$.
 $\chi(C, F) := \sum_{i=0}^d (-1)^i \dim H^i(C, F) = \rho(C, F) - h(F) = \deg D + \chi(C, \mathcal{O}_C)$
 $\leftarrow \rho^m = \dim_k H^0(C, \mathcal{O}_C(m))$ $\leftarrow \chi(C, \mathcal{O}_C) = 1 - \text{genus}(C)$

8.9 Čech cohomology computations on \mathbb{P}^n

Recall the key example in Sec. 8.8:

$\mathbb{P}^n = \bigcup_{i=0}^n U_i$ where $U_i = D_+ x_i = \text{Spec } \mathbb{Z}[\frac{x_0}{x_i}, \dots, \frac{x_{i-1}}{x_i}, \frac{x_{i+1}}{x_i}, \dots, \frac{x_n}{x_i}] \cong \mathbb{A}^n$

Line bundle $\mathcal{L} = \mathcal{O}_{\mathbb{P}^n}(d)$ for $d \in \mathbb{Z}$ has:

$\Gamma(U_i, \mathcal{L}) = (\mathbb{Z}[x_0, \dots, x_n][\frac{1}{x_i}])_d$ \leftarrow so poly. in x 's of degree $n+d$ and $N \geq 0$.
 says localize at x_i .

example: $d=0$ gives $\mathcal{L} = \mathcal{O}_{\mathbb{P}^n}$ and $\Gamma(U_i, \mathcal{L}) = \mathbb{Z}[x_0, \dots, x_n]$ \leftarrow the classical functions on \mathbb{A}^n , $\deg P = N, N \geq 0$.
 U_i well-defined when rescale homogeneous coords.

Use ordered Čech cohomology using obvious ordering of $i \in \{0, 1, \dots, n\}$.

$\Gamma(U_{i_0, \dots, i_k}, \mathcal{L}) = (\mathbb{Z}[x_0, \dots, x_n][\frac{1}{x_{i_0} \dots x_{i_k}}])_d$ $\leftarrow (U_{i_0, \dots, i_k} = U_{i_0} \cap \dots \cap U_{i_k} \quad 0 \leq i_0 < \dots < i_k \leq n)$

Warm-up example $\check{H}^1(\mathbb{P}^2, \mathcal{L}) = 0$

Proof $C_{ij} \in \check{C}^1$ is \mathbb{Z} -combo of terms $\frac{x_0^{m_0} x_1^{m_1} x_2^{m_2}}{(x_i x_j)^n}$ where total degree $\sum m_i = -2N = d$

C cocycle $\Rightarrow (dC)_{012} = 0 \stackrel{*}{=} C_{12} - C_{02} + C_{01} \in \Gamma(U_{012}, \mathcal{L})$ \leftarrow (e.g. $C_{12} \in \Gamma(U_{12}, \mathcal{L})$ and we restrict to U_{012})

Want to show cocycle C is a coboundary i.e. $\exists b_i \in \Gamma(U_i, \mathcal{L})$, $(dC)_{ij} = b_j - b_i = C_{ij}$.

Want $b_i \in \Gamma(U_i, \mathcal{L})$ so only x_i denominators allowed. (e.g. $\frac{x_1^2 x_2}{x_1 x_2} = \frac{x_1 x_2}{x_2}$)

Key observation: C_{12} cannot have both x_1, x_2 arising as a denominator (after simplify) because C_{02} has no x_2 's at denom, C_{01} has no x_1 's at denom.

Expand terms depending on denominators: e.g. C_{12}, x_1 = terms of C_{12} which have x_1 denominators

$C_{12} = \frac{C_{12, x_2}}{x_2} + \frac{C_{12, x_1}}{x_1} + P_{12}$ $\leftarrow P_{12}$ are leftover terms, so no denominators (so polys of degree d if $d \geq 0$, otherwise 0)

$-C_{02} = \frac{-C_{02, x_2}}{x_2} - P_{02}$

$C_{01} = \frac{C_{01, x_1}}{x_1} + P_{01}$

\Rightarrow calling $b_2 = C_{12, x_2}$, $b_1 = -C_{12, x_1}$, $b_0 = -C_{02}$, x_0 get

$\begin{cases} C_{12} = b_2 - b_1 + P_{12} \\ -C_{02} = -b_2 + b_0 - P_{02} \\ C_{01} = b_1 - b_0 + P_{01} \end{cases}$ \leftarrow (boundary so does not change cohomology)

\Rightarrow replacing C by $C - dB$ remains to consider the case $C_{ij} = P_{ij}$ (no denominators)

Trick! Let $q_2 = \alpha$, $q_0 = 1$, $q_1 = 0$ then $(dq)_{ij} = q_j - q_i = \begin{cases} \alpha & \text{if } (i,j) = (0,2) \\ 0 & \text{if } (i,j) = (1,2) \\ 0 & \text{else} \end{cases}$

Taking $\alpha = P_{12}$ we can replace C by $C - dq$ and assume $P_{12} = 0$ (and redefine P_{02} due to) \leftarrow since $dc = 0, P_{12} = 0$

Whereas for $q_2 = 1, q_1 = 0, q_0 = -\beta$ get $(dq)_{ij} = \begin{cases} \beta & \text{if } (0,1) \\ 0 & \text{else} \end{cases}$ \leftarrow (since $dc = 0, P_{12} = 0$)

Taking $\beta = P_{02}$ replacing C by $C - dq$ we can assume $P_{02} = 0$, so also $P_{01} = 0$, so $C = 0$

Lemma $\check{H}^1(\mathbb{P}^n, \mathcal{L}) = 0 \quad \forall n \geq 2$ \leftarrow (n=1 fails because don't have triple overlaps \leftarrow TRY ON YOUR OWN FIRST! \leftarrow we computed the n=1 case in Sec. 8.7)

Proof The first part of proof of $n=2$ case is same: replace $0, 1, 2$ by indices i_0, i_1, i_2 . So reduce to case of cocycle $C \in \check{C}^1$ with C_{ij} having no denominators (agree $d \geq 0$)

Doing Trick 1 now is messy I think, so I'll use another trick first.

9. Sheaf Cohomology

9.1 Resolutions

← (Reference for more details: Lang, Algebra, Chapter XX §4-6)

Motivation: "represent" an object in an abelian category \mathcal{A} by "nicer objects" at the cost of using a chain complex (Sec. 1.8)

right resolution of MEA means an exact sequence $0 \rightarrow M \rightarrow I^0 \rightarrow I^1 \rightarrow I^2 \rightarrow \dots$ in \mathcal{A} abbreviated as $M \rightarrow I^\bullet$

left resolution $\dots \rightarrow P_2 \rightarrow P_1 \rightarrow P_0 \rightarrow M \rightarrow 0$, or $P_\bullet \rightarrow M$

Def I injective if $\text{Hom}(\cdot, I)$ exact \Rightarrow (both always left exact)

P projective if $\text{Hom}(P, \cdot)$ exact

Exercise I injective is equivalent to: $\forall \text{inj } A \hookrightarrow B, \exists \text{ "extend" } \psi: A \xrightarrow{\varphi} I \xrightarrow{\psi} B$

Fact injective resolution $M \rightarrow I^\bullet$ means I^n are injective

projective resolution $P_\bullet \rightarrow M$ " P_n " projective

$f, g: A \rightarrow B$ additive functors of abelian cats (see 1.7)

f left \Rightarrow right-derived functor $R^n f(M) = H^n(f(I^\bullet))$ (see 1.8)

g right \Rightarrow left-derived functor $L_n g(M) = H_n(g(P_\bullet))$

Warning f left exact only implies $0 \rightarrow fM \rightarrow fI^0 \rightarrow fI^1 \rightarrow \dots$ exact. Deduce: $R^0 f(M) \cong fM$

Similarly $L_0 g \cong g$, so $R^0 f, L_0 g$ remember the functors f, g .

Classical Examples $A = S\text{-Mod}$, $f = \text{Hom}(M, \cdot), N \rightarrow I^\bullet$ inj. res. $(\text{Ext}_S^n(M, N) \cong \text{Hom}_S(M, I^n))$

(Similarly: $f = \text{Hom}(\cdot, N): S\text{-Mod} \rightarrow \text{Ab}, \text{Ext}_S^n(M, N) = (R^n f)(M) = H_n(\text{Hom}(P_\bullet, N))$)

$g = M \otimes_S \cdot$ right exact $\Rightarrow \text{Tor}_S^n(M, N) = (L_n g)(N) = H_n(M \otimes_S P_\bullet)$ ($\text{Tor}_0^S(M, N) \cong M \otimes_S N$)

(Similarly: $g = \cdot \otimes_S N, \text{Tor}_S^n(M, N) = (L_n g)(M) = H_n(P_\bullet \otimes_S N)$ for $P_\bullet \rightarrow M$ proj. res.)

For $R\text{-mods}$: I injective \Leftrightarrow if $I \subseteq \text{any mod } M$ then $\exists \text{ mod } J: I \oplus J = M \leftarrow \text{compare linear algebra "extending a basis"}$

Fact $M \rightarrow I^\bullet$ inj. res., \downarrow morph \Rightarrow can extend $M \rightarrow I^\bullet$ and any 2 choices $\Rightarrow f(M) \rightarrow H^i(f(I^\bullet))$

Key idea: I inj $\Rightarrow \text{Hom}(\cdot, I)$ right exact \Rightarrow if $A \xrightarrow{m} B$ then any $A \rightarrow I$ can be extended to $B \rightarrow I$. E.g. $M \rightarrow I^\bullet \Rightarrow M \rightarrow I^0 \rightarrow I^1 \rightarrow \dots$

then consider $\text{Coker}(M \rightarrow I^0) \hookrightarrow I^1$ and continue inductively. Try proving the rest.

Cor 1) $R^n f(M) = H^n(fI^\bullet)$ independent of choice of inj. res. $M \rightarrow I^\bullet$

2) $M \rightarrow N$ induces $R^n f(M) \rightarrow R^n f(N)$, indeed $R^n f: \mathcal{A} \rightarrow \mathcal{A}$ is functor.

Pf 1) Apply fact to $M=N$, get $H^i(fI^\bullet) \rightarrow H^i(fJ^\bullet) \rightarrow H^i(fI^\bullet) \rightarrow H^i(fI^\bullet)$ composite is id by uniqueness.

2) By Fact, $R^n f(M) = H^n(fI^\bullet) \rightarrow H^n(fJ^\bullet) = R^n f(N)$. Exercise: check functor. \square

Trick 2 $\sum_{i,j} c_{ij} \in (\mathbb{Z}[\alpha_0, \dots, \alpha_n] / \langle \alpha_i^2 \rangle)_0 = \mathbb{Z}[\alpha_1, \dots, \alpha_n] =$ global sections on $U_0 \cong \mathbb{A}^n$

This is a 1-cycle on \mathbb{A}^n and we know $H^1(\mathbb{A}^n, \mathcal{O}_{\mathbb{A}^n}) = 0$ (by Sec. 8.3 since \mathbb{A}^n affine)

So $\exists \beta_i \in \mathbb{Z}[\alpha_1, \dots, \alpha_n] / \langle \alpha_i^2 \rangle$ with $(d\beta)_i = \sum_{j \neq i} c_{ij} \alpha_j$ for $1 \leq i \leq n$

Since c_{ij} has no denominators, β_i cannot have any α_i denominator.

Since c_{ij} is homog. of deg $= d$ in the α 's, wlog β_i is homogeneous of deg $= d$ in α 's

\Rightarrow Take $b_i = \alpha_0^d \beta_i =$ homog. deg d poly in α 's with $(db)_i = c_{ij}$ for $1 \leq i \leq n$.

\Rightarrow Replace c by $c - db$, can assume $c_{ij} = 0$ for $i \neq 0$.

Final trick $(dc)_{0,i} = 0 = \sum_{j \neq 0} c_{0j} \alpha_j + c_{00} \alpha_0$ so all c_{0i} are the same say $= \beta$, so we

Trick 1 with $q_i = 0$ for $i \neq 0, q_0 = -\beta$ then $(dq)_i = \begin{cases} \beta & \text{if } i=0 \\ 0 & \text{if } i \neq 0 \end{cases}$ so $c = dq$. \square

Theorem For $\mathcal{L} = \mathcal{O}(d)$, $d \in \mathbb{Z}, n \geq 2$ degree d homog. polys (so $\{0, j$ if $d < 0$)

$H^i(\mathbb{P}^n, \mathcal{L}) = \begin{cases} \mathbb{Z}[\alpha_0, \dots, \alpha_n]_d & \text{for } * = 0 \leftarrow \text{Hwk 4, global sections of } \mathcal{O}_{\mathbb{P}^n}(d) \\ 0 & \text{for } 0 < * \leq n \\ \mathbb{Z} \left\{ \frac{1}{\alpha_0 \alpha_1 \dots \alpha_n} \cdot \frac{1}{x_n^m} \right\} & \text{of total degree } d \text{ for } * = n \leftarrow \text{free } \mathbb{Z}\text{-module with that basis} \\ 0 & \text{for } * > n \leftarrow \text{no } n+2 \text{ overlaps or higher since } n+1 \text{ sets } U_i \text{ cover} \end{cases}$

Proof $0 < * = k < n$ is same as for H^1 : exercise for you.

(Hint: $\pm b_{i_0} \dots \pm b_{i_k} \dots \pm b_{i_n} =$ terms in $c_{i_0 \dots i_k} \alpha_{i_0} \dots \alpha_{i_k}$ with no α_{i_j} at denominator)

(notice) those must cancel with similar terms in $c_{i_0 \dots i_k} \alpha_{i_0} \dots \alpha_{i_k}$

Pick sign it has as a term in $(db)_{i_0 \dots i_k} \alpha_{i_0} \dots \alpha_{i_k} \leftarrow$ since want this to give $c_{i_0 \dots i_k} \alpha_{i_0} \dots \alpha_{i_k}$

Case $* = n$: only one possible overlap: $U_{0,1,2,\dots,n}$, any chain $c \in \mathbb{Z}^n$ is cocycle since

no higher overlaps. Question becomes what are possible $(db)_{0,1,\dots,n}$ for $b_i \in \Gamma(U_{0,1,\dots,n}, \mathcal{L})$.

$(db)_{0,1,\dots,n} = b_{1,2,\dots,n} - b_{0,2,\dots,n} + b_{0,1,3,\dots,n} - \dots$ so can get all \mathbb{Z}^m with some $m_i \geq 0$

(i.e. some α_i not in denom) \leftarrow no α_0 at denom

$\Rightarrow H^n = \mathbb{Z} \{ \frac{1}{\alpha_0^m} : \sum m_i = d \} / \mathbb{Z} \{ \frac{1}{\alpha_0^m} : \sum m_i = d, \text{ some } m_i \geq 0 \}$

$\cong \mathbb{Z} \{ \frac{1}{\alpha_0^m} : \sum m_i = d, \text{ all } m_i < 0 \}$

$= \frac{1}{\alpha_0^d} \cdot \mathbb{Z} \{ \frac{1}{\alpha_1^{m_1} \dots \alpha_n^{m_n}} : \sum m_i = -d - n - 1, \text{ all } m_i \geq 0 \}$. \square

Exercise deduce the ranks $\beta^i = \text{rank}_{\mathbb{Z}} H^i$ are $\beta^i(\mathbb{P}^n, \mathcal{O}(d)) = \begin{cases} \binom{n+d}{n} & \text{if } i=0 \\ \binom{d-n}{n} & \text{if } i=n \\ 0 & \text{else} \end{cases}$

Motivation for chapter 9: Now that we know $H^*(\mathbb{P}^n, \mathcal{O}(d))$, one might hope to compute $H^*(\mathbb{P}^n, F)$ for other $F \in \text{Coh}(\mathbb{P}^n)$ by first finding a resolution $\dots \rightarrow \mathcal{L}_2 \rightarrow \mathcal{L}_1 \rightarrow F \rightarrow 0$ with $\mathcal{L}_i = \bigoplus \mathcal{O}(d_{ij})$ and exploiting LES.

8.10 Product on Čech cohomology (Non-examinable section)

(X, \mathcal{O}_X) any ringed space

$H^p(X, F) \times H^q(X, G) \rightarrow H^{p+q}(X, F \otimes G)$

$\{U_i\} \rightarrow \{U_i\} \rightarrow (S_{\mathbb{Z}} \otimes t_{\mathbb{Z}})$

Rank 1h 8.6 where we took constant coefficients $F=G=\mathbb{Z}$ (note: $\mathbb{Z} \otimes_{\mathbb{Z}} \mathbb{Z} \cong \mathbb{Z}$)

we recover the cup product on singular cohomology (respectively on de Rham cohomology)

using $F=G=\mathbb{R}$ for \mathbb{R}^n

\mathbb{R}^n smooth real functions

so $\mathbb{R} \otimes_{\mathbb{R}} \mathbb{R} \cong \mathbb{R}$

Lemma f left exact, $0 \rightarrow M_1 \rightarrow M_2 \rightarrow M_3 \rightarrow 0$ SES $\Rightarrow \exists$ canonical & functorial LES

$$0 \rightarrow R^0 f(M_1) \rightarrow R^0 f(M_2) \rightarrow R^0 f(M_3) \rightarrow R^1 f(M_1) \rightarrow R^1 f(M_2) \rightarrow R^1 f(M_3) \rightarrow R^2 f(M_1) \rightarrow \dots$$

Sketch Pf $0 \rightarrow I_1^0 \rightarrow I_2^0 \oplus I_3^0 \rightarrow I_3^0 \rightarrow 0$ ← first pick inj. res. I_1^0, I_3^0 then define I_2^0 that way so get obvious SES.
 where these triples are just Rⁿf applied to the SES

$$0 \rightarrow M_1 \xrightarrow{f_{M_1}} M_2 \xrightarrow{f_{M_2}} M_3 \rightarrow 0$$

use obvious map $M_2 \rightarrow M_3 \rightarrow I_3^0$ and $M_1 \hookrightarrow I_1^0$ extends via $M_1 \rightarrow M_2 \rightarrow M_3 \rightarrow I_3^0$
 Exercise: $M_2 \hookrightarrow I_2^0 = I_1^0 \oplus I_3^0$ is injective.
 Then take cokernels $M_1' = \text{Coker}(M_1 \rightarrow I_1^0)$, check that $0 \rightarrow M_1' \rightarrow M_2' \rightarrow M_3' \rightarrow 0$ exact, and repeat construction.

(Fact additive functors preserve \oplus)

$$\Rightarrow 0 \rightarrow fI_1^0 \rightarrow fI_2^0 \oplus fI_3^0 \rightarrow fI_3^0 \rightarrow 0 \leftarrow f \text{ may only be left exact, but here clearly } fI_2^0 \text{ surjects onto } fI_3^0 \text{ since have projection onto } fI_3^0 \text{ summand.}$$

$$0 \rightarrow fM_1 \rightarrow fM_2 \rightarrow fM_3 \rightarrow 0$$

Finally take the LES associated to the SES of complexes $0 \rightarrow fI_1^0 \rightarrow fI_2^0 \rightarrow fI_3^0 \rightarrow 0 \rightarrow 0$

Rmk Indeed $R^0 f$ satisfies universal property that " $R^0 f = f$ and Lemma holds", then it follows that $R^0 f(M) = H^0(f(I^\bullet))$ for any inj. res. $M \rightarrow I^\bullet$ (see end of next section)

Hwk 4 $\text{Ab}(X) \rightarrow \text{Ab}$ left exact \Rightarrow can define sheaf cohomology $H^n(X, F) = R^n \Gamma(X, F)$ (Sec. 1.9)

We now ask how this relates to $H^n(X, F)$ for $F \in \text{QCoh}(X) \subseteq \text{Ab}(X)$ and X scheme.

9.2 Acyclic resolutions (in an abelian cat.)

Rmk If I inj. object \Rightarrow resolution $0 \rightarrow I \rightarrow I^0 \rightarrow I^1 \rightarrow \dots \Rightarrow R^n f(I) = 0 \forall n \geq 1$

So for sheaf cohomology: $H^n(X, I) = 0 \forall n \geq 1$ if I injective sheaf.

Def An acyclic resolution of F is an exact sequence $0 \rightarrow F \rightarrow J^0 \rightarrow J^1 \rightarrow \dots$ with $H^n(X, J^k) = 0 \forall n \geq 1$ ← (so we weakened the condition ← being an inj. resolution)

Claim Any acyclic resolution can be used to compute sheaf cohomology, i.e. $H^n(X, F) = \text{Cohomology of chain complex } \Gamma(X, J^0) \rightarrow \Gamma(X, J^1) \rightarrow \dots$

Pf Trick "break down into SES and take LES":

Let $C_1 = \text{Coker}(F \rightarrow J_0) \cong \text{Im}(J_0 \rightarrow J_1)$ so \exists natural monomorph. $C_1 \hookrightarrow J_1$
 $C_{n+1} = \text{Coker}(C_n \rightarrow J_n) \cong \text{Im}(J_n \rightarrow J_{n+1})$ " " $C_{n+1} \hookrightarrow J_{n+1}$
 $0 \rightarrow F \rightarrow J_0 \rightarrow C_1 \rightarrow 0$
 $0 \rightarrow C_1 \rightarrow J_1 \rightarrow C_2 \rightarrow 0$ exact, and $0 \rightarrow F \rightarrow J_0 \rightarrow J_1 \rightarrow J_2 \rightarrow J_3 \rightarrow \dots$
 $0 \rightarrow C_n \rightarrow J_n \rightarrow C_{n+1} \rightarrow 0$...

Technical Lemma $0 \rightarrow F \rightarrow I \rightarrow G \rightarrow 0$ SES $\Rightarrow H^n(F) \cong H^{n-1}(G) \forall n \geq 2$
 (only uses LES in H^*) with $H^n(I) = 0 \forall n \geq 1$
 $H^1(F) \cong \text{Coker}(H^0 I \rightarrow H^0 G)$
 $H^0(F) \cong H^0 I \oplus H^0 G \rightarrow H^0 I \rightarrow H^0 G \rightarrow H^1(I) \rightarrow H^1(G) \rightarrow H^2(F) \rightarrow H^2(I) \rightarrow \dots \rightarrow 0$
 so surj. so $H^1 F = \text{Coker} \circledast$ so \cong

Finish proof, abbreviate $H^n(F) = H^n(X, F)$, $\Gamma(F) = \Gamma(X, F)$:

$$H^n(F) \cong H^{n-1}(C_1) \cong H^{n-2}(C_2) \cong \dots \cong H^1(C_{n-1}) \cong \text{Coker}(H^0(J_{n-1}) \rightarrow H^0(C_n))$$

$$\Gamma \text{ left exact} \dots \rightarrow \Gamma(J_{n-1}) \xrightarrow{\alpha_n} \Gamma(J_n) \xrightarrow{\alpha_n} \Gamma(J_{n+1}) \rightarrow \dots$$

$$\downarrow \text{exactness of:} \quad \downarrow \text{in} \quad \downarrow \text{in} \quad \downarrow \text{in}$$

$$0 \rightarrow \Gamma(C_n) \rightarrow \Gamma(J_n) \rightarrow \Gamma(C_{n+1}) \rightarrow \Gamma(C_n) \rightarrow \Gamma(C_{n+1})$$

$$\text{hence } \text{Ker } \rho_n = \text{Im } \text{in}_n \quad \text{Ker } \alpha_n / \text{Im } \alpha_{n-1} = \text{Ker } \rho_n / \text{Im } \text{in}_{n-1} = \text{Im } \text{in}_n / \text{Im } \text{in}_{n-1} \cong \Gamma(C_n) / \text{Im } \rho_{n-1} = \text{Coker } \rho_{n-1} = H^n(F).$$

Non-examinable:

Rmk For a left-exact functor $f: A \rightarrow B$ of abelian cats, a resolution $0 \rightarrow M \rightarrow I^\bullet$ is f -acyclic if $R^n(f(I^\bullet)) = 0 \forall n \geq 1$. Similarly for right-exact functors, for $P \rightarrow M \rightarrow 0$ says $L_n(g(P_n)) = 0 \forall n \geq 1$.
Fact Injective resolutions are acyclic resolutions for left exact functors
 Projective " " " right " "

9.3 Čech cohomology vs sheaf cohomology

Theorem X separated, quasi-compact scheme. Suppose $H^n: \text{QCoh}(X) \rightarrow \mathcal{A}$ are functors s.t.

- i) $H^0(X, F) = \Gamma(X, F)$. ← $\in \text{QCoh}(X)$ by Sec. 7.4 Rmk
- ii) $\varphi: U \hookrightarrow X \Rightarrow H^n(X, \varphi_* F) = 0 \forall n \geq 1, \forall F \in \text{QCoh}(U)$. holds for Čech cohomology since $H^n(X, \varphi_* F) = H^n(\varphi^{-1} X, F) = H^n(U, F) = 0, n \geq 1$ affine
- iii) SES induces a LES on H^* $H^n(X, \varphi_* F) = H^n(\varphi^{-1} U, F)$

Then $H^* \cong \check{H}^*$

Pf $X = \cup U_i$ finite affine open covers (use X quasi-compact)
 U_i affine since X separated (using ordered I)

Notice that the Čech complex

$$\check{C}^n = \prod_{|I|=n} F(U_I) = \prod_{|I|=n} \Gamma(U_I, F) = \prod_{|I|=n} \Gamma(X, \varphi_{I,*}(F|_{U_I})) = \Gamma(X, \prod_{|I|=n} \varphi_{I,*}(F|_{U_I}))$$

where $\varphi_I: U_I \hookrightarrow X$ is the inclusion

$\Rightarrow \check{C}^n = \Gamma(X, J^n)$ and have sequence $0 \rightarrow F \rightarrow J^0 \rightarrow J^1 \rightarrow \dots$ ← call this J^n
 By Sec. 9.2 it is enough to check this is an acyclic resolution, since then $H^n(X, F) = H^n(\check{C}^{\bullet}(X, F)) = H^n(X, F)$
 other maps are defined on any open $V \subseteq X$ by the Čech maps $F \rightarrow \varphi_{I,*}(F|_{U_I})$ differential on V for cover $\{U_i\}$

By (ii): $H^n(X, \varphi_{I,*}(F|_{U_I})) = 0 \forall n \geq 1$

$\prod_{|I|=n}$ is a finite product so \cong finite \oplus .

So $H^n(X, J^k) = 0 \forall n \geq 1$ follows by induction by following Trick:

10. Qcoh(P^n), GRADED MODULES, PROJ(R) (Non-examinable chapter)

10.1 Graded modules and Qcoh(P^n)

Def graded ring means a ring R s.t.

$$R = R_0 \oplus R_1 \oplus R_2 \oplus \dots \text{ as abelian groups (so a graded abelian gp graded by } \mathbb{N})$$

$$R_i \cdot R_j \subseteq R_{i+j} \leftarrow \text{link } R_0 \subseteq R \text{ subring since } R_0 \cdot R_0 \subseteq R_0$$

The elements of R_n are called homogeneous elements of degree n

Graded module means R-mod M s.t.

$$M = \dots \oplus M_{-2} \oplus M_{-1} \oplus M_0 \oplus M_1 \oplus M_2 \oplus \dots \text{ as abelian groups (so graded by } \mathbb{Z})$$

$R_i \cdot M_j \subseteq M_{i+j}$ \leftarrow often write M_0 to emphasize \exists grading \circ

A morphism of graded R-mods is R-mod hom $M \rightarrow N$, with $\varphi(M_n) \subseteq N_n \quad \forall n$

From now on: $R = k[x_0, \dots, x_n]$ R_m = homogeneous polys of deg = m (so $R_0 = k$)

$$X = \mathbb{P}^n_k = A_0 \cup A_1 \cup \dots \cup A_n \text{ for}$$

$$A_i := \text{Spec } k \left[\begin{matrix} x_0 \\ \vdots \\ x_i \\ \vdots \\ x_n \end{matrix} \right] = \text{Spec}(k[x_0, \dots, x_n]_{(x_i)})$$

means take 0-th graded part so $P(x_0, \dots, x_n) \leftarrow \text{poly}$
 $x_i \cdot \text{deg}(P)$

$$A_i \cap A_j = \text{Spec } k \left[\begin{matrix} x_0 \\ \vdots \\ x_i \\ \vdots \\ x_j \\ \vdots \\ x_n \end{matrix} \right] = \text{Spec}(k[x_0, \dots, x_n]_{(x_i, x_j)})$$

omit x_i

Claim \exists exact, full & faithful functor

$$\{\text{graded R-mods}\} \longrightarrow \text{Qcoh}(\mathbb{P}^n)$$

$M \longmapsto \tilde{M}$

Pf Let $M_i = (M_{x_i})_{0 \leq k \leq i}$ and $M_{ij} = (M_{x_i, x_j})_0$

Define $\tilde{M}|_{A_i} = \tilde{M}_i$ these glue since $\tilde{M}_i|_{A_i \cap A_j} \cong \tilde{M}_{ij} \cong \tilde{M}_j|_{A_i \cap A_j}$ \leftarrow using $\left(\begin{matrix} M_{x_i, 0} \\ \vdots \\ M_{x_i, i} \end{matrix} \right)_{x_i} \cong (M_{x_i, i})_0$

Exactness is a local condition, so it holds since it holds in affine case.

Full & faithful: $\text{Hom}(\tilde{M}|_{A_i}, \tilde{N}|_{A_i}) = \text{Hom}(\tilde{M}_i, \tilde{N}_i) = \text{Hom}_{(R_{x_i})_0\text{-mods}}((M_{x_i, 0}, \dots, (N_{x_i, 0}))$

this reduces the problem to an exercise in graded R-mods. (omitted here) \square

Warning Not an equivalence of categories because:

$$\text{HWK4 if } M_n = N_n \text{ for } n > N \text{ then } \tilde{M} \cong \tilde{N}$$

Fact If work with graded R-mods "modulo" identifying those which would give rise to "same" \tilde{M} , then get equivalence of categories. So work with $\{\text{R-mods } M\} / \{R\text{-mods } M : \tilde{M} = 0\}$.

For $X = \mathbb{P}^n$, $\tilde{M} = 0 \Leftrightarrow M$ is locally nilpotent, i.e. $\forall m \in M, \exists d \text{ s.t. } x_i^d \cdot m = 0 \quad \forall i$.

If M is f.g., then $\tilde{M} = 0 \Leftrightarrow M$ is finite dim v.s./k.

In reverse direction: $\{\text{graded R-mods}\} \longleftarrow \text{Qcoh}(\mathbb{P}^n)$

$\Gamma_*(F) := \bigoplus_{d \geq 0} \Gamma(\mathbb{P}^n, F(d)) \longleftarrow F$ where $F(d) = F \otimes_{\mathcal{O}_{\mathbb{P}^n}} \mathcal{O}(d) \leftarrow$ called twisting

Trick If $G_1, G_2 \in \text{Qcoh } X$, $H^i(X, G_i) = 0 \quad \forall n \geq 1 \Rightarrow G_1 \oplus G_2$ also, since:

$$0 \rightarrow G_1 \rightarrow G_1 \oplus G_2 \rightarrow G_2 \rightarrow 0 \text{ SES} \Rightarrow \text{take LES get } H^i(X, G_1 \oplus G_2) = 0, \quad n \geq 1 \checkmark$$

$0 \rightarrow F \rightarrow J^* \text{ exact} \Leftrightarrow \text{exact on stalks} \Leftrightarrow 0 \rightarrow \Gamma(U, F) \rightarrow \Gamma(U, J^*) \text{ exact } \forall \text{ affine open } U$

$$0 \rightarrow \Gamma(U, F) \rightarrow \Gamma(U, J_0) \rightarrow \Gamma(U, J_1) \rightarrow \dots$$

exact since $H^i(U, F) = 0$ for $n \geq 1$ \leftarrow for cover $U = U_i$
since U affine, using sec. 8.3 \square

Cor X separated, Noetherian \Rightarrow sheaf cohomology $H^i(X, F) \cong \check{H}^i(X, F) \quad \forall F \in \text{Qcoh}(X)$

\leftarrow Non-examinable

Pf Sheaf cohomology $H^i(X, F) =$ cohomology of $\Gamma(X, I^0) \rightarrow \Gamma(X, I^1) \rightarrow \dots$ for $F \rightarrow I^*$ any injective resolution.

Check the conditions of Theorem:

i) $\Gamma(X, \cdot)$ left exact $\Rightarrow H^0(X, F) \cong \Gamma(X, F) \leftarrow$ general consequence see 9.1, or explicitly: $0 \rightarrow \Gamma(X, F) \rightarrow \Gamma(X, I^0) \rightarrow \Gamma(X, I^1)$

iii) Lemma in 9.1 proves \exists LES exact, so $\text{im } \Gamma$ is ker of J which is H^0

ii) by the Theorem below. \square

Theorem R Noeth., $F \in \text{Qcoh}(\text{Spec } R) \Rightarrow H^n(\text{Spec } R, F) = 0 \quad \forall n \geq 1$

Non-examinable proof ideas The cleanest proof is to build machinery:

1) A sheaf F is flasque if all restrictions $F(U) \rightarrow F(V)$ are surjective.

2) \forall flasque F on a top. space X , have $H^n(X, F) = 0 \quad \forall n \geq 1$ (Hartshorne III.3.4)

3) \forall injective R-module I , and R Noeth. $\Rightarrow \bar{I}$ on spec R is flasque (Hartshorne III.3.4)

Cor Flasque resolutions are acyclic by (2), so can be used to compute $H^n(X, F)$ by 9.2

Pf Thm $F \in \tilde{M} \rightarrow \bar{I}^*$ exact, each \bar{I}^n flasque, so can use this to compute $H^n(X, F)$ by Cor $\Rightarrow H^n(X, \tilde{M}) = H^n(\Gamma(X, \bar{I}^*)) = H^n(\bar{I}^*) \stackrel{\cong}{=} H^n(I^*) \stackrel{\cong}{=} H^n(I^*)$ (in deg = 0 get M , and $H^0(X, \tilde{M}) = \tilde{H}^0(X, M)$)

Rmk Injective \mathcal{O}_X -mods are flasque (Hartshorne III.2.4)

Cultural Rmk For any scheme X and sheaf F of abelian groups have $H^0(X, F) \cong H^0(X, F) = \Gamma(X, F)$ but also in degree 1: $\exists H^1(X, F) \cong H^1(X, F)$. So for example $\text{Pic}(X) \cong \check{H}^1(X, \mathcal{O}^*) \cong H^1(X, \mathcal{O}^*)$ in 8.7.

9.4 Product on sheaf cohomology

(Non-examinable section) (X, \mathcal{O}_X) any ringed space

Fact \exists product $H^i(X, F) \times H^j(X, G) \rightarrow H^{i+j}(X, F \otimes_{\mathcal{O}_X} G)$

idea $0 \rightarrow F \rightarrow I^* \Rightarrow 0 \rightarrow F \otimes G \rightarrow I^* \otimes J^* \rightarrow I^* \otimes J^* \rightarrow 0$

Unfortunately not a resolution \leftarrow non-exact

bi-complex (compare 8.4) with maps $d \otimes \text{id}$, $\text{id} \otimes d$

then take total complex: total degree is sum of degrees \leftarrow non-exact

need $I^* J^*$ to be "pure acyclic resolutions" to ensure this \leftarrow non-exact

is resolution. Then given any inj. res. $F \otimes G \rightarrow K^*$, the identity $F \otimes G \xrightarrow{\text{id}} F \otimes G$ extends to $I^* \otimes J^* \rightarrow K^*$.

Taking $\Gamma(X, \cdot)$ yields the result. (see key idea under the Fact in 9.1)

Fact $F \cong \Gamma_*(F)$

When we mod out by the M with $\tilde{M} = 0$ as in \otimes , this functor together with the functor of claim define an equivalence of cats.

$\text{Coh}(\mathbb{P}^n)$ corresponds to the f.g. graded modules under the equivalence.

Rmk The preferred representative of M in the quotient \otimes is the saturation $\Gamma_*(\tilde{M})$ of M . Call M a saturated module if $M \cong \Gamma_*(\tilde{M})$. (Think of this like a sheafification)

Def $M[d]$ new graded R -mod with $M[d]_i = M_{d+i}$

Example $\mathcal{L} = \widehat{R[d]}$ on $\mathbb{P}^n \leftarrow (S_0 k[x_0, \dots, x_n])[d]$

$\mathcal{L}(A_i) = (R[d]_{x_i})_0 = x_i^d k[\frac{x_0}{x_i}, \dots, \frac{x_n}{x_i}] = x_i^d \cdot (R_{x_i})_0$

line bundle with $\alpha_{ij} = (x_i/x_j)^d$. Hence $\mathcal{L} = \mathcal{O}(d)$.

$(\mathbb{P}^n)_{A_{ij}} \xrightarrow{\cong} \mathcal{L}_{A_{ij}} \xrightarrow{\cong} \mathcal{O}_{\mathbb{P}^n}(d)$, $f \mapsto x_i^d f \mapsto x_j^{-d} x_i^d f$

Exercise $\widehat{M[d]} \cong \tilde{M}(d) (= \tilde{M} \otimes_{\mathcal{O}_{\mathbb{P}^n}} \mathcal{O}(d)) \leftarrow (e.g. \widehat{R[d]} = \tilde{R}(d) = \theta \otimes_{\mathcal{O}_0} \mathcal{O}(d) = \mathcal{O}(d))$

Rmk $k[x_0, \dots, x_n] = \bigoplus_{d \geq 0} \Gamma(\mathbb{P}^n, \mathcal{O}(d))$ (but this does not generalise due to above issue about cats)

The construction of \tilde{M} is so similar to the $\text{Spec } R$ case of \tilde{M} , because \exists analogue of $\text{Spec } R: \text{Proj } R$

10.2 Proj(R) and QCoh(Proj R)

$\text{Proj}(R) = \{ \text{graded prime ideals } I \subseteq R \text{ not containing the irrelevant ideal} \}$

R any graded ring means $I = \bigoplus_{n \geq 0} (I \cap R_n)$

$\mathbb{V}(I) = \{ p \in \text{Proj } R : p \supseteq I \}$ define closed sets of Zariski topology

f homogeneous of degree $> 0 \Rightarrow D_f = \text{Proj } R \setminus \mathbb{V}(f) = \{ p \in \text{Proj } R : f \notin p \}$ basis of open sets

Warning $\text{Proj } R = \bigcup_{D_f} \Leftrightarrow R \subseteq \sqrt{\text{call } f}$ (example: $(\mathbb{P}^n = D_{x_0} \cup \dots \cup D_{x_n}$ and $(x_0, \dots, x_n) = k[x_0, \dots, x_n] +$

Fact $D_f \cong \text{Spec}((R_f)_0)$ as topological spaces (inverse map: $p_0 \mapsto \bigoplus_{k \geq 0} \{ a_k \in R_k : \frac{a_k}{f^k} \in p_0 \}$)

Sheaf $\mathcal{O} := \mathcal{O}_{\text{Proj}(R)} : \mathcal{O}|_p = \mathcal{O}_{\text{Spec}((R_f)_0)}$ then glue. (on $D_{f_0} = D_f \cap D_{g_0}$ get $\mathcal{O}_{\text{Spec}((R_{f_0 g_0})_0)}$)

Examples

1) $S = R[x_0, \dots, x_n]$ with usual grading $\Rightarrow \text{Proj } R = \mathbb{P}^n_R$ (or $\mathbb{P}^n_{\text{Spec } R}$)

2) $R^{(d)} := \bigoplus_{n \geq 0} R_{d+n}$ then the inclusion $R^{(d)} \rightarrow R$ induces an iso $\text{Proj } R \cong \text{Proj } R^{(d)}$

3) S graded ring generated as an S_0 -algebra by $n+1$ elements $s_0, \dots, s_n \in S_1 \Rightarrow S_0[s_0, \dots, s_n] \xrightarrow{\varphi} S \Rightarrow S \cong S_0[x_0, \dots, x_n] \Rightarrow \text{Proj } S \cong \mathbb{V}(I) \subseteq \mathbb{P}^n_{S_0}$ closed subscheme

Example $k[x, y]^{(2)} = k[x^2, xy, y^2]$ $X \mapsto xz, Y \mapsto xz, Z \mapsto yz$ closed sub scheme of \mathbb{P}^2

$\Rightarrow \mathbb{P}^1 = \text{Proj } k[x, y] \cong \text{Proj } k[x, y]^{(2)} \cong \text{Proj } k[X, Y, Z]/(XZ - Y^2)$ closed sub scheme of \mathbb{P}^2

is the Veronese embedding $\nu_2: \mathbb{P}^1 \hookrightarrow \mathbb{P}^2$. Similarly get $\nu_d: \mathbb{P}^1 \hookrightarrow \mathbb{P}^{\binom{n+d}{d}}$ is the Veronese embedding $\nu_d: \mathbb{P}^1 \hookrightarrow \mathbb{P}^{\binom{n+d}{d}}$ some graded ideal I .

4) Every closed subscheme of $\text{Proj } R$ arises as $\text{Proj}(R/I)$

Fact $R = \bigoplus_{n \geq 0} R_n$ graded ring \Rightarrow get line bundles $\mathcal{O}(d) = \tilde{R}(d)$ on $\text{Proj } R$, and \exists exact, full & faithful functor

$\{ \text{graded } R\text{-mods} \} \rightarrow \text{QCoh}(\text{Proj } R)$

$M \mapsto \tilde{M}$

$\Gamma_*(F) \leftarrow F$

where $\Gamma_d(F) := \Gamma(\text{Proj } R, F(d)) \leftarrow (F(d) = F \otimes_{\mathcal{O}_X} \mathcal{O}(d)$ and $\mathcal{O}_X = \tilde{R}$ on $X = \text{Proj } R$)

again, not an equivalence of cats, but $\Gamma_*(F) \cong F$ and the two functors define an equivalence of cats if we work with saturated graded R -mods ($M_i \cong \Gamma_*(M)$)

Fact If R_0 Noetherian, R generated as R_0 -algebra by finitely many elts $\in R_1$ then $\{ \text{f.g. } R\text{-mods} \} / \{ \text{f.g. torsion } R\text{-mods} \} \rightarrow \text{Coh}(\text{Proj } R)$ is equiv. of cats.

Here "torsion" means $\forall m \in M, \exists n \in \mathbb{N}: (R_+)^n \cdot m = 0$. For M f.g. A -mod: this holds $\Leftrightarrow M_k = 0$ for large k

So \otimes same as working with f.g. R -mods modulo identifying those that "agree" in large degrees.

Exercise M "torsion" $\Rightarrow M_f = 0$ \forall homogeneous $f \in R \Rightarrow \tilde{M}(D_f) = M(D_f) = 0 \Rightarrow \tilde{M} = 0$. (homogeneous localisation of f)

Now assume only R Noeth. graded ring. Exercise Show R_0 Noeth., and R generated as R_0 -alg. by finitely many $f_1, \dots, f_n \in R_1$. Let $d := \text{lcm}(\text{deg } f_i)$. Call homogeneous $m \in M$ irrelevant if $(R_+ \cdot m)_{n \cdot d} = 0$ for all large N . M called irrelevant if all m are irrelevant. Fact \otimes holds if replace "torsion" by "irrelevant".

any ring

(recall $R_0 \hookrightarrow R$ subring)

call this I

$N = \#$ degree d monomials in x_0, \dots, x_n

Note: this tells us $\text{QCoh}(\cdot) \forall$ Proj: variety!

\tilde{M} built by gluing as in 10.1 namely $\tilde{M}(D_f) = M(f)$ is homogeneous localisation of f (so localise at f and take \mathcal{O}_X -graded part) stalk $\tilde{M}_I = M_I$ at the homogeneous localisation $I = \mathcal{O}_X$ -th graded part of M_I

Example: $R = k[x_0, \dots, x_n]$ / I the $x_0, \dots, x_n \in R_1$ generate.

$M \mapsto \tilde{M}$ and quasi-inverse $\Gamma_*(F) \leftarrow F$

So \otimes same as working with f.g. R -mods modulo identifying those that "agree" in large degrees.

Exercise M "torsion" $\Rightarrow M_f = 0$ \forall homogeneous $f \in R \Rightarrow \tilde{M}(D_f) = M(D_f) = 0 \Rightarrow \tilde{M} = 0$.

Now assume only R Noeth. graded ring.

Exercise Show R_0 Noeth., and R generated as R_0 -alg. by finitely many $f_1, \dots, f_n \in R_1$.

Let $d := \text{lcm}(\text{deg } f_i)$. Call homogeneous $m \in M$ irrelevant if $(R_+ \cdot m)_{n \cdot d} = 0$ for all large N . M called irrelevant if all m are irrelevant. Fact \otimes holds if replace "torsion" by "irrelevant".