

C7.5: General Relativity 1

University of Oxford: Part C

Chris Couzens

Office N1.06

Mathematical Institute, University of Oxford

Andrew Wiles Building, Radcliffe Observatory Quarter

Woodstock Road

Oxford, OX2 6GG

Disclaimer: There are almost certainly typos in the notes, if something does not look correct or needs further explanation please let me know.

Please send any comments/corrections to christopher.couzens@maths.ox.ac.uk

Contents

| | | |
|----------|--|-----------|
| 1 | Introduction | 8 |
| 2 | Special relativity and Newtonian gravity | 10 |
| 2.1 | Special relativity | 10 |
| 2.1.1 | Lorentz transformations | 11 |
| 2.1.2 | Causal structure and worldlines | 14 |
| 2.1.3 | Some more formal aspects: vectors, one-forms and tensors | 17 |
| 2.1.4 | Newton's law in special relativity and energy momentum | 23 |
| 2.2 | Newtonian gravity | 25 |
| 2.2.1 | Equivalence Principles | 26 |
| 2.2.2 | Planetary orbits in Newtonian mechanics | 27 |
| 2.3 | Problems with Newtonian gravity and why we need GR | 30 |
| 3 | Differential Geometry | 33 |
| 3.1 | Manifolds | 33 |
| 3.2 | Calculus on manifolds | 46 |
| 3.2.1 | Differentiable maps | 46 |
| 3.2.2 | Tangent Vectors | 47 |
| 3.2.3 | Vector fields | 50 |
| 3.2.4 | One-forms | 50 |
| 3.2.5 | Tensors | 51 |
| 3.2.6 | Induced maps | 53 |
| 3.3 | Flows and Lie derivatives | 55 |
| 3.3.1 | One-parameter group of transformations | 57 |
| 3.3.2 | Lie Derivatives | 59 |
| 3.4 | Differential forms | 63 |
| 3.4.1 | Exterior product | 64 |
| 3.4.2 | Exterior derivative | 65 |
| 3.4.3 | Interior product | 69 |
| 3.4.4 | Integration | 71 |
| 3.5 | Summary | 75 |

| | | |
|----------|--|------------|
| 4 | Riemannian geometry | 77 |
| 4.1 | The metric | 77 |
| 4.1.1 | Riemannian metric | 78 |
| 4.1.2 | Lorentzian manifolds | 80 |
| 4.1.3 | Why is the metric useful? | 81 |
| 4.2 | Connections and curvature | 84 |
| 4.2.1 | Differentiating other tensors | 86 |
| 4.3 | Torsion and curvature | 87 |
| 4.3.1 | Component form of the torsion | 88 |
| 4.3.2 | Component form of the Riemann tensor | 89 |
| 4.3.3 | The Ricci identity | 89 |
| 4.3.4 | Levi–Civita connection | 89 |
| 4.4 | Parallel transport and geodesics | 91 |
| 4.4.1 | Geodesics | 92 |
| 4.4.2 | Normal coordinates | 95 |
| 4.4.3 | Path dependence: Curvature and Torsion | 97 |
| 4.4.4 | Explicit Example of path dependence | 101 |
| 4.4.5 | Geodesic deviation | 105 |
| 4.5 | Riemann tensor and its symmetries | 107 |
| 4.5.1 | Ricci and Einstein tensors | 107 |
| 4.6 | Summary | 108 |
| 5 | Einstein’s equations | 110 |
| 5.1 | The Einstein–Hilbert action | 110 |
| 5.1.1 | Newton’s constant | 113 |
| 5.1.2 | Cosmological constant | 114 |
| 5.1.3 | Higher derivative terms | 115 |
| 5.1.4 | Diffeomorphisms | 115 |
| 5.1.5 | Coupling to matter | 117 |
| 5.2 | Summary | 120 |
| 6 | Schwarzschild solution | 122 |
| 6.1 | The Schwarzschild black hole | 122 |
| 6.1.1 | Birkhoff’s theorem | 123 |
| 6.2 | Geodesics of Schwarzschild | 128 |

| | | |
|----------|---|------------|
| 6.2.1 | Perihelion precession | 132 |
| 6.2.2 | Bending of light | 135 |
| 6.2.3 | Gravitational red shift | 138 |
| 6.3 | Schwarzschild solution as a black hole | 141 |
| 6.3.1 | Kruskal spacetime | 147 |
| 6.4 | Summary | 150 |
| A | Some worked examples on Special Relativity | 153 |
| A.1 | Proper time along an accelerated worldline | 153 |
| A.2 | Null curves in Minkowski space | 154 |
| A.3 | Ladders and barns | 154 |
| B | Euler–Lagrange equations in field theory | 158 |
| B.1 | Classical mechanics and Euler–Lagrange | 158 |
| B.2 | Classical field theory and Euler–Lagrange | 159 |
| C | Solutions to exercises | 161 |
| C.1 | Solutions to chapter 2 | 161 |
| C.1.1 | Exercise 2.1 | 161 |
| C.1.2 | Exercise 2.2 | 163 |
| C.1.3 | Exercise 2.3 | 163 |
| C.1.4 | Exercise 2.4 | 163 |
| C.1.5 | Exercise 2.5 | 164 |
| C.1.6 | Exercise 2.6 | 164 |
| C.1.7 | Exercise 2.7 | 165 |
| C.2 | Solutions to chapter 3 | 165 |
| C.2.1 | Exercise 3.1 | 165 |
| C.2.2 | Exercise 3.2 | 166 |
| C.2.3 | Exercise 3.3 | 166 |
| C.2.4 | Exercise 3.4 | 166 |
| C.2.5 | Exercise 3.5 | 166 |
| C.2.6 | Exercise 3.6 | 167 |
| C.2.7 | Exercise 3.7 | 168 |
| C.2.8 | Exercise 3.8 | 170 |
| C.2.9 | Exercise 3.9 | 170 |

| | |
|--------------------------------------|-----|
| C.2.10 Exercise 3.10 | 171 |
| C.3 Solutions to chapter 4 | 172 |
| C.3.1 Exercise 4.1 | 172 |
| C.3.2 Exercise 4.2 | 172 |
| C.3.3 Exercise 4.3 | 173 |

Recommended books and resources

There are a large variety of good textbooks and lecture notes on general relativity. This course borrows from a number of them, in various different places. An assortment of textbooks that have been used in writing these notes are:

- Wald, General Relativity

A very thorough introduction to the subject.

- Weinberg, Gravitation and cosmology
- Carroll, An introduction to general relativity, spacetime and geometry.

Aimed more at particle physicists. We will follow this in the cosmology section and borrow bits for elsewhere.

- Hartle, Gravity, an introduction to Einstein's general relativity
- Misner, Thorne and Wheeler, Gravitation

It is a very big book.

- Nakahara, Geometry, Topology and Physics

An excellent book for learning about geometry and topology and will be useful for the differential geometry section of the notes.

There are also a number of useful lecture notes online. In particular:

- Joe Kier's lecture notes from 2020
- David Tong's lecture notes
- Sean Carroll's lecture notes
- Harvey Reall's lecture notes

Conventions

- We will use the god-given signature convention of mostly plus $(-, +, +, +)$. This may differ with the convention you have used in other courses, especially field theory courses. This convention is preferable when thinking about geometry as it gives positive spatial distances. For quantum field theory the other convention is preferable since it ensures that energies and frequencies are positive. You may map between the two conventions through *Wick rotation*, essentially allowing the coordinates to become complex.
- Spacetime indices will be taken to be greek letters from the middle of the alphabet: μ, ν, ρ, \dots and run over 0, 1, 2, 3. Latin indices i, j, k, \dots run over the spatial directions and take values 1, 2, 3.
- We employ Einstein summation convention, repeated indices are summed over, unless otherwise stated.
- We work in units where the speed of light c is set to 1. Occasionally it is instructive to reintroduce c which can be done by dimensional analysis.
- The Minkowski metric will be denoted by $\eta_{\mu\nu} = \text{diagonal}(-1, 1, 1, 1)_{\mu\nu}$.
- After introducing curvature we will take the metric to be $g_{\mu\nu}$ and the determinant will be $\det(g_{\mu\nu}) \equiv g$.

Useful formulae

- The Lagrangian for the geodesic equation of a massive test particle is

$$\mathcal{L}\left(\frac{dx^\mu}{d\lambda}, x^\mu\right) = \sqrt{-g_{\mu\nu}(x) \frac{dx^\mu}{d\lambda} \frac{dx^\nu}{d\lambda}},$$

with λ an arbitrary parameter along the worldline.

- The geodesic equation for a massive particle is

$$\frac{d^2 x^\mu}{d\tau^2} + \Gamma^\mu_{\nu\rho} \frac{dx^\nu}{d\tau} \frac{dx^\rho}{d\tau} = 0, \quad g_{\mu\nu}(x) \frac{dx^\nu}{d\tau} \frac{dx^\rho}{d\tau} = -1,$$

where τ is the proper time. For light, the first equation takes the same form just replacing τ with an affine parameter. The second is modified by $-1 \rightarrow 0$.

- The Christoffel symbols (Levi–Civita connection) are

$$\Gamma^\mu{}_{\nu\rho} = \frac{1}{2}g^{\mu\sigma} \left(\partial_\nu g_{\sigma\rho} + \partial_\rho g_{\sigma\nu} - \partial_\sigma g_{\nu\rho} \right).$$

- The Riemann tensor is

$$R^\mu{}_{\nu\rho\sigma} = \partial_\rho \Gamma^\mu{}_{\nu\sigma} - \partial_\sigma \Gamma^\mu{}_{\nu\rho} + \Gamma^\mu{}_{\rho\lambda} \Gamma^\lambda{}_{\nu\sigma} - \Gamma^\mu{}_{\sigma\lambda} \Gamma^\lambda{}_{\nu\rho}.$$

- Symmetries

$$R_{\mu\nu\rho\sigma} = -R_{\mu\nu\sigma\rho},$$

$$R_{\mu\nu\rho\sigma} = R_{\rho\sigma\mu\nu}.$$

- Bianchi identity 1

$$R^\mu{}_{\nu\rho\sigma} + R^\mu{}_{\rho\sigma\nu} + R^\mu{}_{\sigma\nu\rho} = 0.$$

- Bianchi Identity 2

$$\nabla_\mu R^\sigma{}_{\lambda\nu\rho} + \nabla_\nu R^\sigma{}_{\lambda\rho\mu} + \nabla_\rho R^\sigma{}_{\lambda\mu\nu} = 0.$$

- Ricci tensor

$$R_{\mu\nu} = R^\rho{}_{\mu\rho\nu}$$

- Ricci scalar

$$R = R_{\mu\nu} g^{\mu\nu}.$$

- Einstein tensor

$$G^{\mu\nu} = R^{\mu\nu} - \frac{1}{2}Rg^{\mu\nu}.$$

- Einstein–Hilbert action plus cosmological constant,

$$S = \frac{1}{16\pi G} \int d^4x \sqrt{-g} (R + \Lambda).$$

- Under a variation $g_{\mu\nu} \rightarrow g_{\mu\nu} + \delta g_{\mu\nu}$ we have

$$\delta g^{\mu\nu} = -g^{\mu\rho} g^{\nu\sigma} \delta g_{\rho\sigma},$$

$$\delta g = g g^{\mu\nu} \delta g_{\mu\nu},$$

$$\delta R_{\mu\nu} = \nabla_\rho \delta \Gamma^\rho{}_{\mu\nu} - \nabla_\mu \delta \Gamma^\rho{}_{\rho\nu}.$$

1 Introduction

Gravity is one of the four¹ fundamental forces alongside electromagnetism, the strong nuclear force and the weak nuclear force. Of these forces gravity is by far the weakest force, the ratio of the gravitational force to electric force acting on an electron is 10^{-36} .² Despite this gravity plays a dominant role in shaping the large scale structure of the universe. This is because the strong and weak forces have a very short range, while, though electromagnetism is a long range force, it is both attractive and repulsive. For bodies of macroscopic dimensions the repulsion of like charges is approximately balanced by the attraction of opposite charges, you don't see big clumps of highly charged bodies in nature. On the other hand, gravity is only an attractive force, thus for sufficiently large bodies the gravitational field of the sum of all its constituents adds up to become the dominant force.

The leading candidate for a theory of gravity for some time was Newton's theory of gravitation. This however, is a non-relativistic theory of gravity and therefore is incompatible with special relativity: it is not invariant under Lorentz transformations. One can see this with a little thought experiment. What would happen if the sun suddenly disappeared? For 8 minutes, the time it takes for light to travel from the sun to Earth, we would be completely oblivious. This is because special relativity tells us that no signal can travel faster than light: the Earth must continue on its orbit for these 8 minutes, after which, it is flung out of the solar system leading to almost certain death for all life on Earth. However, Newton's theory of gravity acts instantaneously, we would be flung out of the solar system immediately. In Newton's theory, the force on one mass depends on the location of the other mass at the same time.

Einstein's breakthrough led to a conceptual revolution in the way that we view spacetime and the role that differential geometry plays. The fact that objects with the same initial conditions follow the same curve, independent of their mass, hints that the curve that is followed is a property of the geometry of spacetime rather than a force acting on the body. General relativity (GR) understands gravity as the curvature of spacetime and the trajectories within spacetime as geodesics on this curved space. Or as John Wheeler once said, "*Mass tells space how to curve, while curved space tells matter how to move*".

The aim of this course is to introduce you to General relativity and by the end of it to

¹One could add *currently known to physics* at this point for safety.

²You can see this very clearly by holding two magnets together, gravity is not strong enough to pull one magnet to the floor. A current research direction, known as the *weak gravity conjecture*, conjectures that in any quantum theory of gravity, the strength of gravity is weaker relative to any other gauge force [1].

allow you to perform calculations. Among other topics we will see how gravity bends light, the corrections to the motion of the planets and a black hole. This is a large topic and we will therefore omit many interesting directions, but this will lay the foundation for further study and for the follow up course General Relativity II.

The notes are organised as follows. We begin by reviewing special relativity and Newtonian gravity in section 2. To understand general relativity properly we need to understand the underlying geometry of spacetime. This requires knowledge of the sophisticated tools of *differential geometry* to describe curved spacetime, which we will study in sections 3 and 4. With these new tools we are finally in a position to introduce Einstein's equations and physics in curved spacetime in section 5. The Schwarzschild solution is the go to solution of general relativity and we will use it as a testing ground for studying many interesting topics in GR including black holes, the motion of the planets and the bending of light in section 6.

2 Special relativity and Newtonian gravity

We begin with a whirlwind exploration of special relativity. This section is by no means meant to be an introduction to special relativity, more a refresher on the subject and to emphasise the pertinent points. We will immediately assume that the reader is familiar with the Einstein summation convention, i.e. that repeated indices, one upper and one lower, are summed over. For readers in need of a more thorough introduction there are a number of excellent texts to consult. For example the notes by [Joe Minahan](#), the book ‘Special Relativity: An Introduction with 200 Problems and Solutions’ by Michael Tsamparlis and Bernard Schutz’s book, ‘A First Course in General Relativity’.

By the end of the 18th century two areas of physics that were in conflict had emerged: Newtonian mechanics and Electromagnetism. Newtonian mechanics has a notion of absolute time and the equations of motion are invariant under Galilean coordinate transformations. The transformation law between two coordinate systems, the latter moving at a uniform speed v in the x direction of the former, is

$$(t', x', y', z') = (t, x - vt, y, z). \quad (2.1)$$

Galilean transformations imply that the speed of light should change in different reference frames moving with respect to each other. This is incompatible with Maxwell’s equations describing electromagnetism where the speed of light is fixed. A resolution to this problem was proposed by conjecturing a preferred frame, the frame of the physical medium in which light propagates, called the *Ether*. The speed of light in any other rest frame would then be modified by the Newtonian addition of velocities. An experiment by Michelson and Morley in 1887 to detect the Ether failed, the speed of light does not obey the Newtonian law of addition of velocities. Either Newtonian mechanics or Maxwell’s equations required modification.

2.1 Special relativity

Einstein gave the resolution to this problem in 1905 with the introduction of special relativity. In special relativity a key role is played by the so-called *inertial reference frame*. Such a frame satisfies three key properties:

1. There is a universal time coordinate which can be synchronised everywhere in the inertial frame.
2. The spatial slices are Euclidean space, satisfying the usual Euclidean axioms.

3. A body with no external forces will move with constant velocity within the frame with respect to the clocks and measuring sticks of the frame.

We can construct an infinite number of such frames and we need to understand how to map between them. This is where Einstein's two postulates for special relativity are needed:

1. The laws of physics in any inertial reference frame are identical.
2. The speed of light in a vacuum in two different reference frames does not change and is given by ³ $c = 299792458 \text{ m s}^{-1}$.

For the second postulate it is important to note the word vacuum. The speed of light through different mediums is less than c , this is why we have refraction when light travels through glass for example. In [2] they even managed to stop light propagating temporarily before letting it propagate again.

The group of spacetime coordinate transformations which map inertial reference frames to inertial reference frames are the Lorentz transformations. One may then rephrase the principle of special relativity to be that the laws of nature are invariant under Lorentz transformations. This requires the abandonment of the Newtonian idea of absolute time. Events which are simultaneous in one inertial reference frame need not be simultaneous in another frame (see problem sheets 0 and 1 for examples to work through).

2.1.1 Lorentz transformations

A Lorentz transformation is a linear transformation from one spacetime coordinate system $x^\mu = (ct, x, y, z)$ to another x'^μ , of the form

$$x'^\mu = \Lambda^\mu{}_\nu x^\nu, \quad (2.2)$$

where Λ is a constant matrix, i.e. spacetime independent, and satisfies

$$\Lambda^\mu{}_\rho \Lambda^\nu{}_\sigma \eta_{\mu\nu} = \eta_{\rho\sigma}. \quad (2.3)$$

The matrix η is the famed Minkowski metric, which in our signature, is given by

$$\eta_{\mu\nu} = \text{diag}(-1, 1, 1, 1)_{\mu\nu} \equiv \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}_{\mu\nu}. \quad (2.4)$$

³There is a nice mnemonic (the number of letters in each word is the value here) to remember this: *We guarantee certainty clearly referring to this light mnemonic.*

The set of matrices satisfying (2.3) is the group $O(1, 3)$.⁴ We could also add in constant shifts of the coordinates, $x'^{\mu} = \Lambda^{\mu}_{\nu} x^{\nu} + a^{\mu}$, with a^{μ} a constant four-vector. This would enhance the Lorentz group to the Poincaré group. For our purposes we will only need to consider the Lorentz group though and so we set $a^{\mu} = 0$ from now on.

From the definition of the Lorentz group it follows that it leaves the line element, (sometimes also called length element, invariant interval)

$$ds^2 = \eta_{\mu\nu} dx^{\mu} dx^{\nu}, \quad (2.5)$$

invariant.⁵ This is merely the proper distance between the two spacetime events: (t, x, y, z) and $(t + dt, x + dx, y + dy, z + dz)$. Here, d stands for an infinitesimal displacement, you also see δ and Δ to mean the same thing.

Aside: The group $O(1, 3)$ described above is sometimes called the *homogeneous Lorentz group*. It admits a proper subgroup defined by imposing

$$\Lambda^0_0 \geq 1, \quad \det \Lambda = 1. \quad (2.6)$$

The proper subgroup restricts to all transformations which can be smoothly joined to the identity. The *improper* Lorentz transformations involve either space inversion $\det \Lambda = -1$, $\Lambda^0_0 \geq 1$, or time reversal $\det \Lambda = 1$, $\Lambda^0_0 \leq 1$. Space and time inversions are known not to be exact symmetries of nature and therefore when we say Lorentz transformation what we really mean is the proper Lorentz transformations.

We now want to understand what types of transformations the proper Lorentz group admits. There is a further subgroup consisting of spatial rotations taking the form:

$$\Lambda^0_0 = 1, \quad \Lambda^0_i = \Lambda^i_0 = 0, \quad \Lambda^i_j = R_{ij}, \quad (2.7)$$

with R an $SO(3)$ matrix: $RR^T = 1$, $\det R = 1$. The remaining transformations are known as *boosts* which mix the space and time directions. Examples of the two types of transformation are⁶

$$\Lambda^{\text{Rotation}} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos \theta & \sin \theta & 0 \\ 0 & -\sin \theta & \cos \theta & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad \Lambda^{\text{Boost}} = \begin{pmatrix} \cosh \phi & -\sinh \phi & 0 & 0 \\ -\sinh \phi & \cosh \phi & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}. \quad (2.8)$$

⁴More generally one could consider $O(p, q)$ which satisfy (2.3) but now with η having $p - 1$'s and $q + 1$'s.

⁵One can show that the Lorentz transformations are the only non-singular coordinate transformations that leave ds^2 invariant. Here non-singular means that both $x'(x)$ and $x(x')$ are well behaved differential functions and thus $\frac{\partial x^{\mu}}{\partial x'^{\nu}}$ has an inverse. When we consider $ds^2 = 0$ there is an enhancement of the symmetry group. You will show this in problem sheet 1.

⁶Note that we only work with the proper Lorentz group.

The first is a rotation in the x, y directions and the second is a boost in the x direction. The rotation parameter is compact $\theta \in [0, 2\pi)$ while the boost parameter, known as the *rapidity* is non-compact $\phi \in (-\infty, \infty)$. Altogether the Lorentz group has six parameters, split evenly between boost and rotations. Rotations do not commute with boosts, thus the Lorentz group is non-abelian.

Rather than considering the matrices Λ in (2.8) it is useful to consider their generators.⁷ These are matrices T^a which satisfy

$$\Lambda = e^{i\theta_a T^a}, \quad (2.9)$$

where θ_a are constant parameters. The generators for the two matrices appearing in (2.8) are

$$T^{\text{Rotation}} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & -i & 0 \\ 0 & i & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad T^{\text{Boost}} = \begin{pmatrix} 0 & i & 0 & 0 \\ i & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad (2.10)$$

with similar expressions for the other generators.

Exercise 2.1: Addition of rapidity

1. Compute the addition of the rapidity under two successive boosts along the x axis.
2. Show that the generators in (2.10) give the matrices in equation (2.8)
3. Compute the commutator of the generators (2.10) of a boost along x and y .
4. Compute the commutator of the generators (2.10) for a boost along x and rotation in the x - y -plane.
5. Compute the commutator of the generators (2.10) of a boost along x and rotation in the y - z -plane.

The interpretation of the rotations is clear from our understanding of Galilean symmetries but what is the interpretation of the boosts? This corresponds to changing coordinates to that of a frame moving with a constant velocity with respect to the first.

Under a boost in the x -direction the transformed coordinates are

$$t' = t \cosh \phi - x \sinh \phi, \quad x' = -t \sinh \phi + x \cosh \phi. \quad (2.11)$$

⁷We are considering the Lie Algebra of the Lie Group.

The point $x' = 0$ is then moving, as viewed from the original frame, with velocity

$$v = \frac{x}{t} = \tanh \phi. \quad (2.12)$$

Motivated by this it is useful to perform the replacement $\phi = \operatorname{arctanh} v$ in the transformations to obtain

$$\begin{aligned} t' &= \gamma(t - vx), \\ x' &= \gamma(x - vt), \end{aligned} \quad \text{with } \gamma = (1 - v^2)^{-1/2}. \quad (2.13)$$

Exercise 2.2: Addition of velocity

Compute the addition of the velocity under two successive boosts along the x axis. You should contrast this with the addition of rapidities.

Understanding these transformation lead to a whole range of interesting phenomena from time dilation to length contraction. *In problem sheet 0 and 1 you will review some of these problems. There will also be some worked examples given in these notes in section A.*

2.1.2 Causal structure and worldlines

The invariance of the proper distance between spacetime events allows us to make the following definition:

Definition 1 (Timelike, Null, Spacelike) *The interval between two spacetime events x^μ and $x^\mu + dx^\mu$ is*

$$\begin{aligned} \text{Timelike separated if } ds^2 &< 0, \\ \text{Null or Lightlike separated if } ds^2 &= 0, \\ \text{Spacelike separated if } ds^2 &> 0. \end{aligned} \quad (2.14)$$

This can be visualised by studying a spacetime diagram. Focussing only on the t, x coordinates and suppressing the y, z directions we may associate to every point p in spacetime a light cone, see figure 1. For simplicity let the point p be the origin, this is of course just a shift in the coordinate, remember our a^μ 's that we removed earlier. Then the path light takes passing through the point p is given by $x = \pm t$. These straight lines moving at the speed of light define the light cone. Points lying on the cone are *null/lightlike separated* from our point p . We can divide the light cone into two cones, the future and past light cones. Those points residing in the light cones are *timelike separated* while those outside the light cones are *spacelike separated*. To fully complete the diagram into a cone we should add in the




Figure 1: The lightcone diagram. The pink areas are time-like separated from the point at the centre, while points in the blue area are space-like separated. The dotted lines are light-like separated.

y, z directions, from which we can see the conical structure. In figure 1 we have sketched the light cone in the t, x -plane

There is a slight subtlety about this diagram which we should point out. If we perform a boost of our coordinates in the x -direction and map this onto the light-cone it seems that the light-cone begins to close up, see figure 1. This is a Euclidean viewpoint however, whereas this properly resides in Lorentzian space. From a Lorentzian viewpoint the axes remain orthogonal and it is not difficult to see that the lines $t = \pm x$ get mapped precisely to the lines $t' = \pm x'$. Thus, we see that our classification in terms of timelike, null and spacelike separated points

is also preserved under Lorentz transformations from this viewpoint.

Note that when we are looking at the distance between points we are drawing straight lines between the points and computing the distance of this line. This works in Minkowski space but when we introduce curved spacetime this is no longer correct since the notion of a straight line in this sense ceases to exist. We should really think about computing the tangent of a path between the two points, for straight lines the tangent lies along the curve but for more general curves this is no longer true. Let us define a path γ in our spacetime and parametrise it by $\lambda \in [\lambda_1, \lambda_2]$. We may choose local inertial coordinates so that we may define our curve as $x^\mu(\lambda)$, note that λ is arbitrary and need not be identified with the time coordinate. We can then compute the tangent to the curve, $\frac{dx^\mu}{d\lambda}$. For a spacelike curve, that is one which for every infinitesimal interval it is spacelike, we define the (change of) *proper length* to be

$$\Delta s = \int_{\lambda_1}^{\lambda_2} \sqrt{\eta_{\mu\nu} \frac{dx^\mu}{d\lambda} \frac{dx^\nu}{d\lambda}} d\lambda. \quad (2.15)$$

There is no such analogue along a null curve since $ds^2 = 0$. For a timelike curve, one in which the infinitesimal intervals are all time like, we define the *proper time* τ via:

$$\Delta\tau = \int_{\lambda_1}^{\lambda_2} \sqrt{-\eta_{\mu\nu} \frac{dx^\mu}{d\lambda} \frac{dx^\nu}{d\lambda}} d\lambda, \quad (2.16)$$

where this is understood to be the change in proper time after following the curve from $x^\mu(\lambda_1)$ to $x^\mu(\lambda_2)$. One may worry that this depends on the parametrisation of the curve, however it is a simple exercise (exercise 2.3) to show that the above is independent of the parametrisation of the curve.

Exercise 2.3: Reparameterisation invariance

Show that the definition of the change in proper time in equation 2.16 is independent of the parametrisation of the curve, i.e. under the change $\lambda \rightarrow \lambda(\sigma)$.

The proper time is useful because of the *clock postulate*.

Clock Postulate *An accurate clock moving along a timelike worldline measures the proper time along the worldline.*

This point of view makes the “twin paradox” and similar puzzles clear. Two worldlines which have two intersections at different events will have proper times which measure their respective proper times, however these numbers in general will be different since the paths are different.

It is often convenient to parametrise a timelike curve by the proper time since for a timelike path we have:

$$\eta_{\mu\nu} \frac{dx^\mu(\tau)}{d\tau} \frac{dx^\nu(\tau)}{d\tau} = -1. \quad (2.17)$$

Exercise 2.4: Parametrising a timelike curve

Show that it is always possible to find a parametrisation of a timelike curve so that it satisfies (2.17), and moreover that it is unique up to constant shifts.

Massive paths Let us now consider the worldlines of massive particles, these follow timelike paths. From our earlier discussion we will use the proper time as the parameter along the path with the path starting at $\tau = 0$ for simplicity. The tangent vector is known as the *four-velocity* U^μ :

$$U^\mu = \frac{dx^\mu}{d\tau}. \quad (2.18)$$

This is automatically normalised, $\eta_{\mu\nu} U^\mu U^\nu = -1$ since we parametrised the curve using the proper time! We may define the *energy-momentum four-vector* as

$$p^\mu = mU^\mu, \quad (2.19)$$

with m the mass of the particle. The mass is a fixed quantity independent of inertial frame, this is what you may have been used to calling the rest mass. The energy is defined simply to be p^0 , and as one component of a four-vector is not invariant under Lorentz transformations. Note that in the particle's rest frame we have $p^0 = m$ (recall $c = 1$) and so this is the celebrated $E = mc^2$. Note that the energy in the rest frame is the norm of the energy momentum four vector. In a general frame we have

$$E^2 - p^i p_i = m^2, \quad (2.20)$$

which is the general version of Einstein's famous formula and is Lorentz invariant and hence holds in any inertial frame.

2.1.3 Some more formal aspects: vectors, one-forms and tensors

Vectors and Vector fields To probe the structure of Minkowski space it is necessary to introduce the concepts of vectors and tensors. We will give a full treatment of this subject later in section 3 introducing only the necessary notation for the moment. You may be used to thinking of a vector as something stretching from one point to another and which can be freely moved around. In relativity this is no longer true and so we must be more careful by what we mean by a vector.

To each point p in spacetime we associate the set of all possible vectors located at that point. A useful class of vectors to consider are the tangent vectors to curves going through the point p . In an n -dimensional spacetime there are n independent such vectors and they span a *vector space* called the *tangent space* at p , and denoted by T_p .

Definition 2 (Vector space) *A vector space, V is a set of elements, $v \in V$, which we call vectors, that may be added together or multiplied by elements of a field, \mathbb{F} (e.g real or complex numbers). The operations of the vector addition and multiplication must satisfy the following axioms:*

1. *Associativity*

$$u + (v + w) = (u + v) + w, \quad \forall u, v, w, \in V, \quad (2.21)$$

2. *Commutativity*

$$u + v = v + u \quad \forall u, v, \in V, \quad (2.22)$$

3. *Identity element. There exists a $0 \in V$ such that*

$$v + 0 = v, \quad \forall v \in V, \quad (2.23)$$

4. *Inverse elements. For every $v \in V$ there exists a $-v \in V$ such that*

$$v + (-v) = 0 = (-v) + v, \quad (2.24)$$

5. *Compatibility of scalar multiplication with field multiplication.*

$$a(bv) = (ab)v, \quad \forall a, b \in \mathbb{F}, \text{ and } v \in V \quad (2.25)$$

6. *Identity element of scalar multiplication. There exists a $1 \in \mathbb{F}$ such that*

$$1v = v, \quad \forall v \in V, \quad (2.26)$$

7. *Distributivity of scalar multiplication.*

$$a(u + v) = au + av, \quad \forall a \in \mathbb{F}, \text{ and } u, v \in V, \quad (2.27)$$

8. *Distributivity of scalar multiplication.*

$$(a + b)v = av + bv, \quad \forall a, b \in \mathbb{F} \text{ and } v \in V \quad (2.28)$$

A vector is a perfectly well-defined geometric object defined at the point p . We may also define a *vector field* to be a set of vectors with exactly one defined at each point in spacetime. The set of all the tangent spaces T_p of a manifold⁸ M is known as the *tangent bundle* $T(M)$. This is a $2n$ -dimensional manifold which is an example of a fiber bundle. *It is important to emphasise that neither the vector nor the vector field transform under Lorentz transformations, they are invariant.*

You may, however, be familiar with the transformation property of the components of a vector. We decompose vectors into components in terms of some basis of the tangent space. Recall that a *basis* is a set of vectors which both spans the vector space and is linearly independent. There are an infinite number of possible bases, but each will have the same number of basis elements, the dimension of the manifold here. Let us imagine that at every point in our n -dimensional space we set up a basis with n vectors \hat{e}_μ . Then any vector V can be expanded in terms of this basis as $V = V^\mu \hat{e}_\mu$. Here V^μ are known as the *components of the vector*. This is sometimes sloppily called a vector or contravariant vector, however this is not correct, V is the vector and V^μ are the components of the vector.⁹

A standard example of a vector in spacetime, and one that will appear frequently, is the tangent to a curve. We can specify a curve by specifying the coordinates in terms of a parameter, $x^\mu(\lambda)$. The tangent vector has components

$$V^\mu = \frac{dx^\mu(\lambda)}{d\lambda}, \quad (2.29)$$

and the vector is

$$V = V^\mu \hat{e}_\mu. \quad (2.30)$$

Under a Lorentz transformation the coordinates transform according to (2.2), and from this we may deduce the transformation of the components of the four-vector V^μ . When the coordinate system is transformed as in (2.2), the components transform as

$$V^\mu \rightarrow V'^\mu = \Lambda^\mu_\nu V^\nu. \quad (2.31)$$

Since the geometric vector itself is invariant; only its components change under a Lorentz transformation, and the parametrisation with λ is unaltered, it follows that the basis vectors transform according to

$$\hat{e}_\mu = \Lambda^\nu_\mu \hat{e}'_\nu. \quad (2.32)$$

⁸We will define a manifold later in section 3.

⁹Despite this we will sometimes call the components the vector, we are however being sloppy and they are just the components of the vector in some basis.

This is just multiplication by the inverse of the Lorentz transformation which transforms the coordinates, therefore

$$\hat{e}'_\mu = \Lambda_\mu{}^\nu \hat{e}_\nu. \quad (2.33)$$

To summarise, we have introduced a set of coordinates labelled by upper indices which transform in a certain way under Lorentz transformations. We then considered vector components with upper indices which transformed in the same way as the coordinates. The basis vectors associated with the coordinate system transformed via the inverse matrix and were labelled by a lower index. These transformations leave invariant the vector, that is summing over the vector components with the basis vectors.

Co-vectors and one-forms Once we have a vector space we can define an associated vector space known as the *dual vector space*. It is usually denoted with an asterisk, so that the dual vector space of the Tangent space T_p is T_p^* . The dual space is the space of all linear maps from the original vector space to the real numbers, so that if $\omega \in T_p^*$ then by the definition of a linear map¹⁰

$$\omega(aV + bW) = a\omega(V) + b\omega(W) \in \mathbb{R}, \quad (2.34)$$

for all $V, W \in T_p$ and $a, b \in \mathbb{R}$. It follows that T_p^* is a vector space itself and since it is finite dimensional its dual vector space is T_p . We may introduce a basis of dual vectors $\hat{\theta}^\mu$ by fixing

$$\hat{\theta}^\mu(\hat{e}_\nu) = \delta_\nu^\mu. \quad (2.35)$$

Every dual vector can be written in components in terms of this basis as

$$\omega = \omega_\mu \hat{\theta}^\mu. \quad (2.36)$$

Typically one refers to the elements of T_p as *contravariant* four vectors and elements of T_p^* as *covariant vectors*, or even *one-forms*, (a name that will make more sense after we have introduced differential geometry in section 3). The set of all cotangent spaces over M is called the *cotangent bundle* $T^*(M)$. The action of a dual vector field on a vector field is no longer a single number but a *scalar*, depending on the spacetime position. A scalar has no indices and is left invariant under Lorentz transformations.

The component notation is useful when considering the action of a dual vector on a vector:

$$\omega(V) = \omega_\mu V^\nu \hat{\theta}^\mu(\hat{e}_\nu) = \omega_\mu V^\nu \delta_\nu^\mu = \omega_\mu V^\mu. \quad (2.37)$$

¹⁰We require a linear map, $f : V \rightarrow W$ to be additive $f(u+v) = f(u) + f(v)$ for all $u, v \in V$ and homogeneous of degree 1 so that $f(cu) = cf(u)$ for all $u \in V$ and $c \in \mathbb{F}$.

Since the action of a co-vector on a vector is a constant it is invariant under Lorentz transformations and we must have

$$\omega'_\mu V'^\mu = \omega'_\mu \Lambda^\mu{}_\sigma V^\sigma \equiv \omega_\mu V^\mu. \quad (2.38)$$

If we were dealing with vector and co-vector fields it would be a *scalar*. It is from here that we can obtain the transformation of the dual vector: a *covariant* four-vector is a quantity which transforms as

$$\omega_\mu \rightarrow \omega'_\mu = \Lambda_\mu{}^\nu \omega_\nu, \quad (2.39)$$

where

$$\Lambda_\mu{}^\nu \equiv \eta_{\mu\rho} \eta^{\nu\sigma} \Lambda^\rho{}_\sigma, \quad (2.40)$$

with $\eta^{\mu\nu}$ the inverse of $\eta_{\mu\nu}$, which are numerically the same.

Exercise 2.5: The inverse of the Lorentz transformation

Using the properties of the Lorentz transformation show that $\Lambda_\mu{}^\nu$ is the inverse of $\Lambda^\mu{}_\nu$.

The simplest example of a co-vector is the *gradient* of a scalar function,

$$d\phi = \frac{\partial\phi}{\partial x^\mu} \hat{\theta}^\mu. \quad (2.41)$$

You can see that the components indeed transform in the correct way.

Observe that the metric η can be used to raise and lower indices. This allows us to transform a contravariant vector into a co-vector and vice-versa. The metric then acts as a map from $T_p(M) \rightarrow T_p^*(M)$, while the inverse metric acts as a map from $T_p^*(M) \rightarrow T_p(M)$.

Exercise 2.6: Raising and lowering indices

- Show that $\eta_{\mu\nu} V^\nu$ transforms as the components of a co-vector if V^ν transforms as the components of a vector.
- Show that $\eta^{\mu\nu} V_\nu$ transforms as the components of a vector if V_ν transforms as the components of a co-vector.

Note that because of the map between contravariant and covariant vectors via the Minkowski metric we can define an *inner product* on two vectors as

$$\eta(V, W) = \eta_{\mu\nu} V^\mu W^\nu. \quad (2.42)$$

Two vectors whose inner product vanishes are called *orthogonal*. Since it is a scalar the dot product is left invariant under Lorentz transformations and therefore orthogonality is basis

and frame independent. We can define the *norm* of a vector to be the inner product with itself. Unlike in Euclidean geometry this is not positive definite, instead

$$\text{if } \eta_{\mu\nu}V^\mu V^\nu \text{ is } \begin{cases} < 0, & V^\mu \text{ is timelike,} \\ = 0, & V^\mu \text{ is lightlike or null,} \\ > 0, & V^\mu \text{ is spacelike,} \end{cases} \quad (2.43)$$

This is the more mathematical definition of these concepts from our earlier discussion.

Tensors One may extend the notion of a vector and co-vector to a *tensor*. A tensor of type (rank) (k, l) , is a multilinear map from a collection of dual vectors and vectors to \mathbb{R} :

$$T : T_p^* \times \dots \times T_p^* \times T_p \times \dots \times T_p \rightarrow \mathbb{R}, \quad (2.44)$$

for example a scalar is a tensor of rank $(0,0)$, a co-vector is a rank $(0,1)$ tensor and a contravariant vector is a tensor of rank $(1,0)$. The space of all tensors of a fixed rank (k, l) forms a vector space. To construct a basis for this space it is useful to define the *tensor product* \otimes . If T is a (k, l) -tensor and S a (m, n) -tensor then $T \otimes S$ is a $(k + m, l + n)$ tensor defined to be

$$\begin{aligned} T \otimes S(\omega^{(1)}, \dots, \omega^{(k)}, \dots, \omega^{(k+m)}, V^{(1)}, \dots, V^{(l)}, \dots, V^{(l+n)}) \\ = T(\omega^{(1)}, \dots, \omega^{(k)}, V^{(1)}, \dots, V^{(l)})S(\omega^{(k+1)}, \dots, \omega^{(k+m)}, V^{(l+1)}, \dots, V^{(l+n)}). \end{aligned} \quad (2.45)$$

Let T be a tensor of rank (k, l) , then under a Lorentz transformation its components transform as

$$T'^{\mu_1 \dots \mu_k}_{\nu_1 \dots \nu_l} = \Lambda^{\mu_1}_{\mu'_1} \dots \Lambda^{\mu_k}_{\mu'_k} \Lambda_{\nu_1}^{\nu'_1} \dots \Lambda_{\nu_l}^{\nu'_l} T^{\mu'_1 \dots \mu'_k}_{\nu'_1 \dots \nu'_l}. \quad (2.46)$$

One can use tensors to construct additional tensors either by taking linear combinations of tensors with the same upper and lower indices, direct products, contraction, or differentiation. Note that the order of the indices of a tensor matters.

Some tensors that will appear regularly are: the metric which is a $(0, 2)$ tensor, with the inverse being a $(2, 0)$ tensor, the *Kronecker delta* δ^μ_ν which is a $(1, 1)$ tensor, and finally the Levi-Civita tensor which is a $(0, 4)$ tensor. Not only can the metric be used to raise and lower indices of a tensor, it can also be used to contract indices. *Contraction* takes a (k, l) tensor to a $(k - 1, l - 1)$ tensor by

$$T^{\mu\nu\rho}_{\mu\sigma} \equiv S^{\nu\rho}_\sigma. \quad (2.47)$$

2.1.4 Newton's law in special relativity and energy momentum

We now want the special relativity version of Newton's second law. The requirement that it be tensorial puts some stringent constraints on the possible form, we must introduce a force four-vector f^μ satisfying

$$f^\mu = m \frac{d^2}{d\tau^2} x^\mu(\tau) = \frac{d}{d\tau} p^\mu(\tau). \quad (2.48)$$

For electromagnetism and the Lorentz force law ($f = q(E + v \times B)$) we find

$$f^\mu = q U^\nu F_\nu{}^\mu, \quad (2.49)$$

where F is the field strength of the electromagnetism gauge field and q is the charge of the particle and U^μ its four-velocity.

Although p^μ provides a complete description of the energy and momentum of a particle for extended systems it is necessary to go further and define the *energy-momentum tensor*, or *stress tensor*, $T^{\mu\nu}$. This is a symmetric $(2, 0)$ tensor which tells us all we need to know about the energy like aspects of a system: energy density, pressure, stress etc.. Consider a *fluid*. This is a continuum of matter described macroscopic quantities such as temperature, pressure, entropy, viscosity, etc. We will work with *perfect fluids* which are completely characterised by their pressure and density. This in particular means that they are isotropic (same in every direction) in the rest frame.

To understand this let us first consider *dust*. This is a collection of particles which are at rest with respect to each other, as a perfect fluid they have zero pressure. Since all the particles have an equal velocity in any fixed inertial frame we can imagine a four-velocity field $U^\mu(x)$ defined over all spacetime. We can define the *number-flux four-vector*

$$N^\mu = n U^\mu, \quad (2.50)$$

where n is the number density of the particles as measured in their rest frame. Then N^0 is the number density of particles as measured in any other frame, while N^i is the flux of particles in the i 'th direction. Let us imagine each of the particles have the same mass m . Then in the rest frame the energy density of the dust is given by

$$\rho = nm. \quad (2.51)$$

This completely specifies the dust, however this only measures the energy density in the rest frame, how do we measure it in other frames? Notice that both n and m are 0-components of four-vectors in their rest frame: $N^\mu = (n, 0, 0, 0)$ and $p^\mu = (m, 0, 0, 0)$. Therefore ρ is the

$\mu = 0, \nu = 0$ component of the tensor $p \otimes N$ as measured in the rest frame. We are therefore lead to define the energy momentum tensor for dust

$$T_{\text{dust}}^{\mu\nu} = p^\mu N^\nu = nmU^\mu U^\nu = \rho U^\mu U^\nu, \quad (2.52)$$

where ρ is the energy density as measured in the rest frame.

We can now consider other perfect fluids. The key point is the isotropic in the rest frame property which implies that the energy momentum tensor must take a diagonal form in the rest frame, since there cannot be a net flux of momentum in an orthogonal direction. Moreover the spacelike components must all be equal $T^{11} = T^{22} = T^{33}$, there are only two independent components. We will take the two independent parameters to be the energy density ρ and the pressure p (note that p is also used for momentum but will always come with a superscript or subscript). In the rest frame the energy momentum tensor takes the form

$$T^{\mu\nu} = \begin{pmatrix} \rho & 0 & 0 & 0 \\ 0 & p & 0 & 0 \\ 0 & 0 & p & 0 \\ 0 & 0 & 0 & p \end{pmatrix}. \quad (2.53)$$

We want a formula which is good in any frame and therefore we want to write this in terms of tensors. For dust we had $T^{\mu\nu} = \rho U^\mu U^\nu$, so we may guess that there should be $(\rho + p)U^\mu U^\nu$, which gives $\rho + p$ in the 00 component and zero elsewhere in the rest frame. To include the remainder we should find something which is of the form $p \text{diag}(-1, 1, 1, 1)$ this is of course given by the Minkowski metric! The general form of the energy momentum tensor for a perfect fluid is

$$T^{\mu\nu} = (\rho + p)U^\mu U^\nu + p\eta^{\mu\nu}. \quad (2.54)$$

This will be important when we consider the cosmology section of the course.

Some worked examples of the concepts of special relativity can be found in appendix [A](#).

Key Points

- The laws of physics are invariant under Lorentz transformations.
- The speed of light is the same in all inertial frames.
- Spacetime intervals classify events into timelike, null and spacelike.
- The quantity $ds^2 = \eta_{\mu\nu} dx^\mu dx^\nu$ is invariant under Lorentz transformations.
- The components of a tensor transform according to equation (2.46) under a Lorentz transformation.

2.2 Newtonian gravity

Above we have reviewed special relativity, now we want to understand why special relativity is incompatible with Newtonian gravity, in particular we will see that it is not Lorentz invariant. To do this we can cast Newtonian gravity in terms of a field theory. The force acting on a particle of mass m is

$$F = -m\nabla\Phi(t, \vec{x}), \quad (2.55)$$

where the gravitational field $\Phi(t, \vec{x})$ is determined by the surrounding matter distribution $\rho(t, \vec{x})$, through

$$\nabla^2\Phi(t, \vec{x}) = 4\pi G_N\rho(t, \vec{x}), \quad (2.56)$$

where G_N is Newton's constant with approximate value

$$G_N \sim 6.67 \times 10^{-11} \text{m}^3 \text{kg}^{-1} \text{s}^{-2}. \quad (2.57)$$

This is simply a rewriting into field theory language of the inverse square law of Newton. For example if there is a mass M concentrated at a single point at $(t, \vec{0})$, then the mass density is

$$\rho(t, \vec{x}) = M\delta^{(3)}(\vec{x}), \quad (2.58)$$

which gives the gravitational field,

$$\Phi(\vec{x}) = -\frac{GM}{r}, \quad r^2 = \vec{x} \cdot \vec{x}. \quad (2.59)$$

This can be extended to more complicated matter distributions, either summing up contributions from the location of point-like particles or more generally by using the Greens function for the Laplacian and the mass density

$$\Phi(\vec{x}) = -\int d^3x' \frac{G\rho(\vec{x}')}{|\vec{x} - \vec{x}'|}. \quad (2.60)$$

Exercise 2.7: Newton's theorem

Newton's theorem states that the gravitational field outside of a spherically symmetric mass distribution depends only on its total mass. Show this by using (2.55), (2.56) and Gauss' theorem.

We can now insert the gravitational force law into Newton's second law of motion $F = ma$. At this point one should ask oneself whether the inertial mass appearing in Newton's second

law is the same as the one appearing in the gravitational force law (2.55), there is no reason that they need to be the same. Application of Newton's second law gives

$$\vec{a} = -\frac{m_G}{m_i} \nabla \Phi, \quad (2.61)$$

with \vec{a} the acceleration. Starting with work by Galileo, Christaan Huygens and more recent experimental data we have seen that $m_i = m_G$ to an accuracy of 10^{-13} . This is known as the *weak equivalence principle*. In the Newtonian theory this appears as an isolated unexplained fact, however it is this experimental fact that underlies general relativity. Since all bodies with the same initial conditions fall along the same curve regardless of their composition, we can interpret that curve to be a property of the geometry of the spacetime not of a force acting on the body.

2.2.1 Equivalence Principles

The *Weak equivalence principle* was one of the starting points for the development of GR. It is motivated by thought experiments using Newtonian gravity. The exact equality of $m_i = m_G$ is one version of the weak equivalence principle. Newtonian gravity gives no explanation for why this should be true. A theory of gravity should be able to explain this. Another way to formulate the weak equivalence principle is:

The weak equivalence principle

The trajectory of a freely falling test body depends only on its initial position and initial velocity and is independent of the composition of the body.

A consequence of the weak equivalence principle is that it is not possible to tell the difference between constant acceleration and a constant gravitational field. Suppose that you are in a closed box and consider the two situations 1) you are on earth, 2) you are in a spaceship undergoing constant acceleration.¹¹ Within Newtonian mechanics there is no local experiment that you can perform which distinguishes the two. One of the important words is *local*. You can use tidal forces to distinguish between the two. Roughly if you drop two masses on Earth they will ever so slightly come together because the direction gravity acts on them is slightly different, they are pulled to the centre of the Earth. On a spaceship this is not the case and they fall down never getting closer together. This however, is a non-local experiment, you need to watch the masses fall for a while and for a distance. Note that it is not the mutual gravitational force between the two test bodies that is bringing them together, rather it the

¹¹Another version of this is 1) the box is in free fall 2) you are floating in deep space. Again there is no local experiment that you can conduct to tell the difference.

fact that they experience a non-uniform gravitational force. This is measured by tidal forces and we will see the mathematics behind this in section 4.4.5.

This motivated Einstein's equivalence principle:

Equivalence Principle

1. The weak equivalence principle is valid.
2. In a local inertial frame the results of all non-gravitational experiments will be indistinguishable from the results of the same experiments in an inertial frame in Minkowski spacetime.

The weak equivalence principle implies that 2) is valid for test bodies. The fact that test bodies which include ordinary matter which is held together by the three other forces, gives evidence that the electromagnetic and nuclear forces also obey 2).

Implications The equivalence principle has many implications. One is that it implies that light is bent in a gravitational field. Consider a uniform gravitational field and a freely falling lab. Inside the lab the Einstein equivalence principle says that light rays must move on straight lines. But a straight line with respect to the lab corresponds to a curved path with respect to a frame at rest relative to the Earth. The effect is small so we cannot demonstrate this in the classroom but with more sensitive equipment in a lab this has been experimentally verified.

2.2.2 Planetary orbits in Newtonian mechanics

We now have all the technology we need to study the orbits of the planets within Newtonian mechanics. Let us set up a coordinate system where the massive body of mass M is at $r = 0$ and the planet of mass m is a distance r from that point. The Lagrangian describing the system is

$$\mathcal{L} = \frac{m}{2}(\dot{x}(t)^2 + \dot{y}(t)^2 + \dot{z}(t)^2) - V(r), \quad V(r) = -\frac{mMG}{r}. \quad (2.62)$$

To make this more tractable it is useful to change coordinates to polar coordinates rather than Cartesian coordinates:

$$x = r \sin \theta \cos \phi, \quad y = r \sin \theta \sin \phi, \quad z = r \cos \theta. \quad (2.63)$$

The Lagrangian becomes

$$\mathcal{L} = \frac{m}{2} \left(\dot{r}^2 + r^2 (\dot{\theta}^2 + \sin^2 \theta \dot{\phi}^2) \right) + \frac{mMG}{r}. \quad (2.64)$$

We can now compute the equations of motion via the Euler–Lagrange equations: we find

$$\begin{aligned}\ddot{r} - r(\dot{\theta}^2 + \sin^2 \theta \dot{\phi}^2) + \frac{MG}{r^2} &= 0, \\ \frac{d}{dt}(r^2 \dot{\theta}) - r^2 \sin \theta \cos \theta \dot{\phi}^2 &= 0, \\ \frac{d}{dt}(r^2 \sin^2 \theta \dot{\phi}) &= 0.\end{aligned}\tag{2.65}$$

First let us consider the $\dot{\theta}$ equation. If we kick the particle off in the $\theta = \frac{\pi}{2}$ plane with $\dot{\theta} = 0$ then it will remain in that plane. We will make this choice from now on. The coordinate ϕ is an ignorable coordinate since it does not appear explicitly in the Lagrangian. Recall that for every ignorable coordinate there is an associated conserved charge, in this case it will be the angular momentum. We may define

$$l = r^2 \dot{\phi},\tag{2.66}$$

which is conserved. We have now solved the last two equations of (2.65) and only the first remains. Then we have

$$\ddot{r} - \frac{l^2}{r^3} + \frac{MG}{r^2} = 0.\tag{2.67}$$

To proceed further it is useful to note that there is one more conserved quantity, the Energy of the system. This follows since the Lagrangian is explicitly time independent, thus

$$Em = \frac{m}{2} \left(\dot{r}^2 + r^2 (\dot{\theta}^2 + \sin^2 \theta \dot{\phi}^2) \right) + V(r),\tag{2.68}$$

is conserved. We can now substitute $\dot{\theta}$ and $\dot{\phi}$ into this final condition to obtain an equation for \dot{r} only:

$$E = \frac{1}{2} \dot{r}^2 + \frac{l^2}{2r^2} - \frac{MG}{r} \equiv \frac{1}{2} \dot{r}^2 + V_N(r).\tag{2.69}$$

We can now study the orbits by looking at the Newtonian potential. At large distances the attractive $-r^{-1}$ dominates, while the angular momentum prohibits the particle from getting too close to the origin, see figure 2.

The potential has a minimum when

$$V'(r_*) - \frac{MG}{r_*^2} - \frac{l^2}{r_*^3} = 0 \quad r_* = \frac{l^2}{MG}.\tag{2.70}$$

The planet can happily sit at $r = r_*$ for all time on a circular orbit, note that $E < 0$ in this case. The planet could also oscillate back and forth around the minima. This happens when $E < 0$ so that the planet cannot escape off to infinity. This describes an orbit where the distance to the massive body varies, as you may expect this is the usual elliptic orbit. For




Figure 2: A representative example of the Newtonian potential. There are three interesting behaviours to consider. When $E > 0$ and the motion describes a fly-by coming from infinity and heading back to infinity. For $E = V_N(r_*) < 0$ the motion is a circular orbit. Finally for $0 > E > V_N(r_*)$ we obtain elliptic orbits.

$E > 0$ the motion describes a flyby, the planet gets close to the massive body, never reaching it, before being flung off to infinity. Clearly such a planet would be dead and inhospitable for life.

So far we have discussed the radial motion of the planet, this does not tell us about the full motion however. We consider the orbit trajectory, the flyby motion is not so interesting for us. We now need to solve the angular momentum equations (2.66). To solve the coupled equations we start by employing a change of coordinates

$$u = r^{-1}, \quad (2.71)$$

and then view this as a function of ϕ . This works nicely because

$$\dot{u} = \frac{du}{d\phi} \dot{\phi} = lu^2 \frac{du}{d\phi}, \quad (2.72)$$

where we have used (2.66). We have

$$\dot{r} = -\frac{1}{u^2} \dot{u} = -l \frac{du}{d\phi}. \quad (2.73)$$

The conservation of energy equation (2.69) becomes

$$\left(\frac{du}{d\phi}\right)^2 + \left(u - \frac{GM}{l^2}\right)^2 = \frac{2E}{l^2} + \frac{G^2 M^2}{l^4}. \quad (2.74)$$

This turns out to be straightforward to solve, the solution is

$$u(\phi) = \frac{GM}{l^2}(1 + e \cos \phi). \quad (2.75)$$

In the original radial coordinate we have

$$r(\phi) = \frac{l^2}{GM} \frac{1}{1 + e \cos \phi}. \quad (2.76)$$

This is the equation for a conic section with the eccentricity given by

$$e = \sqrt{1 + \frac{2El^2}{G^2M^2}}. \quad (2.77)$$

The shape of the orbit depends on the eccentricity. Motion with $E > 0$ is not in a bounded orbit, tracing out a hyperbola for $e > 1$ and a parabola for $e = 1$. Objects in orbit have $e < 1$ with elliptical orbits. An important thing to note about this solution is that the orbit does not *precess*, its closest approach to the origin, known as the *perihelion*¹² is always at the same point it never moves and nor does the furthest point of the orbit, the known as the *aphelion*. This disagrees with observations of Mercury's orbit and is the first observational discrepancy of Newtonian gravity.¹³ We will compare this result with the result from general relativity in section 6.2.1.

2.3 Problems with Newtonian gravity and why we need GR

Newton's theory of gravitation is successful in explaining the motions of the moon and planets. Some irregularities in the orbit of Uranus remained unexplained until the irregularities were used independently by John Couch Adams and Jean Joseph Le Verrier in 1846, to predict the existence and position of Neptune. There were still issues with predictions from Newtonian gravity and experimental data however. The precession of the perihelion of Mercury was one such problem. It was shown to be out by $43''/\text{century}$ ¹⁴, recall that in the section above we showed that the perihelion does not precess in Newtonian gravity. We will see later how GR

¹²Strictly this is for the closest approach to the sun. *Helios* is the word for the sun in greek, while *peri* means around.

¹³To perform a more accurate computation one should also take into account the effect of the gravitation fields of the other planets. This is notoriously difficult since one has to study a multi-body problem. Instead, what one can do is imagine that the other planets form a shell of mass along their orbit, acting equally. One can then evaluate the force due to this mass shell. This approximation works if one considers the problem over a long enough time, but is poor if taken for a short time scale. Since planets closer to the sun have quicker orbits over a long enough time this approximation will give a reasonable result and we can avoid trying to solve this multi-body computation.

¹⁴The " stands for arcseconds, with $3600 \text{ arcseconds}(=3600'')$ in a degree.

corrects this. A more obvious (and mathematical) problem arose after Einstein's work on special relativity in 1905. Newtonian gravity is incompatible with special relativity. A body can, in principle, be accelerated to a speed greater than the speed of light. Moreover, effects are instantaneous in Newtonian gravity clearly this is not allowed in special relativity where the speed of light gives an upper bound on the transfer of information.

Despite Newtonian gravities' failings it is sufficient for studying a large range of phenomena. To understand when a relativistic theory is needed let us consider a circular orbit around a star of mass M . The speed of the planet is easily computed by equating the centripetal force with the gravitational force giving,

$$\frac{v^2}{r} = \frac{GM}{r^2}. \quad (2.78)$$

Relativistic effects become important when $v \sim c$ and therefore the dimensionless parameter which governs corrections to Newtonian gravity is

$$\frac{GM}{rc^2}. \quad (2.79)$$

There is a convenient length scale which one can construct from a mass and the fundamental constants known as the *Schwarzschild radius*,¹⁵

$$r_s = \frac{2GM}{c^2}. \quad (2.80)$$

Relativistic corrections to gravity are then necessary when $r_S \sim 2r$. By this measure the earth is not a relativistic system $r_S \sim 10^{-2}m$ and the corrections on the surface of the Earth are of the order 10^{-8} . For satellites in orbit this is even smaller $\sim 10^{-9}$ however for GPS satellites clocks with such high precision are needed that this effect can be seen and if GR was not taken into account would stop working very soon. The sun has $r_S \sim 3\text{km}$ and for Mercury the corrections are of order 10^{-7} , clearly very small but over a century the precession of Mercury's perihelion adds up to the previously quoted $43''$.

General relativity is the theory that replaces both Newtonian gravity and special relativity. However, general relativity is not the final theory of gravity, one eventually needs a theory of quantum gravity. General relativity breaks down for very extreme phenomena where quantum effects become important, e.g. the Big Bang and inside black holes. If one views gravity as a classical field theory and attempts to quantise it one finds that it is perturbatively non-renormalizable (if you do the QFT courses these words will reappear). Essentially this

¹⁵We will see this appear later when we consider the Schwarzschild solution in section 6.

means that to obtain sensible observable results we must absorb infinities in computations by introducing new parameters. For a renormalizable theory we need to introduce only a finite number of these new parameters but for a non-renormalizable theory we need to introduce an infinite number, rendering the theory unable to give meaningful predictions. A candidate theory for quantum gravity, but no means the only candidate, is string theory. We still do not know what quantum gravity really is! We should emphasise that a theory of quantum gravity is only needed for these extreme phenomena and so for the large part general relativity is sufficient.

3 Differential Geometry

The study of general relativity is the study of curved spacetime and so to make progress we need to learn some differential geometry. Our discussion will not be all-encompassing, there will be both topics and proofs that we omit. Instead we will build up all the necessary mathematical structure, that we will need, in a logical order. As we proceed, many of the objects that we will introduce may already be familiar to you; they will however take a different guise in places. This section will have different levels of mathematical rigour, in some sections we will be good mathematicians defining everything while in other places we will give more physics inspired definitions. For a more formal course one should consult the Manifolds and Riemannian geometry courses.

This section closely follows the excellent book by Nakahara [3] with some more physics explanations and examples.

3.1 Manifolds

Before we begin to define a manifold we need to define a topological space. *Do not worry about these possibly scary sounding words, they are actually simple once the fear has cleared.*

Definition 3 (Topological space) *Let X be any set and $\mathcal{T} = \{U_i | i \in I\}$ denote a certain collection of subsets of X . The pair (X, \mathcal{T}) is called a topological space if \mathcal{T} satisfies*

1. *Both the set X and the empty set \emptyset are open subsets: $X \in \mathcal{T}$ and $\emptyset \in \mathcal{T}$.*
2. *If J is any, possibly infinite, sub-collection of I , then the family $\{U_j | j \in J\}$ satisfies $\cup_{j \in J} U_j \in \mathcal{T}$.*
3. *If K is any finite sub-collection of I then the set $\{U_k | k \in K\}$ satisfies $\cap_{k \in K} U_k \in \mathcal{T}$.*

Sometimes X alone is called a topological space, i.e. without associating to it a topology, here we mean that a topology is associated. The sets U_i are called *open sets* (we may sometimes refer to them as coordinate patches, the reason why will become obvious later) and \mathcal{T} gives a *topology* to X . So when we talk about the open sets, what we really mean is *does this set appear in the topology?*. Therefore what we call an open set depends on what topology we have given the space.

Physical Explanation

We have a set of objects/elements X , for example the people in the lecture room. A topology gives us a way to group the different objects in X with some requirements. Let us now group people in the lecture room into subsets U_i . For example one subset could be

the set of all people who were born on a Thursday another could be those who woke up before 7am this morning and so forth. Now we can define a topology \mathcal{T} by taking a choice of these different subsets, this is the $i \in I$. This space needs to satisfy some properties though.

1. Both the empty set (nothingness) and the set X (everything) must be in \mathcal{T} . These sets are then the trivial ones. I pick none of the room occupants or all of them.
2. If we take any subset of our J of the open sets U_i then the combination (union) of them all is one of the open sets U_i in our topology. Essentially what this is saying is that if we call all the people in the room who are in at least one of the subsets defined by the J open sets then this also has to be an open set.
3. If we take a subset K and intersect them all then the intersection (elements common to all sets) is also an open set U_i in our topology. That is we take some subset K of the open sets and ask who is in all of them. The remainder needs to also be an open set in the topology.

The key point is that a topology specifies which subsets are open.

Example 3.1: Topologies

- a) If X is a set and \mathcal{T} a collection of all subsets of X then this is a topological space, and is known as the *discrete topology*.
- b) Let X be a set and take $\mathcal{T} = \{\emptyset, X\}$. This is then a topological space and the topology is known as the *trivial topology*. While the discrete topology is too stringent, this topology is too trivial.
- c) Take $X = \mathbb{R}$. All open subsets (a, b) (a, b may be $\mp\infty$ respectively) and their unions define a topology known as the *usual topology*.

Exercise 3.1: Usual topology vs discrete topology

Consider the usual topology on \mathbb{R} and show that if we allow for an infinite number of open sets in condition 3 for the definition of a topological space, then the usual topology reduces to the discrete topology.

Definition 4 (Metric) A metric $d : X \times X \rightarrow \mathbb{R}$ is a function that for any $x, y, z \in X$ satisfies:

1. $d(x, y) = d(y, x)$,
2. $d(x, y) \geq 0$ with equality iff $x = y$,
3. $d(x, y) + d(y, z) \geq d(x, z)$.

If X is endowed with a metric then X is made a topological space whose open sets are given by open discs

$$U_\epsilon(x) = \{y \in X \mid d(x, y) < \epsilon\}, \quad (3.1)$$

and all possible unions. The topology \mathcal{T} is called the *metric topology* determined by d .

Physical Explanation

A metric in this abstract sense does exactly what we would expect of a metric: it gives us a way of understanding which elements are “close” together. This seems a bit strange if we take a finite set but we can define such a thing. Let us take a set with two elements: $\{x, y\}$. We take the discrete topology on the space. We can then define a metric via:

$$g(x, x) = 0 = g(y, y), \quad g(x, y) = g(y, x) = 5, \quad (3.2)$$

This then satisfies the axioms of a metric above.

Definition 5 (Neighbourhood) *Suppose \mathcal{T} gives a topology to X . Then N is a neighbourhood of the point $x \in X$ if N is a subset of X and N contains at least one open set U_i which contains x . Note that there is no requirement for N to be open, in the case where it is open it is called an open neighbourhood.*

Physical Explanation

We want to have a notion of which elements are “close together”. Let us pick an element x in X that we are interested in. Our usual intuition of close together does not really work with a finite group of elements. Instead we say that some subset N of X is a neighbourhood (i.e. “close by”) of x if the element x appears in any of the open subsets making up N .

Definition 6 (Hausdorff space) *A topological space (X, \mathcal{T}) is a Hausdorff space if for an arbitrary pair of distinct points $x, y \in X$, there always exists neighbourhoods U_x and U_y such that $U_x \cap U_y = \emptyset$.*

Example 3.2: A non-Hausdorff example

Let $X = \{A, B, C, D\}$ define the sets

$$U_0 = \emptyset, \quad U_1 = \{A\}, \quad U_2 = \{A, B\}, \quad U_3 = \{A, B, C, D\}. \quad (3.3)$$

Then the topology $\mathcal{T} = \{U_0, U_1, U_2, U_3\}$ makes X a topological space but it is not Hausdorff. First note that both the empty set and X are in the topology \mathcal{T} , satisfying point (1) of the definition of a topological space. Note that the union of these sets is within \mathcal{T} thereby satisfying point (2). Finally the intersection of any of the sets is within \mathcal{T} and therefore it is a topological space. To see why it is not Hausdorff it suffices to show that we can pick two points which have no open sets in which one of the points is in and that the intersection of these open sets is not the empty set. There are a few choices we could make but an obvious one is C, D . They both appear in only one open set and therefore the space cannot be Hausdorff.

Most examples in physics that one encounters are Hausdorff spaces. We will *assume* this is the case throughout in this course since it is a form of apotropaic magic: it protects us from evil (=bad things occurring).

Definition 7 (Continuous) *Let X and Y be topological spaces. A map $f : X \rightarrow Y$ is continuous if the inverse image of an open set in Y is an open set in X .*

Physical Explanation

We have two spaces and we want to have a good way of mapping between them. We want to impose some nice properties on this map, the one we are interested in here is it being continuous.

Note that a continuous function does *not* need to map an open set in X to an open set in Y . For example take $f : \mathbb{R} \rightarrow \mathbb{R}$ with $f(x) = x^2$ and the usual topology. This is a continuous function that would fail this since $(-\epsilon, \epsilon)$ gets mapped to the closed set $[0, \epsilon^2)$.

Definition 8 (Closed, closure, interior, boundary) *Let (X, \mathcal{T}) be a topological space.*

- *A subset A of X is closed if its complement $X - A \in \mathcal{T}$ in X is an open set.*
- *The closure of the subset A is the smallest closed set that contains A and is denoted by \bar{A} .*
- *The interior of A is the largest open subset of A and is denoted by A° .*

- The boundary $b(A)$ of A is the complement of A° in \bar{A} : $b(A) = \bar{A} - A^\circ$.

An open set is always disjoint from its boundary while a closed set always contains its boundary. To make this a little more clear let us consider a concrete example.

Example 3.3: Open and closed sets

- Let us consider \mathbb{R}^2 , the two-dimensional plane, with the metric topology and let A be the following open set, $A = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 < 1\}$. Then the closure of A is

$$\bar{A} = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 \leq 1\}. \quad (3.4)$$

The interior of A is itself $A^\circ = A$. The boundary is then the complement of A° in \bar{A} , thus

$$b(A) = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 = 1\}. \quad (3.5)$$

It therefore agrees with our usual understanding of these concepts.

- Consider the set $X = \{a, b, c, d, e, f\}$ with topology

$$\mathcal{T} = \{X, \emptyset, \{a\}, \{c, d\}, \{a, c, d\}, \{b, c, d, e, f\}\}. \quad (3.6)$$

By definition the sets, $\{a\}, \{c, d\}, \{a, c, d\}, \{b, c, d, e, f\}$ are all open in (X, \mathcal{T}) .

Now let us consider the following sets and whether they are open or closed (or both). First it is useful to work out the closed sets in our topology, they are $X - \mathcal{T}$ and thus we have $\{X, \emptyset, \{b, c, d, e, f\}, \{a, b, e, f\}, \{b, e, f\}, \{a\}\}$.

1. $\{a\}$. This is both open, since it appears in \mathcal{T} and closed since $X - \{a\} = \{b, c, d, e, f\} \in \mathcal{T}$. We can see that its closure is itself since it is closed and its interior is itself since it is open.
2. $\{b, c\}$. This is not open since it does not appear in \mathcal{T} . We have $X - \{b, c\} = \{a, d, e, f\}$ which does not appear in \mathcal{T} either and is therefore this set is not closed either. The closure of $\{b, c\}$ is the smallest closed sets that contains $\{b, c\}$, and from the set of closed sets above we see that it is $\{b, c, d, e, f\}$. The interior is the largest open subset of A , we see that this is necessarily the empty set \emptyset . The boundary is therefore $\{b, c, d, e, f\}$.
3. $\{c, d\}$. This is open since it appears in \mathcal{T} , but is not closed since $\{a, b, e, f\}$ does not appear in \mathcal{T} . The closure is $\{b, c, d, e, f\}$, while the interior is itself. The boundary is therefore $\{b, e, f\}$.

4. $\{a, b, e, f\}$. This is not open since it does not appear in \mathcal{T} , it is closed however since $X - \{a, b, e, f\} = \{c, d\}$ does appear in \mathcal{T} . The closure is itself since it is a closed set while the interior is $\{a\}$. Therefore the boundary is $\{b, e, f\}$.

For completeness we give the following two definitions, though we will not call on them much:

Definition 9 (Covering) Let (X, \mathcal{T}) be a topological space. A family $\{A_i\}$ of subsets of X is called a covering of X if

$$\bigcup_{i \in I} A_i = X. \quad (3.7)$$

If all the A_i happen to be the open sets of the topology \mathcal{T} then the covering is called an open covering.

Definition 10 (Compact) Consider a set X and all possible coverings of X . The set X is compact if for every open covering $\{U_i | i \in I\}$ there exists a finite subset J of I such that $\{U_j | j \in J\}$ is also a covering of X .

Theorem 1 Let X be a subset of \mathbb{R}^n , then X is compact if and only if it is closed and bounded.

Definition 11 (Connected, Arcwise connected, Simply connected) Let X be a topological space.

- i) X is connected if it cannot be written as $X = X_1 \cup X_2$ where X_1 and X_2 are both open and $X_1 \cap X_2 = \emptyset$. Otherwise X is called disconnected.
- ii) A topological space is called arcwise connected if for any points $x, y \in X$ there exists a continuous map $f : [0, 1] \rightarrow X$ such that $f(0) = x$ and $f(1) = y$. Only in a few pathological cases is arcwise connectedness not equivalent to connectedness.
- iii) A loop in a topological space X is a continuous map $f : [0, 1] \rightarrow X$ such that $f(0) = f(1)$. If every loop in X can be continuously shrunk to a point, X is called simply connected.

Some simple examples are:

Example 3.4: Connectedness

- $\mathbb{R}^2 - \mathbb{R}$ is not arcwise connected.
- $\mathbb{R}^2 - \{0\}$ is arcwise connected but not simply connected.

- $\mathbb{R}^3 - \{0\}$ is arcwise connected and simply connected.
- The n -dimensional torus is arcwise connected but not simply connected.

The main purpose of topology is to classify spaces. Suppose we have several figures, we want to be able to say which are equal and which are different, and probably more fundamentally what does being equal or different mean. In topology two figures are equivalent if it is possible to deform them continuously into each other. We therefore construct an equivalence relation under which geometrical objects are classified according to whether it is possible to deform one into the other. Of course these are just words and we should define this more mathematically. To wit let us define a *homeomorphism*

Definition 12 (Homeomorphism) *Let X_1 and X_2 be two topological spaces. A map $f : X_1 \rightarrow X_2$ is a homeomorphism if it is continuous and has an inverse $f^{-1} : X_2 \rightarrow X_1$ which is also continuous. If there exists a homeomorphism between X_1 and X_2 we say that X_1 and X_2 are homeomorphic to each other.*

The classic example of two homeomorphic spaces are a donut and a coffee mug, see [here](#) for a gif of this classic example courtesy of wikipedia.

One would like a quick way to understand whether two spaces are homeomorphic to each other. Even today we cannot fully characterise the equivalence classes between spaces. One modest statement that we can make is that if two spaces have different *topological invariants* then they are not homeomorphic to each other. A topological invariant is conserved under homeomorphisms. It may be a number such as the number of connected components of the space, an algebraic structure such as a group or a ring which can be constructed from the space, or something like connectedness, compactness or the Hausdorff property. If we knew the complete class of topological invariants we could specify the equivalence classes easily, however so far we only know a partial list. As such even if all the known topological invariants of two spaces coincide these spaces may still not be homeomorphic.

We are now finally in a position to define a *manifold*. An n -dimensional manifold is a space which looks locally like \mathbb{R}^n . Globally it need not be \mathbb{R}^n but we may glue local patches, each of which look like \mathbb{R}^n together to get the full global space. A manifold is then homeomorphic to \mathbb{R}^n locally. The local homeomorphism allows us to give each point on the manifold a set of n numbers called local *coordinates*. If the manifold is not homeomorphic to \mathbb{R}^n then we need to cover it in more than one patch, and so we need to introduce several local coordinates. We will require that the transition functions between these coordinates on

the overlapping region are *smooth*. In this way we can develop the usual notion of calculus on a manifold. Topology is based on continuity, while manifolds is based on smoothness. With that let us begin with our definitions again.

Definition 13 (Differentiable manifold) *M* is an *n*-dimensional differentiable manifold if it satisfies:

1. *M* is a Hausdorff topological space,
2. *M* is provided with a family of pairs $\{(U_i, \varphi_i)\}$;
3. $\{U_i\}$ is a family of open sets which covers *M*: $\cup_i U_i = M$.
4. φ_i is a homeomorphism from U_i onto an open subset U'_i of \mathbb{R}^n ,
5. Given U_i and U_j such that $U_i \cap U_j \neq \emptyset$, then the map $\psi_{ij} = \varphi_i \circ \varphi_j^{-1}$ from $\varphi_j(U_i \cap U_j)$ to $\varphi_i(U_i \cap U_j)$ is infinitely differentiable. ψ_{ij} is known as a transition function.

In figure 3 we have represented (well copied the image from Nakahara) the ideas above.

Physical Explanation

An *n*-dimensional manifold is therefore a space which locally looks like \mathbb{R}^n , even if the global structure is more complicated. The Earth is an excellent example. As almost everyone knows, the Earth is round, however locally around us it looks flat and the surface looks like \mathbb{R}^2 . Despite this the global properties of the Earth are very different to \mathbb{R}^2 .

The pair (U_i, φ_i) are called a *chart* and the collection of charts is called an *atlas*. The subsets U_i are called the *coordinate neighbourhood* while the φ_i is called the *coordinate function*, or simply the *coordinate*. The homeomorphism φ_i is represented by *n* functions $\varphi_i(p) = \{x^1(p), \dots, x^n(p)\}$, with this set $\{x^\mu(p)\}$ and $\mu = 1, \dots, n$ also called the *coordinate*. A point $p \in M$ exists independently of its coordinates, however we will often be sloppy and denote the point p through its coordinates.

If U_i and U_j overlap, two coordinate systems are assigned to the same point in $U_i \cap U_j$. Axiom 5 asserts that the transition function from one coordinate system to another be smooth \mathbb{C}^∞ . One may be alarmed by this but there is no reason for trepidation, it is analogous to labelling a point by Euclidean coordinates and polar coordinates. The map φ_i assigns *n* coordinates values x^μ , ($1 \leq \mu \leq n$) to a point $p \in U_i \cap U_j$, while φ_j assigns coordinates y^μ to the same point. The transition function from y to x , $x^\mu = x^\mu(y)$ is given by *n* functions of *n* variables, and is the explicit form of the map $\psi_{ji} = \varphi_j \circ \varphi_i^{-1}$. The differentiability in the




Figure 3: Here we see the manifold M and two coordinate charts. The homeomorphisms φ_i maps U_i onto an open set of $U'_i \subset \mathbb{R}^n$ providing coordinates for the point $p \in M$. If $U_i \cap U_j \neq \emptyset$ the transition functions from one coordinate system to another is smooth.

definition is then in the usual sense we are familiar from calculus. All this leads to us being able to move over M so long as we choose coordinates which vary in a smooth way over the manifold.

If the union of two atlases $\{(U_i, \varphi_i)\}$ and $\{(V_j, \psi_j)\}$ is again an atlas, then these two atlases are said to be *compatible*. The compatibility is an equivalence relation. This equivalence class is called the *differentiable structure*. Mutually compatible atlases define the same differentiable structure on M .

Let us briefly comment on manifolds with a boundary. We have assumed that the coordinate neighbourhood U_i is homeomorphic to an open set of \mathbb{R}^n . In some cases this is too restrictive. If a topological space M is covered by a family of open sets $\{U_i\}$ each of which is homeomorphic to an open set $H^n \equiv \{(x^1, \dots, x^n) \in \mathbb{R}^n | x^n \geq 0\}$, M is said to be a manifold with boundary. The analogous plot of figure 3 for the manifold with a boundary is given in

figure 4.




Figure 4: A manifold with a boundary. The point p is on the boundary. Note the subtle difference, for a manifold without a boundary the left figure would be extended below $x^n=0$.

The set of points which are mapped to points with $x^n = 0$ is called the *boundary* of M and is denoted by ∂M . The coordinates on ∂M are given by $n - 1$ numbers $(x^1, \dots, x^{n-1}, 0)$. We now need to be careful when we define smoothness on the overlaps. The map $\psi_{ij} : \varphi_j(U_i \cap U_j) \rightarrow \varphi_i(U_i \cap U_j)$ is defined on an open set of H^n in general, and ψ_{ij} is said to be smooth if it is C^∞ in an open set of \mathbb{R}^n which contains $\varphi_j(U_i \cap U_j)$.

Example 3.5: Charts on some manifolds

- \mathbb{R}^n is a differentiable manifold trivially. A single chart covers the whole space and we take φ to be the identity map.
- Let $n = 1$ and let us impose connectedness. Then there are two choices, either \mathbb{R} or the circle S^1 . Let us work out an atlas for S^1 . For concreteness let us embed the circle in \mathbb{R}^2 via $x^2 + y^2 = 1$. We will need at least two charts. We can take them as in figure 5. Define $\varphi_1^{-1} : (0, 2\pi) \rightarrow S^1$ by^a

$$\varphi_1^{-1} : \theta \rightarrow (\cos \theta, \sin \theta), \tag{3.8}$$

whose image is $S^1 - \{(1, 0)\}$. Similarly define $\varphi_2^{-1} : (-\pi, \pi) \rightarrow S^1$ by

$$\varphi_2^{-1} : \theta \rightarrow (\cos \theta, \sin \theta), \quad (3.9)$$

whose image is $S^1 - \{(-1, 0)\}$. Clearly both φ_i^{-1} are invertible and all the maps are continuous, thus the φ_i 's are homeomorphisms. The transition functions seem trivial for this example but one must be careful to end up in the correct domain. The two charts overlap on the upper and lower hemispheres and therefore we have

$$\varphi_2(\varphi_1^{-1}(\theta)) = \begin{cases} \theta & \text{if } \theta \in (0, \pi) \\ \theta - 2\pi & \text{if } \theta \in (\pi, 2\pi) \end{cases}. \quad (3.10)$$

The transition function isn't defined at $\theta = 0$ or $\theta = \pi$, nonetheless it is smooth on each of the two overlapping open sets as required.




Figure 5: Two charts on S^1 .

- We have seen that we can use two charts to describe an S^1 , but this is a choice that we have made. To show this let us consider a different Atlas on the unit S^1 consisting of the four charts:

$$\begin{aligned} U_1 &\equiv \{(x, y) \in S^1 | x > 0\}, & \varphi_1(x, y) &= y, \\ U_2 &\equiv \{(x, y) \in S^1 | x < 0\}, & \varphi_2(x, y) &= y, \\ U_3 &\equiv \{(x, y) \in S^1 | y > 0\}, & \varphi_3(x, y) &= x, \\ U_4 &\equiv \{(x, y) \in S^1 | y < 0\}, & \varphi_4(x, y) &= x, \end{aligned} \quad (3.11)$$

Note that the circle is defined by $x^2 + y^2 = 1$. Pictorially the charts are given in figure 6. We can see that the patches cover the S^1 from figure 6. Moreover we can see that the φ 's are continuous on $\varphi_i(U_i)$ (since it is just x or y) and that they are invertible. One

needs to be careful with mapping the point into the correct domain, for the inverse, for example for φ_1 we have

$$\varphi_1^{-1}(y) = (\sqrt{1-y^2}, y), \quad (3.12)$$

which is continuous on U_1 . Note that when we took the inverse, working out the point on the circle, we needed to work out the value of x by solving the quadratic $x^2 = 1 - y^2$. We fix the sign by making sure we end up in the correct patch. Note that if we were mapping into U_2 we would have included an additional minus sign. It remains to see that the transition functions are infinitely differentiable. Consider $U_1 \cap U_3$ which is the upper right quadrant in figure 6, with

$$y = \sqrt{1-x^2} \quad \text{with } 0 < y < 1 \text{ and } 0 < x < 1. \quad (3.13)$$

Then

$$\varphi_3 \circ \varphi_1^{-1}(y) = \sqrt{1-y^2}, \quad (3.14)$$

which is indeed infinitely differentiable. This shows that S^1 is a differentiable manifold and we have seen this for two different atlases.




Figure 6: Four charts on S^1 .

- Let us consider a slightly less trivial example, the n -dimensional sphere S^n . We may realise it by embedding it in \mathbb{R}^{n+1} . (Note that embedding it in a higher-dimensional

space is just for convenience and not a necessary requirement for being a manifold, in fact some n -dimensional spaces cannot be embedded in \mathbb{R}^{n+1} , for example hyperbolic space.)

We can realise the n -dimensional sphere S^n in \mathbb{R}^{n+1} as

$$\sum_{i=0}^n (x^i)^2 = 1. \quad (3.15)$$

We can introduce coordinate neighbourhoods

$$\begin{aligned} U_{i+} &\equiv \{(x^0, x^1, \dots, x^n) \in S^n \mid x^i > 0\}, \\ U_{i-} &\equiv \{(x^0, x^1, \dots, x^n) \in S^n \mid x^i < 0\}. \end{aligned} \quad (3.16)$$

Next define the coordinate map $\varphi_{i+} : U_{i+} \rightarrow \mathbb{R}^n$ to be

$$\varphi_{i+}(x^0, \dots, x^n) = (x^0, \dots, x^{i-1}, x^{i+1}, \dots, x^n), \quad (3.17)$$

and $\varphi_{i-} : U_{i-} \rightarrow \mathbb{R}^n$ to be

$$\varphi_{i-}(x^0, \dots, x^{i-1}, x^{i+1}, \dots, x^n). \quad (3.18)$$

Note that the domains of φ_{i+} and φ_{i-} are different and they have no overlap. Instead they are the projections of the hemispheres $U_{i\pm}$ to the plane $x^i = 0$. The transition functions can be obtained simply from the above maps. As an example let us take S^2 , then we have six coordinate neighbourhoods: $U_{x\pm}, U_{y\pm}, U_{z\pm}$. The transition function $\psi_{(y-)(x+)} \equiv \varphi_{y-} \circ \varphi_{x+}^{-1}$ is given by

$$\psi_{(y-)(x+)} : (y, z) \rightarrow (\sqrt{1 - y^2 - z^2}, z). \quad (3.19)$$

This is infinitely differentiable on $U_{x+} \cap U_{y-}$.

^aUntil now we would just have taken the range to be $\theta \in [0, 2\pi)$ and been happy with this. However this does not meet our requirement of being a chart since it is not an open set. This would present problems later when we try to differentiate anything at $\theta = 0$. Recall that the derivative requires us to be able to take limits from both sides, and since there is nothing smaller than 0 we are stuck.

We have seen that to describe n -dimensional spheres we need more than one chart. The need to deal with multiple charts arises when we consider manifolds of non-trivial topology. When we come to discuss general relativity we will care a lot about changing coordinates and the limitations of the coordinate systems. In almost all situations that we will consider a

single set of coordinates generally covers enough of the space to tell us everything we need to know. However as one progresses in physics, topology becomes more important. We will not see much of this but you may see this in some of your other physics/mathematics courses.

Key Points

An n -dimensional manifold is a space which locally looks like \mathbb{R}^n at each point. We can introduce charts within an Atlas which cover the manifold. In order for this to be well defined, and our description of the space to not depend on the choice of charts, we impose the conditions in definition 13.

3.2 Calculus on manifolds

The reason why differentiable manifolds are useful is because it allows us to use the usual calculus we have developed on \mathbb{R}^n for curved backgrounds. One of the key requirements is smoothness of the transition functions, which implies that the calculus is independent of the chosen coordinates.

3.2.1 Differentiable maps

Let $f : M \rightarrow N$ be a map from an m -dimensional manifold M to an n -dimensional manifold N . A point $p \in M$ is mapped to a point $f(p) \in N$. We may take a chart (U, φ) on M and a chart (V, ψ) in N where for all $p \in U$, $f(p) \in V$. Moreover let $\varphi(U) = U' \subset \mathbb{R}^m$ and $\psi(V) = V' \subset \mathbb{R}^n$. Then f has the following coordinate presentation:

$$\psi \circ f \circ \varphi^{-1} : \mathbb{R}^m \rightarrow \mathbb{R}^n. \quad (3.20)$$

If we write $\varphi(p) = \{x^\mu\}$ and $\psi(f(p)) = \{y^\alpha\}$ then, $\psi \circ f \circ \varphi^{-1}$ is just the usual vector-valued function $y = \psi \circ f \circ \varphi^{-1}(x)$ of m variables. Sometimes it is useful to abuse notation and write $y = f(x)$ or $y^\alpha = f^\alpha(x^\mu)$ when we know the coordinate systems on M and N that are in use.

Definition 14 (Smooth function) *We say that a function $\hat{f} : M \rightarrow \mathbb{R}$ is smooth if the map $\hat{f} \circ \varphi^{-1} : U' \rightarrow \mathbb{R}$ is smooth for all charts. We let the set of all smooth functions on M be denoted by $\mathcal{F}(M)$.*

Recall that smoothness in Calculus is defined by being infinitely differentiable, C^∞ . that is it has continuous derivatives over some domain.

Definition 15 (Smooth map, Differentiable map) *We say that a map $f : M \rightarrow N$ between two manifolds is smooth if the map $\psi \circ f \circ \varphi^{-1} : U' \rightarrow V'$ is smooth for all charts $\varphi : M \rightarrow \mathbb{R}^m$ and $\psi : N \rightarrow \mathbb{R}^n$. If $y = \psi \circ f \circ \varphi^{-1}(x)$ is C^∞ then we say that f is differentiable at p . This is actually independent of the coordinate system.*

Note that our definition of a smooth function is a particular case of a smooth map as defined directly above. There we have taken $N = \mathbb{R}$ and therefore we do not need the second coordinate map ψ .

Definition 16 (Diffeomorphism) *Let $f : M \rightarrow N$ be a homeomorphism and ψ and φ coordinate functions. If $\psi \circ f \circ \varphi^{-1}$ is invertible and both, $\psi \circ f \circ \varphi^{-1}$ and its inverse $\varphi \circ f^{-1} \circ \psi^{-1}$ are C^∞ , then f is called a diffeomorphism and M is said to be diffeomorphic to N and vice-versa. This is denoted by $M \equiv N$.*

Since the map is invertible it follows that if $M \equiv N$ then $\dim M = \dim N$. Homeomorphisms classify spaces according to whether it is possible to deform one space into another *continuously*. *Diffeomorphisms* classify spaces into equivalence classes according to whether it is possible to deform one space into the other *smoothly*. As such a diffeomorphism is stronger than a homeomorphism, it requires that both the map and its inverse are smooth. Two diffeomorphic manifolds are viewed as the same manifold.

The question as to whether a homeomorphism is a diffeomorphism is quite subtle and far beyond the scope of this course, but let us give a small taste. Given a differentiable structure on your manifold, this is defined as an equivalence of atlases, one must then ask is it possible that a manifold admits different differentiable structures. This would mean that we can pick two atlases on a manifold which are *not compatible* (the union of the two Atlases is not another Atlas). Finding such examples is non-trivial and it is known that it is only possible if $\dim(M) \geq 4$. In 1956 Milnor showed that the seven-sphere S^7 admits different 28 differentiable structures. More bizarrely it has been shown that \mathbb{R}^4 has uncountably many pairwise non-diffeomorphic open subsets each of which is homeomorphic to \mathbb{R}^4 . Throughout this course we will assume that our manifold admits a unique differentiable structure for simplicity.

3.2.2 Tangent Vectors

Having defined maps on a manifold we can define other objects on the manifold. The elementary notion of a vector no longer works: where is the origin? what is a straight line? etc. On a manifold a vector is defined to be a *tangent vector* to a curve in M .

Definition 17 (Tangent vector) *A tangent vector X_p is an object that differentiates functions at a point $p \in M$, $X_p : C^\infty(M) \rightarrow \mathbb{R}$ which satisfies*

1. *Linearity: $X_p[f + g] = X_p[f] + X_p[g]$ for all $f, g \in C^\infty(M)$.*

2. $X_p[f] = 0$ when f is a constant function.

3. *Leibnitz identity*: $X_p[fg] = f(p)X_p[g] + X_p[f]g(p)$ for all $f, g \in C^\infty(M)$.

To define a tangent vector we need a curve $\gamma : (a, b) \rightarrow M$ and a function $f : M \rightarrow \mathbb{R}$. For simplicity let $0 \in (a, b)$ and let us parametrise our curve by t . We define the *tangent vector at $\gamma(0)$* to be the directional derivative of a function $f(\gamma(t))$ along the curve $\gamma(t)$ at $t = 0$. The tangent vector to γ at $\gamma(0)$ is the linear map X_p from the space of smooth functions of M to \mathbb{R} defined by

$$X_p[f] = \left. \frac{df(\gamma(t))}{dt} \right|_{t=0}. \quad (3.21)$$

In terms of local coordinates we have

$$X_p[f] = \left. \frac{\partial f}{\partial x^\mu} \frac{dx^\mu(\gamma(t))}{dt} \right|_{t=0}. \quad (3.22)$$

Notice the abuse of notation, the first term should really be

$$\frac{\partial(f \circ \varphi^{-1}(x))}{\partial x^\mu} \equiv \frac{\partial f}{\partial x^\mu}, \quad (3.23)$$

and the second should be

$$\frac{d}{dt}(\varphi \circ \gamma)^\mu \equiv \frac{dx^\mu(\gamma(t))}{dt}, \quad (3.24)$$

we will often employ this abuse of notation.

Exercise 3.2: Tangent vectors at p

Show that the set of all tangent vectors at p forms an n -dimensional vector space $T_p(M)$.

We have just found that the tangent vector can be written as

$$X_p = X_p^\mu \left(\frac{\partial}{\partial x^\mu} \right), \quad X_p^\mu = \left. \frac{dx^\mu(\gamma(t))}{dt} \right|_{t=0} \quad (3.25)$$

then

$$\left. \frac{df(\gamma(t))}{dt} \right|_{t=0} = X_p^\mu \frac{\partial f}{\partial x^\mu} \equiv X_p[f]. \quad (3.26)$$

We define X_p to be the tangent vector to M at $p = \gamma(0)$ along the direction given by the curve $\gamma(t)$. There is then a natural basis of $T_p(M)$ for us to take given by

$$\left\{ \left. \frac{\partial}{\partial x^\mu} \right|_p, \mu = 1, \dots, \dim(M) \right\}. \quad (3.27)$$

This is chart dependent since we chose a chart in the neighbourhood of p . Choosing a different chart would give a different basis of $T_p(M)$. The basis defined in this way is called the *coordinate basis*.

Let us see how the coordinate basis changes when we choose different coordinates. Let $p \in U_i \cap U_j$ and let $x = \varphi_i(p)$ and $x' = \varphi_j(p)$ be two charts defined in the neighbourhood of the point p . For a smooth function f we have:

$$\begin{aligned} \left(\frac{\partial}{\partial x^\mu}\right)\Big|_p [f] &= \frac{\partial}{\partial x^\mu} (f \circ \varphi^{-1})\Big|_{\varphi(p)} \\ &= \frac{\partial}{\partial x^\mu} [(f \circ \varphi'^{-1}) \circ (\varphi' \circ \varphi^{-1})]\Big|_{\varphi(p)}. \end{aligned} \quad (3.28)$$

Let $F' = f \circ \varphi'^{-1}$, this is a function in the x' coordinates. Moreover note that $\varphi' \circ \varphi^{-1}$ is simplify the functions $x'^\mu(x)$, that is the primed coordinates in terms of the unprimed ones. Hence it follows that we have:

$$\begin{aligned} \left(\frac{\partial}{\partial x^\mu}\right)\Big|_p [f] &= \frac{\partial}{\partial x^\mu} [F'(x'(x))]\Big|_{\varphi(p)} \\ &= \frac{\partial x'^\nu}{\partial x^\mu}\Big|_{\varphi(p)} \frac{\partial F'(x')}{\partial x'^\nu}\Big|_{\varphi'(p)} \\ &= \frac{\partial x'^\nu}{\partial x^\mu}\Big|_{\varphi(p)} \frac{\partial}{\partial x'^\nu}\Big|_p [f], \end{aligned} \quad (3.29)$$

and therefore we have

$$\frac{\partial}{\partial x^\mu}\Big|_p = \frac{\partial x'^\nu}{\partial x^\mu}\Big|_{\varphi(p)} \frac{\partial}{\partial x'^\nu}\Big|_p. \quad (3.30)$$

This then defines the transformation of one basis into another. It is now straightforward to work out the transformation of the components of a vector X_p^μ into the components of another basis $X_p'^\mu$ by using that the vector X_p is invariant. It follows that:

$$X_p'^\nu = X_p^\mu \frac{\partial x'^\nu}{\partial x^\mu}\Big|_{\varphi(p)}. \quad (3.31)$$

Components of vectors that transform in this way are known as *contravariant*.

We have defined a tangent vector X as a differential operator acting on functions along a curve passing through the point p , but there is some redundancy in this since two curves passing can give the same tangent vector at p . This leads us to define an equivalence class of curves on M .

Definition 18 (Equivalence class of curves) *If two curves $\gamma_1(t)$ and $\gamma_2(t)$ satisfy*

$$(i) \quad \gamma_1(0) = \gamma_2(0) = p,$$

$$(ii) \quad \frac{dx^\mu(\gamma_1(t))}{dt}\Big|_{t=0} = \frac{dx^\mu(\gamma_2(t))}{dt}\Big|_{t=0},$$

then $\gamma_1(t)$ and $\gamma_2(t)$ yield the same differential operator X at p . This allows us to define the equivalence relation between curves at the point p , $\gamma_1(t) \sim \gamma_2(t)$. We identify the tangent vector X with the equivalence class of curves

$$[\gamma(t)] = \left\{ \tilde{\gamma}(t) \mid \gamma(0) = \tilde{\gamma}(0) \text{ and } \left. \frac{dx^\mu(\gamma(t))}{dt} \right|_{t=0} = \left. \frac{dx^\mu(\tilde{\gamma}(t))}{dt} \right|_{t=0} \right\} \quad (3.32)$$

rather than a particular representative of the curve.

All the equivalence classes of curves at a point $p \in M$, i.e. all the tangent vectors at p , form a vector space called the *tangent space* of M at p , $T_p(M)$. From our discussion above we can take the basis of vectors for $T_p(M)$ to be $e_\mu = \frac{\partial}{\partial x^\mu}$. It follows that $\dim T_p(M) = \dim(M)$.

It is often convenient to use the coordinate basis, however the basis clearly depends on the coordinates that we are using. At times it is convenient to use other bases, one such basis which is a ‘non-coordinate bases’ is known as vielbeins. These are necessary when defining spinors on curved backgrounds and can often make computations simpler. We will introduce these later in the problem sheets.

Note that for two distinct points p and q the tangent spaces $T_p(M)$ and $T_q(M)$ are different. We cannot add vectors from one to a vector in the other. In fact even to compare the vectors in $T_p(M)$ with the vectors in $T_q(M)$ we need to introduce the notion of *parallel transport*, which will appear shortly.

3.2.3 Vector fields

So far we have defined tangent vectors at a single point, but what we really want is an object with a choice of tangent vector for every point in our manifold. These objects are called fields.

A vector field X is defined to be a smooth assignment of tangent vectors X_p to each point $p \in M$. Here we mean that if we feed a function into the vector field then we obtain another function which is the differentiation of the first. The vector field is then smooth if, starting with our smooth function f we obtain another smooth function $X[f]$. Therefore a vector field defines a map $X : C^\infty(M) \rightarrow C^\infty(M)$. To evaluate $X[f]$ at a point p we have

$$X[f] \Big|_p = X_p[f]. \quad (3.33)$$

We denote the space of all vector fields on M to be $\mathcal{X}(M)$.

3.2.4 One-forms

Since $T_p(M)$ is a vector space, there exists a dual vector space to $T_p(M)$ whose element is a linear function from $T_p(M) \rightarrow \mathbb{R}$. The dual space is called the *cotangent space* at p , and

is denoted by $T_p^*(M)$. An element ω_p of $T_p^*(M)$ is a map $\omega_p : T_p(M) \rightarrow \mathbb{R}$ and is called a *dual vector/cotangent vector* or in the context of differential forms a *one-form*. The simplest example of a one-form is the differential df for a smooth function f on M . The action of a vector V_p on f is $V_p[f] = V_p^\mu \frac{\partial f}{\partial x^\mu} \in \mathbb{R}$. The action of $df \in T_p^*(M)$ on $V_p \in T_p(M)$ is defined by

$$\langle df, V_p \rangle \equiv V_p[f] = V_p^\mu \frac{\partial f}{\partial x^\mu} \in \mathbb{R}. \quad (3.34)$$

This is then \mathbb{R} -linear in both V_p and f . In terms of the coordinate basis we have

$$df = \frac{\partial f}{\partial x^\mu} dx^\mu, \quad (3.35)$$

and it is natural to regard $\{dx^\mu\}$ as a basis of $T_p^*(M)$. This is a dual basis since

$$\left\langle dx^\mu, \frac{\partial}{\partial x^\nu} \right\rangle = \frac{\partial x^\mu}{\partial x^\nu} = \delta_\nu^\mu. \quad (3.36)$$

We can then write an arbitrary one-form as

$$\omega_p = \omega_{p,\mu} dx^\mu. \quad (3.37)$$

If we take a vector V_p and a one-form ω_p we may define the *inner product* between one-forms and vectors $\langle \cdot, \cdot \rangle : T_p^*(M) \times T_p(M) \rightarrow \mathbb{R}$ to be

$$\langle \omega_p, V_p \rangle = \omega_\mu V^\mu \left\langle dx^\mu, \frac{\partial}{\partial x^\nu} \right\rangle = \omega_\mu V_p^\nu \delta_\nu^\mu = \omega_\mu V^\mu. \quad (3.38)$$

The inner product is defined between a vector and a covector. Since ω_p is defined without reference to any coordinate system for a point $p \in U_i \cap U_j$ we have

$$\omega_p = \omega_\mu dx^\mu = \omega'_\mu dx'^\mu, \quad (3.39)$$

with x and x' as before. Then we have

$$\omega'_\nu = \omega_\mu \frac{\partial x^\mu}{\partial x'^\nu}, \quad (3.40)$$

which is the transformation of the components of a co-vector.

We will often denote the set of one-forms to be $\Omega^{(1)}(M)$.

3.2.5 Tensors

Definition 19 (Tensor) A tensor at the point p of type (q, r) is a multilinear object which maps q elements of $T_p^*(M)$ and r elements of $T_p(M)$ to \mathbb{R} :

$$T : \otimes^q T_p^*(M) \otimes^r T_p(M) \rightarrow \mathbb{R}. \quad (3.41)$$

We define $\mathcal{T}_p^{(q,r)}(M)$ to be the set of (q, r) tensors at $p \in M$.

An element of $\mathcal{T}^{(q,r)}(M)$ can be written in terms of the coordinate bases described above as

$$T = T^{\mu_1 \dots \mu_q}_{\nu_1 \dots \nu_r} \frac{\partial}{\partial x^{\mu_1}} \dots \frac{\partial}{\partial x^{\mu_q}} dx^{\nu_1} \dots dx^{\nu_r}. \quad (3.42)$$

Let $V_i = V_i^\mu \frac{\partial}{\partial x^\mu}$ with $1 \leq i \leq r$ and $\omega_j = \omega_{j\mu} dx^\mu$ with $1 \leq j \leq q$ then the action of T is

$$T(\omega_1, \dots, \omega_q; V_1, \dots, V_r) = T^{\mu_1 \dots \mu_q}_{\nu_1 \dots \nu_r} \omega_{1\mu_1} \dots \omega_{q\mu_q} V_1^{\nu_1} \dots V_r^{\nu_r}. \quad (3.43)$$

The transformation of the components of a tensor under a change of coordinates follows simply from the transformation of the components of a contravariant vector and a covariant vector. We have

$$T^{\mu'_1 \dots \mu'_q}_{\nu'_1 \dots \nu'_r} = T^{\mu_1 \dots \mu_q}_{\nu_1 \dots \nu_r} \frac{\partial x^{\mu'_1}}{\partial x^{\mu_1}} \dots \frac{\partial x^{\mu'_q}}{\partial x^{\mu_q}} \frac{\partial x^{\nu_1}}{\partial x^{\nu'_1}} \dots \frac{\partial x^{\nu_r}}{\partial x^{\nu'_r}}. \quad (3.44)$$

Note that in all cases given the placement of indices this is really the only type of transformation that we can really have.

As before we can define a *Tensor field* of type (q, r) to be a smooth assignment of an element of $\mathcal{T}_p^{q,r}(M)$ at each point $p \in M$. The set of tensor fields of type (q, r) on M is denoted by $\mathcal{T}^{q,r}(M)$.

Operations on Tensor fields There are a variety of different operations we can do on tensor fields to generate new tensor fields. We can add or subtract them, multiply by smooth functions. This is just the statement that tensors at a point p form a vector space. These operations preserve the rank of the tensor but there are operations we can do which do not preserve the rank.

Given a tensor field S of rank (q, r) and a tensor field T of rank (s, t) we may take the tensor product of the two tensor fields together to form a tensor field $S \otimes T$ of rank $(q+s, r+t)$. The *tensor product* $S \otimes T$ is defined by

$$S \otimes T(\omega_1, \dots, \omega_q, \eta_1, \dots, \eta_s, X_1, \dots, X_r, Y_1, \dots, Y_t) = S(\omega_1, \dots, \omega_q, X_1, \dots, X_r)T(\eta_1, \dots, \eta_s, Y_1, \dots, Y_t). \quad (3.45)$$

In terms of components we have:

$$(S \otimes T)^{\mu_1 \dots \mu_q \nu_1 \dots \nu_s}_{\rho_1 \dots \rho_r \sigma_1 \dots \sigma_t} = S^{\mu_1 \dots \mu_q}_{\rho_1 \dots \rho_r} T^{\nu_1 \dots \nu_s}_{\sigma_1 \dots \sigma_t}. \quad (3.46)$$

Given an (q, r) tensor we may define a $(q-1, r-1)$ tensor through *contraction*. To do this we replace one of the $T_p^*(M)$ entries with a basis vector $\hat{\theta}^\mu$ and the corresponding $T_p(M)$

entry with the dual vector \hat{e}_μ and then sum over the indices. For example we have

$$S(\omega_1, \dots, \omega_{q-1}, X_1, \dots, X_{r-1}) = \sum_{\mu=1}^n T(\omega_1, \dots, \omega_{q-1}, \hat{\theta}^\mu, X_1, \dots, X_{r-1}, \hat{e}_\mu). \quad (3.47)$$

The placement of either of the basis vector/co-vector is arbitrary and moving it around we obtain a typically different tensor, again of rank $(q-1, r-1)$. In terms of components we have

$$S^{\mu_1 \dots \mu_{q-1}}{}_{\nu_1 \dots \nu_{r-1}} = T^{\mu_1 \dots \mu_{q-1} \sigma}{}_{\nu_1 \dots \nu_{r-1} \sigma}. \quad (3.48)$$

Another operation we can do is to symmetrise and anti-symmetrise the tensor. For example given a rank $(0, r)$ tensor T we can define the symmetrisation of T to be

$$S[T(X_1, \dots, X_r)] = \frac{1}{r!} \sum_{\sigma \in S_r} T(X_{\sigma(1)}, \dots, X_{\sigma(r)}), \quad (3.49)$$

and the anti-symmetrisation to be

$$A[T(X_1, \dots, X_r)] = \frac{1}{r!} \sum_{\sigma \in S_r} \text{sign}(\sigma) T(X_{\sigma(1)}, \dots, X_{\sigma(r)}), \quad (3.50)$$

where S_r is the permutation group of r objects. In components we have

$$S[T]_{\mu_1 \dots \mu_r} = \frac{1}{r!} \sum_{\sigma \in S_r} T_{\mu_{\sigma(1)} \dots \mu_{\sigma(r)}} \equiv T_{(\mu_1 \dots \mu_r)}, \quad (3.51)$$

and

$$A[T]_{\mu_1 \dots \mu_r} = \frac{1}{r!} \sum_{\sigma \in S_r} \text{sign}(\sigma) T_{\mu_{\sigma(1)} \dots \mu_{\sigma(r)}} \equiv T_{[\mu_1 \dots \mu_r]} \quad (3.52)$$

Note that it only makes sense to symmetrise and anti-symmetrise over objects of the same type, or the same placement of indices. We also divide by the symmetry factor $r!$ which is the number of permutations, this ensures that if an object is (anti-)symmetric then (anti-)symmetrisation acts as the identity.

3.2.6 Induced maps

Definition 20 (Push-forward/differential map) *A smooth map $f : M \rightarrow N$ naturally induces a map f_* called the differential map or push-forward,*

$$f_* : T_p(M) \rightarrow T_{f(p)}(N). \quad (3.53)$$

The explicit form of f_ is obtained by the definition of the directional derivative along a curve. Let $g \in \mathcal{F}(N)$ then $g \circ f \in \mathcal{F}(M)$. A vector $V \in T_p(M)$ acts on $g \circ f$ to give a number $V[g \circ f]$. We can now define $f_*V \in T_{f(p)}(N)$ by*

$$(f_*V)[g] \equiv V[g \circ f]. \quad (3.54)$$

See figure 7 for a pictorial representation.




Figure 7: A map $f : M \rightarrow N$ induces the differential map $f_* : T_p(M) \rightarrow T_{f(p)}(N)$. Note that the mapping is performed by mapping the curve $c(t)$ between the two manifolds using the map f .

[**Aside:** Note that we do not require the map f to be invertible here, just smooth, so it need not be a diffeomorphism. We will however require this later so let us just assume it from now on.]

Physical Explanation

The map f is telling us how we would move points in M to points in N , and in particular how we would move a curve in M to a curve in N . Instead the push forward tells us how the tangent vectors of M are mapped to tangent vectors of N .

We can write this more explicitly by introducing coordinates. Let us introduce the charts (U, φ) on M and (V, ψ) on N , then

$$(f_* V)[g \circ \psi^{-1}(y)] = V[g \circ f \circ \varphi^{-1}(x)], \quad (3.55)$$

where $x = \varphi(p)$ and $y = \psi(f(p))$. Let $V = V^\mu \frac{\partial}{\partial x^\mu}$ and $f_* V = W^\alpha \frac{\partial}{\partial y^\alpha}$, then in components it reads

$$W^\alpha = V^\mu \frac{\partial y^\alpha(x)}{\partial x^\mu}. \quad (3.56)$$

This is nothing but the Jacobian of the map $f : M \rightarrow N$ that we are all familiar with. So far we have done this for vector fields, but this can easily be extended to tensors of type $(q, 0)$ without additional thought. Let us consider an example to put this into practice:

Example 3.6: Push forward

Let (x^1, x^2) and (y^1, y^2, y^3) be coordinates on M and N respectively, and let $V = a \frac{\partial}{\partial x^1} + b \frac{\partial}{\partial x^2}$. Take the map $f : M \rightarrow N$ whose coordinate representation is

$$f(x^1, x^2) = (x^1, x^2, \sqrt{1 - (x^1)^2 - (x^2)^2}). \quad (3.57)$$

Then

$$f_*V = V^\mu \frac{\partial y^\alpha}{\partial x^\mu} \frac{\partial}{\partial y^\alpha} = a \frac{\partial}{\partial y^1} + b \frac{\partial}{\partial y^2} - \left(a \frac{y^1}{y^3} + b \frac{y^2}{y^3} \right) \frac{\partial}{\partial y^3}. \quad (3.58)$$

Definition 21 (Pull-back) A smooth map $f : M \rightarrow N$ also induces a map between cotangent space

$$f^* : T_{f(p)}^*(N) \rightarrow T_p^*(M), \quad (3.59)$$

which is called the pull-back. If we take $V \in T_p(M)$ and $\omega \in T_{f(p)}^*(N)$ then the pull-back of ω by f^* is defined to be

$$\langle f^*\omega, V \rangle = \langle \omega, f_*V \rangle. \quad (3.60)$$

In components we have

$$(f^*\omega)_\mu = \omega_\alpha \frac{\partial y^\alpha}{\partial x^\mu}. \quad (3.61)$$

Physical Explanation

The push forward allowed us to map vectors of M to vectors of N . The pull-back allows us to map the dual vectors (one-forms) between the two spaces. In contrast to the push-forward it takes one-forms from N to one-forms in M and not from M to N .

The pull-back can be extended to tensors of type $(0, r)$.

We have defined the push-forward and pull-back for tensors of type $(q, 0)$ and $(0, r)$ respectively. Note that the push-forward induces a map in the same direction as the map f while the pull-back induces a map in the opposite direction to the original map. If f is a diffeomorphism $f : M \rightarrow N$, then we also have the inverse $f^{-1} : N \rightarrow M$ and therefore we can transport any object from M to N and back to our hearts content.

3.3 Flows and Lie derivatives

In this section we introduce flows generated by vector fields and use them to define the Lie derivative, which measures how tensor fields change along these flows.

Definition 22 (Integral Curve) Let X be a vector field on M . An integral curve $x(t)$ of X is a curve in M whose tangent vector at $x(t)$ is $X|_x$. Given a chart (U, φ) , this means that

$$\frac{dx^\mu(t)}{dt} = X^\mu(x(t)), \quad (3.62)$$

where $x^\mu(t)$ is the μ 'th component of $\varphi(x(t))$ and $X = X^\mu \frac{\partial}{\partial x^\mu}$.

As always we have very much abused notation, using x to denote a point in M as well as its coordinates. Finding an Integral curve is equivalent to solving the ODE with initial conditions $x^\mu(0) = x_0^\mu$. The existence and uniqueness theorems for ODEs implies that there is always a unique solution, at least locally, with the given initial data.

Physical Explanation

Given a vector field X the integral curve is the curve through your manifold which has tangent vector X at every point.

Definition 23 (Flow) Let $\sigma(t, x_0)$ be an integral curve of X which passes through the point x_0 at $t = 0$, and denote the coordinate by $\sigma^\mu(t, x_0)$. The flow equation becomes

$$\frac{d}{dt}\sigma^\mu(t, x_0) = X^\mu(\sigma(t, x_0)), \quad (3.63)$$

with the initial condition

$$\sigma^\mu(0, x_0) = x_0^\mu. \quad (3.64)$$

Then the map $\sigma : \mathbb{R} \times M \rightarrow M$ is called a flow generated by $X \in \mathcal{X}(M)$.

Physical Explanation

The flow gives a way for us to move points on the manifold along the curves defined by our vector field X . We can think of it as given an initial point x_0^μ we move it to the point $\sigma^\mu(t, x_0)$ along the integral curve of the vector field X . By construction this new point is on the integral curve defined by X which passes through the initial point x_0^μ .

From the definition one can show that a flow satisfies the rule

$$\sigma(t, \sigma^\mu(s, x_0)) = \sigma(t + s, x_0), \quad (3.65)$$

for any $s, t \in \mathbb{R}$. This follows from the uniqueness of the ODE with fixed initial condition.

Theorem 2 For any point $x \in M$, there exists a differentiable map $\sigma : \mathbb{R} \times M \rightarrow M$ such that

(i) $\sigma(0, x) = x$,

(ii) $t \mapsto \sigma(t, x)$ is a solution of (3.63) and (3.64),

(iii) $\sigma(t, \sigma^\mu(s, x)) = \sigma(t + s, x)$.

Note that the initial point is denoted by x to emphasise that σ is a map $\mathbb{R} \times M \rightarrow M$.

Example 3.7: Integral Curves

- Let $M = \mathbb{R}^2$ and let $X(x, y) = -y \frac{\partial}{\partial x} + x \frac{\partial}{\partial y}$ be a vector field in M . Then

$$\sigma(t, (x_0, y_0)) = (x_0 \cos t - y_0 \sin t, x_0 \sin t + y_0 \cos t), \quad (3.66)$$

is a flow generated by X . The flow through (x_0, y_0) is a circle whose centre is at the origin. Clearly $\sigma(t, (x_0, y_0)) = (x_0, y_0)$ if $t = 2\pi n, n \in \mathbb{Z}$. If $(x_0, y_0) = (0, 0)$, the flow stays at $(0, 0)$.

- Consider the sphere S^2 in polar coordinates with the vector field $X = \partial_\phi$. The integral curves are:

$$\frac{d\phi}{dt} = 1, \quad \frac{d\theta}{dt} = 0, \quad (3.67)$$

and this has solution:

$$\theta = \theta_0, \quad \phi = \phi_0 + t. \quad (3.68)$$

The associated diffeomorphism is $\sigma_t : (\theta, \phi) \rightarrow (\theta, \phi + t)$, and the flow lines are simply the lines of constant latitude on the sphere.

3.3.1 One-parameter group of transformations

For fixed $t \in \mathbb{R}$ a flow $\sigma(t, x)$ is a diffeomorphism from M to M which we denote by $\sigma_t : M \rightarrow M$. This map is made into a commutative group by the following rules:

- $\sigma_t(\sigma_s(x)) = \sigma_{t+s}(x)$ i.e. $\sigma_t \circ \sigma_s = \sigma_{t+s}$, (Associative)
- $\sigma_0 = \text{identity map}$ (Unit element),
- $\sigma_{-t} = (\sigma_t)^{-1}$ (Inverse).

Exercise 3.3: Flow defines a commutative group

Show that a flow defines a commutative group.

This group is called the *one-parameter group of transformations*. Locally the group looks like the additive group \mathbb{R} , although they may not be isomorphic globally. For example in the example above (see equation (3.66)) we had that $\sigma_{2\pi n+t} = \sigma_t$ and we find that the one-parameter group is isomorphic to $\text{SO}(2)$ the multiplicative group of 2×2 real matrices of the form;

$$\begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \quad (3.69)$$

or $\text{U}(1)$ the multiplicative group of complex numbers of unit modulus $e^{i\theta}$.

We can consider an infinitesimal transformation and see where it maps the point x . Using (3.63) and (3.64) we find

$$\sigma_\epsilon^\mu(x) = \sigma^\mu(\epsilon, x) = x^\mu + \epsilon X^\mu(x). \quad (3.70)$$

The vector field X in this context is called the *infinitesimal generator* of the transformation σ_t .

Given a vector field X the corresponding flow σ is often referred to as the *exponentiation* of X and is denoted by

$$\sigma^\mu(t, x) = \exp(tX)x^\mu. \quad (3.71)$$

To see why this is so, let us take a parameter t and evaluate the coordinate of a point which is separated from the initial point $x = \sigma(0, x)$ by the parameter distance t along the flow σ . The coordinate corresponding to the point $\sigma(t, x)$ is

$$\begin{aligned} \sigma^\mu(t, x) &= x^\mu + t \frac{d}{ds} \sigma^\mu(s, x) \Big|_{s=0} + \frac{t^2}{2!} \left(\frac{d}{ds} \right)^2 \sigma^\mu(s, x) \Big|_{s=0} + \dots \\ &= \left[1 + t \frac{d}{ds} + \frac{t^2}{2!} \left(\frac{d}{ds} \right)^2 + \dots \right] \sigma^\mu(s, x) \Big|_{s=0} \\ &\equiv \exp \left(t \frac{d}{ds} \right) \sigma^\mu(s, x) \Big|_{s=0}. \end{aligned} \quad (3.72)$$

The last expression can also be written as $\sigma^\mu(t, x) = \exp(tX)x^\mu$ as in the definition above. Then the flow satisfies the following exponential properties:

1. $\sigma(0, x) = x = \exp(0X)x$,
2. $\frac{\sigma(t, x)}{dt} = X \exp(tX)x = \frac{d}{dt} \left(\exp(tX)x \right)$,
3. $\sigma(t, \sigma(s, x)) = \sigma(t, \exp(sX)x) = \exp(tX) \exp(sX)x = \exp((t+s)X)x = \sigma(t+s, x)$.

3.3.2 Lie Derivatives

We have now defined maps using flows, but what are these good for? One use is to construct the Lie derivative. This is a derivative which essentially tells us how something changes along the integral curve of a vector. We have already seen that for a curve with tangent vector field X the change of a function f , along the curve is given by $X[f]$. Motivated by this we define the Lie derivative of the function along the curve with tangent X to be

$$\mathcal{L}_X f = X[f]. \quad (3.73)$$

We now want to ask how vector fields and other tensors transform along curves.

Let $\sigma(t, x)$ and $\tau(t, x)$ be two flows generated by the vector fields X and Y respectively:

$$\frac{d\sigma^\mu(s, x)}{ds} = X^\mu(\sigma(s, x)), \quad \frac{d\tau^\mu(t, x)}{dt} = Y^\mu(\tau(t, x)). \quad (3.74)$$

Let us evaluate the change of the vector field Y along $\sigma(s, x)$. To do this we need to compare the vector Y at a point x with Y at a nearby point $x' = \sigma_\epsilon(x)$, see figure 8. We cannot simply take the difference between the components of Y at the two points since they belong to different tangent spaces: $T_x(M)$ and $T_{\sigma_\epsilon(x)}(M)$, and so the difference between the two vectors is ill-defined. To define a sensible derivative, we first map $Y|_{\sigma_\epsilon(x)}$ to $T_x(M)$ by using the push-forward $(\sigma_{-\epsilon})_* : T_{\sigma_\epsilon(x)}(M) \rightarrow T_x(M)$, after which the two vectors are in the same tangent space and we can take the difference between them, see figure 8.

Definition 24 (Lie Derivative) *The Lie derivative of a vector field Y along the flow σ of the vector field X is defined by*

$$\mathcal{L}_X Y = \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} \left[(\sigma_{-\epsilon})_* Y|_{\sigma_\epsilon(x)} - Y|_x \right]. \quad (3.75)$$

This is still somewhat abstract, since it is a coordinate free expression, so let us write it in components. By writing this in components we obtain another expression for the Lie derivative of a vector field. Let (U, φ) be a chart with the coordinates x and let $X = X^\mu \frac{\partial}{\partial x^\mu}$ and $Y = Y^\mu \frac{\partial}{\partial x^\mu}$ be vector fields defined on U . Then $\sigma_\epsilon(x)$ has the coordinates $x^\mu + \epsilon X^\mu(x)$ and

$$\begin{aligned} Y|_{\sigma_\epsilon(x)} &= Y^\mu(x^\nu + \epsilon X^\nu(x)) e_\mu|_{x+\epsilon X} \\ &\simeq \left[Y^\mu(x) + \epsilon X^\nu(x) \partial_\nu Y^\mu(x) \right] e_\mu|_{x+\epsilon X}, \end{aligned} \quad (3.76)$$

with $e_\mu = \frac{\partial}{\partial x^\mu} \equiv \partial_\mu$. Mapping this vector at $\sigma_\epsilon(x)$ to x using $(\sigma_{-\epsilon}(x))_*$ we obtain

$$(\sigma_{-\epsilon}(x))_* Y|_{\sigma_\epsilon(x)} = \left[Y^\mu(x) + \epsilon X^\lambda(x) \partial_\lambda Y^\mu(x) \right] \partial_\mu(x^\nu - \epsilon X^\nu(x)) e_\nu|_x$$




Figure 8: To compare a vector $Y|_x$ with the vector $Y|_{\sigma_\epsilon(x)}$ the latter must be transported back to x by the differential map $(\sigma_{-\epsilon})_*$, that is we use the push-forward.

$$\begin{aligned}
 &= \left[Y^\mu(x) + \epsilon X^\lambda(x) \partial_\lambda Y^\mu(x) \right] \left[\delta_\mu^\nu - \epsilon \partial_\mu X^\nu(x) \right] e_\nu|_x & (3.77) \\
 &= Y^\mu(x) e_\mu|_x + \epsilon \left[X^\mu(x) \partial_\mu Y^\nu(x) - Y^\mu(x) \partial_\mu X^\nu(x) \right] e_\nu|_x + O(\epsilon^2),
 \end{aligned}$$

and therefore we find

$$\mathcal{L}_X Y = (X^\mu \partial_\mu Y^\nu - Y^\mu \partial_\mu X^\nu) e_\nu. \quad (3.78)$$

This motivates the introduction of the Lie bracket.

Definition 25 (Lie bracket) *The Lie bracket, $[\ , \]$. For vector fields X, Y on M we have*

$$[X, Y]f = X[Y[f]] - Y[X[f]], \quad (3.79)$$

for all $f \in \mathcal{F}(M)$.

In components $[X, Y]$ reads

$$(X^\mu \partial_\mu Y^\nu - Y^\mu \partial_\mu X^\nu) e_\nu, \quad (3.80)$$

and in terms of the Lie bracket the Lie derivative of Y along X is

$$\mathcal{L}_X Y = [X, Y]. \quad (3.81)$$

Exercise 3.4: Some properties of the Lie bracket

Show that the Lie bracket defines a vector field. In addition show that it satisfies the following properties:

1. Bilinearity

$$\begin{aligned} [X, c_1 Y_1 + c_2 Y_2] &= c_1 [X, Y_1] + c_2 [X, Y_2], \\ [c_1 X_1 + c_2 X_2, Y] &= c_1 [X_1, Y] + c_2 [X_2, Y], \end{aligned} \quad (3.82)$$

for any constants c_1 and c_2 .

2. Skew symmetry

$$[X, Y] = -[Y, X]. \quad (3.83)$$

3. Jacobi Identity

$$[[X, Y], Z] + [[Z, X], Y] + [[Y, Z], X] = 0, \quad (3.84)$$

4. For X, Y vector fields and f a smooth function on M then

$$\begin{aligned} \mathcal{L}_{fX} Y &= f[X, Y] - Y[f]X, \\ \mathcal{L}_X(fY) &= f[X, Y] + X[f]Y \end{aligned} \quad (3.85)$$

5. For $f : M \rightarrow N$ then

$$f_*[X, Y] = [f_*X, f_*Y]. \quad (3.86)$$

Exercise 3.5: Algebra of Lie derivative

Show that for vector fields $X, Y, Z \in \mathcal{X}(M)$ we have

$$\mathcal{L}_X \mathcal{L}_Y Z - \mathcal{L}_Y \mathcal{L}_X Z = \mathcal{L}_{[X, Y]} Z. \quad (3.87)$$

Geometrically the Lie bracket shows the non-commutativity of two flows. Let us take the flows $\sigma(s, x)$ and $\tau(t, x)$ generated by X and Y respectively. If we first move a small parameter distance ϵ along the flow σ and then by δ along the second flow τ we end up at a point whose coordinates are

$$\begin{aligned} \tau^\mu(\delta, \sigma(\epsilon, x)) &\simeq \tau^\mu(\delta + x^\nu + \epsilon X^\nu(x)) \\ &\simeq x^\mu + \epsilon X^\mu(x) + \delta Y^\mu(x^\nu + \epsilon X^\nu) \\ &\simeq x^\mu + \epsilon X^\mu(x) + \delta Y^\mu(x) + \epsilon \delta X^\nu(x) \partial_\nu Y^\mu(x). \end{aligned} \quad (3.88)$$

If we instead first move along τ and then move along σ we find

$$\sigma^\mu(\epsilon, \tau(\delta, x)) \simeq x^\mu + \delta Y^\mu(x) + \epsilon X^\mu(x) + \epsilon \delta Y^\nu(x) \partial_\nu X^\mu(x). \quad (3.89)$$

The difference between the two points is proportional to the Lie bracket

$$\tau^\mu(\delta, \sigma(\epsilon, x)) - \sigma^\mu(\epsilon, \tau(\delta, x)) = \epsilon \delta [X, Y]^\mu. \quad (3.90)$$

The Lie bracket measures the failure of the parallelogram in figure 9 to close and therefore for the two flows to commute. It is easy to see that

$$\mathcal{L}_X Y = [X, Y] = 0 \iff \sigma(s, \tau(t, x)) = \tau(t, \sigma(s, x)). \quad (3.91)$$




Figure 9: Moving first along the flow σ and then the flow τ or first along τ and then along σ we find that we may not end up at the same point. The difference is measured by the failure of the Lie bracket to vanish.

Lie Derivative for one-forms: We now want to define the Lie derivative of a one-form $\omega \in \Omega^1(M)$ along X . This time we need to use the pull-back, and the Lie derivative of the one-form ω is

$$\mathcal{L}_X \omega \equiv \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} \left[(\sigma_\epsilon)^* \omega|_{\sigma_\epsilon(x)} - \omega|_x \right], \quad (3.92)$$

where $\omega|_x \in T_x^*(M)$ is ω at x . Introducing coordinates such that $\omega = \omega_\mu dx^\mu$, we have

$$(\sigma_\epsilon)^* \omega|_{\sigma_\epsilon(x)} = \omega_\mu(x) dx^\mu + \epsilon \left[X^\nu(x) \partial_\nu \omega_\mu(x) + \partial_\mu X^\nu(x) \omega_\nu(x) \right] dx^\mu, \quad (3.93)$$

which leads to

$$\mathcal{L}_X \omega = (X^\nu \partial_\nu \omega_\mu + \partial_\mu X^\nu \omega_\nu) dx^\mu. \quad (3.94)$$

This remains a one-form, that is $\mathcal{L}_X \omega \in T_x^*(M)$ since it is the difference of two one-forms at the same point.

This may also be extended to functions f on M . Then

$$\begin{aligned}\mathcal{L}_X f &\equiv \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} [f(\sigma_\epsilon(x)) - f(x)] \\ &= \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} [f(x^\mu + \epsilon X^\mu(x)) - f(x^\mu)] \\ &= X^\mu(x) \frac{\partial f}{\partial x^\mu} = X[f],\end{aligned}\tag{3.95}$$

which is just the usual directional derivative of f along X .

To extend this to more general tensors we need the following result:

Exercise 3.6: Properties of the Lie Derivative on tensors

1. Show that the Lie derivative satisfies:

$$\mathcal{L}_X(t_1 + t_2) = \mathcal{L}_X t_1 + \mathcal{L}_X t_2,\tag{3.96}$$

where t_1 and t_2 are tensor fields of the same type.

2. Show that

$$\mathcal{L}_X(t_1 \otimes t_2) = (\mathcal{L}_X t_1) \otimes t_2 + t_1 \otimes (\mathcal{L}_X t_2),\tag{3.97}$$

with t_1 and t_2 tensors of arbitrary type.

3.4 Differential forms

Not all tensors are created equally, some will play a more prominent role than others. One class of interesting tensors are the p -forms, these are totally anti-symmetric $(0, p)$ tensor fields. To define them we must recall the definition of the anti-symmetrisation of a tensor. Consider $\omega \in \mathcal{T}_p^{(0,r)}(M)$, then its (total) anti-symmetrisation is given by:

$$A[\omega(X_1, \dots, X_r)] = \frac{1}{r!} \sum_{\sigma \in S_r} \text{sign}(\sigma) \omega(X_{\sigma(1)}, \dots, X_{\sigma(r)}).\tag{3.98}$$

with $\text{sign}(\sigma) = +1$ for an even permutation and -1 for an odd permutation.

Definition 26 (Differential form) *A differential form of order r , or more succinctly an r -form, is a totally anti-symmetric tensor of type $(0, r)$.*

Definition 27 (Wedge product) *The Wedge product \wedge of r one-forms is defined to be the totally anti-symmetric tensor product of the one-forms*

$$dx^{\mu_1} \wedge dx^{\mu_2} \wedge \dots \wedge dx^{\mu_r} \equiv \sum_{\sigma \in S_r} \text{sign}(\sigma) dx^{\mu_{\sigma(1)}} \otimes dx^{\mu_{\sigma(2)}} \otimes \dots \otimes dx^{\mu_{\sigma(r)}}.\tag{3.99}$$

Thus

$$dx^\mu \wedge dx^\nu = dx^\mu \otimes dx^\nu - dx^\nu \otimes dx^\mu. \quad (3.100)$$

The wedge product satisfies the following conditions

- $dx^{\mu_1} \wedge \dots \wedge dx^{\mu_r} = 0$ if some index is repeated.
- $dx^{\mu_1} \wedge \dots \wedge dx^{\mu_r} = \text{sign}(\sigma) dx^{\mu_{\sigma(1)}} \wedge \dots \wedge dx^{\mu_{\sigma(r)}}$.
- $dx^{\mu_1} \wedge \dots \wedge dx^{\mu_r}$ is linear in each dx^μ .

We will denote the vector space of r -forms at the point $p \in M$ by $\Omega_p^r(M)$, (another common notation is $\Lambda_p^{(r)}(M)$), a basis is provided by the set of all wedge products in (3.99). We can then expand an element of $\Omega_p^r(M)$ as

$$\omega = \frac{1}{r!} \omega_{\mu_1 \dots \mu_r} dx^{\mu_1} \wedge \dots \wedge dx^{\mu_r}, \quad (3.101)$$

where $\omega_{\mu_1 \dots \mu_r}$ are taken to be totally anti-symmetric. Since there are $\binom{m}{r}$ choices of the set $\{\mu_1, \dots, \mu_r\}$ out of $(1, 2, \dots, m = \dim(M))$ the dimension of the vector space $\Omega_p^{(r)}(M)$ is

$$\binom{m}{r} = \frac{m!}{r!(m-r)!}. \quad (3.102)$$

We take $\Omega_p^0(M) = \mathbb{R}$ and it follows that $\Omega_p^1(M) = T_p^*(M)$ from before. Moreover, since we are anti-symmetrising all the indices if r exceeds $m = \dim(M)$ then it vanishes identically. Furthermore we have the identity $\binom{m}{r} = \binom{m}{m-r}$ and it follows that $\dim \Omega_p^r(M) = \dim \Omega_p^{m-r}(M)$. Since $\Omega_p^r(M)$ is a vector space it is isomorphic to $\Omega_p^{(r-m)}(M)$.¹⁶

3.4.1 Exterior product

Definition 28 (Exterior product) *The exterior product, \wedge , is defined to be the following map $\wedge : \Omega_p^q(M) \times \Omega_p^r(M) \rightarrow \Omega_p^{q+r}(M)$. Its action follows by trivial extension of the wedge product defined above. Let $\omega \in \Omega_p^q(M)$ and $\xi \in \Omega_p^r(M)$ be an q -form and an r -form respectively. The action of the $(q+r)$ -form $\omega \wedge \xi$ on $q+r$ vectors V_i is*

$$(\omega \wedge \xi)(V_1, \dots, V_{q+r}) = \frac{1}{q!r!} \sum_{\sigma \in S_{q+r}} \text{sign}(\sigma) \omega(V_{\sigma(1)}, \dots, V_{\sigma(q)}) \xi(V_{\sigma(q+1)}, \dots, V_{\sigma(q+r)}). \quad (3.103)$$

¹⁶When the manifold is equipped with a metric the isomorphism is provided by the Hodge star operation \star . We will see the Hodge star later in section 4.1.3.

It follows that if $q + r > m = \dim(M)$ then $\omega \wedge \xi$ vanishes. With this product we can define and algebra

$$\Omega_p^*(M) \equiv \Omega_p^0(M) \oplus \Omega_p^1(M) \oplus \dots \oplus \Omega_p^m(M). \quad (3.104)$$

Example 3.8: Wedge product

Let us take \mathbb{R}^3 with coordinates (x, y, z) and consider the forms $\omega_1 = f(x)dx + g(x)dy$, $\omega_2 = \sin(x - z)dy \wedge dz$ and $\omega_3 = e^{x+y+z}dz$ then we have:

$$\begin{aligned} \omega_1 \wedge \omega_2 &= f(x) \sin(x - z)dx \wedge dy \wedge dz, \\ \omega_1 \wedge \omega_3 &= e^{x+y+z}f(x)dx \wedge dz + e^{x+y+z}g(x)dy \wedge dz, \\ \omega_2 \wedge \omega_3 &= 0. \end{aligned} \quad (3.105)$$

Exercise 3.7: Properties of the Wedge product

From the properties of the wedge product show that for $\xi \in \Omega_p^q(M)$, $\eta \in \Omega_p^r(M)$ and $\omega \in \Omega_p^s(M)$ that

$$\begin{aligned} \xi \wedge \eta &= (-1)^{qr} \eta \wedge \xi, \\ \xi \wedge \xi &= 0 \quad \text{if } q \text{ odd}, \\ (\xi \wedge \eta) \wedge \omega &= \xi \wedge (\eta \wedge \omega). \end{aligned} \quad (3.106)$$

We may assign an r -form smoothly at each point on a manifold M . We denote the space of smooth r -forms on M by $\Omega^r(M)$, and take $\Omega^0(M) = \mathcal{F}(M)$ to be the space of smooth functions.

3.4.2 Exterior derivative

A useful map between p -forms and $p + 1$ -forms is the exterior derivative:

Definition 29 (Exterior Derivative) *The exterior derivative d_r is a map $\Omega^r(M) \rightarrow \Omega^{r+1}(M)$, whose action on an r -form is:*

$$\begin{aligned} d\omega(X_1, \dots, X_{r+1}) &= \sum_{i=1}^r (-1)^{i+1} X_i \omega(X_1, \dots, \hat{X}_i, \dots, X_{r+1}) \\ &\quad + \sum_{i < j} (-1)^{i+j} \omega([X_i, X_j], X_1, \dots, \hat{X}_i, \dots, \hat{X}_j, \dots, X_{r+1}), \end{aligned} \quad (3.107)$$

where the hats denote that this term should be removed.

We can write this in coordinates, consider the r -form

$$\omega = \frac{1}{r!} \omega_{\mu_1 \dots \mu_r} dx^{\mu_1} \wedge \dots \wedge dx^{\mu_r}, \quad (3.108)$$

then the exterior derivative

$$d_r \omega = \frac{1}{r!} \left(\frac{\partial}{\partial x^\nu} \omega_{\mu_1 \dots \mu_r} \right) dx^\nu \wedge dx^{\mu_1} \wedge \dots \wedge dx^{\mu_r}. \quad (3.109)$$

It is common to drop the r subscript and simply write d , we will do this from now on. The wedge product automatically anti-symmetrises the coefficient so it is indeed a $(r + 1)$ -form that we obtain.

Exercise 3.8: Exterior derivative property

Show that for $\xi \in \Omega_p^q(M)$, $\eta \in \Omega_p^r(M)$ we have

$$d(\xi \wedge \eta) = d\xi \wedge \eta + (-1)^q \xi \wedge d\eta. \quad (3.110)$$

Example 3.9: Exterior Derivative

Let us take \mathbb{R}^3 with coordinates (x, y, z) . The generic r -forms are

$$\begin{aligned} \omega_0 &= f(x, y, z), \\ \omega_1 &= \omega_x(x, y, z)dx + \omega_y(x, y, z)dy + \omega_z(x, y, z)dz, \\ \omega_2 &= \omega_{xy}(x, y, z)dx \wedge dy + \omega_{yz}(x, y, z)dy \wedge dz + \omega_{zx}(x, y, z)dz \wedge dx, \\ \omega_3 &= \omega_{xyz}(x, y, z)dx \wedge dy \wedge dz. \end{aligned} \quad (3.111)$$

The exterior derivative of these forms is

$$\begin{aligned} d\omega_0 &= \frac{\partial}{\partial x} f(x, y, z)dx + \frac{\partial}{\partial y} f(x, y, z)dy + \frac{\partial}{\partial z} f(x, y, z)dz, \\ d\omega_1 &= \left(\frac{\partial}{\partial x} \omega_y - \frac{\partial}{\partial y} \omega_x \right) dx \wedge dy + \left(\frac{\partial}{\partial y} \omega_z - \frac{\partial}{\partial z} \omega_y \right) dy \wedge dz + \left(\frac{\partial}{\partial z} \omega_x - \frac{\partial}{\partial x} \omega_z \right) dz \wedge dx, \\ d\omega_2 &= \left(\frac{\partial}{\partial x} \omega_{yz} + \frac{\partial}{\partial y} \omega_{zx} + \frac{\partial}{\partial z} \omega_{xy} \right) dx \wedge dy \wedge dz, \\ d\omega_3 &= 0. \end{aligned} \quad (3.112)$$

In the usual 3d vector calculus you may identify these as ‘grad’ for d acting on the scalar, ‘curl’ for the one-form and the ‘divergence’ for the two-form.

From either the coordinate free expression (3.107) or the one using the coordinates in (3.109), we can prove the important result that

$$d^2 = 0, \quad (d_{r+1} d_r = 0). \quad (3.113)$$

Using the coordinate form (3.109) we find

$$d^2 \omega = \frac{1}{r!} \frac{\partial^2}{\partial x^\nu \partial x^\sigma} \omega_{\mu_1 \dots \mu_r} dx^\nu \wedge dx^\sigma \wedge dx^{\mu_1} \wedge \dots \wedge dx^{\mu_r}. \quad (3.114)$$

Using that the derivative term is symmetric in $\nu\sigma$ while the wedge product is anti-symmetric in these indices it follows that this vanishes. Since $d^2 = 0$ it follows that an exact form is always closed, though the converse need not be true. The failure of a closed form to be exact tells us interesting information about the topology of the underlying manifold.

Aside: Cohomology

The exterior derivative induces the sequence

$$0 \xrightarrow{i} \Omega^0(M) \xrightarrow{d_0} \Omega^1(M) \xrightarrow{d_1} \dots \xrightarrow{d_{m-1}} \Omega^m(M) \xrightarrow{d_m} 0, \quad (3.115)$$

with i the inclusion map. This is known as the *de Rahm complex*. We let the set of all closed r -forms on M be denoted by $Z^r(M)$, so that for $d_r : \Omega^r(M) \rightarrow \Omega^{r+1}(M)$, $\ker(d_r) = Z^r(M)$, and denote the set of all exact r -forms to be $B^r(M)$, i.e. the B^r is the image of $\Omega^{r-1}(M)$ under $d^{r-1} : \Omega^{r-1}(M) \rightarrow \Omega^r(M)$. Then the r th de-Rahm cohomology group is defined to be

$$H^r(M) = Z^r(M)/B^r(M). \quad (3.116)$$

This is the dual space of the *homology group*, though we will not have time to consider either of these. The cohomology groups tell us important information about a manifold, their dimensions are topological invariants. Let $b^r = \dim(H^r(M))$, these are known as the *Betti numbers* of the manifold and are always finite. For a connected manifold one always has $b_0 = 1$, these are just the constant functions. The higher Betti numbers are non-zero when the manifold has some interesting topology, for example on a round two-sphere we have: $b^0(S^2) = 1 = b^2(S^2)$ and $b^1(S^2) = 0$. The Euler characteristic of a manifold can be given in terms of the Betti numbers by

$$\chi(M) = \sum_{r=0}^m (-1)^r b^r(M). \quad (3.117)$$

For the S^2 we find $\chi(S^2) = 1 - 0 + 1 = 2$ which is the correct result!

We have seen that every exact form is closed, however not every closed form is exact, instead we have:

Theorem 3 (Poincaré’s lemma) *If a coordinate neighbourhood U of a manifold M is contractible to a point $p \in M$, any closed form on U is also exact. In particular on $M = \mathbb{R}^m$, closed implies exact.*

Since we have been mapping our manifolds to \mathbb{R}^m this says that for a general manifold any closed form is *locally exact*. That is if ω is a closed r -form, then in any neighbourhood $U \subset M$ it is always possible to find $\eta \in \Omega^{r-1}(M)$ such that $\omega = d\eta$ on U . Since we cannot generally cover the manifold with a single coordinate patch, it may not be possible to find

such an η everywhere on M . It is for this reason that we say the form is only *locally* exact rather than globally exact.

Example 3.10: Exact forms

Let us consider some examples.

- Consider $M = \mathbb{R}$. We can take a generic one-form to be $\omega = f(x)dx$, with $f(x)$ some function. This is trivially closed since it is a top form, it is also exact since we can write

$$g(x) = \int_0^x dx' f(x'), \quad (3.118)$$

such that $\omega = dg(x)$. This is all very boring because of the Poincaré's lemma.

- Now consider a circle, S^1 . We can obtain a circle by looking at the phase $e^{i\phi} \in \mathbb{C}$. We can introduce the one-form $\omega = d\phi$. Clearly this is once again closed since it is a top form, and from the way that it is written it seems that it must be exact once again, this however is not correct. The caveat is that ϕ is *not* a good coordinate everywhere on S^1 , since it is not single valued, remember that we needed at least 2 patches on S^1 . As such ϕ is not a good smooth function and so it is not a zero-form. Therefore $d\phi$ is closed but not exact.

- Next consider $M = \mathbb{R}^2$. The Poincaré lemma ensures that all closed forms are exact. What happens if we remove a point? Consider instead $\mathbb{R}^2 - \{0, 0\}$ and the one-form

$$\omega = -\frac{y}{x^2 + y^2} dx + \frac{x}{x^2 + y^2} dy. \quad (3.119)$$

This is not a smooth one-form on \mathbb{R}^2 , since it blows up at the origin, however by restricting to $\mathbb{R}^2 - \{0, 0\}$ we obtain a smooth one-form. A simple computation shows that ω is closed, but is it exact? If such a smooth function exists such that $\omega = df$ then the function f must satisfy:

$$\frac{\partial f}{\partial x} = -\frac{y}{x^2 + y^2}, \quad \frac{\partial f}{\partial y} = \frac{x}{x^2 + y^2}. \quad (3.120)$$

The solution is

$$f(x, y) = \arctan\left(\frac{y}{x}\right) + \text{constant}, \quad (3.121)$$

so have we found an exact form? The answer is no, this is not a smooth function everywhere on $\mathbb{R}^2 - \{0, 0\}$, along the line $x = 0$ it is ill-defined, and so ω is *not* exact. We see that removing a single point makes a big difference and closed no longer implies exact. A similar story holds for \mathbb{R}^3 and this is how magnetic monopoles sneak back into physics despite being forbidden by Maxwell's equations. See your favourite course on electromagnetism.

3.4.3 Interior product

We can now go from $\Omega^r(M) \rightarrow \Omega^{r+1}(M)$, what about the other way around? To do this we have to define the *Interior product*.

Definition 30 (Interior product) *Let Y be a vector field and $\omega \in \Omega^r(M)$ then*

$$i_Y \omega(X_1, \dots, X_{r-1}) \equiv \omega(Y, X_1, \dots, X_{r-1}). \quad (3.122)$$

If we introduce coordinates: $Y = Y^\mu \frac{\partial}{\partial x^\mu}$ then

$$i_Y \omega = \frac{1}{(r-1)!} Y^\nu \omega_{\nu \mu_1 \dots \mu_{r-1}} dx^{\mu_1} \wedge \dots \wedge dx^{\mu_{r-1}}. \quad (3.123)$$

Example 3.11:

Let us take \mathbb{R}^3 again with coordinates (x, y, z) , and the usual coordinate basis, then we have

$$i_{e_x}(dx \wedge dy) = dy, \quad i_{e_x}(dy \wedge dz) = 0, \quad i_{e_x}(dz \wedge dx) = -dz. \quad (3.124)$$

Using the interior product and exterior derivative gives a simple way of computing the Lie derivative of a form along the vector field X . We have for any r -form ω and vector field X that:

$$\mathcal{L}_X \omega = (di_X + i_X d)\omega. \quad (3.125)$$

Exercise 3.9: Interior product identities

Show that the interior product satisfies the following:

$$\begin{aligned} i_X^2 &= 0, \\ i_X(\omega \wedge \eta) &= i_X \omega \wedge \eta + (-1)^r \omega \wedge i_X \eta, \\ \mathcal{L}_X i_X \omega &= i_X \mathcal{L}_X \omega. \end{aligned} \quad (3.126)$$

Hamiltonian mechanics in differential geometry We can now combine some of the differential geometry we have learnt so far to reformulate classical Hamiltonian mechanics. Recall that in classical mechanics the phase space is a manifold M parametrised by coordinates (q^i, p_j) where q^i are the positions of particles and p_j their momenta. Note that M must be even dimensional here. The Hamiltonian $H(q, p)$ is a function on M and Hamilton's equations are

$$\dot{q}^i = \frac{\partial H}{\partial p_i}, \quad \text{and} \quad \dot{p}_i = -\frac{\partial H}{\partial q^i}. \quad (3.127)$$

Phase space comes equipped with the *Poisson bracket*, defined on a pair of functions f, g to act as

$$\{f, g\} = \frac{\partial f}{\partial q^j} \frac{\partial g}{\partial p_j} - \frac{\partial f}{\partial p_j} \frac{\partial g}{\partial q^j}, \quad (3.128)$$

from which the time evolution of a function is

$$\dot{f} = \{f, H\}, \quad (3.129)$$

with H the Hamiltonian. To obtain Hamilton's equations one should input $f = q^i$ and $f = p_i$ into the above.

Underlying this structure are forms. The key idea behind this is to convert the scalar function H into a vector field X_H on M . Particles will then follow trajectories which are the integral curves generated by X_H . To convert the scalar into a vector we introduce the *symplectic two-form* ω . This is a two-form which is closed $d\omega = 0$ and is non-degenerate, $\omega \wedge \omega \wedge \dots \wedge \omega \neq 0$. A manifold equipped with such a two-form is called a *symplectic manifold*.

Any two-form provides a map $\omega : T_p(M) \rightarrow T_p^*(M)$, since given a vector field X we can simply take the inner product with ω to obtain a one-form, $i_X\omega$. For our purposes we want to go in the opposite direction, we want to convert a scalar function into a vector field. This is possible if the map $\omega : T_p(M) \rightarrow T_p^*(M)$ is an isomorphism. This is equivalent to ω being non-degenerate. In this case we can define a vector field X_H via

$$i_{X_H}\omega = -dH. \quad (3.130)$$

In coordinate notation we have

$$X_H^\mu \omega_{\mu\nu} = -\partial_\nu H. \quad (3.131)$$

If we take the inverse to be $\omega^{\mu\nu}$ so that $\omega^{\mu\nu}\omega_{\nu\rho} = \delta_\rho^\mu$, then

$$X_H^\mu = \omega^{\mu\nu} \partial_\nu H. \quad (3.132)$$

The integral curves generated by X_H obey

$$\frac{dx^\mu(t)}{dt} = X_H^\mu = \omega^{\mu\nu} \partial_\nu H. \quad (3.133)$$

These are the general form of Hamilton's equations, just written without reference to canonical coordinates. If we let $x^\mu = (q^i, p_j)$ and choose the symplectic form to have block diagonal form

$$\omega^{\mu\nu} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \quad \Leftrightarrow \quad \omega = dp^i \wedge dq_i \quad (3.134)$$

then the integral curves reduce precisely to Hamilton's equations (3.127).

To define the Poisson structure, we first note that we can repeat the map for obtaining a vector from a scalar for any function f , to obtain a vector field X_f . Then

$$\{f, g\} = \omega(X_f, X_g) = -\omega(X_g, X_f). \quad (3.135)$$

This may be written in a multitude of different ways, we have

$$\{f, g\} = -i_{X_f}\omega(X_g) = df(X_g) = X_g(f). \quad (3.136)$$

It follows that the equation of motion in Poisson bracket structure is then

$$\dot{f} = \{f, H\} = X_H(f) = \mathcal{L}_{X_H}f. \quad (3.137)$$

We see that the Lie derivative along X_H generates time evolution!

So far we have not explained why the symplectic two-form was taken to be closed. This is required in order for the Poisson bracket to obey the Jacobi identity. It is also a necessary (and sufficient) condition for the symplectic form to be invariant under Hamiltonian flow.

3.4.4 Integration

We have learnt how to differentiate on a manifold using a vector field X , we now turn our attention to integration? The key question we want to ask is what can we integrate on a manifold M and how? The answer turns out to be our friends the differential forms.

Definition 31 (Orientation) *To begin we need to define an orientation on a manifold. Let M be a connected m -dimensional differentiable manifold. At a point $p \in M$ the tangent space $T_p(M)$ is spanned by the basis $\{e_\mu\} = \{\frac{\partial}{\partial x^\mu}\}$ where x^μ is the local coordinate on the chart U_i which contains p . Take U_j to be another chart such that $U_i \cap U_j \neq \emptyset$ and such that $p \in U_i \cap U_j$. Then the tangent space $T_p(M)$ is spanned by both $\{e_\mu\}$ or $\{\tilde{e}_\nu\} = \{\frac{\partial}{\partial y^\nu}\}$. The change of basis is*

$$\tilde{e}_\nu = \frac{\partial x^\mu}{\partial y^\nu} e_\mu \equiv \Lambda^\mu{}_\nu e_\mu. \quad (3.138)$$

If $\det(\Lambda) > 0$ on $U_i \cap U_j$, the two bases $\{e_\mu\}$ and $\{\tilde{e}_\nu\}$ are said to define the same orientation on $U_i \cap U_j$. If on the other hand $\det(\Lambda) < 0$ then they define the opposite orientation.

Definition 32 (Orientable) *Let M be a connected manifold covered by $\{U_i\}$. The manifold M is orientable if for any overlapping charts U_i, U_j there exist local coordinates $\{x^\mu\}$ for U_i and $\{y^\nu\}$ for U_j such that $\det \Lambda = \det(\frac{\partial x^\mu}{\partial y^\nu}) > 0$. If M is non-orientable, Λ cannot be made positive in all intersections of charts.*

An example of a non-orientable manifold is the Möbius strip, see figure 10. To construct a Möbius strip take two rectangles and glue them together with a twist of π on one of the edges to glue.




Figure 10: To construct the Möbius strip we glue two rectangles together: A with A' and B with B' . When joining A with A' we twist by π . The coordinate transformation on the A, A' intersection $y^1 = x^1$ and $y^2 = -x^2$, which has Jacobian -1 and is thus not orientable. We see that the cyclist going around the Möbius strip end up “up-side down” as they travel around the strip.

If an m -dimensional manifold M is orientable there exists an m -form ω which is nowhere vanishing, called the *volume form* or *volume element*. It plays the role of the measure when we integrate a function $f \in \mathcal{F}(M)$ over M . Two volume elements are said to be *equivalent* if there exists a strictly positive function $h \in \mathcal{F}(M)$ such that $\omega = h\omega'$. A negative-definite function $h' \in \mathcal{F}(M)$ gives an inequivalent orientation to M . Therefore for any orientable manifold there are two inequivalent orientations, we may refer to one of them as right-handed and the other as left-handed.

Since the volume form is a top form locally it can be written as

$$\omega = h(x)dx^1 \wedge \dots \wedge dx^m, \quad (3.139)$$

with the requirement that $h(x) \neq 0$. For the volume form to not flip orientation we must be able to patch this over the whole manifold without the handedness changing. Suppose that we have two sets of coordinates x^μ and y^ν on the charts U_i and U_j respectively, then in the

new coordinates we have

$$\omega = h(x) \frac{\partial x^1}{\partial y^{\nu_1}} dy^{\nu_1} \wedge \dots \wedge \frac{\partial x^n}{\partial y^{\nu_m}} dy^{\nu_m} = h(x) \det \left(\frac{\partial x^\mu}{\partial y^\nu} \right) dy^1 \wedge \dots \wedge dy^m, \quad (3.140)$$

which makes clear that we may only define a volume form when the manifold is orientable, since the determinant appears. For the Möbius strip we see that we begin with volume form $\omega = dx \wedge dy$ but as we change charts this becomes $\omega = -dx \wedge dy$ and so ω is not of definite handedness on the Möbius strip.

With our volume form in tow we can now define integration of a function $f : M \rightarrow \mathbb{R}$ over an orientable manifold M . Let us take the volume form to be ω . Then in a coordinate neighbourhood U_i with coordinates x^μ we define the integration of an m -form $f\omega$ to be

$$\int_{U_i} f\omega \equiv \int_{\varphi(U_i)} f(\varphi_i^{-1}(x)) h(\varphi_i^{-1}(x)) dx^1 \dots dx^m. \quad (3.141)$$

Notice that the right-hand side is just ordinary integration we are familiar with, albeit in m variables. Once the integral of f over U_i is defined it can be extended to an integration over all of M by making use of a *partition of unity*.

Definition 33 (Partition of unity) *Take an open covering $\{U_i\}$ on M such that each point of M is covered with a finite number of U_i . If this is always possible we call M paracompact.¹⁷ If a family of differentiable functions $\epsilon_i(p)$ satisfies*

1. $0 \leq \epsilon_i(p) \leq 1$,
2. $\epsilon_i(p) = 0$ if $p \notin U_i$,
3. $\epsilon_1(p) + \epsilon_2(p) + \dots = 1$ for every point $p \in M$.

The family $\{\epsilon_i(p)\}$ is called a *partition of unity for the covering $\{U_i\}$* .

From condition (3) it follows that

$$f(p) = \sum_i f(p) \epsilon_i(p) = \sum_i f_i(p), \quad f_i(p) \equiv \epsilon_i(p) f(p). \quad (3.142)$$

Hence given a point $p \in M$ assumed paracompactness ensures that there are only a finite number of terms in the summation over i , this was one of the magical properties we imposed but forgot about. For each of the $f_i(p)$ we may define the integral over U_i via (3.141), and therefore we have

$$\int_M f\omega \equiv \sum_i \int_{U_i} f_i\omega. \quad (3.143)$$

¹⁷We will assume this is the case whenever we integrate something in this course.

Though a different choice of atlas gives a different set of coordinates and a different partition of unity the integral as defined above stays the same.

Example 3.12: Integrating on a circle

Let us consider integrating a function on the circle. Let us take the atlas as given in (3.8) and (3.9). Let $U_1 = S^1 - \{(1, 0)\}$ and $U_2 = S^1 - \{(-1, 0)\}$. Then we may give a partition of unity by fixing $\epsilon_1(\theta) = \sin^2 \frac{\theta}{2}$ and $\epsilon_2(\theta) = \cos^2 \frac{\theta}{2}$. Note that $\epsilon_1(0) = 0$ and $\epsilon_2(\pi) = 0$ and therefore they vanish at the removed points as required. Moreover $\epsilon_1(\theta) + \epsilon_2(\theta) = 1$ as required. Thus $\{\epsilon_i(\theta)\}$ furnishes us with a partition of unity for the atlas $\{U_i\}$. Let us integrate the function $f = \cos^2 \theta$. Of course we know

$$\int_0^{2\pi} d\theta \cos^2 \theta = \pi, \tag{3.144}$$

but we should check with our partition of unity that we obtain the same result. We find

$$\int_{S^1} d\theta \cos^2 \theta = \int_0^{2\pi} d\theta \sin^2 \frac{\theta}{2} \cos^2 \theta + \int_{-\pi}^{\pi} d\theta \cos^2 \frac{\theta}{2} \cos^2 \theta = \frac{1}{2}\pi + \frac{1}{2}\pi = \pi. \tag{3.145}$$

So far we have left the function $h(x)$ appearing in the volume-form arbitrary. Since this gets multiplied by the Jacobian it changes between different coordinate patches and therefore there is no canonical way to pick this. Once we endow the manifold with a metric, as we are required to do in GR, there is a canonical choice that we can make.

We can also integrate forms over sub-manifolds of M , rather than the full manifold. A manifold Σ with dimension $k < n$ is a *sub-manifold* of M if we can find a map $\sigma : \Sigma \rightarrow M$ which is one-to-one and $\sigma_* : T_p(\Sigma) \rightarrow T_{\sigma(p)}(M)$ is also one-to-one. We can then integrate a k -form ω on M over a k -dimensional sub-manifold Σ by pulling the form back to Σ :

$$\int_{\sigma(\Sigma)} \omega = \int_{\Sigma} \sigma^* \omega. \tag{3.146}$$

For example consider a one-form A living on M and take C to be a one-dimensional manifold in M . We can introduce a map $\sigma : C \rightarrow M$ which defines a non-intersecting curve $\sigma(C)$ which is a sub-manifold of M . We can then pull-back A onto the curve and integrate to obtain,

$$\int_{\sigma(C)} A = \int_C \sigma^* A. \tag{3.147}$$

Let the curve trace out a path $x^\mu(\tau)$ in M then, in coordinates this reads

$$\int_C \sigma^* A = \int d\tau A_\mu(x) \frac{dx^\mu}{d\tau}, \tag{3.148}$$

which is precisely the way in which a worldline of a particle couples to the electromagnetic field.

Until now our focus has been on smooth manifolds without boundary. We saw that this can be extended to manifolds with a boundary in section 3.1. There we have charts $\varphi : M \rightarrow U_i$ where U_i is an open subset of $\mathbb{R}^m = \{(x^1, \dots, x^m) | x^m \geq 0\}$. The boundary is denoted by ∂M , and is the sub-manifold fixed by $x^m = 0$.

Theorem 4 (Stokes' Theorem) *For a manifold M with a boundary, for any $(m-1)$ -form ω we have*

$$\int_M d\omega = \int_{\partial M} \omega. \quad (3.149)$$

Stokes' theorem is the mother of all integral theorems. You may be familiar with the divergence theorem, Green's theorem, etc., this is the generalisation of those.

Exercise 3.10: Stokes' Theorem

Show that this reduces to Stokes' theorem on \mathbb{R}^3 .

We will see integrals later in the course when we compute the mass of the Schwarzschild black hole.

3.5 Summary

- A **manifold** is a space that locally looks like \mathbb{R}^n , even if its global structure is more complicated.
- A **chart** assigns coordinates to a region of the manifold, and different charts are related by smooth transition functions.
- At each point $p \in M$, the **tangent space** $T_p(M)$ is a vector space consisting of tangent vectors at that point.
- A **vector field** assigns a tangent vector to each point on the manifold.
- A **one-form** is a linear map $\omega : T_p(M) \rightarrow \mathbb{R}$; the dual basis $\{dx^\mu\}$ satisfies

$$dx^\mu \left(\frac{\partial}{\partial x^\nu} \right) = \delta^\mu_\nu. \quad (3.150)$$

- A **tensor** is a multilinear map built from vectors and one-forms; its components depend on coordinates, but the tensor itself is geometric.
- The **pushforward** ϕ_* maps vectors between manifolds, while the **pullback** ϕ^* maps one-forms in the opposite direction.

- A **flow** generated by a vector field describes how points move along its integral curves.
- The **Lie derivative** \mathcal{L}_X measures how a tensor field changes along the flow generated by a vector field X :

$$\mathcal{L}_X f = X(f), \quad \mathcal{L}_X Y = [X, Y]. \quad (3.151)$$

- **Differential forms** are totally antisymmetric tensors that can be integrated over manifolds.
- The **exterior derivative** d generalises the gradient and satisfies $d^2 = 0$.
- **Stokes' theorem** relates integrals over a manifold and its boundary:

$$\int_M d\omega = \int_{\partial M} \omega, \quad (3.152)$$

and is the mother of all integral theorems (Divergence, Greens' Stokes').

Key Points

Differential geometry allows us to do calculus on curved spaces by combining local coordinate descriptions with coordinate-independent geometric objects. We have generalisations of all the usual Euclidean operators (divergence, curl etc.) along with other new operators like the Lie derivative.

4 Riemannian geometry

We now have all the necessary pre-requisites to introduce the most valuable player of general relativity: the metric. The introduction of a metric brings a whole slew of new objects that we can define. Here we will continue to talk about Riemannian geometry, this is spaces with Euclidean signature, whereas what we really want to consider is Lorentzian geometry.

4.1 The metric

Definition 34 (Riemannian metric) *Let M be a differentiable manifold.*

A Riemannian metric g on M is a type $(0, 2)$ tensor field on M which at each point $p \in M$ satisfies:

- *Symmetric: $g_p(X, Y) = g_p(Y, X)$,*
- *$g_p(X, X) \geq 0$ with equality iff $X = 0$,*

with $X, Y \in T_p(M)$.

A tensor field g of type $(0, 2)$ is a pseudo-Riemannian metric if it satisfies the first condition and

- *Non-degenerate. If for any $p \in M$ $g_p(X, Y) = 0$ for all $Y \in T_p(M)$ then $X|_p = 0$,*

We may extend the tensor g_p over the full manifold. With a choice of coordinates we can write the metric as

$$g = g_{\mu\nu}(x)dx^\mu \otimes dx^\nu . \quad (4.1)$$

This defines the metric to be a smooth tensor field over our whole manifold, that is a multi-linear map from $T(M) \times T(M) \rightarrow \mathcal{F}(M)$. We will often write this as the line element ds^2 ,

$$ds^2 = g_{\mu\nu}(x)dx^\mu dx^\nu , \quad (4.2)$$

in particular removing the tensor product. We can do this unambiguously because of the symmetry property of the metric. This also captures our intuitive understanding of the infinitesimal distance being measured by the infinitesimal coordinate separations dx^μ weighted by the metric.

One can extract out the components by evaluating the metric on a pair of basis elements

$$g_{\mu\nu}(x) = g\left(\frac{\partial}{\partial x^\mu}, \frac{\partial}{\partial x^\nu}\right). \quad (4.3)$$

We may view $g_{\mu\nu}$ as a matrix, which by the symmetry property above is symmetric. This implies that the matrix can be diagonalised with real eigenvalues. If there are i positive eigenvalues and j negative eigenvalues the pair (i, j) is called the *index* of the metric. If $j = 1$ the metric is called a *Lorentz metric*, for $j = 0$ we have a *Euclidean* metric. The number of negative entries is called the *signature* and by Sylvester's law of inertia¹⁸, this is independent of the choice of basis. By an abuse of notation we will often call this symmetric matrix the metric when really this is just components of the metric tensor field in some coordinate basis.

For most applications of differential geometry, we are interested in manifolds with signature 0, i.e. a Riemannian manifold. The simplest example which you are probably familiar with, though maybe not in this language, is the metric on Euclidean space \mathbb{R}^m , which in Cartesian coordinates has the metric

$$g = dx^1 \otimes dx^1 + \dots + dx^m \otimes dx^m, \quad (4.4)$$

which in components reads $g_{\mu\nu} = \delta_{\mu\nu}$.

4.1.1 Riemannian metric

A general Riemannian metric is a useful object to have in one's tool belt. It gives us a way of measuring the length of a vector X at each point

$$|X| = \sqrt{g(X, X)}. \quad (4.5)$$

Moreover we may measure the angle between two vectors

$$g(X, Y) = |X||Y| \cos \theta. \quad (4.6)$$

Furthermore, it can be used to measure the distance between two points p and q along a curve in M . For the curve $\sigma : [a, b] \rightarrow M$ with $\sigma(a) = p$ and $\sigma(b) = q$ the distance between the two points along the curve is

$$d(p, q) = \int_a^b dt \sqrt{g(X, X)|_{\sigma(t)}}, \quad (4.7)$$

where X is the tangent vector field to the curve. If the curve has coordinates $x^\mu(t)$ then $X^\mu = \frac{dx^\mu}{dt}$ and the distance is

$$d(p, q) = \int_a^b dt \sqrt{g_{\mu\nu} \frac{dx^\mu(t)}{dt} \frac{dx^\nu(t)}{dt}}. \quad (4.8)$$

Importantly this distance does not depend on the parametrisation of the curve, and only on the curve itself.

¹⁸This has nothing to do with inertia, Sylvester just wanted a law of inertia like Newton.

Example 4.1: Metric on a two sphere

A less trivial example is the metric on a unit round two-sphere, denoted S^2 ,

$$ds^2(S^2) = d\theta^2 + \sin^2 \theta d\phi^2. \quad (4.9)$$

Here $\theta \in (0, \pi)$ and $\phi \in [0, 2\pi)$. This is written in a chart which does not cover the full S^2 , to show that this is a smooth tensor field one must define a second patch whose union with the above one covers S^2 . To see this one can first realise the S^2 by embedding the two sphere in \mathbb{R}^3 via:

$$x^2 + y^2 + z^2 = 1. \quad (4.10)$$

First introduce polar coordinates as

$$x = \sin \theta \cos \phi, \quad y = \sin \theta \sin \phi, \quad z = \cos \theta, \quad (4.11)$$

which define $\theta \in (0, \pi)$ and $\phi \in (0, 2\pi)$ uniquely. This covers all of the S^2 but for the line of longitude $y = 0, x > 0$ and the points $(0, 0, \pm 1)$. We can define a second chart using a different set of polar coordinates:

$$x = -\sin \theta' \cos \phi', \quad y = \cos \theta', \quad z = \sin \theta' \sin \phi', \quad (4.12)$$

where $\theta' \in (0, \pi)$ and $\phi' \in (0, 2\pi)$. The points $(0, \pm 1, 0)$ and the line $z = 0, x < 0$ is not covered however one can see that the union of these two charts covers the S^2 . One can check that this makes S^2 a manifold. Now in the chart the metric we obtain is

$$ds^2(S^2) = d\theta'^2 + \sin^2 \theta' d\phi'^2. \quad (4.13)$$

One can then check this defines a smooth tensor field.

We can use the metric to work out the circumference of the unit circle. Let us first compute the path along the equator. This is the integral curve of the vector field $X = \partial_\phi$ which we computed in example 3.7. The form of the integral curve is

$$\theta(\lambda) = \theta_0, \quad \phi(\lambda) = \lambda, \quad (4.14)$$

and we can compute the length of the curve. We end up back where we are after $\lambda = 2\pi$ and so the path length is:

$$d = \int_0^{2\pi} \sqrt{\dot{\theta}^2 + \sin^2 \theta \dot{\phi}^2} d\lambda = \int_0^{2\pi} \sin \theta_0 d\lambda = 2\pi \sin \theta_0. \quad (4.15)$$

The equator has $\theta = \frac{\pi}{2}$ and therefore we find that the circumference is 2π which is the correct value for a unit sphere. If we wanted to work with a non-unit sphere we should multiply the metric by a factor l^2 with l the radius. It is simple to see that the result in this case would then be $2\pi l$.

Exercise 4.1: Metric on S^2 from pull back

This metric in (4.9) is the pull back of the metric on Euclidean space to the S^2 defined by the polar coordinates. Show this.

The possible issues with the tensor field (4.9) came from the need to change charts since we cannot cover the S^2 with a single chart. The metric on S^2 will appear later in this course and we will simply ignore any subtleties of the regularity of the metric in (4.9) since, as we have just seen, these can be removed by properly considering the different patches of the S^2 .

4.1.2 Lorentzian manifolds

For General Relativity we need to consider Lorentzian manifolds. The simplest example is Minkowski space. This is $\mathbb{R}^{1,m-1}$ equipped with the metric

$$\eta = -dx^0 \otimes dx^0 + dx^1 \otimes dx^1 + \dots + dx^{m-1} \otimes dx^{m-1}, \quad (4.16)$$

which has components $\eta_{\mu\nu} = \text{diag}(-1, 1, \dots, 1)$. Note that on a Lorentzian manifold we take the index to run over $0, 1, \dots, m-1$.

At any point p on a general Lorentzian manifold it is always possible to find an orthonormal basis $\{e_\mu\}$ of $T_p(M)$ such that locally the metric looks like the Minkowski metric

$$g_{\mu\nu}|_p = \eta_{\mu\nu}. \quad (4.17)$$

This is closely related to the equivalence principle we discussed in section 2.2.1, we will discuss the coordinates, known as *normal coordinates* that allow us to do this shortly. The fact that locally the metric looks like Minkowski space allows us to import some of the ideas of special relativity, namely we can classify the elements of $T_p(M)$ into three classes

- $g(X, X) > 0 \rightarrow X$ is *spacelike* ,
- $g(X, X) = 0 \rightarrow X$ is *lightlike* or *null* ,
- $g(X, X) < 0 \rightarrow X$ is *timelike* .

At each point on M we can then draw light cones which are the null tangent vectors at that point. The novelty is that the directions of these light cones can vary smoothly as we move around the manifold. This specifies the causal structure of spacetime which determines which regions of spacetime can interact together.

As in the Riemannian case we can use the metric to determine the length of curves. The nature of a curve is inherited from the nature of its tangent vector. A curve is called *timelike* if its tangent vector is everywhere timelike. We then measure the proper time

$$\tau = \int_a^b d\lambda \sqrt{-g_{\mu\nu} \frac{dx^\mu}{d\lambda} \frac{dx^\nu}{d\lambda}}. \quad (4.18)$$

4.1.3 Why is the metric useful?

The existence of a metric comes with a large number of benefits.

The metric as an isomorphism The metric gives a natural isomorphism between vectors and covectors, $g : T_p(M) \rightarrow T_p^*(M)$ for each p . In a coordinate basis we can write $X = X^\mu \partial_\mu$, and map it to a one-form $X = X_\mu dx^\mu$, with the components given by

$$X_\mu = g_{\mu\nu} X^\nu. \quad (4.19)$$

We will usually say that we use the metric to lower (or raise) an index. What we really mean is that the metric provides an isomorphism between a vector space and its dual. Since g is non-degenerate and is thus invertible we also have the inverse map. We take the inverse of $g_{\mu\nu}$ to be $g^{\mu\nu}$ so that $g^{\mu\nu} g_{\nu\rho} = \delta_\rho^\mu$. This can then be thought of as the components of a symmetric $(2,0)$ tensor

$$\hat{g} = g^{\mu\nu} \partial_\mu \otimes \partial_\nu. \quad (4.20)$$

Then

$$X^\mu = g^{\mu\nu} X_\nu. \quad (4.21)$$

The Volume form The metric also gives a natural volume form on the manifold M rather than the variety of volume forms we had previously. On a Riemannian manifold we take the volume form to be

$$\text{vol}(M) = \sqrt{\det(g_{\mu\nu})} dx^1 \wedge \dots \wedge dx^m, \quad (4.22)$$

and we use the shorthand $\sqrt{\det(g_{\mu\nu})} = \sqrt{g}$. On a Lorentzian manifold the determinant is negative and therefore we take the volume form to be

$$\text{vol}(M) = \sqrt{-g} dx^0 \wedge dx^1 \wedge \dots \wedge dx^{m-1}. \quad (4.23)$$

As it is written it looks coordinate dependent however it is not. To see this recall that if we change coordinates $y = y(x)$ we have (see (3.40))

$$dx^\mu = \frac{\partial x^\mu}{\partial y^\nu} dy^\nu \equiv \Lambda^\mu{}_\nu dy^\nu. \quad (4.24)$$

Then

$$\begin{aligned}
dx^1 \wedge \dots \wedge dx^m &= \Lambda^1_{\nu_1} \dots \Lambda^m_{\nu_m} dy^{\nu_1} \wedge \dots \wedge dy^{\nu_m} \\
&= \sum_{\sigma \in S_m} \Lambda^1_{\sigma(1)} \dots \Lambda^m_{\sigma(m)} dy^{\sigma(1)} \wedge \dots \wedge dy^{\sigma(m)} \\
&= \sum_{\sigma \in S_m} \text{sign}(\sigma) \Lambda^1_{\sigma(1)} \dots \Lambda^m_{\sigma(m)} dy^1 \wedge \dots \wedge dy^m \\
&= \det(\Lambda) dy^1 \wedge \dots \wedge dy^m,
\end{aligned} \tag{4.25}$$

where in the penultimate line we have used the properties of the wedge product and in the last line used the definition of the determinant. The metric components transform as

$$g_{\mu\nu} = \frac{\partial y^\rho}{\partial x^\mu} \frac{\partial y^\sigma}{\partial x^\nu} \tilde{g}_{\rho\sigma}, \tag{4.26}$$

and therefore

$$\det(g_{\mu\nu}) = \det(\tilde{g}_{\mu\nu}) (\det(\Lambda))^{-2}, \tag{4.27}$$

and therefore this cancels with the transformation of the wedge product leaving

$$\text{vol}(M) = \sqrt{|g|} dy^1 \wedge \dots \wedge dy^m, \tag{4.28}$$

which is therefore invariant.

We may rewrite the volume form as

$$\text{vol}(M) = \frac{1}{m!} v_{\mu_1 \dots \mu_m} dx^{\mu_1} \wedge \dots \wedge dx^{\mu_m}, \quad \text{where} \quad v_{\mu_1 \dots \mu_m} = \sqrt{|g|} \epsilon_{\mu_1 \dots \mu_m}. \tag{4.29}$$

Here $\epsilon_{\mu_1 \dots \mu_m}$ is the Levi–Civita symbol which is m -dimensional totally anti-symmetric tensor giving 1 for an even permutation of the indices, -1 for an odd permutation and 0 when an index is repeated. It follows that $v_{\mu_1 \dots \mu_m}$ is a tensor, while $\epsilon_{\mu_1 \dots \mu_m}$ is not, instead it is a tensor density (one needs to multiply by the square root of the determinant to obtain a tensor). Note that we define $\epsilon^{\mu_1 \dots \mu_m}$ to again be the totally anti-symmetric tensor with $\epsilon^{1 \dots m} = 1$, i.e. we do not raise the indices on ϵ with the metric. Instead we have

$$v^{\mu_1 \dots \mu_m} = g^{\mu_1 \nu_1} \dots g^{\mu_m \nu_m} v_{\nu_1 \dots \nu_m} = \pm \frac{1}{\sqrt{|g|}} \epsilon^{\mu_1 \dots \mu_m}. \tag{4.30}$$

Hodge dual On an oriented manifold M we can use the totally anti-symmetric tensor density to define a map which takes a p -form $\omega \in \Omega^p(M)$ to a $(m-p)$ -form $\star\omega \in \Omega^{m-p}(M)$.

We define this map to be

$$(\star\omega)_{\mu_1 \dots \mu_{m-p}} = \frac{1}{p!} \sqrt{|g|} \epsilon_{\mu_1 \dots \mu_{m-p} \nu_1 \dots \nu_p} \omega^{\nu_1 \dots \nu_p}. \tag{4.31}$$

This is called the *Hodge dual* and is independent of coordinates. One can see that it satisfies

$$\star(\star\omega) = \pm(-1)^{p(m-p)}\omega, \quad (4.32)$$

with $+$ for a Riemannian metric and $-$ for a Lorentzian.¹⁹

With the Hodge dual in tow we can define an inner product on each vector space $\Omega^r(M)$. If $\omega, \eta \in \Omega^r(M)$ then we define the inner product

$$\langle \eta, \omega \rangle \equiv \int_M \eta \wedge \star\omega. \quad (4.33)$$

With such an inner product one can look at operators on $\Omega^r(M)$ and their adjoints. The differential operator we have introduced on r -forms is the exterior derivative. For $\omega \in \Omega^r(M)$ and $\alpha \in \Omega^{r-1}(M)$ the adjoint is defined via

$$\langle d\alpha, \omega \rangle = \langle \alpha, d^\dagger\omega \rangle, \quad (4.34)$$

where the adjoint operator $d^\dagger : \Omega^r(M) \rightarrow \Omega^{r-1}(M)$ is given by

$$d^\dagger = \pm(-1)^{m(r+1)-1} \star d \star. \quad (4.35)$$

One can then define a Laplacian $\square : \Omega^r(M) \rightarrow \Omega^r(M)$ defined as²⁰

$$\square = (d + d^\dagger)^2 = dd^\dagger + d^\dagger d. \quad (4.36)$$

It can be defined on both Riemannian manifolds and Lorentzian, however it is only positive definite on Riemannian manifolds. On a function f the Laplacian acts as

$$\square f = -\frac{1}{\sqrt{|g|}} \partial_\nu \left(\sqrt{|g|} g^{\mu\nu} \partial_\mu f \right). \quad (4.37)$$

¹⁹One has actually seen the Hodge dual before, it was just hidden from view. Consider two vectors \vec{a} and \vec{b} in \mathbb{R}^3 , We can take the cross product to obtain a third vector \vec{c} as $\vec{a} \times \vec{b} = \vec{c}$. This however mixes a lot of different objects. This is equivalent in our new language to first use the metric to relate the vectors to one-forms. The cross product it really the wedge product of the two one-forms to give a two-form. We then take the Hodge dual of this two-form to obtain a one-form and then use the metric once again to extract out a vector. This more complicated route is hidden since the metric is just the Kronecker delta and so we can raise and lower indices with impunity. Going to curved space and a non-trivial metric these subtleties become relevant.

²⁰You may also see the Laplacian denoted by Δ rather than \square .

Aside: There is a beautiful interplay between the Eigenforms and Eigenvalues of the Laplacian and the topology of the space that we will not cover. If one defines a harmonic form to be one which is annihilated by the Laplacian $\square\omega = 0$, then there is an isomorphism between the set of all harmonic forms and the cohomology group:

$$\text{Harm}^r(M) \cong H^r(M). \quad (4.38)$$

The Betti numbers which were the dimensions of the cohomology groups are then just the dimension of the group of harmonic r -forms on the manifold.

4.2 Connections and curvature

A vector field X is a directional derivative acting on a function $f \in \mathcal{F}(M)$. However so far we have not introduced such a derivative for tensors of type (q, r) . The Lie derivative is not quite what we want since it also involves derivatives of the vector defining the direction and so we want to introduce something additional. This other derivative is more useful than the Lie derivative, but requires the introduction of a *connection* to map the vector spaces at one point to vector spaces at another. The resultant object is known as the *covariant derivative* and is distinct from the Lie derivative that we introduced previously.

Definition 35 (Affine connection) *An affine connection, which we denote by ∇ is a map $\nabla : \mathcal{X}(M) \times \mathcal{X}(M) \rightarrow \mathcal{X}(M)$, that is $(X, Y) \mapsto \nabla_X Y$ which satisfies*

$$\nabla_X(Y + Z) = \nabla_X Y + \nabla_X Z, \quad (4.39)$$

$$\nabla_{(fX+gY)}Z = f\nabla_X Z + g\nabla_Y Z, \quad (4.40)$$

$$\nabla_X(fY) = X[f]Y + f\nabla_X Y, \quad (4.41)$$

for vector fields $X, Y, Z \in \mathcal{X}(M)$ and functions $f, g \in \mathcal{F}(M)$.

Key Points

The connection provides a way to compare vectors at different points and to define differentiation on a curved manifold.

Let us take a chart (U, φ) with coordinate $x = \varphi(p)$ and define m^3 functions $\Gamma^\mu_{\nu\rho}$ called the *connection coefficients* by

$$\nabla_\nu e_\mu \equiv \nabla_{e_\nu} e_\mu = e_\lambda \Gamma^\lambda_{\nu\mu}, \quad (4.42)$$

where $\{e_\mu\} = \{\frac{\partial}{\partial x^\mu}\}$ is the coordinate basis in $T_p(M)$. The connection coefficients specify how the basis vectors change from point to point, i.e. how to map the tangent space $T_p(M)$

to $T_q(M)$. Using the properties of the connection we can work out the general covariant derivative of a vector field

$$\begin{aligned}
\nabla_X Y &= \nabla_X (Y^\mu e_\mu) \\
&= X[Y^\mu]e_\mu + Y^\mu \nabla_X e_\mu \\
&= X^\nu \partial_\nu (Y^\mu) e_\mu + X^\nu Y^\mu \nabla_\nu e_\mu \\
&= X^\nu \left(\partial_\nu Y^\mu + \Gamma^\mu_{\nu\rho} Y^\rho \right) e_\mu .
\end{aligned} \tag{4.43}$$

We can strip off the overall X^ν to write

$$\left(\nabla_\nu Y \right)^\mu = \frac{\partial Y^\mu}{\partial x^\nu} + \Gamma^\mu_{\nu\rho} Y^\rho , \tag{4.44}$$

so that

$$\left(\nabla_X Y \right)^\mu = X^\nu \nabla_\nu Y^\mu \tag{4.45}$$

On a function the covariant derivative coincides with both the Lie derivative and the regular partial derivative, however its action on vectors differs. While the Lie derivative $\mathcal{L}_X Y$ depends on both X and its first derivative, the covariant derivative depends only on X . This is the natural generalisation of the partial derivative on curved space.

We will often be sloppy and write

$$\left(\nabla_X Y \right)^\mu = \nabla_X Y^\mu . \tag{4.46}$$

Typically in older books, though some still like to use this stupid convention, one may see the semi-colon notation

$$\nabla_\nu Y^\mu = Y^\mu_{;\nu} . \tag{4.47}$$

We will refrain from using this convention to preserve our sanity.

At the moment the connection $\Gamma^\mu_{\nu\rho}$ is somewhat abstract. One may guess that it is a tensor however this is not correct. To see this let us consider how it transforms under a change of coordinates. Recall that the basis elements transform as

$$\tilde{e}_\nu = \Lambda^\mu_{\nu} e_\mu , \quad \text{with} \quad \Lambda^\mu_{\nu} = \frac{\partial x^\mu}{\partial y^\nu} . \tag{4.48}$$

Recall that a $(1, 2)$ tensor $T^\mu_{\nu\rho}$ transforms as

$$\tilde{T}^{\mu_1}_{\nu_1\rho_1} = (\Lambda^{-1})^{\mu_1}_{\mu_2} \Lambda^{\nu_2}_{\nu_1} \Lambda^{\rho_2}_{\rho_1} T^{\mu_2}_{\nu_2\rho_2} . \tag{4.49}$$

We can compute the transformation of the connection. In the basis $\{\tilde{e}_\mu\}$ we have

$$\begin{aligned}
\nabla_{\tilde{e}_\rho} \tilde{e}_\nu &= \tilde{\Gamma}^\mu_{\nu\rho} \tilde{e}_\mu \\
&= \nabla_{\Lambda^\sigma_\rho e_\sigma} \left(\Lambda^\tau_\nu e_\tau \right) \\
&= \Lambda^\sigma_\rho \left(\nabla_\sigma (\Lambda^\tau_\nu) e_\tau + \Lambda^\tau_\nu \nabla_\sigma e_\tau \right) \\
&= \Lambda^\sigma_\rho \left(\Lambda^\tau_\nu \Gamma^\kappa_{\sigma\tau} + \partial_\sigma \Lambda^\kappa_\nu \right) e_\kappa \\
&= \Lambda^\sigma_\rho \left(\Lambda^\tau_\nu \Gamma^\kappa_{\sigma\tau} + \partial_\sigma \Lambda^\kappa_\nu \right) (\Lambda^{-1})^\mu_\kappa \tilde{e}_\mu.
\end{aligned} \tag{4.50}$$

From this we obtain

$$\tilde{\Gamma}^\mu_{\nu\rho} = (\Lambda^{-1})^\mu_\kappa \Lambda^\sigma_\rho \Lambda^\tau_\nu \Gamma^\kappa_{\sigma\tau} + (\Lambda^{-1})^\mu_\kappa \Lambda^\sigma_\rho \partial_\sigma \Lambda^\kappa_\nu. \tag{4.51}$$

The first term is the expected transformation term of a $(1,2)$ tensor, however there is an additional piece. This additional piece is independent of Γ and depends only on the $\partial\Lambda$. This is the characteristic transformation of a connection coefficient.

Key Points

To see why the connection is necessary let us see what happens if we had tried to just use the usual differentiation for a vector field. Consider the transformation of $\partial_\mu V^\nu$ under a change of coordinates:

$$\begin{aligned}
\frac{\partial}{\partial y^\mu} \tilde{V}^\nu &= \frac{\partial x^\sigma}{\partial y^\mu} \frac{\partial}{\partial x^\sigma} \left(\frac{\partial y^\nu}{\partial x^\tau} V^\tau \right) \\
&= \frac{\partial x^\sigma}{\partial y^\mu} \frac{\partial y^\nu}{\partial x^\tau} \frac{\partial}{\partial x^\sigma} V^\tau + \frac{\partial x^\sigma}{\partial y^\mu} \frac{\partial^2 y^\nu}{\partial x^\sigma \partial x^\tau} V^\tau.
\end{aligned} \tag{4.52}$$

The first term has exactly the correct transformation of a $(1,1)$ -tensor, the second term however should not be there. Fortunately, the transformation of the connection coefficients precisely cancels off this extra term. The two terms, derivative plus connection coefficients, despite neither giving a well defined tensor, do give a tensor when combined. This is the utility of the connection it gives a way of differentiating tensors on a curved manifold.

4.2.1 Differentiating other tensors

We can use the properties of the covariant derivative to extend its action to any tensor field. Consider a one-form ω , we want the covariant derivative to take the one-form and return another one-form, $\nabla_X \omega$, as such we should check its action on a vector field $Y \in \mathcal{X}(M)$. We impose that the connection obeys the (generalised) Leibniz identity, so that

$$\nabla_X(\omega(Y)) = (\nabla_X \omega)(Y) + \omega(\nabla_X Y). \tag{4.53}$$

Since $\omega(Y)$ is a function we know that

$$\nabla_X(\omega(Y)) = X[\omega(Y)]. \quad (4.54)$$

Using the Leibniz condition we have

$$(\nabla_X\omega)(Y) = X(\omega(Y)) - \omega(\nabla_X Y), \quad (4.55)$$

and reducing to coordinates we find

$$\begin{aligned} X^\mu(\nabla_\mu\omega)_\nu Y^\nu &= X^\mu\partial_\mu(\omega_\nu Y^\nu) - \omega_\nu X^\mu(\partial_\mu Y^\nu + \Gamma^\nu_{\mu\rho} Y^\rho) \\ &= X^\mu(\partial_\mu\omega_\rho - \Gamma^\nu_{\mu\rho}\omega_\nu)Y^\rho. \end{aligned} \quad (4.56)$$

We may then write

$$(\nabla_\mu\omega)_\rho \equiv \nabla_\mu\omega_\rho = \frac{\partial}{\partial x^\mu}\omega_\rho - \Gamma^\nu_{\mu\rho}\omega_\nu. \quad (4.57)$$

We can now extend this argument to an arbitrary tensor of rank (q, r) , again imposing the generalised Leibniz identity, and we find

$$\begin{aligned} \nabla_\mu T^{\nu_1 \dots \nu_q}_{\rho_1 \dots \rho_r} &= \frac{\partial}{\partial x^\mu} T^{\nu_1 \dots \nu_q}_{\rho_1 \dots \rho_r} + \Gamma^{\nu_1}_{\mu\sigma} T^{\sigma \dots \nu_q}_{\rho_1 \dots \rho_r} + \dots + \Gamma^{\nu_q}_{\mu\sigma} T^{\nu_1 \dots \nu_{q-1} \sigma}_{\rho_1 \dots \rho_r} \\ &\quad - \Gamma^\sigma_{\mu\rho_1} T^{\nu_1 \dots \nu_q}_{\sigma \dots \rho_r} - \dots - \Gamma^\sigma_{\mu\rho_r} T^{\nu_1 \dots \nu_q}_{\rho_1 \dots \rho_{r-1} \sigma}. \end{aligned} \quad (4.58)$$

In words, you first differentiate the tensor and then for each upper index you add in a $+\Gamma T$ and for every down index a $-\Gamma T$.

Key Points

By using the property that the connection acting on a tensor gives another tensor we can work out how the connection actions on an arbitrary tensor by knowing the action on a scalar (just usual differentiation) and a vector.

4.3 Torsion and curvature

Even though the connection is not a tensor we can use it to construct two tensors. The first is a rank $(1, 2)$ tensor T known as *Torsion*, the second is a rank $(1, 3)$ tensor known as *curvature* or the *Riemann tensor*.

Definition 36 (Torsion tensor) *The torsion tensor acts on $X, Y \in \mathcal{X}(M)$ and $\omega \in \Omega^1(M)$ by*

$$T(\omega : X, Y) = \omega(\nabla_X Y - \nabla_Y X - [X, Y]). \quad (4.59)$$

We may equivalently think of this as a map $T : \mathcal{X}(M) \times \mathcal{X}(M) \rightarrow \mathcal{X}(M)$ defined by

$$T(X, Y) = \nabla_X Y - \nabla_Y X - [X, Y]. \quad (4.60)$$

Definition 37 (Curvature tensor) *The curvature acts on $X, Y, Z \in \mathcal{X}(M)$ and $\omega \in \Omega^1(M)$ as*

$$R(\omega : X, Y, Z) = \omega(\nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X, Y]} Z) \quad (4.61)$$

As for the torsion we may think of this as a map $\mathcal{X}(M) \times \mathcal{X}(M)$ to a differential operator acting on $\mathcal{X}(M)$ as

$$R(X, Y) = \nabla_X \nabla_Y - \nabla_Y \nabla_X - \nabla_{[X, Y]}. \quad (4.62)$$

Exercise 4.2: Torsion and Curvature are tensors

Check that both the Torsion and Curvature tensors are actually tensors. There are two ways of doing this.

1. Show that it is multi-linear in all arguments. For example show $T(\omega : fX, Y) = fT(\omega : X, Y)$ for all $f \in \mathcal{F}(M)$ and so forth.
2. Show that it transforms as a tensor should under a change of coordinates.

We can evaluate the tensors in a basis to obtain the component form. Let $\{\theta^\rho\} = \{dx^\rho\}$ be a basis of the cotangent space and $\{e_\mu\} = \{\partial_\mu\}$ be our basis for the tangent space.

4.3.1 Component form of the torsion

In this basis the components of the torsion tensor are

$$\begin{aligned} T^\rho_{\mu\nu} &= T(\theta^\rho : e_\mu, e_\nu) \\ &= \theta^\rho(\nabla_\mu e_\nu - \nabla_\nu e_\mu - [e_\mu, e_\nu]) \\ &= \theta^\rho(\Gamma^\sigma_{\mu\nu} - \Gamma^\sigma_{\nu\mu})e_\sigma \\ &= \Gamma^\rho_{\mu\nu} - \Gamma^\rho_{\nu\mu}. \end{aligned} \quad (4.63)$$

We therefore end up with

$$T^\rho_{\mu\nu} = \Gamma^\rho_{\mu\nu} - \Gamma^\rho_{\nu\mu}. \quad (4.64)$$

So despite $\Gamma^\sigma_{\mu\nu}$ not being a tensor, the anti-symmetrised part is! The torsion tensor is clearly anti-symmetric in the two lowered indices.

Definition 38 (Torsion free) *We see that connections $\Gamma^\sigma_{\mu\nu}$ which are symmetric in the lowered indices have $T^\rho_{\mu\nu} = 0$ and are called torsion-free.*

4.3.2 Component form of the Riemann tensor

A similar computation for the Riemann tensor gives

$$R^\sigma{}_{\rho\mu\nu} = \partial_\mu \Gamma^\sigma{}_{\nu\rho} - \partial_\nu \Gamma^\sigma{}_{\mu\rho} + \Gamma^\lambda{}_{\nu\rho} \Gamma^\sigma{}_{\mu\lambda} - \Gamma^\lambda{}_{\mu\rho} \Gamma^\sigma{}_{\nu\lambda}. \quad (4.65)$$

4.3.3 The Ricci identity

Consider the commutator of covariant derivatives acting on a vector field, we have

$$\begin{aligned} \nabla_{[\mu} \nabla_{\nu]} X^\sigma &= \partial_{[\mu} (\nabla_{\nu]} X^\sigma) + \Gamma^\sigma{}_{[\mu|\lambda]} \nabla_{\nu]} X^\lambda - \Gamma^\rho{}_{[\mu\nu]} \nabla_\rho X^\sigma \\ &= \partial_{[\mu} \partial_{\nu]} X^\sigma + (\partial_{[\mu} \Gamma^\sigma{}_{\nu]\rho}) X^\rho + (\partial_{[\mu} X^\rho) \Gamma^\sigma{}_{\nu]\rho} + \Gamma^\sigma{}_{[\mu|\lambda]} \partial_{\nu]} X^\lambda \\ &\quad + \Gamma^\sigma{}_{[\mu|\lambda]} \Gamma^\lambda{}_{\nu]\rho} X^\rho - \Gamma^\rho{}_{[\mu\nu]} \nabla_\rho X^\sigma. \end{aligned} \quad (4.66)$$

The first term of the second line vanishes, while the third and fourth cancel. The second term on the second line and first on the third line combine to give the Riemann tensor while the last gives the torsion. Putting everything together we have the *Ricci identity*

$$2\nabla_{[\mu} \nabla_{\nu]} X^\sigma = R^\sigma{}_{\rho\mu\nu} X^\rho - T^\rho{}_{\mu\nu} \nabla_\rho X^\sigma. \quad (4.67)$$

Similar identities hold when acting on other tensors and can be shown following similar steps to the above.

4.3.4 Levi–Civita connection

So far the discussion has not required a metric, however when a metric exists we have the following theorem.

Theorem 5 (Levi–Civita Connection) *There exists a unique, torsion free, connection that is compatible with the metric g :*

$$\nabla_X g = 0, \quad (4.68)$$

for all vector fields X . This connection is called the Levi–Civita connection.

Proof: To prove this we first show uniqueness before constructing the connection. Suppose that such a connection exists, then we have

$$X(g(Y, Z)) = \nabla_X(g(Y, Z)) = (\nabla_X g)(Y, Z) + g(\nabla_X Y, Z) + g(Y, \nabla_X Z). \quad (4.69)$$

Since $\nabla_X g = 0$ we have

$$X(g(Y, Z)) = g(Y, \nabla_X Z) + g(\nabla_X Y, Z). \quad (4.70)$$

We may use our favourite trick and cyclically permute X, Y, Z to find

$$\begin{aligned} Y(g(Z, X)) &= g(Z, \nabla_Y X) + g(\nabla_Y Z, X), \\ Z(g(X, Y)) &= g(X, \nabla_Z Y) + g(\nabla_Z X, Y). \end{aligned} \tag{4.71}$$

By the no torsion condition we have

$$\nabla_X Y - \nabla_Y X = [X, Y], \tag{4.72}$$

and therefore²¹

$$\begin{aligned} X(g(Y, Z)) &= g(\nabla_X Z, Y) + g(\nabla_Y X, Z) + g([X, Y], Z), \\ Y(g(Z, X)) &= g(\nabla_Z Y, X) + g(\nabla_Y X, Z) + g([Y, Z], X), \\ Z(g(X, Y)) &= g(\nabla_Z Y, X) + g(\nabla_X Z, Y) + g([Z, X], Y), \end{aligned} \tag{4.73}$$

Adding the first and second and subtracting the third we find

$$\begin{aligned} g(\nabla_Y X, Z) &= \frac{1}{2} \left[X(g(Y, Z)) + Y(g(Z, X)) - Z(g(X, Y)) \right. \\ &\quad \left. - g([X, Y], Z) - g([Y, Z], X) + g([Z, X], Y) \right] \end{aligned} \tag{4.74}$$

With a non-degenerate metric this specifies the connection uniquely.

It remains to be seen that the connection as defined does satisfy the properties of a connection. We will present one of the terms to check. The most finicky one is $\nabla_{fX} Y = f\nabla_X Y$, so let us present that one

$$\begin{aligned} g(\nabla_{fX} Y, Z) &= \frac{1}{2} \left[X(g(fY, Z)) + fY(g(Z, X)) - Z(g(X, fY)) \right. \\ &\quad \left. - g([X, fY], Z) - g([fY, Z], X) + g([Z, X], fY) \right] \\ &= \frac{1}{2} \left[fX(g(Y, Z)) + X(f)g(Y, Z) + fY(g(Z, X)) - fZ(g(X, Y)) - Z(f)g(X, Y) \right. \\ &\quad \left. - fg([X, Y], Z) - X(f)g(Y, Z) - fg([Y, Z], X) + Z(f)g(Y, X) + fg([Z, X], Y) \right] \\ &= g(f\nabla_X Y, Z). \end{aligned} \tag{4.75}$$

The coloured terms in the penultimate line cancel amongst themselves, leaving just the black terms as required. The other properties follow similarly. This then proves the uniqueness and has explicitly constructed such a connection.

²¹Note that in the X equation we eliminate the term $g(\nabla_X Y, Z)$ using the identity (4.72). Similarly for the Y term we eliminate the $g(\nabla_Y Z, X)$ and for Z we eliminate the $g(\nabla_Z X, Y)$ term.

In components we can evaluate

$$g(\nabla_\nu e_\mu, e_\rho) = \Gamma^\lambda_{\nu\mu} g_{\lambda\rho} = \frac{1}{2}(\partial_\mu g_{\nu\rho} + \partial_\nu g_{\mu\rho} - \partial_\rho g_{\mu\nu}). \quad (4.76)$$

Multiplying by the inverse metric we have

$$\Gamma^\lambda_{\mu\nu} = \frac{1}{2}g^{\lambda\rho}(\partial_\mu g_{\nu\rho} + \partial_\nu g_{\mu\rho} - \partial_\rho g_{\mu\nu}). \quad (4.77)$$

The connection compatible with the metric is called the *Levi-Civita connection* while the components of the Levi-Civita connection are called the *Christoffel symbols*.

There is a nice expression if you contract two indices of the Christoffel symbols, we have

$$\Gamma^\mu_{\mu\nu} = \frac{1}{\sqrt{|g|}}\partial_\nu \sqrt{|g|} \quad (4.78)$$

To see this note

$$\Gamma^\mu_{\mu\nu} = \frac{1}{2}g^{\mu\rho}\partial_\nu g_{\mu\rho} = \frac{1}{2}\text{tr}(g^{-1}\partial_\nu g) = \frac{1}{2}\text{tr}(\partial_\nu \log g), \quad (4.79)$$

for diagonalisable matrices we have $\text{tr} \log A = \log \det(A)$ and therefore we find

$$\Gamma^\mu_{\mu\nu} = \frac{1}{2}\partial_\nu \log \det(g) = \frac{1}{\sqrt{|\det(g)|}}\partial_\nu \sqrt{|\det(g)|}. \quad (4.80)$$

This implies that

$$\sqrt{|g|}\nabla_\mu X^\mu = \sqrt{|g|}(\partial_\mu X^\mu + \Gamma^\mu_{\mu\nu}X^\nu) + \sqrt{|g|}\left(\partial_\mu X^\mu + X^\nu \frac{1}{\sqrt{|g|}}\partial_\nu \sqrt{|g|}\right) = \partial_\mu(\sqrt{|g|}X^\mu). \quad (4.81)$$

Using this result we can prove the divergence theorem:

$$\int_M d^m x \sqrt{|g|}\nabla_\mu X^\mu = \int_{\partial M} d^{n-1}x \sqrt{\gamma}n_\mu X^\mu, \quad (4.82)$$

where γ_{ij} is the pull-back of the metric to ∂M , $\gamma = \det(\gamma_{ij})$ and n_μ is an outward pointing unit vector orthogonal to ∂M . On a Lorentzian manifold this holds provided that ∂M is either purely spacelike or purely timelike, which guarantees that $\gamma \neq 0$.

4.4 Parallel transport and geodesics

We have introduced the connection but we are yet to explain what it connects. It connects tangent spaces, or more generally any vector space at different points of the manifold. This map is called *parallel transport*. Take a vector field X with some associated integral curve γ with coordinates $x^\mu(\lambda)$ such that

$$X^\mu|_\gamma = \frac{dx^\mu(\lambda)}{d\lambda}. \quad (4.83)$$

We say that a tensor field T is parallel transported along γ if

$$\nabla_X T = 0. \quad (4.84)$$

Physical Explanation

This describes how to move the tensor along the curve whilst keeping it as constant as possible. As we will see shortly, parallel transporting a tensor along a closed curve (loop), does not necessarily leave the tensor invariant, it can differ at the start and end-points. This is due to the curvature of spacetime.

Let γ connect two points $p, q \in M$. The condition (4.84) provides a map from the vector space defined at p to the vector space defined at q . Consider a second vector field Y . In components (4.84) reads

$$X^\nu (\partial_\nu Y^\mu + \Gamma^\mu_{\nu\rho} Y^\rho) = 0. \quad (4.85)$$

If we evaluate it on the curve γ , we can write $Y^\mu = Y^\mu(x(\lambda))$ and therefore the condition is

$$\frac{dY^\mu}{d\lambda} + X^\nu \Gamma^\mu_{\nu\rho} Y^\rho = 0. \quad (4.86)$$

This defines a set of coupled ordinary differential equations, given an initial condition at $p = \gamma(\lambda = 0)$ for example these can be solved to find a unique vector field at each point along the curve. This is path dependent and depends on the connection and the underlying path which was characterised by X here.

There is a subtle difference between what we are doing here and what we did with the push-forward and pull-back, which we used to define the Lie derivative. Here X only appears to define the map, there are no derivatives applied to X^μ as was for those maps. The connection does the work of relating the vector spaces along the curve and not the vector X .

4.4.1 Geodesics

Definition 39 (Affinely parametrised geodesic) *An affinely parametrised geodesic is an integral curve with tangent vector field X that obeys*

$$\nabla_X X = 0. \quad (4.87)$$

Physical Explanation

Geodesics are the analogue of straight lines in flat space.

Along the curve γ with coordinates x^μ and tangent vector field X this implies

$$\frac{d^2 x^\mu}{d\lambda^2} + \Gamma^\mu_{\nu\rho} \frac{dx^\nu}{d\lambda} \frac{dx^\rho}{d\lambda} = 0. \quad (4.88)$$

We have thrown around the phrase affinely parametrised geodesic but what does this really mean? Consider a curve parametrised by λ and with tangent vector field X that satisfies (4.88). Let us parametrise it by some other parameter τ with $\lambda(\tau)$ and take $f = \frac{d\lambda}{d\tau} > 0$. The change in parametrisation leads to a different tangent vector to the curve since

$$Y^\mu \equiv \frac{dx^\mu}{d\tau} = \frac{dx^\mu}{d\lambda} \frac{d\lambda}{d\tau} = X^\mu \frac{d\lambda}{d\tau}. \quad (4.89)$$

We therefore have

$$\nabla_Y Y = \nabla_{fX} fX = f\nabla_X(fX) = fX[f]X + f^2\nabla_X X = X[f]Y. \quad (4.90)$$

We see that (4.90) defines the same geodesic however it is *not* affinely parametrised, the right-hand side of (4.88) no longer vanishes but is instead proportional to an arbitrary function multiplied by the vector field.

We see conversely that if we work in the opposite direction, with a non-affinely parametrised geodesic, that is a Y satisfying (4.90) then we can always perform a reparametrisation that gives us an affinely parametrised geodesic, therefore there is no loss of generality for us to consider affinely parametrised geodesics.

Observe that we still have some freedom to reparametrise our curve without ruining the affine parametrisation, this requires $X[f] = 0$. This is then equivalent to f being constant along the curve and therefore we may take $\lambda = a\tau + b$ with a, b constants without ruining the affine parametrisation. There is a two-parameter family of affine parameters for any affinely parametrised geodesic.

Exercise 4.3: Geodesic equation from Euler–Lagrange

Consider the action of a particle following the path $x^\mu(\lambda)$:

$$S = \int d\lambda \sqrt{-g_{\mu\nu}(x) \frac{dx^\mu}{d\lambda} \frac{dx^\nu}{d\lambda}}. \quad (4.91)$$

Using the Euler–Lagrange equations show that generically you obtain the non-affinely parametrised geodesics from the action with respect to the Levi–Civita connection.

What is the affine parameter for a time-like geodesic?

If we choose the Levi–Civita connection, since $\nabla_X g = 0$ it follows that for any vector field Y which is parallel transported along a geodesic defined by X we have

$$\frac{d}{d\lambda} g(X, Y) = 0. \quad (4.92)$$

The vector field Y makes the same angle with the tangent vector at each point along the geodesic. Further, this holds true if we replace Y by X in the expression above. Since the

norm of the vector field X tangent to the geodesic classifies the character of the geodesic, (timelike/null/spacelike), if we define a geodesic using a metric compatible connection, then the nature of the geodesic does not change. This statement relies on us using a metric compatible connection though, in this course we will always take such a connection and therefore the nature of a geodesic is preserved throughout all spacetime.

Let us consider a timelike geodesic. When we vary the action (4.91) what are we extremising and is it a maximum of minimum? From our definition of the proper time, (4.18) we see that we are extremising the proper time, and geodesics maximise the proper time. Why is this true? Well given any time-like curve we can approximate it to arbitrary accuracy by a null curve. We should consider jagged null curves that follow the time-like one, see figure 11. As we increase the number of null curves the approximation gets better and better, while still having zero length. Timelike curves cannot therefore be curves with minimal proper time since they are infinitesimally close to curves of zero length (and therefore zero proper time). They must therefore maximise the proper time. This is why the twin who remains home in the twin paradox ages more, they are on a geodesic (for most of the journey). We should really say that this maximises the proper time locally. If we took a sphere, then there is more than one geodesic between two points, we can either go the short way around or the long way around. One is longer than the other (assuming the points are not opposite each other, i.e. picking the poles), but both maximise locally the length functional.




Figure 11: We approximate the time-like curve with null curves. As we increase the number of null curves the approximation gets better and better.

Key Points

A geodesic is the path of a freely falling particle. There are no forces acting on the particle. From the perspective of GR gravity is not a force, rather it is encoded in the curvature of the spacetime geometry.

- For a time-like geodesic the geodesic equation maximises the proper time along the curve.
- For a space-like geodesic, though not physical, on minimises the proper distance.

4.4.2 Normal coordinates

Geodesics allow for the construction of a particularly useful coordinate system. This holds independently of whether the Levi–Civita connection is employed or not, however it takes a particularly simple form when it is used. On a Riemannian manifold, in the neighbourhood of a point $p \in M$ we can always find coordinates such that:

$$g_{\mu\nu}(p) = \delta_{\mu\nu}, \quad \text{and} \quad \Gamma^\mu_{\nu\rho}(p) = 0. \quad (4.93)$$

The same is true for Lorentzian manifolds with $\delta \rightarrow \eta$. These coordinates are known as *normal coordinates*. If one uses the Levi–Civita connection the vanishing of the Christoffel symbols implies that $\partial_\rho g_{\mu\nu}(p) = 0$. As we move away from p this does not need to continue to hold. It should be noted that one cannot generically make the derivative of the connection coefficients vanish at p . This means that it is not possible to find coordinates such that the Riemann tensor vanishes at the given point.

We can brute force this. Start with a metric $\tilde{g}_{\mu\nu}$ in coordinates \tilde{x}^μ and try to find a new set of coordinates $x^\mu(\tilde{x})$ which satisfy the required conditions. In the new coordinates we have

$$\frac{\partial \tilde{x}^\rho}{\partial x^\mu} \frac{\partial \tilde{x}^\sigma}{\partial x^\nu} \tilde{g}_{\rho\sigma} = g_{\mu\nu} = \delta_{\mu\nu}. \quad (4.94)$$

We can take the point p to be the origin of both coordinate systems and Taylor expand around the point

$$\tilde{x}^\rho = 0 + \left. \frac{\partial \tilde{x}^\rho}{\partial x^\mu} \right|_{x=0} x^\mu + \frac{1}{2} \left. \frac{\partial^2 \tilde{x}^\rho}{\partial x^\mu \partial x^\nu} \right|_{x=0} x^\mu x^\nu + \dots \quad (4.95)$$

Inserting the expansion into (4.94) together with the Taylor expansion of $\tilde{g}_{\mu\nu}$ we can try to solve the resulting PDEs. The first order variation implies

$$\left. \frac{\partial \tilde{x}^\rho}{\partial x^\mu} \right|_{x=0} \left. \frac{\partial \tilde{x}^\sigma}{\partial x^\nu} \right|_{x=0} \tilde{g}_{\rho\sigma}(p) = \delta_{\mu\nu}. \quad (4.96)$$

We can always find $\partial \tilde{x} / \partial x$ such that this is true, there are many choices. For $\dim M = m$ there are m^2 independent coefficients of $\partial \tilde{x} / \partial x$. The equation above contains $\frac{1}{2}m(m+1)$

conditions on these, since \tilde{g} is symmetric. This leaves us with $\frac{1}{2}m(m-1)$ parameters which are un-fixed. Notice that this remainder is precisely the same number of components of the rotational group of $\text{SO}(m)$ or $\text{SO}(1, m-1)$, this is of course the group that leaves the flat metric unchanged and is therefore to be expected. Next consider the second order variations. There are $\frac{1}{2}m^2(m+1)$ independent components of $\partial^2\tilde{x}^\rho/\partial x^\mu\partial x^\nu$ which is the same number of components of $\partial_\rho g_{\mu\nu}$ and so we can always choose the first derivative of the metric at p to vanish. Consider now the second derivative term, requiring $\partial_\rho\partial_\sigma g_{\mu\nu} = 0$ imposes $\frac{1}{4}m^2(m+1)^2$ constraints. However the next term in the Taylor expansion is $\partial^3\tilde{x}^\rho/\partial x^\mu\partial x^\nu\partial x^\sigma$ and has only $\frac{1}{6}m^2(m+1)(m+2)$ independent coefficients: there are not enough independent coefficients to cancel all of the terms of the second derivative. The difference is the number of ways of characterising the second derivative of the metric that cannot be undone by coordinate transformations. This is precisely the number of independent components of the Riemann tensor, this is

$$\frac{1}{4}m^2(m+1)^2 - \frac{1}{6}m^2(m+1)(m+2) = \frac{1}{12}m^2(m+1)(m-1). \quad (4.97)$$

One can explicitly construct the normal coordinates using the exponential map and geodesics flowing through the point p . One can consider all affinely parametrised geodesics through p and label the point q at a small fixed distance of the affine parameter by the coordinates of the geodesic flowing through q . One then essentially uses geodesics to construct your basis vectors. We will not consider this construction here.

The Equivalence principle again. Normal coordinates play an important role in GR. Any observer at a point p who parametrises their immediate surroundings using normal coordinates will experience a locally flat metric. This is the mathematics underlying the Einstein equivalence principle. Any freely falling observer, performing local experiments will not experience a gravitational field. Here free falling means following a geodesic and therefore they can use normal coordinates. The lack of gravitational field is the statement that $g_{\mu\nu}(p) = \eta_{\mu\nu}$.

There are limitations to the equivalence principle and the important word is **local**. There is a way to distinguish whether there is a gravitational field at p or not. We simply compute the Riemann tensor. This depends on the second derivative of the metric and will in general be non-vanishing. However to measure the effects of the Riemann tensor one typically has to compare the result of an experiment at p with the result at a nearby point q , this is then a “non-local” observable, according to the equivalence principle.

4.4.3 Path dependence: Curvature and Torsion

We have introduced the curvature and torsion tensors but what are they really measuring? For the Riemann tensor let us consider a vector $Z_p \in T_p(M)$ and parallel transport it along a curve C to some point $r \in M$. In addition consider another curve C' along which we can parallel transport Z_p to r . We will see that the difference between the two vectors at r is determined by the Riemann tensor.

The meaning of curvature Let us construct our curves from two segments, generated by linearly independent vector fields X, Y and let us take $[X, Y] = 0$. (Recall that this implies that the parallelogram constructed from the vectors closes, see section 3.3.2). We take the points to be close and pick normal coordinates $x^\mu = (\tau, \sigma, 0, \dots, 0)$ so that the starting point is at $x^\mu(p) = 0$, and the tangent vectors are aligned along the coordinates $X = \frac{\partial}{\partial \tau}$ and $Y = \frac{\partial}{\partial \sigma}$. The other corner points are $x^\mu(r) = (\delta\tau, 0, 0, \dots)$, $x^\mu(s) = (0, \delta\sigma, 0, \dots)$ and $x^\mu(r) = (\delta\tau, \delta\sigma, 0, \dots)$, with $\delta\tau$ and $\delta\sigma$ small, see figure 12.




Figure 12: Parallel transporting a vector Z_p along two different paths does not give the same answer. From the lecture notes of Tong.

First parallel transport Z_p along X to obtain Z_q . Along the curve, by definition of the parallel transport Z^μ satisfies

$$\frac{dZ^\mu}{d\tau} + X^\nu \Gamma^\mu_{\rho\nu} Z^\rho = 0. \quad (4.98)$$

We can Taylor expand the solution as

$$Z_q^\mu = Z_p^\mu + \left. \frac{dZ^\mu}{d\tau} \right|_p \delta\tau + \frac{1}{2} \left. \frac{d^2 Z^\mu}{d\tau^2} \right|_p \delta\tau^2 + \mathcal{O}(\delta\tau^3). \quad (4.99)$$

We can use normal coordinates at the point p which implies that $\Gamma^\mu_{\rho\nu}(p) = 0$ and therefore $\left. \frac{dZ^\mu}{d\tau} \right|_p = 0$. To calculate the second derivative we differentiate (4.98), to obtain

$$\begin{aligned} \left. \frac{dZ^\mu}{d\tau^2} \right|_{\tau=0} &= - \left(X^\nu Z^\rho \frac{d\Gamma^\mu_{\rho\nu}}{d\tau} + \frac{dX^\nu}{d\tau} Z^\rho \Gamma^\mu_{\rho\nu} + X^\nu \frac{dZ^\rho}{d\tau} \Gamma^\mu_{\rho\nu} \right) \Big|_p \\ &= -X^\nu Z^\rho \left. \frac{d\Gamma^\mu_{\rho\nu}}{d\tau} \right|_p \\ &= -X^\nu X^\sigma Z^\rho \partial_\sigma \Gamma^\mu_{\rho\nu} \Big|_p \end{aligned} \quad (4.100)$$

To get to the second line we have used that we are working in normal coordinates at p and the final line is because τ parametrises the integral curve of X . We find

$$Z_q^\mu = Z_p^\mu - \frac{1}{2} X^\nu X^\sigma Z^\rho \partial_\sigma \Gamma^\mu_{\rho\nu} \Big|_p \delta\tau^2 + \dots \quad (4.101)$$

Now we parallel transport again, this time along Y to the point r with resultant vector Z_r^μ . The Taylor expansion is

$$Z_r^\mu = Z_q^\mu + \left. \frac{dZ^\mu}{d\sigma} \right|_q \delta\sigma + \frac{1}{2} \left. \frac{d^2 Z^\mu}{d\sigma^2} \right|_q \delta\sigma^2 + \mathcal{O}(\delta\sigma^3). \quad (4.102)$$

We can evaluate the first derivative $\left. \frac{dZ^\mu}{d\sigma} \right|_q$ using the analogue of the parallel transport equation (4.98),

$$\left. \frac{dZ^\mu}{d\sigma} \right|_q = -Y^\nu Z^\rho \Gamma^\mu_{\rho\nu} \Big|_q, \quad (4.103)$$

however since our normal coordinates are at p and not q we cannot argue that this term immediately vanishes, instead we can Taylor expand about p to get

$$Y^\nu Z^\rho \Gamma^\mu_{\rho\nu} \Big|_q = Y^\nu Z^\rho X^\sigma \partial_\sigma \Gamma^\mu_{\rho\nu} \Big|_p \delta\tau + \dots \quad (4.104)$$

One should also expand Y^ν and Z^ν however to leading order they multiply $\Gamma^\mu_{\rho\nu}(p) = 0$ ergo, only contribute at the next order. For the second order term in the Taylor expansion (4.102) there is a similar expression to before, we find

$$\begin{aligned} \left. \frac{d^2 Z^\mu}{d\sigma^2} \right|_q &= -Y^\nu Y^\sigma Z^\rho \partial_\sigma \Gamma^\mu_{\rho\nu} \Big|_q + \dots \\ &= -Y^\nu Y^\sigma Z^\rho \partial_\sigma \Gamma^\mu_{\rho\nu} \Big|_p + \dots \end{aligned} \quad (4.105)$$

After the dust settles we have

$$Z_r^\mu = Z_q^\mu - Y^\nu Z^\rho X^\sigma \partial_\sigma \Gamma^\mu_{\rho\nu} \Big|_p \delta\tau \delta\sigma - \frac{1}{2} Y^\nu Y^\sigma Z^\rho \partial_\sigma \Gamma^\mu_{\rho\nu} \Big|_p \delta\sigma^2 + \dots \quad (4.106)$$

and therefore

$$Z_r^\mu = Z_p^\mu - \frac{1}{2} \partial_\sigma \Gamma^\mu_{\rho\nu} \Big|_p \left[X^\nu X^\sigma Z^\rho \delta\tau^2 + 2Y^\nu Z^\rho X^\sigma \delta\sigma \delta\tau + Y^\nu Y^\sigma Z^\rho \delta\sigma^2 \right] \Big|_p + \dots \quad (4.107)$$

with ... cubic and higher terms. We can now consider the same computation for the path C' .

We merely need to swap the role of $\tau \leftrightarrow \sigma$ and $X \leftrightarrow Y$, so that

$$Z_r'^\mu = Z_p^\mu - \frac{1}{2} \partial_\sigma \Gamma^\mu_{\rho\nu} \Big|_p \left[X^\nu X^\sigma Z^\rho \delta\tau^2 + 2X^\nu Z^\rho Y^\sigma \delta\sigma \delta\tau + Y^\nu Y^\sigma Z^\rho \delta\sigma^2 \right] \Big|_p + \dots \quad (4.108)$$

and therefore

$$\begin{aligned} \Delta Z_r^\mu &= Z_r^\mu - Z_r'^\mu = - \left(\partial_\sigma \Gamma^\mu_{\rho\nu} - \partial_\nu \Gamma^\mu_{\rho\sigma} \right) \Big|_p (Y^\nu Z^\rho X^\sigma) \Big|_p \delta\sigma \delta\tau + \dots \\ &= R^\mu_{\rho\sigma\nu} Y^\nu Z^\rho X^\sigma \Big|_p \delta\sigma \delta\tau. \end{aligned} \quad (4.109)$$

The final expression follows from the Riemann tensor expression in normal coordinates. Although our calculation was performed in a certain choice of coordinates since the end result is an equality between tensors it must hold in any coordinate system. This is a common trick, normal coordinates generally simplify expressions.

The Riemann tensor tells us the path dependence of parallel transport. This is related to the concept of *holonomy*. If we transport a vector around a closed loop we can ask how it compares to the original vector. This is captured by the Riemann tensor. A particularly easy example is to consider a two-sphere. We can draw a loop by considering the intersection of three great circles. First go along the equator by $1/4$ of the circumference. Then make a $\pi/2$ turn and head to the north pole. At the north pole go south on another $\pi/2$ angle. You will end up with a triangle with angle $3\pi/2$. Now consider parallel transporting a vector along this loop. You will see that it changes direction when you get back to the start. Of course one could take any path and the direction you end up facing depends on the path. The set of all possible transformations of the vector at p along loops form a group known as the *holonomy group*. For a Riemannian manifold with a metric this is a subgroup of $\text{SO}(m)$ while for a Lorentzian manifold it is a subgroup of $\text{SO}(1, m-1)$.

The meaning of Torsion Torsion will not play a role in this course since we will exclusively use the Levi-Civita connection which is torsion free. Before we completely rid ourselves of the torsion let us first understand its geometric meaning.

Take two vectors $X, Y \in T_p(M)$ and let us use coordinates x^μ . Starting at $p \in M$ we can use these vectors to construct two points infinitesimally close to p , let them be r and s respectively:

$$r : x^\mu + \epsilon X^\mu \quad \text{and} \quad s : x^\mu + \epsilon Y^\mu. \quad (4.110)$$

We can now parallel transport X along Y to give a new vector $X' \in T_s(M)$ and similarly parallel transport Y along X to get a new vector $Y' \in T_r(M)$. The new vectors have components

$$X' = (X^\mu - \epsilon \Gamma^\mu_{\nu\rho} Y^\nu X^\rho) \partial_\mu, \quad Y' = (Y^\mu - \epsilon \Gamma^\mu_{\nu\rho} X^\nu Y^\rho) \partial_\mu. \quad (4.111)$$

Each now defines a new point. Starting from s and moving in the direction X' we get a new point

$$q : x^\mu + (X^\mu + Y^\mu) \epsilon - \epsilon^2 \Gamma^\mu_{\nu\rho} Y^\nu X^\rho. \quad (4.112)$$

Similarly if we sit at r and move in the direction of Y' we get to a typically different point t with coordinates

$$t : x^\mu + (X^\mu + Y^\mu) \epsilon - \epsilon^2 \Gamma^\mu_{\nu\rho} X^\nu Y^\rho. \quad (4.113)$$

The two points are not the same when $\Gamma^\mu_{\nu\rho} \neq \Gamma^\mu_{\rho\nu}$, i.e. when the connection has torsion. Torsion measures the failure of the parallelogram in figure 13 to close.




Figure 13: We transport the two vectors X and Y along each other. The failure for the parallelogram to close is measured by the torsion of the connection.

4.4.4 Explicit Example of path dependence

To bring everything we have learnt in the last few pages let us explicitly show path dependence using the two sphere as an example. We draw a path consisting of four parts, two paths along constant latitude (constant θ) and two paths along constant longitude (constant ϕ) which are joined up into a rectangle on the sphere, see figure 14




Figure 14: We consider parallel transporting a vector field around the rectangular looking path. We will find that we do not end up with the same vector field we started with.

We take the metric to be

$$ds^2 = d\theta^2 + \sin^2 \theta d\phi^2, \quad (4.114)$$

which in components is

$$g_{\mu\nu} = \begin{pmatrix} 1 & 0 \\ 0 & \sin^2 \theta \end{pmatrix}_{\mu\nu}, \quad g^{\mu\nu} = \begin{pmatrix} 1 & 0 \\ 0 & \frac{1}{\sin^2 \theta} \end{pmatrix}^{\mu\nu}. \quad (4.115)$$

Recall from example 4.1 that the metric in these coordinates is not well defined at the two poles. One needs to use another coordinate patch to describe these two points. We will continue to use this metric, staying away from the poles. (In the end we will take a limit to one of the poles which remains well defined.)

The non-trivial Christoffel symbols following from the above metric are

$$\Gamma^{\theta}_{\phi\phi} = -\sin \theta \cos \theta, \quad \Gamma^{\phi}_{\theta\phi} = \Gamma^{\phi}_{\phi\theta} = \cot \theta. \quad (4.116)$$

These can be easily computed by plugging the metric into the explicit formula in equation (4.77). There is a computationally simpler way to do it however which involves using the Euler Lagrange equations and the equation for an affinely parametrised geodesic. Recall that

an affinely parametrised geodesic satisfies (4.88) and are the equations of motion following from the Lagrangian:

$$\mathcal{L} = g_{\mu\nu} \dot{x}^\mu \dot{x}^\nu. \quad (4.117)$$

We have

$$\mathcal{L} = \dot{\theta}^2 + \sin^2 \theta \dot{\phi}^2, \quad (4.118)$$

and therefore the Euler–Lagrange equations give:

$$\begin{aligned} 0 &= \ddot{\theta} - \cos \theta \sin \theta \dot{\phi}^2, \\ 0 &= \ddot{\phi} + 2 \cot \theta \dot{\theta} \dot{\phi}. \end{aligned} \quad (4.119)$$

Comparing with (4.88) we easily read off that the non-trivial Christoffel symbols are as in equation (4.116).

We can now consider the different paths we want to take. For constant longitude (ϕ fixed) we have the path $(\theta, \phi) : (\alpha_0, \beta) \rightarrow (\alpha_1, \beta)$ is given by

$$x^\mu = (\alpha_0 + (\alpha_1 - \alpha_0)\lambda, \beta), \quad \lambda \in [0, 1]. \quad (4.120)$$

This has tangent vector field

$$X^\mu = \frac{dx^\mu}{d\lambda} = (\alpha_1 - \alpha_0, 0)^\mu. \quad (4.121)$$

For the constant latitude (fixed θ) consider the end-points $(\theta, \phi) : (\alpha, \beta_0) \rightarrow (\alpha, \beta_1)$ which has path between them

$$x^\mu = (\alpha, \beta_0 + \lambda(\beta_1 - \beta_0))^\mu, \quad \lambda \in [0, 1], \quad (4.122)$$

with tangent vector field

$$X^\mu = \frac{dx^\mu}{d\lambda} = (0, \beta_1 - \beta_0)^\mu. \quad (4.123)$$

The parallel transport of the vector field Y along the path with tangent vector field X is given by

$$\nabla_X Y = 0, \quad (4.124)$$

which in components can be written as:

$$\frac{d}{d\lambda} Y^\mu + \Gamma^\mu_{\rho\sigma} Y^\rho X^\sigma = 0. \quad (4.125)$$

First consider parallel transport of Y along constant longitude ($\phi = \text{const}$) and take the path in equation (4.120). For ease of notation let $\Omega = \alpha_1 - \alpha_0$. Now we just need to solve the

parallel transport equation. We have the two equations:

$$\begin{aligned} 0 &= \frac{d}{d\lambda} Y^\theta + \Gamma^\theta_{\nu\rho} Y^\nu X^\rho, \\ 0 &= \frac{d}{d\lambda} Y^\phi + \Gamma^\phi_{\nu\rho} Y^\nu X^\rho. \end{aligned} \quad (4.126)$$

Plugging in our X and Christoffel symbols we find the two equations:

$$\begin{aligned} 0 &= \frac{d}{d\lambda} Y^\theta, \\ 0 &= \frac{d}{d\lambda} Y^\phi + \Omega \cot \theta Y^\phi. \end{aligned} \quad (4.127)$$

The first can be solved simply by

$$Y^\theta = Y_0^\theta, \quad (4.128)$$

with Y_0^θ the initial value of Y at the initial point (α_0, ϕ_0) of the path. The second is slightly trickier to solve but one finds

$$Y^\phi = \frac{c}{\sin(\alpha_0 + \lambda\Omega)}, \quad (4.129)$$

with c an integration constant. We need to solve the initial condition which then fixes:

$$Y^\phi = Y_0^\phi \frac{\sin \alpha_0}{\sin(\alpha_0 + \lambda\Omega)}, \quad (4.130)$$

with Y_0^ϕ the initial vector field at (α_0, ϕ_0) . We therefore find that moving along a path of constant longitude we have that the vector field after a parameter time λ is at

$$Y^\theta = Y_0^\theta, \quad Y^\phi = Y_0^\phi \frac{\sin \alpha_0}{\sin(\alpha_0 + \lambda\Omega)}. \quad (4.131)$$

From this we can obtain the new transformed vector field at the end-point of our path by substituting in $\lambda = 1$.

Now consider the constant latitude path. We have $X^\mu = (0, \phi_1 - \phi_0) \equiv (0, \omega)$. The resultant parallel transport equations are:

$$\begin{aligned} 0 &= \frac{d}{d\lambda} Y^\theta - \omega \sin \theta \cos \theta Y^\phi, \\ 0 &= \frac{d}{d\lambda} Y^\phi + \omega \cot \theta Y^\theta. \end{aligned} \quad (4.132)$$

We now have a pair of coupled ODEs. Recall that along the path we are taking we have $\dot{\theta} = 0$ and therefore to uncouple the equations we take a second derivative of the first. This gives:

$$\frac{d^2}{d\lambda^2} Y^\theta = -\omega^2 \cos^2 \theta Y^\theta, \quad (4.133)$$

which has solution:

$$Y^\theta = A \sin(\omega \cos \theta_{\text{in}} \lambda) + B \cos(\omega \cos \theta_{\text{in}} \lambda), \quad (4.134)$$

and therefore for Y^ϕ we find:

$$Y^\phi = \frac{1}{\sin \theta_{\text{in}}} (A \cos(\omega \cos \theta_{\text{in}} \lambda) - B \sin(\omega \cos \theta_{\text{in}} \lambda)). \quad (4.135)$$

We need to fix the initial conditions at $\lambda = 0$ which fixes:

$$B = Y_0^\theta, \quad A = \sin \theta_{\text{in}} Y_0^\phi. \quad (4.136)$$

Therefore we find that the parallel transported vector field is:

$$\begin{aligned} Y^\theta &= \sin \theta_{\text{in}} Y_0^\phi \sin(\omega \cos \theta_{\text{in}} \lambda) + Y_0^\theta \cos(\omega \cos \theta_{\text{in}} \lambda), \\ Y^\phi &= Y_0^\phi \cos(\omega \cos \theta_{\text{in}} \lambda) - \frac{1}{\sin \theta_{\text{in}}} Y_0^\theta \sin(\omega \cos \theta_{\text{in}} \lambda). \end{aligned} \quad (4.137)$$

We now have the transformation of a vector field along both the longitudinal direction and the latitudinal direction. To fix the transformation of the vector field along the loop we need to combine these. We want to perform the map:

$$(\theta_0, \phi_0) \rightarrow (\theta_1, \phi_0) \rightarrow (\theta_1, \phi_1) \rightarrow (\theta_0, \phi_1) \rightarrow (\theta_0, \phi_0). \quad (4.138)$$

For simplicity let the initial vector field at (θ_0, ϕ_0) be $Y_0 = \partial_\theta$, i.e. $Y_0^\theta = 1, Y_0^\phi = 0$ and let us transform this to each of the points using (4.131) and (4.137).

For the first transformation we find:

$$Y_1^\theta = 1, \quad Y_1^\phi = 0. \quad (4.139)$$

For the second, using (4.137) we find:

$$Y_2^\theta = \cos(\omega \cos \theta_1), \quad Y_2^\phi = -\frac{1}{\sin \theta_1} \sin(\omega \cos \theta_1), \quad \omega = \phi_1 - \phi_0. \quad (4.140)$$

For the third path we find:

$$Y_3^\theta = \cos(\omega \cos \theta_1), \quad Y_3^\phi = -\frac{1}{\sin \theta_0} \sin(\omega \cos \theta_1). \quad (4.141)$$

For the final path we have:

$$\begin{aligned} Y_4^\theta &= -\sin(\omega \cos \theta_1) \sin(\tilde{\omega} \cos \theta_0) + \cos(\omega \cos \theta_1) \cos(\tilde{\omega} \cos \theta_0), \\ Y_4^\phi &= -\frac{1}{\sin \theta_0} (\sin \omega \cos \theta_1) \cos(\tilde{\omega} \cos \theta_0) + \cos(\omega \cos \theta_1) \sin(\tilde{\omega} \cos \theta_0), \end{aligned} \quad (4.142)$$

where $\tilde{\omega} = \phi_0 - \phi_1 = -\omega$. We may simplify this using trig identities to find:

$$Y_4^\theta = \cos(\omega(\cos \theta_0 - \cos \theta_1)), \quad Y_4^\phi = \frac{1}{\sin \theta_0} \sin(\omega(\cos \theta_0 - \cos \theta_1)). \quad (4.143)$$

We immediately see that we need not obtain the same vector field once we have gone around the loop! This is a consequence of curvature. We see that after going around the loop the vector field we have obtained has been transformed. Thus this operation gives a map $T_p(M) \rightarrow T_p(M)$. The set of transformations is known as the *Holonomy* group. For our example this is an $SO(2)$ rotation. In general for a metric compatible connection the holonomy is a subgroup of $SO(m)$ for a Euclidean manifold and $SO(1, m-1)$ for a Lorentzian manifold.

Note that we can set $\theta_0 = \frac{\pi}{2}$, $\omega = \frac{\pi}{2}$ and $\theta_1 = 0$ which is then the triangle one can make with three 90 degree angles starting from the equator.²² The resultant vector field is $Y^\mu = (0, -1)^\mu$, which undergoes a 90 degree rotation.

4.4.5 Geodesic deviation

In Euclidean space or in Minkowski spacetime, geodesics which are initially parallel will remain parallel forever. On a general curved manifold this notion of parallel is not possible, instead we can study whether nearby geodesics move together or apart, and characterise their relative acceleration.

Consider a one-parameter family of geodesics with coordinates $x^\mu(\tau : \sigma)$. Here τ is the affine parameter along the geodesics, all of which are tangent to the vector field X . Thus, along the surface spanned by $x^\mu(\tau : \sigma)$ we have

$$\left. \frac{\partial x^\mu}{\partial \tau} \right|_\sigma = X^\mu. \quad (4.144)$$

The parameter σ labels the different geodesics, see figure 15. We can compute the tangent vector in the σ direction to be generated by a second vector field S so that

$$S^\mu = \left. \frac{\partial x^\mu}{\partial \sigma} \right|_\tau. \quad (4.145)$$

This tangent vector is known as the *deviation vector*, its job is to take us from one geodesic to a nearby geodesic with the same affine parameter τ . The family of geodesics sweep out a 2d surface embedded in the manifold. We have freedom to choose coordinates so that on the surface $S = \frac{\partial}{\partial \sigma}$ and $X = \frac{\partial}{\partial \tau}$ consequently we have $[X, S] = 0$.

²²Recall that the metric we took is bad at the poles $\theta = 0, \theta = \pi$. Despite this we can take this limit and still find a sensible result.




Figure 15: The black lines are geodesics generated by X while the red lines label constant τ and are generated by S with $[X, S] = 0$.

We can ask how neighbouring geodesics behave, do they converge, diverge, or remain the same distance apart? Consider a torsion free connection so that

$$\nabla_X S - \nabla_S X = [X, S]. \quad (4.146)$$

Since $[X, S] = 0$, we have

$$\nabla_X \nabla_X S = \nabla_X \nabla_S X = \nabla_S \nabla_X X + R(X, S)X, \quad (4.147)$$

where we have used the expression for the Riemann tensor in (4.62). Since X is tangent to geodesics we have $\nabla_X X = 0$ and therefore

$$\nabla_X \nabla_X S = R(X, S)X. \quad (4.148)$$

In index notation we have

$$X^\nu \nabla_\nu (X^\rho \nabla_\rho S^\mu) = R^\mu{}_{\nu\rho\sigma} X^\nu X^\rho S^\sigma. \quad (4.149)$$

If we take an integral curve γ associated to X as before we have

$$\frac{D^2 S^\mu}{D\tau^2} = R^\mu{}_{\nu\rho\sigma} X^\nu X^\rho S^\sigma, \quad (4.150)$$

with $D/D\tau$ the covariant derivative along the curve γ , $D/D\tau \equiv \frac{\partial x^\mu}{\partial \tau} \nabla_\mu$. The left hand side tells us how the deviation vector S changes as we move along the geodesic and it measures the relative acceleration of neighbouring geodesics. From (4.150) we see that the relative acceleration of neighbouring geodesics is measured by the Riemann tensor. This is nothing other than the *tidal forces* mentioned previously. Note that the relative acceleration vanishes for all families of geodesics if and only if the Riemann tensor vanishes.

4.5 Riemann tensor and its symmetries

We have just seen that the Riemann tensor is responsible for tidal forces, let us now study this tensor in more detail. The components of the Riemann tensor are given in (4.65). It is not hard to see that it is anti-symmetric in the final two indices:

$$R^\sigma{}_{\rho\mu\nu} = -R^\sigma{}_{\rho\nu\mu}. \quad (4.151)$$

This does not exhaust the symmetries however.

Exercise 4.4: Symmetries of the Riemann tensor

Prove that the Riemann tensor satisfies the identities: If we lower an index then we have

$$R_{\mu\nu\rho\sigma} = R_{\rho\sigma\mu\nu}, \quad (4.152)$$

$$R_{\mu[\nu\rho\sigma]} = 0, \quad (4.153)$$

$$\nabla_{[\mu} R_{\sigma\rho]\tau\nu} = 0. \quad (4.154)$$

Hint: These expressions can be proven by using normal coordinates.

4.5.1 Ricci and Einstein tensors

Given a rank $(1, 3)$ tensor we can construct a rank $(0, 2)$ tensor by contraction, for the Riemann tensor the resultant $(0, 2)$ -rank tensor is called the Ricci tensor.

Definition 40 (Ricci tensor) *The Ricci tensor is*

$$R_{\mu\nu} = R^\rho{}_{\mu\rho\nu}. \quad (4.155)$$

It inherits symmetry in its indices from the properties of the Riemann tensor and therefore

$$R_{\mu\nu} = R_{\nu\mu}. \quad (4.156)$$

We can play a similar game to create a scalar by contracting the indices again, the resultant scalar is known as the Ricci-scalar.

Definition 41 (Ricci Scalar) *The Ricci scalar is*

$$R = g^{\mu\nu} R_{\mu\nu}. \quad (4.157)$$

Using the metric compatible connection, the Bianchi identity implies that

$$\nabla^\mu \left(R_{\mu\nu} - \frac{1}{2} R g_{\mu\nu} \right) = 0. \quad (4.158)$$

Definition 42 (Einstein tensor) *The covariantly constant tensor*

$$G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu}, \quad (4.159)$$

is called the Einstein tensor.

This will appear in the next section when we consider GR and its conservation has physical consequences.

4.6 Summary

We have now introduced the basics of Riemannian geometry. You should understand the following:

- A **metric** $g_{\mu\nu}(x)$ is a symmetric $(0, 2)$ tensor that defines distances and angles on a manifold:

$$ds^2 = g_{\mu\nu}(x) dx^\mu dx^\nu. \quad (4.160)$$

- The metric allows us to:

- measure lengths of vectors,
- define angles between vectors,
- compute distances along curves,
- define a canonical volume form.

- A **Lorentzian metric** has signature $(-, +, +, +)$ and allows us to classify vectors as:

$$\begin{cases} g(X, X) < 0 & \text{timelike,} \\ g(X, X) = 0 & \text{null/lightlike,} \\ g(X, X) > 0 & \text{spacelike.} \end{cases} \quad (4.161)$$

- The metric provides a map between vectors and one-forms:

$$V^\mu \leftrightarrow V_\mu = g_{\mu\nu}V^\nu. \quad (4.162)$$

- The **connection** ∇ defines how to differentiate tensor fields on a manifold. It maps tensor fields to tensor fields.

- The **Levi-Civita connection** is uniquely determined by:

- metric compatibility: $\nabla_\rho g_{\mu\nu} = 0$,

– vanishing torsion.

- Its components are the **Christoffel symbols**:

$$\Gamma^\mu{}_{\nu\rho} = \frac{1}{2}g^{\mu\sigma} (\partial_\nu g_{\sigma\rho} + \partial_\rho g_{\sigma\nu} - \partial_\sigma g_{\nu\rho}), \quad (4.163)$$

which is *not* a tensor.

- The **parallel transport** of a tensor T along the integral curve of the vector field X is defined by finding the solution to

$$\nabla_X T = 0. \quad (4.164)$$

It tells you how T evolves as we move it along the curve with tangent X .

- **Geodesics** are curves that parallel transport their own tangent vector:

$$\frac{d^2 x^\mu}{d\tau^2} + \Gamma^\mu{}_{\nu\rho} \frac{dx^\nu}{d\tau} \frac{dx^\rho}{d\tau} = 0. \quad (4.165)$$

- The **Riemann tensor** measures curvature:

$$R^\mu{}_{\nu\rho\sigma} = \partial_\rho \Gamma^\mu{}_{\nu\sigma} - \partial_\sigma \Gamma^\mu{}_{\nu\rho} + \Gamma^\mu{}_{\rho\lambda} \Gamma^\lambda{}_{\nu\sigma} - \Gamma^\mu{}_{\sigma\lambda} \Gamma^\lambda{}_{\nu\rho}. \quad (4.166)$$

- Curvature leads to **geodesic deviation**, describing how nearby geodesics accelerate relative to each other.

- Contractions of the Riemann tensor give:

- the **Ricci tensor** $R_{\mu\nu}$,
- the **Ricci scalar** R .

5 Einstein's equations

After defining all this mathematics we can now use it to introduce general relativity. Like the other forces, gravity is also mediated by some field, in this case it is the metric $g_{\mu\nu}$. It is a dynamical object, not something fixed and therefore there must be some rules as to how it can evolve. These are provided by the equations of motion following from the *Einstein–Hilbert* action.

5.1 The Einstein–Hilbert action

We want to write down an action for the gravity. Differential geometry places some rigid constraints on what this can be. We want the action to be diffeomorphism invariant, since physics should not depend on the choice of coordinates. It should therefore depend on the intrinsic properties of the metric.

Spacetime is a manifold M equipped with a metric of Lorentzian signature. The action is an integral over M and so we require a volume-form. Thankfully the metric provides us with a canonical volume-form, which is reparametrisation invariant with which we can integrate any scalar. Given that we only have a metric there is not really much that we can construct. The simplest non-trivial scalar we can construct is the Ricci scalar, and therefore we can guess the action

$$S_{\text{EH}} = \int d^4x \sqrt{-g} R. \quad (5.1)$$

As a quick check since the Ricci scalar takes the form $R \sim \partial\Gamma + \Gamma\Gamma$ and the Levi–Civita connection is $\Gamma \sim \partial g$ it follows that the action is second derivative in the metric. This is like all other actions that we have considered previously, they all had two derivatives of our fundamental field.

The equations of motion will follow from varying the action with respect to the metric. We start with a fixed metric and see how the action varies as we shift according to

$$g_{\mu\nu}(x) \rightarrow g_{\mu\nu}(x) + \delta g_{\mu\nu}(x). \quad (5.2)$$

Writing the Ricci scalar as $R = g^{\mu\nu} R_{\mu\nu}$ the Einstein–Hilbert action changes as

$$\delta S = \int d^4x \left((\delta\sqrt{-g}) g^{\mu\nu} R_{\mu\nu} + \sqrt{-g} (\delta g^{\mu\nu}) R_{\mu\nu} + \sqrt{-g} g^{\mu\nu} \delta R_{\mu\nu} \right). \quad (5.3)$$

It turns out that it is simpler to consider the variation with respect to the inverse metric, this is of course equivalent to considering the variation with the metric since

$$g_{\mu\nu} g^{\nu\rho} = \delta_\mu^\rho, \quad \Rightarrow \quad (\delta g_{\mu\nu}) g^{\nu\rho} + g_{\mu\nu} \delta g^{\nu\rho} = 0, \quad \Rightarrow \quad \delta g^{\nu\rho} = -g^{\nu\sigma} g^{\rho\mu} \delta g_{\sigma\mu}. \quad (5.4)$$

The second term in the variation of the Einstein–Hilbert action is already proportional to $\delta g^{\mu\nu}$, so we need to consider the first and third terms only. First consider the variation of the determinant term. To do this we must remember a few properties of a diagonalisable matrix A , namely

$$\log \det A = \operatorname{tr} \log A. \quad (5.5)$$

(To prove this use that this is clearly true for a diagonal matrix since the determinant is the product of the eigenvalues while the trace is the sum of the eigenvalues. Since both the determinant and trace are invariant under conjugation it follows for any diagonalisable matrix.) Thus we have

$$\frac{1}{\det A} \delta \det A = \operatorname{tr}(A^{-1} \delta A), \quad (5.6)$$

after recalling the properties of the (matrix) Logarithm. Applying this to the metric we have

$$\delta \sqrt{-g} = \frac{1}{2\sqrt{-g}} (-g) g^{\mu\nu} \delta g_{\mu\nu} = \frac{1}{2} \sqrt{-g} g^{\mu\nu} \delta g_{\mu\nu}. \quad (5.7)$$

Using the identity (5.4) we have

$$\delta \sqrt{-g} = -\frac{1}{2} \sqrt{-g} g_{\mu\nu} \delta g^{\mu\nu}, \quad (5.8)$$

as claimed.

With this the variation of the Einstein–Hilbert action takes the form

$$\delta S = \int d^4x \sqrt{-g} \left(R_{\mu\nu} - \frac{1}{2} R g_{\mu\nu} \right) \delta g^{\mu\nu} + \sqrt{-g} g^{\mu\nu} \delta R_{\mu\nu}, \quad (5.9)$$

and it remains to consider the final term only. We claim that this term is a total derivative and can therefore be neglected by using Stoke’s theorem under suitable assumptions of spacetime (no boundary). To confirm this we need to prove the following identity:

$$\delta R_{\mu\nu} = \nabla_\rho \delta \Gamma^\rho_{\mu\nu} - \nabla_\nu \delta \Gamma^\rho_{\mu\rho}, \quad (5.10)$$

where

$$\delta \Gamma^\rho_{\mu\nu} = \frac{1}{2} g^{\rho\sigma} (\nabla_\mu \delta g_{\sigma\nu} + \nabla_\nu \delta g_{\mu\sigma} - \nabla_\sigma \delta g_{\mu\nu}). \quad (5.11)$$

We start by looking at the variation of the Christoffel symbols $\Gamma^\rho_{\mu\nu}$. Though the Christoffel symbol is not a tensor the variation $\delta \Gamma^\rho_{\mu\nu}$ is a tensor. This is because it is the difference of two Christoffel symbols, one computed using the metric $g_{\mu\nu}$ and one with $g_{\mu\nu} + \delta g_{\mu\nu}$ and the term in the transformation of the Christoffel which shows that it is not a tensor is independent of the metric and therefore cancels in the difference. This observation makes our

lives a lot simpler. It implies that at any point $p \in M$ we can work in normal coordinates such that $\partial_\rho g_{\mu\nu}|_p = 0$ and therefore $\Gamma^\rho{}_{\mu\nu}|_p = 0$. To linear order in the variation the change in the Christoffel symbol evaluated at p is

$$\begin{aligned}\delta\Gamma^\rho{}_{\mu\nu}|_p &= \frac{1}{2}g^{\rho\sigma}(\partial_\mu\delta g_{\sigma\nu} + \partial_\nu\delta g_{\sigma\mu} - \partial_\sigma\delta g_{\mu\nu})|_p \\ &= \frac{1}{2}g^{\rho\sigma}(\nabla_\mu\delta g_{\sigma\nu} + \nabla_\nu\delta g_{\sigma\mu} - \nabla_\sigma\delta g_{\mu\nu})|_p\end{aligned}\tag{5.12}$$

where we have used that in normal coordinates we can replace partial derivatives with covariant derivatives. Both the left and right hand side are tensors and therefore this holds in any coordinate system, moreover the point p was arbitrary and therefore this holds in all coordinate systems at all points $p \in M$.

Key Points

When using normal coordinates it is important that the two objects being compared at the end are tensors. In this case it follows that it must then hold for all choices of coordinates since both sides of the equation transform in the same way.

Next consider the variation of the Riemann tensor. In normal coordinates we have

$$R^\sigma{}_{\rho\mu\nu} = \partial_\mu\Gamma^\sigma{}_{\nu\rho} - \partial_\nu\Gamma^\sigma{}_{\mu\rho},\tag{5.13}$$

and the variation is

$$\delta R^\sigma{}_{\rho\mu\nu} = \partial_\mu\delta\Gamma^\sigma{}_{\nu\rho} - \partial_\nu\delta\Gamma^\sigma{}_{\mu\rho} = \nabla_\mu\delta\Gamma^\sigma{}_{\nu\rho} - \nabla_\nu\delta\Gamma^\sigma{}_{\mu\rho},\tag{5.14}$$

where we have once again used that in normal coordinates we can replace partial derivatives with covariant derivatives. As before we have a tensorial equation and therefore this must hold in any coordinate system not just normal coordinates. Contracting the σ and μ indices we find

$$\delta R_{\rho\nu} = \nabla_\mu\delta\Gamma^\mu{}_{\nu\rho} - \nabla_\nu\delta\Gamma^\mu{}_{\rho\mu}.\tag{5.15}$$

It follows that

$$g^{\mu\nu}\delta R_{\mu\nu} = \nabla_\mu(g^{\rho\nu}\delta\Gamma^\mu{}_{\rho\nu} - g^{\mu\nu}\delta\Gamma^\rho{}_{\nu\rho}) = \nabla_\mu X^\mu,\tag{5.16}$$

with X^μ the bracketed tensor. The variation of the Einstein–Hilbert action can then be written as

$$\delta S = \int d^4x\sqrt{-g}\left[\left(R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu}\right)\delta g^{\mu\nu} + \nabla_\mu X^\mu\right].\tag{5.17}$$

The final term is a total derivative after using the identity (4.81) and with suitable assumptions on spacetime can be neglected. Requiring that the action is extremised, so that $\delta S = 0$, implies the equation of motion

$$G_{\mu\nu} := R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} = 0. \quad (5.18)$$

These are the *Einstein field equations* in the absence of matter. We may further simplify them by first contracting with the inverse metric to find $R = 0$ and therefore in the absence of matter Einstein's equations are simply

$$R_{\mu\nu} = 0. \quad (5.19)$$

A metric obeying this equation is known as Ricci flat. Though this looks deceptively simple this holds a very rich set of solutions, in fact not all solutions to this equation are known. Notice that it does not imply that the Riemann tensor vanishes!

Physical Explanation

The Einstein equation in a vacuum tells us that there is a constraint on the curvature of spacetime.

We threw away the boundary term in the usual cavalier way one does with such variational principles. We may make this more rigorous by introducing the Gibbons–Hawking boundary term to allow for M to admit a boundary. The addition of this term gives the same Einstein field equations as before even for a manifold with a boundary.

5.1.1 Newton's constant

As it stands the action we have given does not have the correct dimension, this will become a problem when we want to couple to matter, so let us remedy this now. We take the coordinates to have dimension of length $[x] = L$ and therefore the metric is dimensionless. The Ricci scalar involves two derivatives and therefore it has dimension $[R] = L^{-2}$. Including the dimension of the integration measure the current action in (5.1) has dimension $[S] = L^2$. An action should have dimension of Energy \times time and therefore we should multiply the action by an appropriate dimensionful constant. This constant is known as Newton's constant. The correct action is:

$$S_{\text{EH}} = \frac{c^3}{16\pi G_N} \int d^4x \sqrt{-g} R, \quad (5.20)$$

where c is of course the speed of light, and G_N is Newton's constant

$$G_N \sim 6.67 \times 10^{-11} m^3 \text{kg}^{-1} s^{-2}. \quad (5.21)$$

This will not change the equation of motion in the vacuum but once we couple to matter this will determine the strength of the coupling of the gravitational field to the matter.

If we are just interested in phenomena related to gravity it is sensible to set $G_N = 1$. Instead if we want to consider other phenomena other than gravity this is not so sensible since it defines the coupling of the forces. Instead the more useful convention is to pick $\hbar = 1$, which equates energy with time. With this convention Newton's constant has dimension $[G] = m^{-2}$. The corresponding energy scale is called the *Planck mass* and is given by

$$M_{pl}^2 = \frac{\hbar c}{8\pi G_N}. \quad (5.22)$$

It is around 10^{18} GeV which is a very high energy scale and far beyond anything we can probe experimentally. This is why the gravitational force is so weak, the coupling constant is much smaller than that of the other forces.

5.1.2 Cosmological constant

We motivated the Einstein–Hilbert action as the simplest action one can write down. There is in fact a simpler term we may write down other than the Einstein–Hilbert term we considered previously. We may simply add a constant to the volume form. The resulting action is

$$S = \frac{1}{16\pi G_N} \int d^4x \sqrt{-g} (R - 2\Lambda). \quad (5.23)$$

The constant Λ is known as the *cosmological constant* and has dimension $[\Lambda] = L^{-2}$. The minus sign in the action comes from thinking of the Lagrangian as $T - V$ with the cosmological constant playing the role of the potential energy V .

Varying the action as before yields the Einstein equations

$$R_{\mu\nu} - \frac{1}{2} R g_{\mu\nu} = -\Lambda g_{\mu\nu}. \quad (5.24)$$

This time if we contract with the inverse metric we get $R = 4\Lambda$. Substituting this back into the vacuum Einstein equations, in the presence of a cosmological constant they become

$$R_{\mu\nu} = \Lambda g_{\mu\nu}. \quad (5.25)$$

Metrics satisfying this property are known as Einstein metrics.

Einstein considered adding the cosmological constant his biggest blunder. He was searching for a static, non-expanding universe which he could not find in the vacuum equations. By adding the Cosmological constant he could counteract the gravitational collapse and achieve a balanced static universe. In 1929 Edwin Hubble had discovered that the universe is actually expanding and so Einstein abandoned the constant calling it his biggest blunder. Modern

cosmologists have revived the cosmological constant to represent dark energy and we believe that there is a small positive cosmological constant (we live in a de Sitter universe). Einstein's addition of the cosmological constant was not a really a blunder, rather not trusting his own theory enough was the true blunder.

5.1.3 Higher derivative terms

The Einstein–Hilbert action with cosmological constant is the simplest thing we can write down. We may construct other scalars from the Riemann tensor, however they will introduce higher derivative terms. For example, there are three terms that we can add at four-derivative level in the metric

$$S_{4\text{-deriv}} = \int d^4x \sqrt{-g} \left(c_1 R^2 + c_2 R_{\mu\nu} R^{\mu\nu} + c_3 R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} \right), \quad (5.26)$$

with the c_i dimensionless constants. Generic choices of the constants will not give rise to higher derivative equations of motion with a well-defined initial value problem. Nonetheless there are certain combinations which conspire to keep the equations second order in derivatives. This goes by the name of *Lovelock's theorem* and says that in four-dimensions the combination

$$\frac{1}{8\pi^2} \int_M d^4x \sqrt{g} (R^2 - 4R_{\mu\nu} R^{\mu\nu} + R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma}) = \chi(M), \quad (5.27)$$

where $\chi(M) \in \mathbb{Z}$, is the Euler character of M . Though not obvious, this is a total derivative, and therefore does not affect the classical equations of motion. Higher derivative terms only become relevant for fast varying fields. For us these terms will not be important and therefore we stick to the 2-derivative action.

5.1.4 Diffeomorphisms

A natural question to ask is how many degrees of freedom are there in the metric? Since it is a 4×4 symmetric matrix the naive guess is $\frac{1}{2} \times 4 \times 5 = 10$ however this is not quite correct. Not all of these 10 components are physical. Two metrics which are related by a change of coordinates $x^\mu \rightarrow \tilde{x}^\mu(x)$ describe the same physical spacetime. This means that there is a redundancy in any given representation of the metric which removes precisely 4 of the 10 degrees of freedom, leaving just 6 actual degrees of freedom.

This redundancy is implemented by diffeomorphisms. Recall that a diffeomorphism is a map $\phi : M \rightarrow M$. We may use it to map all fields, including the metric on M to a new set of fields on M . The end result is physically indistinguishable from the original, it describes the same system just in a different set of coordinates. Such diffeomorphisms are analogous to the gauge transformations of a gauge theory such as electromagnetism.

Let us look at how diffeomorphisms modify the action. Consider a diffeomorphism which takes a point with coordinate x^μ to a nearby point with coordinates

$$x^\mu \rightarrow \tilde{x}^\mu = x^\mu + \delta x^\mu . \quad (5.28)$$

We can view this either as an active change in which one point with coordinates x^μ is mapped to another point with coordinates $x^\mu + \delta x^\mu$ or as a passive transformation in which we use two different coordinate patches to label the same point. Either viewpoint leads to the same conclusion, here we will take the passive viewpoint.

We can think of the change of coordinates as being generated by an infinitesimal vector field X ,

$$\delta x^\mu = -X^\mu(x) . \quad (5.29)$$

Under the change of coordinates the metric transforms as

$$g_{\mu\nu}(x) \rightarrow \tilde{g}_{\mu\nu}(\tilde{x}) = \frac{\partial x^\rho}{\partial \tilde{x}^\mu} \frac{\partial x^\sigma}{\partial \tilde{x}^\nu} g_{\rho\sigma}(x) . \quad (5.30)$$

We can invert the Jacobian matrix to find

$$\frac{\partial \tilde{x}^\mu}{\partial x^\rho} = \delta_\rho^\mu - \partial_\rho X^\mu \quad \Rightarrow \quad \frac{\partial x^\rho}{\partial \tilde{x}^\mu} = \delta_\mu^\rho + \partial_\mu X^\rho , \quad (5.31)$$

where the inverse holds to leading order in the variation X . Continuing to work infinitesimally we have

$$\begin{aligned} \tilde{g}_{\mu\nu}(\tilde{x}) &= (\delta_\mu^\rho + \partial_\mu X^\rho)(\delta_\nu^\sigma + \partial_\nu X^\sigma) g_{\rho\sigma}(x) \\ &= g_{\mu\nu}(x) + g_{\mu\rho}(x) \partial_\nu X^\rho + g_{\nu\rho}(x) \partial_\mu X^\rho . \end{aligned} \quad (5.32)$$

We can also Taylor expand the left-hand side to find

$$\tilde{g}_{\mu\nu}(\tilde{x}) = \tilde{g}_{\mu\nu}(x + \delta x) = \tilde{g}_{\mu\nu}(x) - X^\lambda \partial_\lambda \tilde{g}_{\mu\nu}(x) . \quad (5.33)$$

Comparing the different metrics at the same point x we find that the metric undergoes the infinitesimal change

$$\delta g_{\mu\nu}(x) = \tilde{g}_{\mu\nu}(x) - g_{\mu\nu}(x) = X^\lambda \partial_\lambda g_{\mu\nu} + g_{\mu\rho} \partial_\nu X^\rho + g_{\nu\rho} \partial_\mu X^\rho . \quad (5.34)$$

This is precisely the Lie derivative of the metric. If we act with an infinitesimal diffeomorphism along X then the metric changes as

$$\delta g_{\mu\nu} = (\mathcal{L}_X g)_{\mu\nu} . \quad (5.35)$$

We may rewrite equation (5.34) by lowering the index on X^ρ to find

$$\delta g_{\mu\nu} = \partial_\mu X_\nu + \partial_\nu X_\mu + X^\rho (\partial_\rho g_{\mu\nu} - \partial_\mu g_{\rho\nu} - \partial_\nu g_{\mu\rho}), \quad (5.36)$$

the last term is just the Christoffel symbols and therefore we have

$$\delta g_{\mu\nu} = \nabla_\mu X_\nu + \nabla_\nu X_\mu. \quad (5.37)$$

We may put this together to see how the action changes. Under a general change of the metric the Einstein–Hilbert action changes as

$$\delta S = \int d^4x \sqrt{-g} G^{\mu\nu} \delta g_{\mu\nu}, \quad (5.38)$$

where we have discarded the boundary term. Insisting that $\delta S = 0$ for any variation $\delta g_{\mu\nu}$ gives the equations of motion $G^{\mu\nu} = 0$. In contrast, symmetries of the action are those variations $\delta g_{\mu\nu}$ for which $\delta S = 0$ for any choice of metric. Since diffeomorphisms are symmetries we know that the action is invariant under changes of the form (5.37). Using the fact that $G_{\mu\nu}$ is symmetric we must have

$$\delta S = 2 \int d^4x \sqrt{-g} G^{\mu\nu} \nabla_\mu X_\nu = 0, \quad \text{for all } X_\mu(x). \quad (5.39)$$

After integrating by parts we find that this results in the Bianchi identity

$$\nabla_\mu G^{\mu\nu} = 0. \quad (5.40)$$

We learn that from the path integral perspective the Bianchi identity is a result of diffeomorphism invariance.

5.1.5 Coupling to matter

Until now the action has only involved gravity, and at most we can allow for test particles moving on geodesics. Since they are test particles we ignore any back-reaction they produce on spacetime. However matter is not just an actor doing what gravity says in spacetime, it also back-reacts and affects the dynamics of spacetime. The first question to ask is how does matter couple to the metric? We will consider matter that comes with a Lagrangian in Minkowski spacetime.

Scalar Field Consider first a scalar field $\phi(x)$. In flat spacetime the action takes the form

$$S_{\text{scalar}} = \int d^4x \left(-\frac{1}{2} \eta^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - V(\phi) \right), \quad (5.41)$$

with $\eta^{\mu\nu}$ the inverse Minkowski metric.²³

It is straightforward to generalise this to describe a field moving in curved spacetime, we simply need to replace the Minkowski metric with the curved metric, replace partial derivatives with covariant derivatives and introduce the volume form when we integrate in the action. This means that we take

$$S_{\text{scalar}} = \int d^4x \sqrt{-g} \left(-\frac{1}{2} g^{\mu\nu} \nabla_\mu \phi \nabla_\nu \phi - V(\phi) \right). \quad (5.42)$$

Despite upgrading the partial derivatives to covariant ones this is somewhat redundant here as they act the same on a scalar field: we keep it for later though.

Curved spacetime also introduces new possibilities for us to add to the action, for example we could add a term such as $\xi R \phi^2$ to the action which gives rise to extra couplings. We will not interest ourselves in such terms here however.

Maxwell Theory The action of Maxwell theory from special relativity is

$$S_{\text{Maxwell}} = -\frac{1}{4} \int d^4x \eta^{\mu\rho} \eta^{\nu\sigma} F_{\mu\nu} F_{\rho\sigma}, \quad (5.43)$$

with $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$. The electric and magnetic fields are encoded in F via

$$F_{\mu\nu} = \begin{pmatrix} 0 & -E_1 & -E_2 & -E_3 \\ E_1 & 0 & B_3 & -B_2 \\ E_2 & -B_3 & 0 & B_1 \\ E_3 & B_2 & -B_1 & 0 \end{pmatrix}, \quad (5.44)$$

and the Bianchi identity $dF = d^2A = 0$ yields two of the four Maxwell equations

$$\nabla \cdot \vec{B} = 0, \quad \nabla \times \vec{B} + \frac{\partial \vec{E}}{\partial t} = 0. \quad (5.45)$$

See problem sheet 1.

We may couple to curved space time through the minimal coupling outlined for the scalar theory. The action is

$$S_{\text{Maxwell}} = -\frac{1}{4} \int d^4x \sqrt{-g} g^{\mu\rho} g^{\nu\sigma} F_{\mu\nu} F_{\rho\sigma} = -\frac{1}{2} \int F \wedge \star F. \quad (5.46)$$

We again take $F = dA$, which in components reads $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu = \nabla_\mu A_\nu - \nabla_\nu A_\mu$. Antisymmetry implies that we may replace the covariant derivatives with normal derivatives. The equations of motion are

$$\nabla^\mu F_{\mu\nu} = 0, \quad \Leftrightarrow \quad d \star F = 0. \quad (5.47)$$

²³Note that the minus sign is due to our mostly plus signature convention, you may be more used to the opposite convention when considering a field theory. The Lagrangian will take the form of kinetic energy minus potential energy.

We have now seen the algorithm of how to couple matter to gravity for two examples, for generic matter we follow the exact same rules. It remains to be seen how coupling to matter change the Einstein equations of the previous section. We need to consider the combined action

$$S = \frac{1}{16\pi G_N} \int d^4x \sqrt{-g} (R - 2\Lambda) + S_{\text{Matter}}, \quad (5.48)$$

where S_{Matter} is the action for any matter fields in the theory minimally coupled to gravity. When we vary the Einstein–Hilbert term we know that we will obtain the Einstein tensor, what about S_{Matter} ?

Definition 43 (Energy-Momentum tensor) *We define the Energy-Momentum tensor to be*

$$T_{\mu\nu} = -\frac{2}{\sqrt{-g}} \frac{\delta \mathcal{L}_{\text{Matter}}}{\delta g^{\mu\nu}}. \quad (5.49)$$

By construction $T_{\mu\nu}$ is symmetric. Varying the full action with respect to the metric we have

$$\delta S = \frac{1}{16\pi G_N} \int d^4x \sqrt{-g} (G_{\mu\nu} + \Lambda g_{\mu\nu}) \delta g^{\mu\nu} - \frac{1}{2} \int d^4x \sqrt{-g} T_{\mu\nu} \delta g^{\mu\nu}, \quad (5.50)$$

from which we may read the following equation of motion

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = 8\pi G_N T_{\mu\nu}. \quad (5.51)$$

This is the Einstein equation describing gravity coupled to matter. Note that the presence of the energy-momentum tensor says that the matter distribution sources the curvature of the spacetime.

Example 5.1: Computing the energy momentum tensor

- For the scalar theory above the energy-momentum tensor is

$$T_{\mu\nu} = \nabla_\mu \phi \nabla_\nu \phi - g_{\mu\nu} \left(\frac{1}{2} \nabla^\rho \phi \nabla_\rho \phi + V(\phi) \right). \quad (5.52)$$

To see this observe that

$$\mathcal{L}_{\text{scalar}} = \sqrt{-g} \left(-\frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - V(\phi) \right), \quad (5.53)$$

where we have used that the connection on a scalar field is equivalent to the usual derivative and therefore the variation with respect to the metric of the connection here

is trivial. Therefore

$$\begin{aligned}
T_{\mu\nu} &= -\frac{2}{\sqrt{-g}} \frac{\delta}{\delta g^{\mu\nu}} \left[\sqrt{-g} \left(-\frac{1}{2} g^{\mu\nu} \nabla_\mu \phi \nabla_\nu \phi - V(\phi) \right) \right] \\
&= -\frac{2}{\sqrt{-g}} \left[-\frac{1}{2} \sqrt{-g} \nabla_\mu \phi \nabla_\nu \phi + \frac{\delta \sqrt{-g}}{\delta g^{\mu\nu}} \left(-\frac{1}{2} g^{\rho\sigma} \nabla_\rho \phi \nabla_\sigma \phi - V(\phi) \right) \right] \\
&= \nabla_\mu \phi \nabla_\nu \phi - g_{\mu\nu} \left(\frac{1}{2} \nabla^\rho \phi \nabla_\rho \phi - V(\phi) \right)
\end{aligned} \tag{5.54}$$

If we restrict to flat space then

$$T_{00} = \frac{1}{2} \dot{\phi}^2 + \frac{1}{2} (\nabla \phi)^2 + V(\phi), \tag{5.55}$$

with ∇ the usual 3d spatial derivative. This is the energy density of a scalar field.

- For the Maxwell action we have

$$T_{\mu\nu} = g^{\rho\sigma} F_{\mu\rho} F_{\nu\sigma} - \frac{1}{4} g_{\mu\nu} F^{\rho\sigma} F_{\rho\sigma}, \tag{5.56}$$

see problem sheet 1 and replace $\eta \rightarrow g$. In flat space we have

$$T_{00} = \frac{1}{2} \left[\vec{E}^2 + \vec{B}^2 \right]. \tag{5.57}$$

This is the energy density of the magnetic and electric fields.

5.2 Summary

We have now seen how to derive the Einstein field equations.

- The dynamics of gravity are determined by varying the **Einstein–Hilbert action** which is the simplest, second order in derivatives, diffeomorphism invariant action possible.
- Varying the action with respect to the inverse metric, leads to the **Einstein field equations in a vacuum**:

$$G_{\mu\nu} \equiv R_{\mu\nu} - \frac{1}{2} R g_{\mu\nu} = 0 \quad \Leftrightarrow \quad R_{\mu\nu} = 0. \tag{5.58}$$

- Adding a **cosmological constant** Λ modifies the field equations to be

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = 0. \tag{5.59}$$

- To couple to matter we take the flat space action and **minimally couple**. We replace $\eta_{\mu\nu} \rightarrow g_{\mu\nu}$ $\partial_\mu \rightarrow \nabla_\mu$ and introduce the usual determinant factor $d^4x \rightarrow \sqrt{-g} d^4x$.
- The **Energy momentum tensor** is defined to be

$$T_{\mu\nu} = -\frac{2}{\sqrt{-g}} \frac{\delta \mathcal{L}_{\text{matter}}}{\delta g^{\mu\nu}}. \quad (5.60)$$

- The fully fledged Einstein equations are

$$R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} + \Lambda g_{\mu\nu} = 8\pi G_N T_{\mu\nu}. \quad (5.61)$$

- The Einstein field equations describe how matter and energy curves spacetime. In turn, the Geodesic equations tell us how spacetime makes matter move.
- **Diffeomorphism invariance** of the theory implies that not all the components of the metric are physical. Invariance of the action implies that the Einstein tensor is conserved

$$\nabla_{\mu} G^{\mu\nu} = 0, \quad (5.62)$$

this is known as the *Bianchi identity*.

6 Schwarzschild solution

Black holes are one of the most enigmatic objects and probably the reason why most of you are here. Well the moment is finally upon us and we will take our first steps to understanding black holes.

6.1 The Schwarzschild black hole

In 1915 Einstein published his work on General relativity and made a comment saying that he was not optimistic that the equations he had found could be solved except for Minkowski space. Also in 1915, with the first world war raging in Europe, Karl Schwarzschild was in the German army on the Russian front performing ballistic calculations, and suffering from pemphigus, a rare and painful autoimmune disease. Despite this, in his spare time he worked on finding solutions to general relativity and found the first exact (non-trivial) solution to Einstein's field equations.²⁴ Schwarzschild's breakthrough was to use a convenient system of coordinates, taking a polar like coordinate system as opposed to Einstein's rectangular coordinate system. The metric that bears his name is

$$ds^2 = -\left(1 - \frac{2G_N M}{r}\right) dt^2 + \left(1 - \frac{2G_N M}{r}\right)^{-1} dr^2 + r^2(d\theta^2 + \sin^2 \theta d\phi^2). \quad (6.1)$$

This solves Einstein's equations in a vacuum, $R_{\mu\nu} = 0$. The coordinate ranges are²⁵

$$t \in \mathbb{R}, \quad 0 < \theta < \pi, \quad 0 < \phi < 2\pi. \quad (6.2)$$

The range of r is slightly more subtle. At $r = 2G_N M$ something funky is happening since the prefactor of dt^2 and dr^2 vanish and diverge respectively. For the moment we will keep $2G_N M < r < \infty$ and we then do not need to worry about what is happening at $r = 2G_N M$. This value of the radial coordinate is called the *Schwarzschild radius* and will play a prominent role when we view the Schwarzschild solution as a black hole in section 6.3.

The metric depends on a single parameter M which is interpreted as the mass of the object. Indeed using our results from problem sheet 1 (or the Linearised equations and their Newtonian limit we will see later in GR2) in the weak field limit we have

$$g_{00} = -(1 + 2\Phi), \quad (6.3)$$

²⁴Schwarzschild died in 1916 having left military service due to his illness.

²⁵There are singularities at $\theta = 0, \pi$ however these are just the expected singularities from considering a two-sphere and attempting to use just one coordinate patch, as we studied in example 4.1. We should be careful about this but it is not a problem since we learnt earlier that this problem could be removed.

with Φ the Newtonian potential. For the Schwarzschild metric we have

$$\Phi = -\frac{G_N M}{r}, \quad (6.4)$$

which is the Newtonian potential for a point mass M at the origin.

Not examinable: We can compute the mass of the black hole by using Komar integrals, these will be explained in the GR2 course, but we will give a flavour of it here. The Schwarzschild solution admits a time-like Killing vector $K = \partial_t$: a Killing vector satisfies $\mathcal{L}_K g = 0$ which is equivalent to $\nabla_{(\mu} K_{\nu)} = 0$. To compute the Komar integral we must construct the dual one-form to the time-like Killing vector, (which in an abuse of notation we also call K)

$$K = g_{00} dt = -\left(1 - \frac{2GM}{r}\right) dt. \quad (6.5)$$

The Komar integral is given by

$$M_{\text{Komar}} = -\frac{1}{8\pi G_N} \int_{S^2} \star dK, \quad (6.6)$$

where the S^2 is any sphere with a radius larger than the horizon at $r = 2G_N M$ where the Killing vector has vanishing norm. Then

$$dK = -\frac{2G_N M}{r^2} dr \wedge dt \quad \Rightarrow \quad \star dK = -2G_N M \sin \theta d\theta \wedge d\phi. \quad (6.7)$$

and therefore

$$M_{\text{Komar}} = M. \quad (6.8)$$

Note that $d \star dK = 0$ and therefore it obeys an equation similar to Maxwell's equations $d \star F = 0$. These are Maxwell's equations in the absence of any current and therefore one would expect the electric charge to vanish. Yet this electric charge is precisely the mass and this is non-zero. For the solution the mass is localised at the origin $r = 0$ where the field strength diverges, and it is this singular source at $r = 0$ which gives rise to the mass.

We may thus expect that this describes something physical only when $M > 0$. For $M = 0$ we find Minkowski space while for $M < 0$ the metric becomes unphysical. For $M < 0$ we find a naked singularity (more on this in GR2) and this is generally considered unphysical.

6.1.1 Birkhoff's theorem

The Schwarzschild solution turns out to be the unique spherically symmetric asymptotically flat solution to the vacuum Einstein equations, this fact is known as *Birkhoff's theorem*. This means that the Schwarzschild solution does not just describe the spacetime outside of a black hole but outside any non-rotating, spherically symmetric object such as a star or planet.

We will sketch the proof of this fact since it allows us to get a feel for solving the Einstein equations.

The spherical symmetry of the metric means that it has an $\text{SO}(3)$ isometry. If you hold up a round sphere and rotate it it looks the same no matter which way you rotate it. If instead you did the same with a golf ball, which has dimples then this rotational symmetry is broken. The distinction between these two situations should be captured by the metric. The metric on a round two-sphere will look the same wherever you sit on the sphere whereas the metric on the golf ball will depend on where you are.

To define this mathematically we need to use the concept of a flow that we introduced a number of lectures ago. A flow on a manifold M is a one-parameter family of diffeomorphisms $\sigma_t : M \rightarrow M$, and may be associated to a vector field $K \in \mathcal{X}(M)$ at each point along the flow which is tangent to the flow

$$K^\mu = \frac{dx^\mu(\lambda)}{d\lambda}. \quad (6.9)$$

The flow is said to be an isometry if the metric looks the same at each point along a given flow line, mathematically this means that an isometry satisfies

$$\mathcal{L}_K g = 0, \quad \Leftrightarrow \quad \nabla_\mu K_\nu + \nabla_\nu K_\mu = 0. \quad (6.10)$$

A vector satisfying this equation is known as a *Killing vector field*. Sometimes it is simple to see that a vector generates an isometry, particularly when it is an ignorable coordinate, i.e. the metric does not depend on the coordinate. Sometimes, however, the Killing vectors are not so obvious.

There is a group structure underlying the symmetries, well technically a Lie algebra structure. This follows since the Lie derivative satisfies

$$\mathcal{L}_X \mathcal{L}_Y - \mathcal{L}_Y \mathcal{L}_X = \mathcal{L}_{[X,Y]}. \quad (6.11)$$

Killing vectors form a Lie algebra of the isometry group of the manifold. (See problem sheet 3 where we consider the Killing vectors on the round three-sphere).

One can then prove that the $\text{SO}(3)$ isometry implies that the metric must take the form

$$ds^2 = g_{\tau\tau}(\tau, \rho)d\tau^2 + 2g_{\tau\rho}(\tau, \rho)d\tau d\rho + g_{\rho\rho}(\tau, \rho)d\rho^2 + r^2(\tau, \rho)ds^2(S^2), \quad (6.12)$$

where

$$ds^2(S^2) = d\theta^2 + \sin^2 \theta d\phi^2, \quad (6.13)$$

is the metric on a round two-sphere. The $\text{SO}(3)$ isometry then acts on the two-sphere and leaves τ and ρ untouched. This is called a *foliation* of the space by S^2 leaves.

The size of the sphere is determined by $r(\tau, \rho)$ and it is convenient to redefine the coordinates such that r is a coordinate, we can then eliminate the ρ coordinate in favour of r . The metric becomes

$$ds^2 = g_{\tau\tau}(\tau, r)d\tau^2 + 2g_{\tau r}(\tau, r)d\tau dr + g_{rr}(\tau, r)dr^2 + r^2 ds^2(S^2). \quad (6.14)$$

The only subtlety we could encounter in doing this change of coordinates is if it is not possible to exchange ρ with r , for example r could have been independent of ρ . We can rule out these cases by imposing that asymptotically the spacetime looks like Minkowski space.

The logic now is to remove the cross term $d\tau dr$ by using a change of coordinates. If we define $\tilde{t}(\tau, r)$ then

$$d\tilde{t} = \frac{\partial \tilde{t}}{\partial \tau} d\tau + \frac{\partial \tilde{t}}{\partial r} dr \quad (6.15)$$

and therefore, we can pick a change of coordinates such that we can remove the cross term. The resultant metric is

$$ds^2 = -e^{2\alpha(\tilde{t}, r)} d\tilde{t}^2 + e^{2\beta(\tilde{t}, r)} dr^2 + r^2 ds^2(S^2). \quad (6.16)$$

We have included a minus sign since we are looking for a Lorentzian metric and then we can introduce the exponential terms which are manifestly positive definite. This is the simplest form of the metric that we can achieve just through coordinate transformations and we now need to plug this into Einstein's equations. Observe that we have used symmetries to restrict the form of the metric and then used diffeomorphisms to write the metric in the simplest form possible. This makes solving Einstein's equations simpler, the correct ansatz and choice of coordinates simplifies

We can compute the Christoffel symbols for the metric, the non-trivial ones are²⁶

$$\begin{aligned} \Gamma_{\tilde{t}\tilde{t}}^{\tilde{t}} &= \partial_{\tilde{t}}\alpha, & \Gamma_{\tilde{t}r}^{\tilde{t}} &= \partial_r\alpha, & \Gamma_{rr}^{\tilde{t}} &= e^{2\beta-2\alpha}\partial_{\tilde{t}}\beta, \\ \Gamma_{\tilde{t}\tilde{t}}^r &= e^{2\alpha-2\beta}\partial_r\alpha, & \Gamma_{\tilde{t}r}^r &= \partial_{\tilde{t}}\beta, & \Gamma_{rr}^r &= \partial_r\beta, \\ \Gamma_{r\theta}^\theta &= \frac{1}{r}, & \Gamma_{\theta\theta}^r &= -re^{-2\beta}, & \Gamma_{r\phi}^\phi &= \frac{1}{r}, \\ \Gamma_{\phi\phi}^r &= -re^{-2\beta}, & \Gamma_{\phi\phi}^\theta &= -\sin\theta\cos\theta, & \Gamma_{\theta\phi}^\phi &= \frac{\cos\theta}{\sin\theta}. \end{aligned} \quad (6.17)$$

²⁶You should see that you can do this. There is also a mathematica file on the course webpage where this has been computed for you to check.

It follows that the non-vanishing components of the Riemann tensor are

$$\begin{aligned}
R^{\tilde{t}}_{r\tilde{t}r} &= e^{2\beta-2\alpha} \left(\partial_{\tilde{t}}^2 \beta + (\partial_{\tilde{t}} \beta)^2 - \partial_{\tilde{t}} \alpha \partial_{\tilde{t}} \beta \right) + \left(\partial_r \alpha \partial_r \beta - \partial_r^2 \alpha - (\partial_r \alpha)^2 \right), \\
R^{\tilde{t}}_{\theta\tilde{t}\theta} &= -r e^{-2\beta} \partial_r \alpha, \\
R^{\tilde{t}}_{\phi\tilde{t}\phi} &= -r e^{-2\beta} \sin^2 \theta \partial_r \alpha, \\
R^{\tilde{t}}_{\theta r \theta} &= -r e^{-2\alpha} \partial_{\tilde{t}} \beta, \\
R^{\tilde{t}}_{\phi r \phi} &= -r e^{-2\alpha} \sin^2 \theta \partial_{\tilde{t}} \beta, \\
R^r_{\theta r \theta} &= r e^{-2\beta} \partial_r \beta, \\
R^r_{\phi r \phi} &= r e^{-2\beta} \sin^2 \theta \partial_r \beta, \\
R^\theta_{\phi \theta \phi} &= (1 - e^{-2\beta}) \sin^2 \theta.
\end{aligned} \tag{6.18}$$

From the Riemann tensor we can construct the Ricci tensor finding the non-trivial components

$$\begin{aligned}
R_{\tilde{t}\tilde{t}} &= \left(\partial_{\tilde{t}}^2 \beta + (\partial_{\tilde{t}} \beta)^2 - \partial_{\tilde{t}} \alpha \partial_{\tilde{t}} \beta \right) + e^{2\alpha-2\beta} \left(\partial_r^2 \alpha + (\partial_r \alpha)^2 - \partial_r \alpha \partial_r \beta + \frac{2}{r} \partial_r \alpha \right), \\
R_{rr} &= - \left(\partial_r^2 \alpha + (\partial_r \alpha)^2 - \partial_r \alpha \partial_r \beta - \frac{2}{r} \partial_r \beta \right) + e^{2\beta-2\alpha} \left(\partial_{\tilde{t}}^2 \beta + (\partial_{\tilde{t}} \beta)^2 - \partial_{\tilde{t}} \alpha \partial_{\tilde{t}} \beta \right) \\
R_{\tilde{t}r} &= \frac{2}{r} \partial_{\tilde{t}} \beta, \\
R_{\theta\theta} &= e^{-2\beta} \left(r (\partial_r \beta - \partial_r \alpha) - 1 \right) + 1, \\
R_{\phi\phi} &= R_{\theta\theta} \sin^2 \theta.
\end{aligned} \tag{6.19}$$

Our job is to now solve Einstein's equations in the vacuum, $R_{\mu\nu} = 0$. There is an obvious component to consider first, $R_{\tilde{t}r}$ which implies

$$\partial_{\tilde{t}} \beta = 0. \tag{6.20}$$

If we now take the \tilde{t} derivative of $R_{\theta\theta}$ and use the above condition we find

$$\partial_{\tilde{t}} \partial_r \alpha = 0, \tag{6.21}$$

and therefore we have

$$\beta = \beta(r), \quad \alpha = f(r) + g(\tilde{t}). \tag{6.22}$$

The first term in the metric is then

$$-e^{2f(r)+2g(\tilde{t})} d\tilde{t}^2, \tag{6.23}$$

and by a redefinition of \tilde{t} we can set

$$e^{g(\tilde{t})} d\tilde{t} = dt, \tag{6.24}$$

and we end up with the metric

$$ds^2 = -e^{2f(r)}dt^2 + e^{2\beta(r)}dr^2 + r^2ds^2(S^2). \quad (6.25)$$

We need to solve the remaining Einstein equations. Note that the metric is now independent of t , this naturally comes out of the Einstein equations, we did not impose this! This implies that *any spherically symmetric vacuum metric possesses a timelike Killing vector*. A metric with this property is called *stationary*, in fact the Schwarzschild metric is also *static* we will come back to this shortly.

We can now remove all \tilde{t} derivatives and exchange $\alpha \rightarrow f$ in the Ricci tensor components and where we see \tilde{t} replace with just t . We are free to add components and so we take the combination

$$0 = e^{2\beta-2f(r)}R_{tt} + R_{rr} = \frac{2}{r}(\partial_r f(r) + \partial_r \beta). \quad (6.26)$$

We then have

$$f(r) = -\beta(r) + \text{const}, \quad (6.27)$$

but we may rescale the time coordinate to set the constant to 0. Plugging this into $R_{\theta\theta}$ we find

$$e^{2f(r)}(2r\partial_r f(r) + 1) = 1 \quad \Leftrightarrow \quad \partial_r(re^{2f(r)}) = 1, \quad (6.28)$$

which has solution

$$e^{2f(r)} = 1 - \frac{r_S}{r}, \quad (6.29)$$

with r_S an undetermined constant which we will set to be $r_S = 2G_N M$. There is no remaining freedom except to set r_S to a certain value so the remaining components must vanish, and it turns out that they do, so we have solved Einstein's equations and derived the Schwarzschild solution.

Stationary vs Static There are two different meanings to time independence of a metric.

Definition 44 *Stationary* A spacetime is stationary if it admits an everywhere timelike Killing vector field K . We typically normalise it so that asymptotically $K^2 \rightarrow -1$.

Definition 45 A spacetime is static if, in addition to being stationary, it is invariant under $t \rightarrow -t$, where t is the coordinate along the integral curves of K . This rules out $dt dx$ cross terms in the metric with x any other coordinate except t .

We see that the Schwarzschild solution is static (and therefore also stationary).

Theorem 6 Birkhoff's theorem

The Schwarzschild solution is the unique spherically symmetric, asymptotically flat solution to Einstein's equations in a vacuum.

Key Points

The beauty of Birkhoff's theorem lies in the fact that it must describe the spacetime around any spherically symmetric object: black holes, stars, footballs. Moreover since we did not assume time independence it equally applies to the spacetime around a collapsing star, provided the collapse is spherically symmetric.

6.2 Geodesics of Schwarzschild

We now want to consider the geodesics of the Schwarzschild metric. We have computed the Christoffel symbols above and could just substitute this into the geodesic equation (4.88) however if one did not already have the Christoffel symbols this is not necessarily the quickest method. Instead one should use the Euler–Lagrange equations for the Lagrangian

$$\mathcal{L} = \sqrt{-g_{\mu\nu}\dot{x}^\mu\dot{x}^\nu}, \quad (6.30)$$

and parametrise with an affine parameter. With the choice of an affine parameter we can then compute the Euler–Lagrange equations of \mathcal{L}^2 instead and obtain the same equations of motion. We take

$$\begin{aligned} \mathcal{L} &= g_{\mu\nu} \frac{dx^\mu}{d\lambda} \frac{dx^\nu}{d\lambda} \\ &= -\left(1 - \frac{2G_N M}{r}\right) \dot{t}^2 + \left(1 - \frac{2G_N M}{r}\right)^{-1} \dot{r}^2 + r^2 \dot{\theta}^2 + r^2 \sin^2 \theta \dot{\phi}^2, \end{aligned} \quad (6.31)$$

with $\dot{\bullet} \equiv \frac{d\bullet}{d\lambda}$. Since we are using an affine parameter this is equal to a constant ϵ which we may take to be -1 for time-like geodesics, 0 for null and 1 for space-like geodesics.

Before embarking on a brute force approach we should take a step back to see how we can simplify the problem. The answer is to study the conserved quantities. Ignorable coordinates, ones which do not appear explicitly, give rise to conserved quantities since from the Euler–Lagrange equations²⁷ we find

$$\frac{d\mathcal{L}}{dx^\mu} = 0 \quad \Rightarrow \quad \frac{d}{d\lambda} \frac{d\mathcal{L}}{dx^\mu} = 0. \quad (6.32)$$

²⁷More precisely Killing vectors define conserved quantities for geodesics. Let K be a Killing vector then, $K_\mu p^\mu$ is conserved along a geodesic with $p_\mu = \frac{\partial \mathcal{L}}{\partial \dot{x}^\mu}$. Here the Killing vectors are ∂_t and ∂_ϕ and so we obtain that the conserved quantities are simply p_t and p_ϕ .

The action has two such ignorable coordinates t and ϕ : giving

$$\begin{aligned} 2l &= \frac{d\mathcal{L}}{d\dot{\phi}} = 2r^2 \sin^2 \theta \dot{\phi}, \\ -2E &= \frac{d\mathcal{L}}{dt} = -2 \left(1 - \frac{2G_N M}{r} \right) \dot{t}. \end{aligned} \quad (6.33)$$

Of course these should be identified with the angular momentum and energy respectively. Next consider the equation for θ , we find

$$\frac{d}{d\lambda}(r^2 \dot{\theta}) = r^2 \sin \theta \cos \theta \dot{\phi}^2. \quad (6.34)$$

Recall that in computing the motion in Newtonian gravity we noted that if we started the particle at $\theta = \frac{\pi}{2}$ with $\dot{\theta} = 0$ then it remained in the plane, the same is true here and so we can without loss of generality set $\theta = \frac{\pi}{2}$.

Key Points

The Euler–Lagrange equations are much simpler to compute for the Lagrangian:

$$\mathcal{L} = g_{\mu\nu} \dot{x}^\mu \dot{x}^\nu, \quad (6.35)$$

giving rise to the affinely parametrised geodesic equations. The action for this Lagrangian is not reparametrisation invariant and so we must take an affine parametrisation. For a timelike geodesic the affine parameter is the proper time and therefore we impose:

$$-1 = g_{\mu\nu} \dot{x}^\mu \dot{x}^\nu, \quad (6.36)$$

after applying the Euler–Lagrange equations. It is typically easier to solve this equation than the second order geodesic equations.

One should then look at the symmetries of the metric and ignorable coordinates which give rise to conserved quantities. For the Schwarzschild the ignorable coordinates are t and ϕ giving rise to conserved energy and angular momentum.

Finally, the spherical symmetry implies that we may orient our geodesic to be initially at $\theta = \frac{\pi}{2}$ and also to have initial velocity $\dot{\theta}(0) = 0$. This then means that there is no dynamics in the θ direction and motion is fixed in a plane.

We have therefore solved the motion in the t, θ and ϕ directions and reduced the problem to studying the evolution of the radial coordinate r .

We can now plug this into (6.31) and equate with our constant parameter ϵ giving

$$\epsilon = - \left(1 - \frac{2G_N M}{r} \right)^{-1} E^2 + \left(1 - \frac{2G_N M}{r} \right)^{-1} \dot{r}^2 + r^{-2} \dot{t}^2. \quad (6.37)$$

Rearranging we have

$$\frac{1}{2}\dot{r}^2 + V_{\text{eff}}(r) = \frac{E^2}{2}, \quad (6.38)$$

with

$$V_{\text{eff}}(r) = -\frac{\epsilon}{2} + \frac{\epsilon G_N M}{r} + \frac{l^2}{2r^2} - \frac{l^2 G_N M}{r^3}. \quad (6.39)$$

We should contrast this with the equivalent Newtonian expression in (2.69) for a massive particle which was

$$V_N(r) = -\frac{G_N M}{r} + \frac{l^2}{2r^2}. \quad (6.40)$$

We see that General relativity leads to additional corrections to the potential. The first term is simply a constant shift and so does not play much of a role since we can absorb it into a redefinition of the energy, the r^{-3} term is completely new and changes the Newtonian potential at small distances. Note that the effective potential vanishes at $r = 2G_N M$ which is the Schwarzschild radius.

Let us reinstate the speed of light in the potential, we have

$$V_{\text{eff}}(r) = -\frac{\epsilon c^2}{2} + \frac{\epsilon G_N M}{r} + \frac{l^2}{2r^2} - \frac{l^2 G_N M}{r^3 c^2}, \quad (6.41)$$

then the equation for \dot{r} is

$$\frac{1}{2}\dot{r}^2 + V_{\text{eff}}(r) = \frac{1}{2} \frac{E^2}{c^2}. \quad (6.42)$$

We now want to analyse the different forms of trajectories that are possible. In figure 16 we have plotted the potential for various values of l , with fixed mass M , you should compare the left-hand side one with the plot in figure 2 for the Newtonian potential.

Circular orbits will be at points where the potential has a turning point. Then we are stuck in a circular orbit, which is stable if it corresponds to a minimum of the potential and unstable if it corresponds to a maximum. Differentiating the potential we have

$$V'_{\text{eff}}(r) = \frac{1}{r^4} \left(3G_N l^2 M - l^2 r - G_N M \epsilon r^2 \right) \quad (6.43)$$


which potentially has two zeroes at

$$r_c = -\frac{l^2 \pm \sqrt{l^4 + 12G_N l^2 M \epsilon}}{2G_N M \epsilon}, \quad (6.44)$$

depending on the range of the parameters, for $\epsilon \neq 0$ and

$$r_c = 3G_N M, \quad (6.45)$$

for $\epsilon = 0$.



(a) Plots of the potential for massive particles, $\epsilon = -1$ and for $G_N M = 1$. The different plots correspond to increasing l .

(b) Plots of the potential for massless particles, $\epsilon = 0$ and for $G_N M = 1$. The different plots correspond to increasing l . Note the maximum at $3 \equiv 3G_N M$ in the massless case for all l .

Figure 16: Plots of the potential for massive and massless particles. Note that the plots tends to $-\frac{\epsilon}{2}$ as $r \rightarrow \infty$. Moreover the potentials both vanish at $2 = 2G_N M$ which is the Schwarzschild radius. This should be compared to the corresponding Newtonian plot in figure 2.

For the massless photon the orbit is at a maximum and is therefore unstable. A photon can orbit in a circular orbit forever around the black hole, but any perturbation will send it flying off to either $r = 0$ or $r = \infty$. This is known as the *photon sphere*. The focussing effects mean that much of the light emitted from an accretion disc around a non-rotating black hole emerges from the photon sphere. In practice, it seems likely that the photographs by the Event Horizon Telescope do not have the required resolution to see this yet.

For massive particles there are different regimes depending on the angular momentum. For large l there will be two circular orbits, one stable and one unstable. In the $l \rightarrow \infty$ regime they are at

$$r_c = \left(3G_N M, \frac{l^2}{G_N M} \right). \quad (6.46)$$

The stable circular orbit gets further away while the unstable orbit approaches $3G_N M$ where the photon sphere is located. As we decrease l the two orbits come together and coincide when the discriminant of the quadratic in (6.43) vanishes. This is at

$$l = \sqrt{12}G_N M, \quad (6.47)$$

which gives

$$r_c = 6G_N M. \quad (6.48)$$

For smaller l there are no circular orbits and so $6G_N M$ is the smallest possible radius of a stable circular orbit of the Schwarzschild metric. We have derived that:

The Schwarzschild solution possesses stable circular orbits for massive test masses for $r > 6G_N M$ and unstable circular orbits for $3G_N M < r < 6G_N M$.
 For massless particles/light there are unstable circular orbits at $r_c = 3G_N M$.

We should comment that these are the motions of geodesics. For an accelerating observer such as a rocket ship, there is nothing stopping them from dipping below $r = 3G_N M$ and then reemerging, so long as they stay away from $r = 2G_N M$.

Most experimental tests of general relativity involve the motion of test particles in the solar system. More recently, with the advancements in technology, using gravitational waves to test general relativity has also become possible and will be discussed in GR2. For the moment we will concentrate on three particular tests: the precession of perihelia, the bending of light and gravitational red-shift.

6.2.1 Perihelion precession

We saw when we consider the orbits in Newtonian gravity that the non-circular orbits were closed ellipses. Observation of the orbit of Mercury showed that the closed elliptic orbits of Newtonian gravity were not realised, instead the orbit precessed. This was first noticed in 1859 by Urbain Le Verrier. He analysed the available timed observations of transits of Mercury over the sun's disk from 1697 to 1848. He showed that the actual rate of precession disagreed from that predicted by Newton's theory by 38 " per century. This was later re-estimated by Simon Newcomb in 1882 to be 43". A number of different, and ultimately unsuccessful solutions were proposed, however they tended to introduce new problems. Le Verrier suggested another hypothetical planet might exist, given the previous success in predicting Neptune this was given some faith, and this planet was named Vulcan. No evidence for Vulcan was found. A non-trivial check of General Relativity is then to show that the orbits of the planets precess. We can approximate the metric of the sun to be the Schwarzschild metric and take the planet to follow the geodesic of a massive particle.

The strategy is to describe the evolution of the radial coordinate r as a function of ϕ . If the orbit is a perfect ellipse $r(\phi)$ should be periodic with period 2π , then the perihelion occurs at the same point every orbit. Instead, for a non-closed ellipse the perihelion is shifted after every orbit. We will see that General Relativity gives a slight modification of the Newtonian result such that the orbit precesses.

First consider the radial equation of motion for a massive particle, (6.38), setting $\epsilon = -1$. To get an equation for $\frac{dr}{d\phi}$ we can use the chain rule and multiply the equation by

$$\left(\frac{d\phi}{d\lambda}\right)^{-2} = \frac{r^4}{l^2}, \quad (6.49)$$

yielding

$$\left(\frac{dr}{d\phi}\right)^2 + \frac{r^4}{l^2} - \frac{2G_N M}{l^2} r^3 + r^2 - 2G_N M r = \frac{E^2 r^4}{l^2} \quad (6.50)$$

Define the new variable

$$x = \frac{l^2}{G_N M r}, \quad (6.51)$$

which gives rise to the Newtonian circular orbit when $x = 1$. The equation of motion becomes

$$\left(\frac{dx}{d\phi}\right)^2 + \frac{l^2}{G_N^2 M^2} - 2x + x^2 - \frac{2G_N^2 M^2 x^3}{l^2} = \frac{E^2 l^2}{G_N^2 M^2}. \quad (6.52)$$

Next differentiate with respect to ϕ to obtain

$$\frac{d^2 x}{d\phi^2} - 1 + x = \frac{3G_N^2 M^2 x^2}{l^2}. \quad (6.53)$$

In the Newtonian calculation the term on the right-hand-side would be absent and we could solve for x exactly. Here we will treat this as a perturbation around the Newtonian result.

We expand x into a Newtonian solution plus a small deviation

$$x = x_0 + x_1, \quad (6.54)$$

where the zeroth order part satisfies

$$\frac{d^2 x_0}{d\phi^2} - 1 + x_0 = 0. \quad (6.55)$$

The equation for the first order part becomes

$$\frac{d^2 x_1}{d\phi^2} + x_1 = \frac{3G_N^2 M^2}{l^2} x_0^2. \quad (6.56)$$

A solution to the zeroth order equation is (see (2.75))

$$x_0 = 1 + e \cos \phi, \quad (6.57)$$

which recall describes a perfect ellipse with eccentricity e , $e = 1 - \frac{b^2}{a^2}$ with a the semi-major axis, the distance from the centre to the farthest point on the ellipse and the semi-minor axis

b the distance from the centre to the closest point. Plugging in the Newtonian solution into the first order equation of motion we find

$$\frac{d^2x_1}{d\phi^2} + x_1 = \frac{3G_N^2M^2}{l^2}(1 + e \cos \phi)^2. \quad (6.58)$$

A solution is given by

$$x_1 = \frac{3G_N^2M^2}{l^2} \left[\left(1 + \frac{e^2}{2}\right) + e\phi \sin \phi - \frac{1}{6}e^2 \cos 2\phi \right]. \quad (6.59)$$

The first term is just a constant displacement while the third oscillates around 0. The important effect is contained within the second term which accumulates over successive orbits. Combining only this term with the zeroth-order solution we have

$$x = 1 + e \cos \phi + \frac{3G_N^2M^2e}{l^2} \phi \sin \phi. \quad (6.60)$$

We should emphasise that this is not a full solution, it is an approximation but it encapsulates the part we are interested in. We may write

$$x = 1 + e \cos \left((1 - \alpha)\phi \right), \quad (6.61)$$

where

$$\alpha = \frac{3G_N^2M^2}{l^2}. \quad (6.62)$$

where one should view this as a series expansion around $\alpha = 0$. It follows that during each orbit the perihelion advances by an angle

$$\Delta\phi = 2\pi\alpha = \frac{6\pi G_N^2M^2}{l^2}. \quad (6.63)$$

We may replace the angular momentum in favour of the eccentricity by looking at the Newtonian solution. An ordinary ellipse satisfies

$$r = \frac{(1 - e^2)a}{1 + e \cos \phi}, \quad (6.64)$$

with a the semi-major axis. This leads us to identify

$$l^2 \sim G_N M (1 - e^2) a, \quad (6.65)$$

for the Newtonian orbit. Plugging this in and restoring the speed of light we find

$$\Delta\phi = \frac{6\pi G_N M}{c^2(1 - e^2)a}. \quad (6.66)$$

Historically the precession of mercury was the first test of GR. The apparent discrepancy between observation and Newtonian gravity was known long before the advent of GR, and a number of solutions had been proposed including additional planets. For the motion of Mercury around the sun we have

$$\begin{aligned}\frac{G_N M_\odot}{c^2} &= 1.48 \times 10^3 m, \\ a &= 5.79 \times 10^{10} m, \\ e &= 0.2056.\end{aligned}\tag{6.67}$$

This gives

$$\Delta\phi_{\text{Mercury}} = 5.01 \times 10^{-7} \text{radians/orbit} = 0.103''/\text{orbit}\tag{6.68}$$

with '' denoting arcseconds. Mercury orbits once every 88 days and therefore

$$\Delta\phi_{\text{Mercury}} = 43.0''/\text{century}.\tag{6.69}$$

From our computation we conclude that the major axis of Mercury's orbit precesses at a rate of 43.0 arcseconds every 100 years. The observed value is 5601 arcseconds/100 years. Much of that is due to the precession of equinoxes in our geocentric coordinate system: 5025 arcseconds/100 years. The gravitational perturbations of the other planets contributes an additional 532 arcseconds/100 years leaving a 43 arcseconds/100 years to be explained by GR which it does quite well.

6.2.2 Bending of light

We can now extend these results for null geodesics. We have seen that there is an unstable circular orbit for light. What about other orbits? The fate of other light rays depends on the relative value of their energy E to their angular momentum l . The maximum value of the potential is

$$V_{\text{null}}(r_*) = \frac{l^2}{54G_N^2 M^2},\tag{6.70}$$

and therefore the physics depends on how this compares with the right-hand side of (6.38). There are two possibilities we need to consider

- $E < \frac{l}{\sqrt{27}G_N M}$. The energy of the light is lower than the angular momentum barrier. This means that light emitted from $r < r_*$ cannot escape to infinity; it will orbit the star before falling back towards the origin. For light coming from infinity it will not fall into the star but will instead bounce off the angular momentum barrier and return to infinity: the light will be scattered.




Figure 17: The perihelion precession in blue overlaid with the Newtonian elliptical orbit in red. The parameters have been chosen to exaggerate the precession over a few orbits. A more faithful representation

- $E > \frac{l}{\sqrt{27G_N M}}$. The energy of light is greater than the angular momentum barrier. Light can be emitted from $r < r_*$ and escape to infinity (this is only true for $r_S < r$). Meanwhile light coming from infinity is captured by the star/black hole.

To see this more clearly let us once again use the inverse parameter $u = \frac{1}{r}$. The equation of motion becomes

$$\left(\frac{du}{d\phi}\right)^2 + u^2(1 - 2G_N M u) = \frac{E^2}{l^2}. \quad (6.71)$$

Differentiating again we find

$$\frac{d^2u}{d\phi^2} + u = 3G_N M u^2. \quad (6.72)$$

We may once again work perturbatively. At zeroth order we can ignore the $G_N M$ term on

the right-hand-side. Then to leading order we have

$$\frac{d^2u}{d\phi^2} + u = 0, \quad \Rightarrow \quad u = \frac{1}{b} \sin \phi, \quad (6.73)$$

for b a constant. Reinstating r we have $r \sin \phi = b$: which is the equation of a horizontal straight line, a distance b above the origin, see 18. The distance b is known as the *impact parameter*.




Figure 18: Light bending in the Schwarzschild metric. The dashed line at the top is the constant line $r \sin \phi = b$. The curved line is the geodesic.

With the zeroth order solution we can now solve (6.72) in an expansion around $\frac{G_N M}{b} = \beta$. We have

$$u = u_0 + \beta u_1 + \dots \quad (6.74)$$

At first order we need to solve

$$\frac{d^2u_1}{d\phi^2} + u_1 = \frac{3 \sin^2 \phi}{b} = \frac{3(1 - \cos 2\phi)}{2b}. \quad (6.75)$$

The general solution is

$$u_1 = A \cos \phi + B \sin \phi + \frac{1}{2b}(3 + \cos 2\phi), \quad (6.76)$$

where the first two parts are the solutions of the homogeneous part and A, B two integration constants. We should choose them so that the initial trajectory at $\phi = \pi$ agrees with the straight line u_0 . For this to hold we must take $A = \frac{2}{b}$ and $B = 0$ so that $u_1 \rightarrow 0$ as $\phi \rightarrow \pi$. To leading order in β the solution is

$$u = \frac{1}{b} \sin \phi + \frac{G_N M}{2b^2}(3 + 4 \cos \phi + \cos 2\phi). \quad (6.77)$$

What angle does the particle escape to $r = \infty \Leftrightarrow u = 0$? Before the correction this was at $\phi = 0$, within our perturbative approach we can approximate $\sin \phi \sim \phi$ and $\cos \phi \sim 1$ to find that the particle escapes at

$$\phi \sim -\frac{4G_N M}{b}. \quad (6.78)$$

This means that the light is bent by gravity, this bending of light is known as *gravitational lensing*.

For the sun, $\frac{G_N M_\odot}{c^2} \sim 1.48$ km. If the light rays just graze the surface of the sun, then the impact parameter is the radius of the sun $R_\odot \sim 7 \times 10^5$ km. This gives a scattering angle of $\phi \sim 8.6 \times 10^{-5}$ radians or $\phi \sim 1.8''$. The Newtonian prediction gives only half of this contribution.

There is a difficulty in testing this prediction since things behind the sun are rarely visible. By sheer coincidence, the size of the moon in the sky is about the same size of the Sun which leads to total solar eclipses. This means that during a solar eclipse the light from the sun is blocked allowing for the measurement of stars whose light passes nearby the Sun. This can then be compared with the positions of these stars 6 months later when the Sun is behind the Earth and so the light from the star sources is not lensed by the Sun.

The first measurement was carried out in 1919 by two expeditions lead by Arthur Eddington and Frank Watson Dyson (we will see Eddington's name again shortly).²⁸ Since then our evidence of the bending of light is more impressive. Clusters of galaxies have been seen to distort the light from a background source often revealing a distinct ring-like pattern of multiple copies of the light source. See figure 19.

6.2.3 Gravitational red shift

We have seen two tests of general relativity, next we will now look at time dilation due to strong gravitational fields.

Let us consider an observer with four velocity U^μ who is stationary in Schwarzschild coordinates, i.e. $U^i = 0$.²⁹ The four-velocity is normalised so that $U_\mu U^\mu = 1$, which for our stationary observer in a Schwarzschild background implies

$$U^0 = \left(1 - \frac{2G_N M}{r}\right)^{-1/2}. \quad (6.79)$$

²⁸This result was considered spectacular news and made the front page of most major newspapers.

²⁹We could allow for the observer to be moving, however the difference is just to superimpose the usual Doppler shift on top of the gravitational effect and therefore we consider the simpler example.




Figure 19: A diagram of light lensing picked up by the Hubble telescope. Notice that there are four copies of the distant quasar in the picture obtained by Hubble. Image credited to NASA, ESA and STScI.

Such an observer measures the frequency of a photon following a null geodesic $x^\mu(\lambda)$ to be

$$\omega = -g_{\mu\nu}U^\mu \frac{dx^\nu}{d\lambda}. \quad (6.80)$$

We have

$$\begin{aligned} \omega &= \left(1 - \frac{2G_N M}{r}\right)^{1/2} \frac{dt}{d\lambda} \\ &= \left(1 - \frac{2G_N M}{r}\right)^{-1/2} E, \end{aligned} \quad (6.81)$$

where E was defined to be the conserved quantity associated to time translations when we worked out the geodesics. Since E is conserved it follows that ω will have different values when measured at different radial distances. For a photon emitted at r_1 and an observer at r_2 , the observed frequencies will be related by

$$\frac{\omega_2}{\omega_1} = \sqrt{\frac{1 - 2G_N M/r_1}{1 - 2G_N M/r_2}}. \quad (6.82)$$

This is the exact result for the frequency shift, in the limit $r \gg 2G_N M$ we have

$$\begin{aligned} \frac{\omega_2}{\omega_1} &= 1 - \frac{G_N M}{r_1} + \frac{G_N M}{r_2} \\ &= 1 + \Phi(r_1) - \Phi(r_2), \end{aligned} \quad (6.83)$$

with $\Phi = -G_N M/r$ the Newtonian potential.

We see that the frequency goes down as Φ increases which happens as we climb out of a gravitational field, leading to a red-shift. On the other hand photons which fall towards the gravitating body are blue shifted. Gravitational red-shift was first detected in 1960 by Pound and Rebka using gamma rays travelling a distance of 72-feet (about 22m) which was the height of the physics building at Harvard. Increasingly precise tests have found excellent agreement with GR. There is a cosmological counterpart to this, where light is red-shifted in an expanding universe.

Time delay Since the temporal component of the metric is

$$g_{00}(x) = 1 + 2\Phi(x), \quad (6.84)$$

we see that there is a connection between time and gravity. Let us once again use the Schwarzschild solution. An observer sitting at a fixed distance r from the origin will measure a time interval

$$d\tau^2 = -g_{00}dt^2 = \left(1 - \frac{2G_N M}{r}\right) dt^2. \quad (6.85)$$

For an asymptotic observer at $r \rightarrow \infty$ who measures a time t , an observer at r will measure the time T

$$T(r) = t\sqrt{1 - \frac{2G_N M}{r}}. \quad (6.86)$$

It follows that time goes slower in the presence of a massive gravitating object. Notice that at $r = r_S$ that time seems to stop for the observer at r_S . We will come back to this later.

We can make this more quantitative by considering two observers: Alice and Bob. Bob has gone up in a hot air balloon while Alice is on the surface of the earth at r_A . Bob is at a distance $r_B = r_A + \Delta r$. The time measured by Bob is

$$\begin{aligned} T_B &= t\sqrt{1 - \frac{2G_N M}{(r_A + \Delta r)}} \sim t\sqrt{1 - \frac{2G_N M}{r_A} + \frac{2G_N M \Delta r}{r_A^2}} \\ &\sim t\sqrt{1 - \frac{2G_N M}{r_A}} \left(1 + \frac{G_N M \Delta r}{r_A^2}\right) = T_A \left(1 + \frac{G_N M \Delta r}{r_A^2}\right). \end{aligned} \quad (6.87)$$

A double expansion has been utilised where we assume $\Delta r \ll r_A$ and $\frac{2G_N M}{r_A} \ll 1$. If the hot air balloon flies a distance $\Delta r = 1000m$ above Alice then taking the radius of the Earth to be $r_A \approx 6000km$ the difference in times is about 10^{-12} and therefore over the whole day Bob ages by an extra 10^{-18} seconds or so. Clearly this is a small amount, in the vicinity of a black hole this can be more pronounced. Recall that the smallest stable orbit was at $r = 3G_N M$ and such a person experiences time at a rate of $T = 3^{-1/2}t \approx 0.6t$ compared to an asymptotic

observer at $r \rightarrow \infty$. For more dramatic results one would need to fly closer to the horizon and then return to asymptotic infinity.

This also gives a different perspective on the gravitational redshift. Bob doesn't like Alice and wants to ruin her day so he hovers above Alice and chucks peanuts at her. He throws peanuts at time intervals ΔT_B . Alice, wise to Bob's antics, opens up an umbrella. The peanuts hit the umbrella at time intervals ΔT_A where as above

$$\Delta T_A = \Delta T_B \sqrt{\frac{1 + 2\Phi(r_A)}{1 + 2\Phi(r_B)}} \approx \left(1 + \Phi(r_A) - \Phi(r_B)\right) \Delta T_B. \quad (6.88)$$

We have that $r_A < r_B$ and therefore $\Phi(r_A) < \Phi(r_B) < 0$ and hence $\Delta T_A < \Delta T_B$. Alice receives the peanuts at a higher frequency than Bob threw them.

Having seen the peanuts hitting the umbrella Bob decides to instead shine a light down at Alice with a frequency $\omega_B \sim \Delta T_B^{-1}$. Alice will then receive the light at a frequency ω_A where

$$\omega_A \approx \left(1 + \Phi(r_A) - \Phi(r_B)\right)^{-1} \omega_B. \quad (6.89)$$

This is a higher frequency $\omega_A > \omega_B$ and therefore a shorter wave-length. The light is therefore blue-shifted. In contrast if Alice retaliates and shines a light up to Bob then the frequency decreases and the light is redshifted.

6.3 Schwarzschild solution as a black hole

We have now studied geodesics for the Schwarzschild solution and some phenomena of these geodesics. Each time we have carefully avoided the Schwarzschild radius $r_S = 2G_N M$ and also $r = 0$. At both of these points something funky happens with the metric, at least one of the components of the metric diverges or vanishes. The interpretation of the singularities is different for the two cases. The divergence at $r = 0$ is a genuine *singularity*. General relativity breaks down here and we need a theory of quantum gravity. GR predicts its own death! In contrast the divergence at $r = 2G_N M$ is a result of our choice of coordinates. This surface is referred to as the *event horizon* or simply the *horizon* of the black hole. Many of the surprising properties of a black hole happen here.

There is a simple way to check whether a divergence is due to a singularity or a poor choice of coordinates. We can build scalar quantities, these are then independent of the choice of coordinates and therefore if they diverge in one coordinate system they diverge in all and the spacetime is ill-defined at this point. On the other hand if it does not diverge we cannot say much, one would have to consider all possible scalar quantities to concretely say it is just

a coordinate singularity. Since the Einstein equations in a vacuum set $R_{\mu\nu} = 0$ it follows that the simplest scalar quantities one can construct R and $R_{\mu\nu}R^{\mu\nu}$ both vanish. The next simplest is the *Kretschmann scalar* $R^{\mu\nu\rho\sigma}R_{\mu\nu\rho\sigma}$. For the Schwarzschild metric we find

$$R^{\mu\nu\rho\sigma}R_{\mu\nu\rho\sigma} = \frac{48G_N^2M^2}{r^6}. \quad (6.90)$$

There is no pathology at $r = 2G_NM$ while there is at $r = 0$ where the scalar diverges.

One way to understand the geometry of spacetime is to explore its causal structure as defined by light cones. We therefore consider radial null curves, i.e. those with constant θ, ϕ and $ds^2 = 0$, such that they satisfy

$$ds^2 = 0 = -\left(1 - \frac{2G_NM}{r}\right)dt^2 + \left(1 - \frac{2G_NM}{r}\right)^{-1}dr^2. \quad (6.91)$$

Rearranging gives

$$\frac{dt}{dr} = \pm \left(1 - \frac{2G_NM}{r}\right)^{-1}. \quad (6.92)$$

This measures the slope of the light cones on a spacetime diagram in the t - r plane. For large r the slope is ± 1 as it would be for flat spacetime. On the other hand as we approach $r = 2G_NM$ we get $\frac{dt}{dr} \rightarrow \pm\infty$ and the light cones close up, see figure 20. Thus a light ray which approaches $r = 2G_NM$ never seems to get there, at least in this coordinate system. This apparent inability to get to $r = 2G_NM$ is actually an illusion and an artefact of a bad choice of coordinates. An in-falling light ray or massive particle has no trouble reaching this radius. On the other hand an observer far away would never be able to tell. If we all hovered outside a black hole and one of your class mates jumped in the black hole sending back signals the whole way down we would simply see the signals reach us less frequently, see figure 21.

The fact that we never see them reach $r = 2G_NM$ is a meaningful statement but the fact that their trajectory in the t - r plane never reaches there is not: it is highly dependent on our coordinate system. We want to change coordinates to some that are better behaved at $r = r_S$. Note that we can solve (6.92) for t as a function of r by introducing the *tortoise coordinate* r_*

$$r_* = r + 2G_NM \log\left(\frac{r - 2G_NM}{2G_NM}\right). \quad (6.93)$$

The solution then becomes

$$t = \pm r_* + \text{constant}, \quad (6.94)$$

and we see that this is well adapted to null radial geodesics: they are simply straight lines in these coordinates. The plus sign corresponds to out-going geodesics and the negative to




Figure 20: In Schwarzschild coordinates the light cones appear to close up as we approach the horizon. We will see that this is not quite correct.




Figure 21: A beacon freely falling into a black hole emits signals at intervals of proper time $\Delta\tau_1$. An observer at fixed r receives these signals at a successively longer time intervals $\Delta\tau_2$.

in-going geodesics³⁰. Next we introduce a pair of null coordinates further adapted to the null

³⁰The quick way to see this is to note that as $r \rightarrow \infty$ we have $r_* \rightarrow \infty$ and therefore we need the plus sign for out-going geodesics so that the radial direction increases with time.

geodesics:

$$v = t + r_*, \quad u = t - r_*. \quad (6.95)$$

In these coordinates the null radial geodesics are simply $u = \text{const}$ or $v = \text{const}$. We can write the metric in these new coordinates. First consider the metric in (v, r) coordinates and then we will study (u, r) coordinates before biting the bullet and using (v, u) coordinates.

Ingoing Eddington–Finkelstein coordinates Eliminating t via $t = v - r_*(r)$ we find

$$ds^2 = -\left(1 - \frac{2G_N M}{r}\right)dv^2 + 2dvdr + r^2 ds^2(S^2). \quad (6.96)$$

This is the Schwarzschild solution in *ingoing Eddington–Finkelstein coordinates*. Even though the metric coefficient g_{vv} vanishes at $r = 2G_N M$ there is no real degeneracy. The determinant of the metric is

$$\det g = \det \begin{pmatrix} -\left(1 - \frac{2G_N M}{r}\right) & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & r^2 & 0 \\ 0 & 0 & 0 & r^2 \sin^2 \theta \end{pmatrix} = -r^4 \sin^2 \theta. \quad (6.97)$$

The cross terms stops the metric from being degenerate at the horizon. The metric is still degenerate at $r = 0$ and $\theta = 0, \pi$ however the latter are just the usual pole problems of the S^2 and nothing to worry about. This is the benefit of the Eddington–Finkelstein coordinates, the radial coordinate can be extended beyond the horizon.

To build further intuition we can look at the behaviour of light rays. We saw that the null radial geodesics were given by (6.94). The outgoing geodesics are

$$u = t - r_* = \text{const}. \quad (6.98)$$

Eliminating t in favour of v we have that the outgoing geodesics satisfy $v = 2r_* + \text{const}$. The solutions of this equation have a different behaviour depending on whether they are inside the horizon or outside. For $r > 2G_N M$ we can use the original definition of r_* in (6.93) to get

$$v = 2r + 4G_N M \log \left(\frac{r - 2G_N M}{2G_N M} \right) + \text{const}. \quad (6.99)$$

The Log term goes bad when $r < 2G_N M$, however we can simply modify the coordinate to take the norm of the argument of the log, so that

$$r_* = r + 2G_N M \log \left| \frac{r - 2G_N M}{2G_N M} \right|. \quad (6.100)$$

This means that r_* is multi-valued. Outside the horizon it takes values $r_* \in (-\infty, \infty)$ while inside the horizon it takes values $r_* \in (-\infty, 0)$. The singularity sits at $r_* = 0$. Outgoing geodesics inside the horizon obey

$$v = 2r + 4G_N M \log\left(\frac{2G_N M - r}{2G_N M}\right) + \text{const}. \quad (6.101)$$

Finally note that $r = 2G_N M$ is itself a null geodesic. This information can be captured in a *Finkelstein diagram*. It is designed so that ingoing null rays travel at 45° . This is simple to do if we label the coordinates of the diagram by t and r_* , however since r_* is not single valued we use r instead. We define a new temporal coordinate t_* by the requirement

$$v = t + r_* = t_* + r. \quad (6.102)$$

Thus ingoing null rays travel at 45° in the (t_*, r) -plane. See figure 22




Figure 22: The Finkelstein diagram in in-going coordinates. The ingoing null geodesics are in red while the outgoing are in blue. Inside the horizon the outgoing geodesics never go past the horizon.

The outgoing null geodesics that sit outside the horizon tend to infinity, whereas those inside the horizon don't actually go out, but rather go towards the singularity at $r = 0$. Each

hits the singularity at some finite t_* . We can draw lightcones on the Penrose diagram. These are regions which are bounded by the in-going and out-going future pointing null geodesics. Any massive particle must follow a timelike path and this must then sit within these lightcones. We see that the lightcones get tipped as we get closer to the horizon, and then once inside the horizon there is no way of getting back out. The causal structure of spacetime prevents this. The term black hole really refers to this area inside the horizon $r < 2G_N M$, any observer outside the horizon can never know what is happening inside the black hole.

We can also see what happens if we watch someone fall into a black hole. The person falls through the horizon without realising anything is wrong. However as they fall the light signals that come back to us take longer and longer to reach us. The actions of the in-falling person become increasingly slowed as they approach the horizon. In this way we continue to see the person forever, but we know nothing about their fate past the horizon. Since the light returns to us from a deeper and deeper gravitational well it appears increasingly red-shifted to us.

Out-going Eddington–Finkelstein coordinates We can also extend the exterior of the Schwarzschild black hole by replacing the time coordinate with the null coordinate

$$u = t - r_*. \quad (6.103)$$

Surfaces of constant u correspond to outgoing radial null geodesics. After the change of coordinates we have

$$ds^2 = -\left(1 - \frac{2G_N M}{r}\right) du^2 - 2du dr + r^2 ds^2(S^2). \quad (6.104)$$

This is the Schwarzschild solution in *out-going Eddington–Finkelstein coordinates*. The only difference is in the sign of the cross term. This seemingly trivial modification changes the interpretation drastically.

As before the metric is smooth at the horizon and we can continue the metric down to the singularity at $r = 0$. However the region $r < 2G_N M$ now describes a different part of spacetime from the analogous region in ingoing Eddington–Finkelstein coordinates.

We again look at the ingoing and outgoing null radial geodesics. This time we pick coordinates so that the outgoing geodesics travel at 45° . This means that we take r and $t_* = u + r$ to be the axes.

This time the ingoing null geodesics have the interesting property. Those which start outside are unable to reach the singularity, instead they pile up at the horizon. Those that




Figure 23: The Finkelstein diagram in out-going coordinates. The ingoing null geodesics are in red while the outgoing are in blue. Inside the horizon the ingoing geodesics never go past the horizon.

start behind the horizon move towards the horizon, once again piling up there. What happens to massive particles that sit inside the horizon? Their trajectories must lie inside the future pointing light-cones. They cannot stay inside the horizon and the causal structure of spacetime requires them to be ejected outside of the horizon. This is a *white hole*, an object which expels matter. This is the time reversal of a black hole, indeed the difference is purely a minus sign. Moreover if we flip white-hole upside down we get the black hole.

White holes are perfectly acceptable solutions of general relativity. Indeed they are implied by the time reversal invariance of Einstein's equations. However white holes are not physically relevant since in contrast to a black hole they cannot be formed by collapsing matter.

6.3.1 Kruskal spacetime

We have seen that we can extend the $r \in (2G_N M, \infty)$ coordinate in two ways so that we gain the region $r \in (0, 2G_N M]$ which corresponds to two different parts of spacetime. We can

write the Schwarzschild metric using both null (u, v) -coordinates, the metric is

$$ds^2 = -\left(1 - \frac{2G_N M}{r}\right) du dv + r^2 ds^2(S^2), \quad (6.105)$$

where r is a function of $u-v$. In these coordinates the metric is again degenerate at $r = 2G_N M$ so we need to perform another change of coordinates. We can introduce the *Kruskal-Szekeres coordinates*,

$$U = -\exp\left(-\frac{u}{4G_N M}\right), \quad V = \exp\left(\frac{v}{4G_N M}\right), \quad (6.106)$$

which are both null coordinates. The Schwarzschild black hole in Schwarzschild coordinates (that is the region outside the horizon of the black hole) are covered by $U < 0$ and $V > 0$. Outside the horizon these new coordinates satisfy

$$UV = -\exp\left(\frac{r_*}{2G_N M}\right) = \frac{2G_N M - r}{2G_N M} \exp\left(\frac{r}{2G_N M}\right), \quad (6.107)$$

and similarly

$$\frac{U}{V} = -\exp\left(-\frac{t}{2G_N M}\right). \quad (6.108)$$

The metric is then

$$ds^2 = -\frac{32(G_N M)^3}{r} e^{-\frac{r}{2G_N M}} dU dV + r^2 ds^2(S^2), \quad (6.109)$$

with $r(U, V)$ defined by inverting (6.107). The original Schwarzschild metric covers just $U < 0$ and $V > 0$ however there is no obstruction to extending $U, V \in \mathbb{R}$. Nothing bad happens at $r = 2G_N M$, the metric is smooth and non-degenerate. The Kruskal spacetime is the maximal extension of the Schwarzschild solution.

The Kruskal Diagram To find the location of the horizon in the new coordinates we can use equation (6.107). We see that this is at

$$r = 2G_N M \quad \Rightarrow \quad U = 0 \text{ or } V = 0. \quad (6.110)$$

The horizon is not just one null surface but 2 which intersect at $U = V = 0$. On the other hand the singularity is at

$$r = 0 \quad \Rightarrow \quad UV = 1. \quad (6.111)$$

This hyperbola has two disconnected, one with $U, V > 0$ and the other with $U, V < 0$. The former corresponds to the singularity of the black hole and the latter the singularity of the white hole, see figure 24 We can define $T = \frac{1}{2}(U + V)$ and $X = \frac{1}{2}(V - U)$ as the vertical and




Figure 24: The Kruskal diagram. The U and V axes have been rotated 45° . They are the locations of the horizons at $r = 2G_N M$ and the red lines are the singularities at $r = 0$. Lines of constant r are in green and lines of constant t are in blue.

horizontal lines respectively. Lines of constant r are given by $UV = \text{constant}$ while lines of constant t are $U/V = \text{constant}$.

We see that the singularity is spacelike. Once you pass through the horizon the singularity lies in your future. You cannot avoid the singularity once you cross the horizon. Similarly the singularity of the white hole lies in the past, one could think of this as the singularity of the Big Bang.

We can understand three quadrants of the four. The right quadrant is the exterior of the black hole, the top quadrant is the black hole interior and the bottom quadrant is the interior of the white hole. The left hand quadrant is in fact another copy of the black hole exterior, it is just covered by $U > 0$ and $V < 0$. To see this write

$$U = + \exp\left(-\frac{u}{4G_N M}\right), \quad V = - \exp\left(\frac{v}{4G_N M}\right). \quad (6.112)$$

Undoing all the coordinate transformations we see that this is precisely the metric of the Schwarzschild solution again.

Our spacetime contains two asymptotically flat regions joined by a black hole. Note that it is not possible for an observer to cross from one to the other, nor to send a signal from one

region to the other. The causal structure of spacetime forbids this.

One could ask what the spatial geometry that connects the two regions is. Fix the $t = 0$ slice of Kruskal spacetime ($U = V = 0$). In our original Schwarzschild solution the spatial geometry is

$$ds^2 = \left(1 - \frac{2G_N M}{r}\right)^{-1} dr^2 + r^2 ds^2(S^2), \quad (6.113)$$

which is valid for $r > 2G_N M$. There is another copy of this that describes the geometry of the left-hand side and we can glue these two together at $r = 2G_N M$, giving a worm-hole like geometry. This is known as the *Einstein–Rosen bridge*. Before getting excited about travelling through the black hole you cannot travel through the worm-hole as the paths are space-like not time-like.

6.4 Summary

We have studied the first non-trivial solution to Einstein’s equations: the Schwarzschild solution.

- The **Schwarzschild solution** is the unique, asymptotically flat spherically symmetric solution of Einstein’s equation in a vacuum.
- The metric depends on one parameter which we may identify as the **mass**.
- Geodesics exhibit behaviours which differ from the usual Newtonian particle motion.
- Elliptical orbits are seen to **precess, light is bent and gravitational fields lead to time dilation**.
- The **Schwarzschild solution** has a coordinate singularity at $r = 2G_N M$ which is in fact the location of the **black hole horizon**. This is the boundary of the region in space from which nothing can escape to infinity.
- Changing to **Eddington–Finkelstein coordinates** we may extend the metric inside the horizon. Using these coordinates we can draw Finkelstein diagrams for in-going and out-going coordinates, see figures 22 and 23. These show the light cones formed by looking at in- and out-going null radial geodesics.
- We can make a further change of coordinates to **Kruskal coordinates**, (6.106) to maximally extend the spacetime. The maximal extension contains four regions; two asymptotically flat exterior regions, a black hole interior and a white hole interior.

- There is an **Einstein–Rosen** bridge, also known as a wormhole, connecting the two asymptotically flat exterior regions. However, no causal (time-like or null) curve can pass through the wormhole and so they are causally disconnected regions.

Acknowledgments

It is a pleasure to thank all the students who have attended these lectures. You have made this an enjoyable course to teach and all your questions are much appreciated. I hope that you have enjoyed the course and learnt some of the amazing things that GR has to offer.

A Some worked examples on Special Relativity

We have now completed our review of special relativity and Newtonian gravity. We present some worked examples on special relativity below.

A.1 Proper time along an accelerated worldline

We treat the planets as being at rest relative to each other in this question.

Leia begins at rest on the planet Polis Massa and sets off in a spaceship to visit a distant planet called Alderaan. Alderaan is at rest relative to Polis Massa and is a proper distance D away. Leia's spaceship accelerates during the journey at a constant rate α ,

$$\eta_{\mu\nu} a^\mu a^\nu = \alpha^2, \quad (\text{A.1})$$

where a^μ is the four-acceleration of Leia. We want to answer two questions: 1) what path does Leia take in terms of coordinates centred on Polis Massa? 2) How much time passes, from Leia's point of view until she reaches Alderaan?

We can choose coordinates (t, x, y, z) where the worldline of Polis Massa is simply $(t, 0, 0, 0)$ and the worldline of Alderaan is $(t, D, 0, 0)$ (recall that the two planets are at rest relative to each other). Leia's world line is then of the form

$$(t(\tau), x(\tau), 0, 0), \quad (\text{A.2})$$

where τ is the proper time along Leia's worldline. Since we have parametrised Leia's worldline by the proper time we have

$$-\dot{t}(\tau)^2 + \dot{x}(\tau)^2 = -1 \quad \bullet \equiv \frac{d\bullet}{d\tau}. \quad (\text{A.3})$$

Leia's acceleration is therefore,

$$a = (\ddot{t}(\tau), \ddot{x}(\tau), 0, 0) = \left(\frac{\dot{x}(\tau)\ddot{x}(\tau)}{\sqrt{1 + \dot{x}(\tau)^2}}, \ddot{x}(\tau), 0, 0 \right), \quad (\text{A.4})$$

where for the second equality we have used (A.3) to eliminate $\ddot{t}(\tau)$. Since Leia's acceleration is constant, (A.1), we have

$$\alpha^2 = \frac{\ddot{x}(\tau)^2}{1 + \dot{x}(\tau)^2}. \quad (\text{A.5})$$

We have that $\dot{x}(\tau) > 0$ and therefore the solution for $\dot{x}(\tau)$ is

$$\dot{x}(\tau) = \sinh(\alpha\tau + \beta), \quad (\text{A.6})$$

with β a constant of integration. Since Leia began at rest on Polis Massa, we take $\beta = 0$. Integrating again and using that Leia begins at Polis Massa at $\tau = 0$, i.e. $x(0) = 0$, we have

$$x(\tau) = \frac{1}{\alpha} (\cosh(\alpha\tau) - 1). \quad (\text{A.7})$$

Inserting this into (A.3), solving for $t(\tau)$ and imposing $t(0) = 0$ we find

$$t(\tau) = \frac{1}{\alpha} \sinh(\alpha\tau). \quad (\text{A.8})$$

Leia reaches Alderaan when

$$\tau = \text{arccosh}(1 + \alpha D), \quad (\text{A.9})$$

If αD is large then $\tau \sim \frac{1}{\alpha} \log(\alpha D)$ and therefore no matter how large D is, for a sufficiently large acceleration Leia can reach Alderaan in a "reasonable" proper time. On the other hand, when Leia reaches Alderaan

$$t = \sqrt{D^2 + \frac{2D}{\alpha}}, \quad (\text{A.10})$$

and therefore no matter how large α is it always takes at least a time of D (recall $c = 1$) to reach Alderaan as viewed from Polis Massa.

A.2 Null curves in Minkowski space

By now we have all seen that a straight line is a null curve in Minkowski space but are there more? Note that we are not asking about geodesics. Consider the curve, given in inertial coordinates, by

$$x^\mu = (\lambda, \sin \lambda, \cos \lambda, 0). \quad (\text{A.11})$$

The tangent to the vector is

$$v^\mu = \frac{dx^\mu}{d\lambda} = (1, \cos \lambda, -\sin \lambda, 0), \quad (\text{A.12})$$

and has norm

$$v^\mu v^\nu \eta_{\mu\nu} = -1 + \cos^2 \lambda + \sin^2 \lambda = 0. \quad (\text{A.13})$$

This is a null curve that is not straight, it is not a geodesic however.

A.3 Ladders and barns

Barry and Paul Chuckle have been employed by Albert E. to put a ladder in a barn, a simple feat you would imagine but these are the Chuckle brothers and nothing is simple with them. Albert E. stands outside the barn, and tells Barry and Paul to run very quickly at a constant

speed in a straight line through the barn carrying the ladder. The barn has doors at the front and back, and two apprentices (Jimmy and Brian) stand at either door ready to close or open them. Initially the front door is open and the back door is closed. The proper length of the ladder is l , while the proper length of the barn is b with $b < l$.

Albert E. claims that if Barry and Paul run fast enough, and that there is no slacking, then both doors of the barn can be temporarily closed with both Barry, Paul and the ladder inside the barn. One of the apprentices can then open the back door again so that Barry, Paul and the ladder can pass through the barn safely. The brothers are stumped, “oh dear, oh dear” says Barry, “the ladder is bigger than the barn, it will never work”. To put their minds at rest show that the ladder will fit in a chosen reference frame.

Let us work in inertial coordinates where the barn is at rest, which corresponds to Albert E.’s point of view. In these coordinates the front of the barn is at $x^\mu = (\lambda, 0, 0, 0)$ while the back of the barn is at $x^\mu = (\lambda, b, 0, 0)$.

The worldline of the front of the ladder in this reference frame is $x^\mu = (\lambda, v\lambda, 0, 0)$, where v is the velocity of the ladder. We have chosen coordinates so that the front of the ladder enters the barn at $\lambda = 0$. The back of the ladder follows the worldline $x^\mu = (\lambda, \lambda v - L, 0, 0)$ for some L which is not l !

First we must work out what L is in terms of l . We could of course perform a Lorentz transformation to switch to the rest frame of the ladder, the proper length of the ladder is then the coordinate length in this frame. We will use an alternative approach, staying in the original coordinate frame. How can we measure a length? Well we can define it to be half the proper time along the worldline at one end of the ladder between the emission and reception of a light signal which bounces off the other end of the body. The worldlines of the points making up the ladder are given by $x^\mu = (\lambda, \lambda v - \beta, 0, 0)$ where $\beta \in [0, L]$ and their tangent vectors are

$$\frac{dx^\mu}{d\lambda} = (1, v, 0, 0). \quad (\text{A.14})$$

We now want to find a spacelike straight line orthogonal to this tangent vector. Such a worldline is given by $n^\mu = (-v\tilde{\lambda}, -\tilde{\lambda}, 0, 0)$. This curve meets the front of the ladder at $\tilde{\lambda} = 0$ and the back of the ladder at $\tilde{\lambda} = L(1 - v^2)^{-1}$. We want to calculate the proper length of this curve with $\tilde{\lambda} \in [0, \frac{L}{1-v^2}]$. To do so we should parametrise the curve by the proper length. The norm of the tangent of the above vector is $\eta_{\mu\nu}\dot{n}^\mu(\tilde{\lambda})\dot{n}^\nu(\tilde{\lambda}) = 1 - v^2$. Then the proper length is

$$s = \tilde{\lambda}\sqrt{1 - v^2}. \quad (\text{A.15})$$

The ladder then has proper length

$$l = s \Big|_{\lambda = \frac{L}{1-v^2}} = \frac{L}{\sqrt{1-v^2}}. \quad (\text{A.16})$$

The entire ladder can fit into the barn from Albert E.'s perspective if $b \geq L$ and therefore the Chuckle brothers must run at a speed of

$$v \geq \sqrt{1 - \frac{b^2}{l^2}}. \quad (\text{A.17})$$

Both doors of the barn can be closed if: the front of the ladder is still in the barn, $b > v\lambda$ and the back of the ladder is in the barn $\lambda v - L > 0$. Since $t = \lambda$ the ladder is in the barn for

$$\frac{l\sqrt{1-v^2}}{v} \leq t \leq \frac{b}{v}. \quad (\text{A.18})$$

We see that Albert E. sees the ladder fully inside the barn with the doors closed.

Now consider what happens from the Chuckle brother's perspective. We can do a Lorentz transformation to coordinates in which they are at rest:

$$(t', x', y', z') = (\gamma t - \gamma v x, \gamma x - \gamma v t, y, z), \quad \gamma = \frac{1}{\sqrt{1-v^2}}. \quad (\text{A.19})$$

In the Chuckle brother's coordinates the barn follows the worldline $(\lambda, -v\lambda, 0, 0)$, while the back of the barn follows the worldline $(\lambda, -v\lambda + b\sqrt{1-v^2}, 0, 0)$.

The front door can be closed when the front of the barn passes the back of the ladder, so $-v\lambda < -l$ and therefore the front door of the barn is closed for $t' > \frac{l}{v}$.

The back door must open when the front of the ladder is about to go through it. So it is closed until $b\sqrt{1-v^2} - vt' = 0$ and therefore the back door is closed for $t' \in [0, \frac{b}{v\gamma}]$. In summary we have

$$\begin{cases} \text{Front door closed} & \frac{l}{v} \leq t', \\ \text{Back door closed} & 0 \leq t' \leq \frac{b}{v\gamma}. \end{cases} \quad (\text{A.20})$$

Since the ladder is longer than the barn $l > b$ and $\gamma \geq 1$ it follows that there is no time for which both doors are closed from the point of view of the Chuckle brothers. The entire ladder never fits into the barn from their perspective. The two view-points are depicted in figure 25.

Having been convinced by your arguments the brothers were off with a "to me, to you".³¹

³¹ChuckleVision was a British children's comedy tv show following the antics of the Chuckle brothers Barry and Paul. Carrying a ladder was a common theme.




Figure 25: The two different perspectives of the ladder and barn. On the left from the perspective of Albert E., a stationary observer in the rest frame of the barn. On the right from the perspective of the Chuckle brothers carrying the ladder.

B Euler–Lagrange equations in field theory

General relativity is an example of a classical field theory. We want to understand how to compute the equations of motion for such a field theory. Let us begin with the computation of the equations of motion in classical mechanics before building up to the field theory version.

B.1 Classical mechanics and Euler–Lagrange

Consider a single particle in one dimension with coordinate $q(t)$. The equations of motion are computed by using the *principle of least action*. We need to define a functional of an *action* S

$$S = \int dt L(q, \dot{q}), \quad (\text{B.1})$$

and find the critical points of S . Here $L(q, \dot{q})$ is known as the *Lagrangian*, and typically for point-particle mechanics takes the form:

$$L = \text{Kinetic Energy} - \text{Potential Energy}. \quad (\text{B.2})$$

From your classical mechanics course you will have derived the *Euler–Lagrange equations*:

$$\frac{\partial L}{\partial q} = \frac{d}{dt} \left(\frac{\partial L}{\partial \dot{q}} \right), \quad (\text{B.3})$$

see below for a recap of its derivation.

Consider a trajectory between the points q_1 and q_2 with $q(t_1) = q_1$ and $q(t_2) = q_2$. Let us deform the trajectory by

$$q(t) \rightarrow q(t) + \epsilon \delta q(t), \quad (\text{B.4})$$

whilst keeping the end-points fixed, that is $\delta q(t_1) = \delta q(t_2) = 0$. We have introduced the parameter ϵ which we take to be small. Consider now the action for this shifted path:

$$\begin{aligned} S[q(t) + \epsilon \delta q(t)] &= \int_{t_1}^{t_2} L(q + \epsilon \delta q, \dot{q} + \epsilon \frac{d}{dt} \delta q) dt \\ &= \int_{t_1}^{t_2} \left[L(q, \dot{q}) + \epsilon \left\{ \frac{\partial L}{\partial q} \delta q + \frac{\partial L}{\partial \dot{q}} \frac{d}{dt} \delta q \right\} + \mathcal{O}(\epsilon^2) \right] dt \\ &= S[q(t)] + \epsilon \left[\int_{t_1}^{t_2} \left(\frac{\partial L}{\partial q} - \frac{d}{dt} \left(\frac{\partial L}{\partial \dot{q}} \right) \right) \delta q dt + \left[\frac{\partial L}{\partial \dot{q}} \delta q \right]_{t_1}^{t_2} \right] + \mathcal{O}(\epsilon^2). \end{aligned} \quad (\text{B.5})$$

Note that due to our boundary conditions the last term vanishes. The first order variation

is then

$$\begin{aligned}\delta S &= S(q + \delta q) - S(q) \\ &= \epsilon \int_{t_1}^{t_2} \left(\frac{\partial L}{\partial q} - \frac{d}{dt} \left(\frac{\partial L}{\partial \dot{q}} \right) \right) \delta q dt\end{aligned}\tag{B.6}$$

and the requirement that this be an extremum, $\delta S = 0$ then implies

$$\frac{\partial L}{\partial q} - \frac{d}{dt} \left(\frac{\partial L}{\partial \dot{q}} \right) = 0,\tag{B.7}$$

which is the Euler–Lagrange equation.

B.2 Classical field theory and Euler–Lagrange

Having reviewed the Euler–Lagrange equations for classical mechanics consider the field theory version. We replace the single coordinate $q(t)$ by a set of spacetime dependent fields $\Phi^i(x^\mu)$, and take the action S to be a functional of these fields. The Lagrangian is expressed in terms of a *Lagrange density*, \mathcal{L} , which we take to be a function of the fields Φ^i and their spacetime derivatives, $\partial_\mu \Phi^i$:

$$L = \int d^3x \mathcal{L}(\Phi^i, \partial_\mu \Phi^i).\tag{B.8}$$

In principle the fields could be tensorial but we will suppress these details here, as we will see tensorial objects are necessary when considering the geodesics of GR. Similar to the classical mechanics case reviewed above we consider a small variation of the fields:

$$\Phi^i \rightarrow \Phi^i + \epsilon \delta \Phi^i, \quad \partial_\mu \Phi^i \rightarrow \partial_\mu \Phi^i + \epsilon \partial_\mu (\delta \Phi^i),\tag{B.9}$$

with ϵ our small parameter once again. The boundary conditions are similar to before with $\delta \Phi^i$ vanishing on the boundary of spacetime. The the first order variation of the action is

$$\begin{aligned}\delta S &= \epsilon \int d^4x \left[\frac{\partial \mathcal{L}}{\partial \Phi^i} \delta \Phi^i + \frac{\partial \mathcal{L}}{\partial (\partial_\mu \Phi^i)} \partial_\mu (\delta \Phi^i) \right] \\ &= \epsilon \int d^4x \left[\frac{\partial \mathcal{L}}{\partial \Phi^i} - \partial_\mu \left(\frac{\partial \mathcal{L}}{\partial (\partial_\mu \Phi^i)} \right) \right] \delta \Phi^i + \epsilon \int d^4x \partial_\mu \left(\frac{\partial \mathcal{L}}{\partial (\partial_\mu \Phi^i)} \delta \Phi^i \right).\end{aligned}\tag{B.10}$$

Due to our boundary conditions the last term vanishes and for a stationary point, $\delta S = 0$ we obtain the Euler–Lagrange equations

$$\frac{\partial \mathcal{L}}{\partial \Phi^i} - \partial_\mu \left(\frac{\partial \mathcal{L}}{\partial (\partial_\mu \Phi^i)} \right) = 0.\tag{B.11}$$

Exercise B.1: Euler–Lagrange equation examples

- Compute the Euler–Lagrange equations for a scalar field with Lagrangian density:

$$\mathcal{L} = -\frac{1}{2}\eta^{\mu\nu}\partial_\mu\phi\partial_\nu\phi - V(\phi). \quad (\text{B.12})$$

- Compute the Euler–Lagrange equations for electromagnetism with Lagrangian density:

$$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + A_\mu J^\mu, \quad (\text{B.13})$$

with $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$.

Hint: You should vary with respect to A and not F

C Solutions to exercises

In this appendix we provide solutions to the various exercises which are not covered in the tutorial problems.

C.1 Solutions to chapter 2

C.1.1 Exercise 2.1

- Under two successive boosts along the x direction we have

$$\begin{aligned}
\Lambda_{\text{final}} &= \Lambda_1^{\text{Boost}} \Lambda_1^{\text{Boost}} \\
&= \begin{pmatrix} \cosh \phi_1 & -\sinh \phi_1 & 0 & 0 \\ -\sinh \phi_1 & \cosh \phi_1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} \cosh \phi_2 & -\sinh \phi_2 & 0 & 0 \\ -\sinh \phi_2 & \cosh \phi_2 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \\
&= \begin{pmatrix} \cosh \phi_1 \cosh \phi_2 + \sinh \phi_1 \sinh \phi_2 & -\cosh \phi_1 \sinh \phi_2 - \sinh \phi_1 \cosh \phi_2 & 0 & 0 \\ -\cosh \phi_1 \sinh \phi_2 - \sinh \phi_1 \cosh \phi_2 & \cosh \phi_1 \cosh \phi_2 + \sinh \phi_1 \sinh \phi_2 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \\
&= \begin{pmatrix} \cosh(\phi_1 + \phi_2) & -\sinh(\phi_1 + \phi_2) & 0 & 0 \\ -\sinh(\phi_1 + \phi_2) & \cosh(\phi_1 + \phi_2) & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \\
&= \begin{pmatrix} \cosh \phi_{\text{final}} & -\sinh \phi_{\text{final}} & 0 & 0 \\ -\sinh \phi_{\text{final}} & \cosh \phi_{\text{final}} & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}
\end{aligned} \tag{C.1}$$

We see that we end up with another boost in the x -direction and that the rapidity of two successive boosts is additive: $\phi_{\text{final}} = \phi_1 + \phi_2$.

- We have

$$\begin{aligned}
e^{i\theta T^{\text{Rotation}}} &= 1_{4 \times 4} + \sum_{k=2}^{\infty} \frac{(i\theta)^k}{k!} (T^{\text{Rotation}})^k \\
&= \sum_{m=0}^{\infty} \frac{(-1)^m \theta^{2m}}{(2m)!} (T^{\text{Rotation}})^{2m} + i \sum_{m=0}^{\infty} \frac{(-1)^m \theta^{2m+1}}{(2m+1)!} (T^{\text{Rotation}})^{2m+1}.
\end{aligned} \tag{C.2}$$

Now

$$(T^{\text{Rotation}})^{2k} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad (T^{\text{Rotation}})^{2k+1} = T^{\text{Rotation}} \quad (\text{C.3})$$

and therefore we have

$$\begin{aligned} e^{i\theta T^{\text{Rotation}}} &= 1_{4 \times 4} + \sum_{m=2}^{\infty} \frac{(-1)^m \theta^{2m}}{(2m)!} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} + iT^{\text{Rotation}} \sum_{m=0}^{\infty} \frac{(-1)^m \theta^{2m+1}}{(2m+1)!} \\ &= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos \theta & \sin \theta & 0 \\ 0 & -\sin \theta & \cos \theta & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \end{aligned} \quad (\text{C.4})$$

where in the last step we use the series expansion of the trigonometric functions. A similar computation gives the boost.

3. Now consider the commutator of the generator of a boost along the x -direction with the generator of a boost along the y -direction. We have

$$\begin{aligned} [T_x^{\text{Boost}}, T_y^{\text{Boost}}] &= T_x^{\text{Boost}} T_y^{\text{Boost}} - T_y^{\text{Boost}} T_x^{\text{Boost}} \\ &= \begin{pmatrix} 0 & i & 0 & 0 \\ i & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \cdot \begin{pmatrix} 0 & 0 & i & 0 \\ 0 & 0 & 0 & 0 \\ i & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} - \begin{pmatrix} 0 & 0 & i & 0 \\ 0 & 0 & 0 & 0 \\ i & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \cdot \begin{pmatrix} 0 & i & 0 & 0 \\ i & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \\ &= -iT_{xy}^{\text{Rotation}} \end{aligned} \quad (\text{C.5})$$

4. Compute the commutator of the generators (2.10) for a boost along x and rotation in the x - y -plane.

We have the result:

$$[T_x^{\text{Boost}}, T_{xy}^{\text{Rotation}}] = -iT_y^{\text{Boost}} \quad (\text{C.6})$$

5. Compute the commutator of the generators ((2.10)) of a boost along x and rotation in the y - z -plane.

In this case we find that the commutator vanishes.

Let $K_i \equiv T_i^{\text{Boost}}$ and $L_i = \frac{1}{2} \epsilon_{ijk} T_{jk}^{\text{Rotation}}$ then we find that the commutators are:

$$[K_i, K_j] = -i\epsilon_{ijk} L_k, \quad [K_i, L_j] = i\epsilon_{ijk} K_k, \quad [L_i, L_j] = i\epsilon_{ijk} L_k. \quad (\text{C.7})$$

C.1.2 Exercise 2.2

We have just computed the addition of the rapidity. The velocity, in terms of the rapidity is (2.12)

$$v = \frac{x}{t} = \tanh \phi. \quad (\text{C.8})$$

We saw that the rapidity was additive. We have that after two boosts with rapidity/velocity (ϕ_i/v_i) we have

$$\begin{aligned} v_3 &= \tanh \phi_3 \\ &= \tanh(\phi_1 + \phi_2) \\ &= \frac{v_1 + v_2}{1 + v_1 v_2}. \end{aligned} \quad (\text{C.9})$$

This uses well known trig identities.

C.1.3 Exercise 2.3

We want to take the proper time defined in (2.16)

$$\Delta\tau = \int_{\lambda_1}^{\lambda_2} \sqrt{-\eta_{\mu\nu} \frac{dx^\mu}{d\lambda} \frac{dx^\nu}{d\lambda}} d\lambda, \quad (\text{C.10})$$

and show that under $\lambda \rightarrow \lambda(\sigma)$ the proper time is left invariant.

$$\begin{aligned} \Delta\tau &= \int_{\lambda_1}^{\lambda_2} \sqrt{-\eta_{\mu\nu} \frac{dx^\mu}{d\lambda} \frac{dx^\nu}{d\lambda}} d\lambda \\ &= \int_{\lambda_1(\sigma_1)}^{\lambda_2(\sigma_2)} \sqrt{-\eta_{\mu\nu} \frac{dx^\mu}{d\sigma} \frac{d\sigma}{d\lambda} \frac{dx^\nu}{d\sigma} \frac{d\lambda}{d\sigma}} d\sigma \\ &= \int_{\eta_1}^{\eta_2} \sqrt{-\eta_{\mu\nu} \frac{dx^\mu}{d\sigma} \frac{dx^\nu}{d\sigma}} d\sigma, \end{aligned} \quad (\text{C.11})$$

and therefore it is reparametrisation invariant. Note that the square root is important. Any other power would not be reparametrisation invariant!

C.1.4 Exercise 2.4

Show that it is always possible to find a parametrisation of a timelike curve so that it satisfies (2.17), and moreover that it is unique up to constant shifts.

Let us take a time-like curve. It must therefore satisfy

$$-f^2 = \eta_{\mu\nu} \frac{dx^\mu(\lambda)}{d\lambda} \frac{dx^\nu(\lambda)}{d\lambda}. \quad (\text{C.12})$$

We can make a change of reparametrisation which leads to

$$\begin{aligned} -f(x)^2 &= \eta_{\mu\nu} \frac{dx^\mu(\lambda)}{d\lambda} \frac{dx^\nu(\lambda)}{d\lambda} \\ &= \eta_{\mu\nu} \frac{dx^\mu}{d\sigma} \frac{d\sigma}{d\lambda} \frac{dx^\nu}{d\sigma} \frac{d\sigma}{d\lambda} \end{aligned} \quad (\text{C.13})$$

Bringing the factor of $\frac{d\sigma}{d\lambda}$ to the other side we have

$$-f(x)^2 \left[\frac{d\lambda}{d\sigma} \right]^2 = \eta_{\mu\nu} \frac{dx^\mu}{d\sigma} \frac{dx^\nu}{d\sigma}. \quad (\text{C.14})$$

We see that if we solve

$$f(x(\lambda(\sigma))) \frac{d\lambda}{d\sigma} = 1, \quad (\text{C.15})$$

then we have the correct normalised curve.

This is just a first order ODE which has a unique solution given a boundary condition. The boundary condition only fixes σ up to a constant which is the constant shift.

To see that σ is just the proper time we need to substitute this into (2.16) and everything falls into place.

C.1.5 Exercise 2.5

We take

$$\Lambda_\mu{}^\nu \equiv \eta_{\mu\rho} \eta^{\nu\sigma} \Lambda^\rho{}_\sigma, \quad (\text{C.16})$$

then

$$\begin{aligned} \Lambda_\mu{}^\nu \Lambda^\tau{}_\nu &= \eta_{\mu\rho} \eta^{\nu\sigma} \Lambda^\rho{}_\sigma \Lambda^\tau{}_\nu \\ &= \eta_{\mu\rho} \eta^{\rho\tau} \\ &= \delta_\mu^\tau \end{aligned} \quad (\text{C.17})$$

C.1.6 Exercise 2.6

- If V^ν transforms as a vector then

$$V'^\mu = \Lambda^\mu{}_\nu V^\nu. \quad (\text{C.18})$$

Transforming $\eta_{\mu\nu} V^\nu$ we would have

$$\begin{aligned} (\eta_{\mu\nu} V^\nu)' &= \eta'_{\mu\nu} V'^\nu \\ &= \eta_{\tau\kappa} \Lambda_\mu{}^\tau \Lambda_\nu{}^\kappa \Lambda^\nu{}_\sigma V^\sigma \\ &= \Lambda_\mu{}^\tau \eta_{\tau\kappa} \delta_\sigma^\kappa V^\sigma \\ &= \Lambda_\mu{}^\tau (\eta_{\tau\sigma} V^\sigma) \end{aligned} \quad (\text{C.19})$$

We have used the result from exercise (2.5) in going to the third line.

- This works in the same way as the one above.

C.1.7 Exercise 2.7

Integrating equation (2.56) over the volume in a sphere of radius r , centred about the center of symmetry, and whose surface contains all of the mass one finds

$$\int_{B^3(r)} d^3x \nabla \cdot F = 4\pi G \int_{B^3(r)} d^3x \rho(r) = -4\pi GM, \quad (\text{C.20})$$

where M is the total mass. The divergence theorem allows us to express the left-hand side as a surface integral over the sphere of radius r giving

$$\int_r d\vec{A} \cdot F = -4\pi GM. \quad (\text{C.21})$$

Due to the spherical symmetry F can only depend on r and point only in a radial direction. This implies that the surface integral is simply

$$\int_r d\vec{A} \cdot F = 4\pi r^2 |F|, \quad (\text{C.22})$$

where $|F|$ is the magnitude of F . Therefore if e_r is a unit vector in the radial direction then

$$F = -\frac{GM}{r^2} e_r, \quad (\text{C.23})$$

and depends only on M . We can now solve for Φ , this is of course only defined up to a constant and if we choose the constant so that Φ vanishes at ∞ then we end up with the simple expression

$$\Phi(r) = -\frac{GM}{r}. \quad (\text{C.24})$$

We learn that it does not matter how the mass is distributed in a spherically symmetric configuration. It could be localised at the centre or in spherically symmetric shells. The potential only depends on the total mass.

C.2 Solutions to chapter 3

C.2.1 Exercise 3.1

The discrete topology is the obtained by taking a collection of all subsets of the topological space.

The usual topology is obtained by taking all open sets (a, b) and their unions. Imagine we were to change the 3rd condition in the definition of a topological space to include an infinite subcollection of I . Then we could take the subsets

$$U_n = (a, b + \frac{1}{n}). \quad (\text{C.25})$$

Taking the infinite intersection we have

$$\bigcap_{n=1}^{\infty} (a, b + \frac{1}{n}) = (a, b]. \quad (\text{C.26})$$

This implies that $\{b\}$ is in the topology and this holds for all b . Therefore the topology is reduced to the discrete one.

C.2.2 Exercise 3.2

To show that the Tangent space at p is a vector space we need to show a number of properties. It boils down to show that if $X_1, X_2 \in T_p(M)$ and $\alpha \in \mathbb{R}$ then $X_1 + \alpha X_2 \in T_p(M)$. By definition both X_i are linear transformations on functions. We then need to show $X_1 + \alpha X_2$ satisfied the Leibniz property and is therefore a tangent vector too. For functions f, g we have

$$\begin{aligned} (X_1 + \alpha X_2)(fg) &= X_1(fg) + \alpha X_2(fg) \\ &= fX_1(g) + gX_1(f) + \alpha(gX_2(f) + fX_2(g)) \\ &= f(X_1 + \alpha X_2)(g) + g(X_1 + \alpha X_2)(f), \end{aligned} \quad (\text{C.27})$$

hence $f(X_1 + \alpha X_2)(g) \in T_p(M)$

C.2.3 Exercise 3.3

We want to show that the flow defines a commutative group. This turns out to be simple since

$$\begin{aligned} \sigma_t(\sigma_s(x)) &= \sigma_{t+s}(x) \\ &= \sigma_{s+t}(x) \\ &= \sigma_s(\sigma_t(x)), \end{aligned} \quad (\text{C.28})$$

and therefore it commutes. Note that we have used that \mathbb{R} is commutative. Since it is 1-dimensional there are two options, \mathbb{R} or S^1 .

C.2.4 Exercise 3.4

These are given in problem sheet 2.

C.2.5 Exercise 3.5

Let $X, Y, Z \in \mathcal{X}(M)$ be vector fields. We have

$$\mathcal{L}_X \mathcal{L}_Y Z = \mathcal{L}_X([Y, Z]) = [X, [Y, Z]]. \quad (\text{C.29})$$

The Lie bracket satisfies the Jacobi identity, which after some rearranging can be written in the form

$$[X, [Y, Z]] - [Y, [X, Z]] = -[[X, Y], Z]. \quad (\text{C.30})$$

Using our expression above for the Lie derivative we have

$$\mathcal{L}_X \mathcal{L}_Y Z - \mathcal{L}_Y \mathcal{L}_X Z = \mathcal{L}_{[X, Y]} Z, \quad (\text{C.31})$$

as required.

C.2.6 Exercise 3.6

We want to show:

1. That the Lie derivative satisfies:

$$\mathcal{L}_X(t_1 + t_2) = \mathcal{L}_X t_1 + \mathcal{L}_X t_2, \quad (\text{C.32})$$

where t_1 and t_2 are tensor fields of the same type.

2. That

$$\mathcal{L}_X(t_1 \otimes t_2) = (\mathcal{L}_X t_1) \otimes t_2 + t_1 \otimes (\mathcal{L}_X t_2), \quad (\text{C.33})$$

with t_1 and t_2 tensors of arbitrary type.

For the first by the way we constructed the Lie derivative for vector fields, one forms and scalars this is obvious.

For the second we have to do a little work. We will show it in a single example, but it will be obvious how it generalises. Take a vector field Y and one-form ω and construct the tensor product $Y \otimes \omega$. Then $(Y \otimes \omega)|_{\sigma_\epsilon(x)}$ is mapped onto a tensor at x by the action of $(\sigma_{-\epsilon})^* \otimes (\sigma_\epsilon)^*$. Therefore

$$[(\sigma_{-\epsilon})^* \otimes (\sigma_\epsilon)^*](Y \otimes \omega)|_{\sigma_\epsilon(x)} = [(\sigma_{-\epsilon})_* Y \otimes (\sigma_\epsilon)^* \omega]|_x. \quad (\text{C.34})$$

We therefore have

$$\begin{aligned} \mathcal{L}_X &= \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} [[(\sigma_{-\epsilon})_* Y \otimes (\sigma_\epsilon)^* \omega]|_x - (Y \otimes \omega)|_x] \\ &= \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} [(\sigma_{-\epsilon})_* Y \otimes \{(\sigma_\epsilon)^* \omega - \omega\} + \{(\sigma_{-\epsilon})_* Y - Y\} \otimes \omega] \\ &= Y \otimes \mathcal{L}_X \omega + \mathcal{L}_X Y \otimes \omega. \end{aligned} \quad (\text{C.35})$$

For other tensors the computation works similarly.

C.2.7 Exercise 3.7

From the properties of the wedge product show that for $\xi \in \Omega_p^q(M)$, $\eta \in \Omega_p^r(M)$ and $\omega \in \Omega_p^s(M)$ that

$$\begin{aligned}\xi \wedge \eta &= (-1)^{qr} \eta \wedge \xi, \\ \xi \wedge \xi &= 0 \quad \text{if } q \text{ odd}, \\ (\xi \wedge \eta) \wedge \omega &= \xi \wedge (\eta \wedge \omega).\end{aligned}\tag{C.36}$$

Recall that the action is given by:

$$(\omega \wedge \xi)(V_1, \dots, V_{q+r}) = \frac{1}{q!r!} \sum_{\sigma \in S_{q+r}} \text{sign}(\sigma) \omega(V_{\sigma(1)}, \dots, V_{\sigma(q)}) \xi(V_{\sigma(q+1)}, \dots, V_{\sigma(q+r)}).\tag{C.37}$$

Then if we have two forms ξ, η of degree q, r respectively then

$$\begin{aligned}(\xi \wedge \eta)(V_1, \dots, V_{q+r}) &= \frac{1}{q!r!} \sum_{\sigma \in S_{q+r}} \text{sign}(\sigma) \xi(V_{\sigma(1)}, \dots, V_{\sigma(q)}) \eta(V_{\sigma(q+1)}, \dots, V_{\sigma(q+r)}) \\ &= \frac{1}{q!r!} \sum_{\sigma \in S_{q+r}} \text{sign}(\sigma) \eta(V_{\sigma(q+1)}, \dots, V_{\sigma(q+r)}) \xi(V_{\sigma(1)}, \dots, V_{\sigma(q)})\end{aligned}\tag{C.38}$$

We now want to move the indices on the V 's to be in the canonical order. Assume $q \geq r$, the alternative is easily checked too. If we exchange an index we pick up a sign from the permutation. We can flip the indices in η with the first r indices in the ξ . This will introduce r signs.

$$\begin{aligned}(\xi \wedge \eta)(V_1, \dots, V_{q+r}) &= (-1)^r \frac{1}{q!r!} \sum_{\sigma \in S_{q+r}} \text{sign}(\sigma) \eta(V_{\sigma(1)}, \dots, V_{\sigma(r)}) \xi(V_{\sigma(q+1)}, \dots, V_{\sigma(r+q)}, \dots, V_{\sigma(r+1)}, \dots, V_{\sigma(q)}) \\ &= (-1)^r (-1)^{r(q-r)} \frac{1}{q!r!} \sum_{\sigma \in S_{q+r}} \text{sign}(\sigma) \eta(V_{\sigma(1)}, \dots, V_{\sigma(r)}) \xi(V_{\sigma(r+1)}, \dots, V_{\sigma(r+q)}) \\ &= (-1)^{r(q-r-1)} (\eta \wedge \xi)(V_1, \dots, V_{q+r}) \\ &= (-1)^{rq} (-1)^{-r(r+1)} (\eta \wedge \xi)(V_1, \dots, V_{q+r}) \\ &= (-1)^{rq} (\eta \wedge \xi)(V_1, \dots, V_{q+r},\end{aligned}\tag{C.39}$$

where we used that $(-1)^{-r(r+1)} = 1$.

This proves the first, the second follows simply from the first since

$$\xi \wedge \xi = (-1)^{r^2} \xi \wedge \xi = (-1)^r \xi \wedge \xi = \begin{cases} \xi \wedge \xi & r \in 2\mathbb{Z} \\ -\xi \wedge \xi & r \notin 2\mathbb{Z} \end{cases}\tag{C.40}$$

It then follows for odd dimensional form it vanishes.

The associativity property follows from the associativity of functions and some messing with permutations. Let us use the shorthand $[V] = (V_1, \dots, V_{q+r+s})$ then

$$((\xi \wedge \eta) \wedge \omega)[V] = \frac{1}{s!} \frac{1}{(q+r)!} \sum_{\sigma \in S_{q+r+s}} \text{sign}(\sigma) (\xi \wedge \eta)(V_{\sigma(1)}, \dots, V_{\sigma(q+r)}) \omega(V_{\sigma(q+r+1)}, \dots, V_{\sigma(q+r+s)}) \quad (\text{C.41})$$

We now want to simplify the permutations. We can decompose the permutation group S_{q+r+s} into residual classes of the subgroup $S_{q+r} \subset S_{q+r+s}$. These are permutations which act as the identity on the final s indices. Let us call the set of these residual classes \mathcal{C} and let $R \in \mathcal{C}$ be one of these residual classes. Each $R \in \mathcal{C}$ is isomorphic to S_{q+r} . Let us pick one class R and within the class a particular permutation σ_R . Then each element $\sigma \in R$ can be decomposed as $\sigma = \sigma_R \otimes \pi$ where $\pi \in S_{q+r}$. We therefore have that the sum over S_{q+r+s} may be written as:

$$((\xi \wedge \eta) \wedge \omega)[V] = \frac{1}{s!} \frac{1}{(q+r)!} \sum_{R \in \mathcal{C}} \text{sign}(\sigma_R) \left[\sum_{\sigma \in R} \text{sign}(\pi) (\xi \wedge \eta)(V_{\sigma(1)}, \dots, V_{\sigma(q+r)}) \right] \times \omega(V_{\sigma_R(q+r+1)}, \dots, V_{\sigma_R(q+r+s)}) \quad (\text{C.42})$$

The terms in the brackets are all equal because they are permutations of π from a fixed ordering given by σ_R . Since each residual class R is isomorphic to S_{q+r} there are $(q+r)!$ terms. Thus,

$$\begin{aligned} ((\xi \wedge \eta) \wedge \omega)[V] &= \frac{1}{s!} \sum_{R \in \mathcal{C}} \text{sign}(\sigma_R) (\xi \wedge \eta)(V_{\sigma_R(1)}, \dots, V_{\sigma_R(q+r)}) \omega(V_{\sigma_R(q+r+1)}, \dots, V_{\sigma_R(q+r+s)}) \\ &= \frac{1}{q!r!s!} \sum_{R \in \mathcal{C}} \text{sign}(\sigma_R) \sum_{\tau \in S_{q+r}} \text{sign}(\tau) \xi(V_{\tau(\sigma_R(1))}, \dots, V_{\tau(\sigma_R(q))}) \\ &\quad \times \eta(V_{\tau(\sigma_R(q+1))}, \dots, V_{\tau(\sigma_R(q+r))}) \omega(V_{\sigma_R(q+r+1)}, \dots, V_{\sigma_R(q+r+s)}) \end{aligned} \quad (\text{C.43})$$

We need to use a similar trick to before. All permutations $\sigma \in S_{q+r+s}$ can be decomposed as $\sigma = \tau \circ \sigma_R$ and since τ acts on the last s indices as the identity we have $\sigma = \sigma_R$ for these. Thus we obtain:

$$((\xi \wedge \eta) \wedge \omega)[V] = \frac{1}{q!r!s!} \sum_{\sigma \in S_{q+r+s}} \text{sign}(\sigma) \xi(V_{\sigma(1)}, \dots, V_{\sigma(q)}) \times \eta(V_{\sigma(q+1)}, \dots, V_{\sigma(q+r)}) \omega(V_{\sigma(q+r+1)}, \dots, V_{\sigma(q+r+s)}) \quad (\text{C.44})$$

We can play the same game starting from $(\xi \wedge (\eta \wedge \omega))[V]$. It is clear though that the difference is that we decompose in terms of the residual class of $S_{r+s} \subset S_{q+r+s}$ instead and we end up with the exact same result. We have therefore shown associativity.

C.2.8 Exercise 3.8

Show that for $\xi \in \Omega_p^q(M)$, $\eta \in \Omega_p^r(M)$ we have

$$d(\xi \wedge \eta) = d\xi \wedge \eta + (-1)^q \xi \wedge d\eta. \quad (\text{C.45})$$

We can do this in coordinate free notation or using a coordinate basis, we choose the latter since it is simpler though the former is equally doable.

Let

$$\xi = \frac{1}{q!} \xi_{\mu_1 \dots \mu_q} dx^{\mu_1} \wedge \dots \wedge dx^{\mu_q}, \quad \eta = \frac{1}{r!} \eta_{\mu_1 \dots \mu_r} dx^{\mu_1} \wedge \dots \wedge dx^{\mu_r}, \quad (\text{C.46})$$

then

$$\begin{aligned} d(\xi \wedge \eta) &= d(\xi_{\mu_1 \dots \mu_q} \eta_{\nu_1 \dots \nu_r} dx^{\mu_1} \wedge \dots \wedge dx^{\mu_q} \wedge dx^{\nu_1} \wedge \dots \wedge dx^{\nu_r}) \\ &= \frac{1}{q!r!} \frac{\partial}{\partial x^\sigma} (\xi_{\mu_1 \dots \mu_q} \eta_{\nu_1 \dots \nu_r}) dx^\sigma \wedge dx^{\mu_1} \wedge \dots \wedge dx^{\mu_q} \wedge dx^{\nu_1} \wedge \dots \wedge dx^{\nu_r} \\ &= \frac{1}{q!r!} \left(\frac{\partial}{\partial x^\sigma} \xi_{\mu_1 \dots \mu_q} \right) \eta_{\nu_1 \dots \nu_r} dx^\sigma \wedge dx^{\mu_1} \wedge \dots \wedge dx^{\mu_q} \wedge dx^{\nu_1} \wedge \dots \wedge dx^{\nu_r} \\ &\quad + (-1)^q \frac{1}{q!r!} \xi_{\mu_1 \dots \mu_q} \left(\frac{\partial}{\partial x^\sigma} \eta_{\nu_1 \dots \nu_r} \right) dx^{\mu_1} \wedge \dots \wedge dx^{\mu_q} \wedge dx^\sigma \wedge dx^{\nu_1} \wedge \dots \wedge dx^{\nu_r} \\ &= d\xi \wedge \eta + (-1)^q \xi \wedge d\eta. \end{aligned} \quad (\text{C.47})$$

C.2.9 Exercise 3.9

$$\begin{aligned} i_X^2 &= 0, \\ i_X(\omega \wedge \eta) &= i_X \omega \wedge \eta + (-1)^r \omega \wedge i_X \eta, \\ i_{[X,Y]} \omega &= X(i_Y \omega) - Y(i_X \omega), \\ \mathcal{L}_X i_X \omega &= i_X \mathcal{L}_X \omega. \end{aligned} \quad (\text{C.48})$$

1. For the first if we contract into a q -form ω we have

$$i_X^2 \omega = \frac{1}{(q-2)!} X^{\mu_1} X^{\mu_2} \omega_{\mu_1 \mu_2 \nu_1 \dots \nu_{q-2}} dx^{\nu_1} \wedge \dots \wedge dx^{\nu_{q-2}} \quad (\text{C.49})$$

Since the two X 's are symmetric and the two indices of the p -form are antisymmetric this vanishes.

2. In components we have:

$$\begin{aligned} i_{V_1}(\omega \wedge \eta)(V_2, \dots, V_{q+r}) &= (\omega \wedge \eta)(V_1, \dots, V_{q+r}) \\ &= \frac{1}{q!r!} \sum_{\sigma \in S_{q+r}} \text{sign}(\sigma) \omega(V_{\sigma(1)}, \dots, V_{\sigma(q)}) \eta(V_{\sigma(q+1)}, \dots, V_{\sigma(q+r)}) \end{aligned} \quad (\text{C.50})$$

We now want to identify the ones with the V_1 and the ones without.

$$\begin{aligned} i_{V_1}(\omega \wedge \eta)(V_2, \dots, V_{q+r}) &= \frac{1}{q!r!} \sum_{\sigma \in S_{q+r-1}} \left[q\omega(V_1 V_{\sigma(2)}, \dots, V_{\sigma(q)})\eta(V_{\sigma(q+1)}, \dots, V_{\sigma(q+r)}) \right. \\ &\quad \left. + (-1)^q r\omega(V_{\sigma(2)}, \dots, V_{\sigma(q+1)})\eta(V_1, V_{\sigma(q+2)}, \dots, V_{\sigma(q+r)}) \right] \\ &= (i_{V_1}\omega) \wedge \eta + (-1)^q \omega \wedge (i_{V_1}\eta). \end{aligned} \tag{C.51}$$

3. Using that on a form

$$\mathcal{L}_X \omega = (i_X d + di_X)\omega, \tag{C.52}$$

we have

$$\begin{aligned} \mathcal{L}_X i_X \omega &= (i_X d + di_X)i_X \omega \\ &= i_X(di_X \omega) \\ &= i_X \mathcal{L}_X \omega \end{aligned} \tag{C.53}$$

C.2.10 Exercise 3.10

Stoke's theorem states that in 3d, given a vector field \vec{F} and a smooth oriented surface in \mathbb{R}^3 with boundary $\partial\Sigma = B$ then

$$\int_{\Sigma} (\nabla \times \vec{F}) \cdot d\Sigma = \oint_{\partial\Sigma} \vec{F} \cdot d\Gamma \tag{C.54}$$

We need to identify the vector field F as a 1-form and the curl as a two form given by the exterior derivative of F . We have that

$$\vec{F} \cdot d\Gamma = F_x dx + F_y dy + F_z dz \equiv F, \tag{C.55}$$

similarly

$$dF = (\partial_x F_y - \partial_y F_x)dx \wedge dy + (\partial_y F_z - \partial_z F_y)dy \wedge dz + (\partial_z F_x - \partial_x F_z)dz \wedge dx \tag{C.56}$$

this is nothing other than

$$dF = (\nabla \times \vec{F}) \cdot dS, \tag{C.57}$$

and therefore putting everything together we have the claimed result.

C.3 Solutions to chapter 4

C.3.1 Exercise 4.1

We take the embedding of the sphere into \mathbb{R}^3 given via equation (4.11). Plugging this parametrisation into the metric on \mathbb{R}^3

$$ds^2 = dx^2 + dy^2 + dz^2, \quad (\text{C.58})$$

we find

$$\begin{aligned} ds^2(S^2) &= [d(\sin \theta \cos \phi)]^2 + [d(\sin \theta \sin \phi)]^2 + [d \cos \theta]^2 \\ &= d\theta^2 + \sin^2 \theta d\phi^2. \end{aligned} \quad (\text{C.59})$$

C.3.2 Exercise 4.2

1. Let us take the torsion tensor defined via (4.59). Then

$$\begin{aligned} T(\omega; fX, Y) &= \omega(\nabla_{fX}Y - \nabla_Y(fX) - [fX, Y]) \\ &= \omega(f\nabla_XY - Y[f]X - f[X, Y] + Y[f]X) \\ &= \omega(f(\nabla_XY - \nabla_YX - [X, Y])) \\ &= f\omega(\nabla_XY - \nabla_YX - [X, Y]) \\ &= fT(\omega; X, Y). \end{aligned} \quad (\text{C.60})$$

For f on the other arguments it follows identically to above on Y and trivially on ω . We also need to show that if we take sums of two then it all becomes linear as well.

$$\begin{aligned} T(\omega; X_1 + X_2, Y) &= \omega(\nabla_{X_1+X_2}Y - \nabla_Y(X_1 + X_2) - [X_1 + X_2, Y]) \\ &= \omega(\nabla_{X_1}Y + \nabla_{X_2}Y - \nabla_YX_1 - \nabla_YX_2 - [X_1, Y] - [X_2, Y]) \\ &= \omega(\nabla_{X_1}Y - \nabla_YX_1 - [X_1, Y]) + \omega(\nabla_{X_2}Y - \nabla_YX_2 - [X_2, Y]) \\ &= T(\omega; X_1, Y) + T(\omega; X_2, Y), \end{aligned} \quad (\text{C.61})$$

with a similar result for other arguments. This shows that T is a multi-linear map in all indices. Note that we have used the established results that ω is a linear map, ∇_X a linear map and properties of the connection to show this.

2. We now want to show that this is a tensor by checking that it transforms correctly in coordinate components. In components we have that

$$T^\rho{}_{\mu\nu} = \Gamma^\rho{}_{\mu\nu} - \Gamma^\rho{}_{\nu\mu}, \quad (\text{C.62})$$

we have already seen the transformation of the connection coefficients under a coordinate transformation in (4.51),

$$\tilde{\Gamma}^{\mu}_{\nu\rho} = (\Lambda^{-1})^{\mu}_{\kappa} \Lambda^{\sigma}_{\rho} \Lambda^{\tau}_{\nu} \Gamma^{\kappa}_{\sigma\tau} + (\Lambda^{-1})^{\mu}_{\kappa} \Lambda^{\sigma}_{\rho} \partial_{\sigma} \Lambda^{\kappa}_{\nu} \quad (\text{C.63})$$

We therefore have

$$\begin{aligned} \tilde{T}^{\rho}_{\mu\nu} &= \tilde{\Gamma}^{\rho}_{\mu\nu} - \tilde{\Gamma}^{\rho}_{\nu\mu} \\ &= (\Lambda^{-1})^{\rho}_{\kappa} \Lambda^{\sigma}_{\nu} \Lambda^{\tau}_{\mu} \Gamma^{\kappa}_{\sigma\tau} + (\Lambda^{-1})^{\rho}_{\kappa} \Lambda^{\sigma}_{\nu} \partial_{\sigma} \Lambda^{\kappa}_{\mu} \\ &\quad - [(\Lambda^{-1})^{\rho}_{\kappa} \Lambda^{\sigma}_{\nu} \Lambda^{\tau}_{\mu} \Gamma^{\kappa}_{\tau\sigma} + (\Lambda^{-1})^{\rho}_{\kappa} \Lambda^{\sigma}_{\mu} \partial_{\sigma} \Lambda^{\kappa}_{\nu}] \\ &= (\Lambda^{-1})^{\rho}_{\kappa} \Lambda^{\sigma}_{\nu} \Lambda^{\tau}_{\mu} (\Gamma^{\kappa}_{\tau\sigma} - \Gamma^{\kappa}_{\sigma\tau}) - \frac{\partial x^{\sigma}}{\partial y^{[\nu}} \frac{\partial x^{\kappa}}{\partial y^{\mu]}} \frac{\partial^2 y^{\rho}}{\partial x^{\sigma} \partial x^{\kappa}} \\ &\quad (\Lambda^{-1})^{\rho}_{\kappa} \Lambda^{\sigma}_{\nu} \Lambda^{\tau}_{\mu} T^{\kappa}_{\tau\sigma} \end{aligned} \quad (\text{C.64})$$

which is as claimed. We have used the coordinate expression for the Λ 's to simplify the derivative term and then that the contraction of anti-symmetric indices with symmetric indices vanishes.

C.3.3 Exercise 4.3

Covered in problem sheet 3.

References

- [1] N. Arkani-Hamed, L. Motl, A. Nicolis and C. Vafa, *The String landscape, black holes and gravity as the weakest force*, *JHEP* **06** (2007) 060, [[hep-th/0601001](#)].
- [2] N. S. Ginsberg, S. R. Garner and L. V. Hau, *Coherent control of optical information with matter wave dynamics*, *Nature* **445** (2007) 623–626.
- [3] M. Nakahara, *Geometry, topology and physics*. 2003.