

C3.3 Differentiable Manifolds revision lecture,  
May 2026  
To go over 2023 C3.3 paper

Dominic Joyce

These slides available on course website

## C3.3 2023 question 1

**(a) (6 marks)** Define a *chart*, an *atlas*, and a *maximal atlas* on a topological space  $X$ . Define (*smooth*) *manifolds*.

**(b) (3 marks)** Define  $X$  to be the set of unoriented affine real lines in  $\mathbb{R}^3$ , made into a topological space in the natural way. One way to do this is to note that

$$X \cong \{(\mathbf{u}, \mathbf{v}) : \mathbf{u}, \mathbf{v} \in \mathbb{R}^3 : |\mathbf{u}| = 1, \mathbf{u} \cdot \mathbf{v} = 0\} / (\mathbf{u}, \mathbf{v}) \sim (-\mathbf{u}, \mathbf{v}),$$

where  $(\pm\mathbf{u}, \mathbf{v})$  corresponds to the line  $\{t\mathbf{u} + \mathbf{v} : t \in \mathbb{R}\}$ . Prove that  $X$  has the properties required of the topological space of a manifold.

**(c) (7 marks)** Define three charts  $(U_1, \varphi_1), (U_2, \varphi_2), (U_3, \varphi_3)$  on  $X$  by  $U_1 = U_2 = U_3 = \mathbb{R}^4$  and

$$\varphi_1 : (a_1, b_1, c_1, d_1) \mapsto \{(x, y, z) \in \mathbb{R}^3 : y = a_1x + b_1, z = c_1x + d_1\},$$

$$\varphi_2 : (a_2, b_2, c_2, d_2) \mapsto \{(x, y, z) \in \mathbb{R}^3 : z = a_2y + b_2, x = c_2y + d_2\},$$

$$\varphi_3 : (a_3, b_3, c_3, d_3) \mapsto \{(x, y, z) \in \mathbb{R}^3 : x = a_3z + b_3, y = c_3z + d_3\}.$$

Prove that  $\{(U_1, \varphi_1), (U_2, \varphi_2), (U_3, \varphi_3)\}$  is an atlas on  $X$ . Deduce that  $X$  is a smooth manifold.

[You may assume that  $(U_1, \varphi_1), (U_2, \varphi_2), (U_3, \varphi_3)$  are charts.]

**(d) (6 marks)** Prove that  $X$  is orientable.

[Hint: prove the transition functions are orientation-preserving.]

**(e) (3 marks)** Now let  $Y$  be the set of (unoriented) affine real lines in  $\mathbb{R}^2$ , made into a manifold in a similar way. Is  $Y$  orientable? Give brief justification.

## C3.3 2023 question 1

(a) (6 marks) Define a *chart*, an *atlas*, and a *maximal atlas* on a topological space  $X$ . Define (*smooth*) *manifolds*. [Blue = bookwork.]

(b) (3 marks)] Define  $X$  to be the set of unoriented affine real lines in  $\mathbb{R}^3$ , made into a topological space in the natural way. One way to do this is to note that

$$X \cong \{(\mathbf{u}, \mathbf{v}) : \mathbf{u}, \mathbf{v} \in \mathbb{R}^3 : |\mathbf{u}| = 1, \mathbf{u} \cdot \mathbf{v} = 0\} / (\mathbf{u}, \mathbf{v}) \sim (-\mathbf{u}, \mathbf{v}),$$

where  $(\pm\mathbf{u}, \mathbf{v})$  corresponds to the line  $\{t\mathbf{u} + \mathbf{v} : t \in \mathbb{R}\}$ . Prove that  $X$  has the properties required of the topological space of a manifold.

(c) (7 marks) Define three charts  $(U_1, \varphi_1), (U_2, \varphi_2), (U_3, \varphi_3)$  on  $X$  by  $U_1 = U_2 = U_3 = \mathbb{R}^4$  and [Green = similar to coursework. Easier.]

$$\varphi_1 : (a_1, b_1, c_1, d_1) \mapsto \{(x, y, z) \in \mathbb{R}^3 : y = a_1x + b_1, z = c_1x + d_1\},$$

$$\varphi_2 : (a_2, b_2, c_2, d_2) \mapsto \{(x, y, z) \in \mathbb{R}^3 : z = a_2y + b_2, x = c_2y + d_2\},$$

$$\varphi_3 : (a_3, b_3, c_3, d_3) \mapsto \{(x, y, z) \in \mathbb{R}^3 : x = a_3z + b_3, y = c_3z + d_3\}.$$

Prove that  $\{(U_1, \varphi_1), (U_2, \varphi_2), (U_3, \varphi_3)\}$  is an atlas on  $X$ . Deduce that  $X$  is a smooth manifold.

[You may assume that  $(U_1, \varphi_1), (U_2, \varphi_2), (U_3, \varphi_3)$  are charts.]

(d) (6 marks) Prove that  $X$  is orientable. [Red = new. Harder.]

[Hint: prove the transition functions are orientation-preserving.]

(e) (3 marks) Now let  $Y$  be the set of (unoriented) affine real lines in  $\mathbb{R}^2$ , made into a manifold in a similar way. Is  $Y$  orientable? Give brief justification.

# How to do Oxford exam questions if you are new to them

- Questions usually divided into **bookwork**, **similar**, **new**.
- 2023 q 1: 6 marks bookwork, q 2 11 marks bw, q 3 11 marks bw.
- **Learn** likely definitions, theorems, proofs bookwork. Look at past papers to see what bookwork has been asked before.
- You answer each question in a separate booklet. If you get stuck on one question you can go on to the next, and come back to it.
- If a question has parts (a)–(e) and you can't do (c), you can still attempt (d),(e).
- **Make sure you get all the easy marks on two questions!**
- Past papers are the best guide to what this year's papers are like.
- **Practice** past papers under timed conditions. You want to be able to finish a question in 45 minutes.

(a)[6 marks] Define a *chart*, an *atlas*, and a *maximal atlas* on a topological space  $X$ . Define (*smooth*) *manifolds*.

All bookwork.

Don't forget Hausdorff and second countable conditions on  $X$ .

(b)[3 marks] Define  $X$  to be the set of unoriented affine real lines in  $\mathbb{R}^3$ , made into a topological space in the natural way. One way to do this is to note that

$$X \cong \{(\mathbf{u}, \mathbf{v}) : \mathbf{u}, \mathbf{v} \in \mathbb{R}^3 : |\mathbf{u}| = 1, \mathbf{u} \cdot \mathbf{v} = 0\} / (\mathbf{u}, \mathbf{v}) \sim (-\mathbf{u}, \mathbf{v}),$$

where  $(\pm\mathbf{u}, \mathbf{v})$  corresponds to the line  $\{t\mathbf{u} + \mathbf{v} : t \in \mathbb{R}\}$ . Prove that  $X$  has the properties required of the topological space of a manifold.

Need to show that  $X$  is Hausdorff and second countable.

The space  $\{(\mathbf{u}, \mathbf{v}) : \mathbf{u}, \mathbf{v} \in \mathbb{R}^3 : |\mathbf{u}| = 1, \mathbf{u} \cdot \mathbf{v} = 0\}$  is both as it is a subset of  $\mathbb{R}^6$  with the subspace topology, and  $\mathbb{R}^6$  is both.

Hence  $X$  is both, as it is the quotient of a Hausdorff and second countable space by a finite group.

(c)[7 marks] Define three charts  $(U_1, \varphi_1), (U_2, \varphi_2), (U_3, \varphi_3)$  on  $X$  by  $U_1 = U_2 = U_3 = \mathbb{R}^4$  and

$$\varphi_1 : (a_1, b_1, c_1, d_1) \mapsto \{(x, y, z) \in \mathbb{R}^3 : y = a_1x + b_1, z = c_1x + d_1\},$$

$$\varphi_2 : (a_2, b_2, c_2, d_2) \mapsto \{(x, y, z) \in \mathbb{R}^3 : z = a_2y + b_2, x = c_2y + d_2\},$$

$$\varphi_3 : (a_3, b_3, c_3, d_3) \mapsto \{(x, y, z) \in \mathbb{R}^3 : x = a_3z + b_3, y = c_3z + d_3\}.$$

Prove that  $\{(U_1, \varphi_1), (U_2, \varphi_2), (U_3, \varphi_3)\}$  is an atlas on  $X$ . Deduce that  $X$  is a smooth manifold.

[You may assume that  $(U_1, \varphi_1), (U_2, \varphi_2), (U_3, \varphi_3)$  are charts.]

Need to show the  $(U_i, \varphi_i)$  are pairwise compatible, and cover  $X$ .

The transition function  $\varphi_2^{-1}\varphi_1$  maps

$$\varphi_2^{-1}\varphi_1 : \{(a_1, b_1, c_1, d_1) \in \mathbb{R}^4 : a_1 \neq 0\} \rightarrow \{(a_2, b_2, c_2, d_2) \in \mathbb{R}^4 : c_2 \neq 0\},$$

$$\varphi_2^{-1}\varphi_1 : (a_1, b_1, c_1, d_1) \mapsto \left(\frac{c_1}{a_1}, d_1 - \frac{b_1c_1}{a_1}, \frac{1}{a_1}, \frac{-b_1}{a_1}\right), \quad (1)$$

$$\text{as } y = a_1x + b_1, z = c_1x + d_1 \Leftrightarrow z = \frac{c_1}{a_1}y + \left(d_1 - \frac{b_1c_1}{a_1}\right), x = \frac{1}{a_1}y - \frac{b_1}{a_1}.$$

This is smooth, with smooth inverse

$$\varphi_1^{-1}\varphi_2 : \{(a_2, b_2, c_2, d_2) \in \mathbb{R}^4 : c_2 \neq 0\} \rightarrow \{(a_1, b_1, c_1, d_1) \in \mathbb{R}^4 : a_1 \neq 0\},$$

$$\varphi_1^{-1}\varphi_2 : (a_2, b_2, c_2, d_2) \mapsto \left(-\frac{1}{c_2}, -\frac{d_2}{c_2}, \frac{a_2}{c_2}, b_2 - \frac{a_2 d_2}{c_2}\right).$$

Hence  $(U_1, \varphi_1)$  and  $(U_2, \varphi_2)$  are compatible.

Similarly  $(U_2, \varphi_2), (U_3, \varphi_3)$  and  $(U_3, \varphi_3), (U_1, \varphi_1)$  are compatible, by cyclic permutation of 1, 2, 3 and  $x, y, z$ .

A line in  $\mathbb{R}^3$  lies in  $\varphi_1(U_1), \varphi_2(U_2), \varphi_3(U_3)$  if it is not parallel to the  $(y, z)$  plane, or  $(x, z)$  plane, or  $(x, y)$  plane, respectively. As no line is parallel to all three,

$$X = \varphi_1(U_1) \cup \varphi_2(U_2) \cup \varphi_3(U_3).$$

Hence  $\{(U_1, \varphi_1), (U_2, \varphi_2), (U_3, \varphi_3)\}$  is an atlas on  $X$ . It is contained in a unique maximal atlas.

We know  $X$  is Hausdorff and second countable by (b). Hence  $X$  is a smooth manifold.

(d)[6 marks] Prove that  $X$  is orientable.

[Hint: prove the transition functions are orientation-preserving.]

Differentiate  $\varphi_2^{-1}\varphi_1$  in (1) at  $(a_1, b_1, c_1, d_1)$ . It acts with matrix

$$D(\varphi_2^{-1}\varphi_1) = \begin{pmatrix} -\frac{c_1}{a_1^2} & 0 & \frac{1}{a_1} & 0 \\ \frac{b_1 c_1}{a_1^2} & -\frac{c_1}{a_1} & -\frac{b_1}{a_1} & 1 \\ -\frac{1}{a_1^2} & 0 & 0 & 0 \\ \frac{b_1}{a_1^2} & -\frac{1}{a_1} & 0 & 0 \end{pmatrix}$$

This has determinant  $\frac{1}{a_1^4}$ , as the only nonzero term comes from the product of the four red terms.

As  $D(\varphi_2^{-1}\varphi_1)$  has positive determinant everywhere,  $\varphi_2^{-1}\varphi_1$  is orientation-preserving. Similarly,  $\varphi_3^{-1}\varphi_2$  and  $\varphi_1^{-1}\varphi_3$  are orientation-preserving, by cyclic permutation of 1, 2, 3 and  $x, y, z$ . Hence  $\{(U_1, \varphi_1), (U_2, \varphi_2), (U_3, \varphi_3)\}$  is an oriented atlas, and defines an orientation on  $X$ .

Note: we have several different ways to define orientations:

- as an orientation on  $T_x X$  for  $x \in X$ , varying continuously with  $x$ .
- as an equivalence class  $[\omega]$  of non-vanishing  $n$ -forms  $\omega$  on  $X$ .
- as an atlas with orientation-preserving transition functions.

You can use any of these you like. This question uses the last.

(e)[3 marks] Now let  $Y$  be the set of (unoriented) affine real lines in  $\mathbb{R}^2$ , made into a manifold in a similar way. Is  $Y$  orientable? Give brief justification.

No,  $Y$  is not orientable, as it is topologically the Möbius strip, or equivalently  $\mathbb{RP}^2 \setminus \{[1, 0, 0]\}$ . (Space of projective lines in  $\mathbb{RP}^2$  is  $\mathbb{RP}^2$ .) [You can repeat the above calculations with two charts

$$\varphi_1 : (a_1, b_1) \mapsto \{(x, y) \in \mathbb{R}^2 : y = a_1x + b_1\},$$

$$\varphi_2 : (a_2, b_2) \mapsto \{(x, y) \in \mathbb{R}^2 : x = a_2y + b_2\}.$$

The transition function  $\varphi_2^{-1}\varphi_1$  maps

$$\varphi_2^{-1}\varphi_1 : \{(a_1, b_1) \in \mathbb{R}^2 : a_1 \neq 0\} \rightarrow \{(a_2, b_2) \in \mathbb{R}^2 : a_2 \neq 0\},$$

$$\varphi_2^{-1}\varphi_1 : (a_1, b_1) \mapsto \left(\frac{1}{a_1}, -\frac{b_1}{a_1}\right).$$

We have  $\det D(\varphi_2^{-1}\varphi_1) = \frac{1}{a_1^3}$ , which changes sign at  $a_1 = 0$  and is not orientation-preserving. This in itself doesn't prove  $Y$  not orientable, but going round the circle  $b_1 = b_2 = 0$  in  $Y$ , you cross  $a_1 = 0$  once, so orientations change sign around the circle.

**This much detail not required.]**

(a)[11 marks] (i) Let  $X$  be a manifold and  $v \in \Gamma^\infty(TX)$  a vector field on  $X$ . Define the *maximal integral curve* of  $v$  through a point  $x \in X$ . What is the domain of a maximal integral curve if  $X$  is compact?

(ii) Define *1-parameter groups of diffeomorphisms*  $\varphi : \mathbb{R} \times X \rightarrow X$ . In the case in which  $X$  is compact, describe the 1-1 correspondence between vector fields  $v$  and 1-parameter groups of diffeomorphisms  $\varphi$ , in terms of maximal integral curves.

(iii) If  $v$  is a vector field and  $\alpha$  a tensor on  $X$ , define the *Lie derivative*  $\mathcal{L}_v \alpha$ .

[You may assume the 1-1 correspondence in (ii) applies to  $v$ .]

All bookwork.

For (iii), define  $\mathcal{L}_v \alpha = \frac{d}{dt}(\varphi_t^*(\alpha))|_{t=0}$ . If  $X$  is not compact then  $\varphi_t$  may not be defined if  $v$  is not complete – a ‘local’ definition is possible – but the question allows you to assume  $\varphi_t$  makes sense.

On  $\mathbb{R}^3$  with coordinates  $(x_1, x_2, x_3)$ , define vector fields

$$u = x_1 \frac{\partial}{\partial x_1} + x_2 \frac{\partial}{\partial x_2} + x_3 \frac{\partial}{\partial x_3}, \quad v = x_1^2 \frac{\partial}{\partial x_1} + x_2^2 \frac{\partial}{\partial x_2} + x_3^2 \frac{\partial}{\partial x_3}.$$

(b)[4 marks] Find the maximal integral curves of  $u, v$  through each  $(x_1, x_2, x_3) \in \mathbb{R}^3$ .

Write  $\gamma(t) = (\gamma_1(t), \gamma_2(t), \gamma_3(t))$ . For  $\gamma$  to be a flow-line of  $u$ , need

$$\dot{\gamma}_1 = \gamma_1, \quad \dot{\gamma}_2 = \gamma_2, \quad \dot{\gamma}_3 = \gamma_3,$$

so  $\gamma_i(t) = x_i e^t$ . Domain of maximal integral curve is  $\mathbb{R}$ .

For  $\gamma$  to be a flow-line of  $v$ , need

$$\dot{\gamma}_1 = \gamma_1^2, \quad \dot{\gamma}_2 = \gamma_2^2, \quad \dot{\gamma}_3 = \gamma_3^2,$$

so  $\int \frac{d\gamma_i}{\gamma_i^2} = \int dt$ , and  $-\frac{1}{\gamma_i} = t - \frac{1}{x_i}$ , giving  $\gamma_i(t) = \frac{x_i}{1-x_it}$ .

The domain of the maximal integral curve is  $(a, b)$ , where

$$a = \begin{cases} -\infty, & \text{all } x_i \geq 0, \\ \max(\frac{1}{x_i} : x_i < 0), & \text{otherwise,} \end{cases}$$
$$b = \begin{cases} \infty, & \text{all } x_i \leq 0, \\ \min(\frac{1}{x_i} : x_i > 0), & \text{otherwise.} \end{cases}$$

(c)[6 marks] Prove that the only 2-form  $\alpha$  on  $\mathbb{R}^3$  with  $\mathcal{L}_u\alpha = 0$  is  $\alpha = 0$ .

[Well known formulae may be used if clearly stated.]

Write  $\alpha = \alpha_1 dx_2 \wedge dx_3 + \alpha_2 dx_3 \wedge dx_1 + \alpha_3 dx_1 \wedge dx_2$  for  $\alpha_i : \mathbb{R}^3 \rightarrow \mathbb{R}$  smooth. **Cartan's formula:**  $\mathcal{L}_u\alpha = i_u(d\alpha) + d(i_u\alpha)$ . So

$$\begin{aligned}\mathcal{L}_u\alpha &= i_u\left[\left(\frac{\partial\alpha_1}{\partial x_1} + \frac{\partial\alpha_2}{\partial x_2} + \frac{\partial\alpha_3}{\partial x_3}\right)dx_1 \wedge dx_2 \wedge dx_3\right] \\ &+ d\left[\alpha_1 x_2 dx_3 - \alpha_1 x_3 dx_2 + \alpha_2 x_3 dx_1 - \alpha_2 x_1 dx_3 + \alpha_3 x_1 dx_2 - \alpha_3 x_2 dx_1\right] \\ &= \left(\frac{\partial\alpha_1}{\partial x_1} + \frac{\partial\alpha_2}{\partial x_2} + \frac{\partial\alpha_3}{\partial x_3}\right)(x_1 dx_2 \wedge dx_3 + x_2 dx_3 \wedge dx_1 + x_3 dx_1 \wedge dx_2) \\ &+ \left(\frac{\partial\alpha_1}{\partial x_2} x_2 + \frac{\partial\alpha_1}{\partial x_3} x_3 + 2\alpha_1 - \frac{\partial\alpha_2}{\partial x_2} x_1 - \frac{\partial\alpha_3}{\partial x_3} x_1\right)dx_2 \wedge dx_3 \\ &+ \left(\frac{\partial\alpha_2}{\partial x_3} x_3 + \frac{\partial\alpha_2}{\partial x_1} x_1 + 2\alpha_2 - \frac{\partial\alpha_3}{\partial x_3} x_2 - \frac{\partial\alpha_1}{\partial x_1} x_2\right)dx_3 \wedge dx_1 \\ &+ \left(\frac{\partial\alpha_3}{\partial x_1} x_1 + \frac{\partial\alpha_3}{\partial x_2} x_2 + 2\alpha_3 - \frac{\partial\alpha_1}{\partial x_1} x_3 - \frac{\partial\alpha_2}{\partial x_2} x_3\right)dx_1 \wedge dx_2 \\ &= \left(\frac{\partial\alpha_1}{\partial x_1} x_1 + \frac{\partial\alpha_1}{\partial x_2} x_2 + \frac{\partial\alpha_1}{\partial x_3} x_3 + 2\alpha_1\right)dx_2 \wedge dx_3 \\ &+ \left(\frac{\partial\alpha_2}{\partial x_1} x_1 + \frac{\partial\alpha_2}{\partial x_2} x_2 + \frac{\partial\alpha_2}{\partial x_3} x_3 + 2\alpha_2\right)dx_3 \wedge dx_1 \\ &+ \left(\frac{\partial\alpha_3}{\partial x_1} x_1 + \frac{\partial\alpha_3}{\partial x_2} x_2 + \frac{\partial\alpha_3}{\partial x_3} x_3 + 2\alpha_3\right)dx_1 \wedge dx_2.\end{aligned}$$

Thus  $\mathcal{L}_u \alpha = 0$  provided

$$\frac{\partial \alpha_i}{\partial x_1} x_1 + \frac{\partial \alpha_i}{\partial x_2} x_2 + \frac{\partial \alpha_i}{\partial x_3} x_3 + 2\alpha_i = 0, \quad i = 1, 2, 3.$$

Here is the tricky part:

Along the ray  $(tx_1, tx_2, tx_3)$  for  $t \in \mathbb{R}$  this gives

$$t \frac{d}{dt} (\alpha_i(tx_1, tx_2, tx_3)) + 2\alpha_i(tx_1, tx_2, tx_3) = 0,$$

with solution  $\alpha_i(tx_1, tx_2, tx_3) = Ct^{-2}$ .

But this is only continuous at  $t = 0$  if  $C = 0$ , so when  $t = 1$ ,  $\alpha_i(x_1, x_2, x_3) = 0$ . Thus  $\alpha = 0$ .

(d)[4 marks] Find all vector fields  $w$  on  $\mathbb{R}^3$  with  $\mathcal{L}_u w = 0$ , that is,  $[u, w] = 0$ .

[Well known formulae may be used if clearly stated.]

Write  $u = u_1 \frac{\partial}{\partial x_1} + u_2 \frac{\partial}{\partial x_2} + u_3 \frac{\partial}{\partial x_3}$  and  $w = w_1 \frac{\partial}{\partial x_1} + w_2 \frac{\partial}{\partial x_2} + w_3 \frac{\partial}{\partial x_3}$ .

Then  $[u, w] = \sum_{i,j=1}^3 (u_i \frac{\partial w_j}{\partial x_i} - w_i \frac{\partial u_j}{\partial x_i}) \frac{\partial}{\partial x_j}$ . Learn this.

As  $u_i = x_i$  we see that  $[u, w] = 0$  iff

$$\frac{\partial w_i}{\partial x_1} x_1 + \frac{\partial w_i}{\partial x_2} x_2 + \frac{\partial w_i}{\partial x_3} x_3 - w_i = 0, \quad i = 1, 2, 3.$$

(Another tricky part.) Along the ray  $(tx_1, tx_2, tx_3)$  for  $t \in \mathbb{R}$  this gives

$$t \frac{d}{dt} (w_i(tx_1, tx_2, tx_3)) - w_i(tx_1, tx_2, tx_3) = 0,$$

with solution  $w_i(tx_1, tx_2, tx_3) = Ct$ . Thus  $w_i$  is linear along each ray in  $\mathbb{R}^3$ . For  $w_i$  to be smooth at  $(0, 0, 0)$ , this forces  $w_i$  to be linear,  $w_i = \sum_{j=1}^3 a_{ij} x_j$ . So the vector fields  $w$  with  $\mathcal{L}_u w = 0$  are  $w = \sum_{i,j=1}^3 a_{ij} x_j \frac{\partial}{\partial x_i}$  for real matrices  $(a_{ij})_{i,j=1}^3$ .

(a)[6 marks] Define the *de Rham cohomology groups*  $H^k(X)$  of an  $n$ -manifold  $X$ . Show that if  $X$  is compact and oriented then there is a well-defined, surjective linear map  $\Phi : H^n(X) \rightarrow \mathbb{R}$  with  $\Phi([\omega]) = \int_X \omega$ .

[*Standard results about integration of exterior forms may be used if clearly stated.*]

In the rest of the question you may assume that  $\Phi$  is an isomorphism if  $X$  is connected.

All bookwork.

To show  $\Phi$  is surjective, make an  $n$ -form  $\omega$  with nonzero integral, supported in a small coordinate ball, using a 'bump function'.

(b)[5 marks] Let  $f : X \rightarrow Y$  be a smooth map between compact, connected, oriented  $n$ -manifolds  $X, Y$ . Define the *degree*  $\deg f$  of  $f$ , using de Rham cohomology. State an alternative definition in terms of preimages of points (you need not prove they are equivalent).

All bookwork.

(c)[9 marks] Show that the cohomology of  $X = \mathcal{S}^2 \times \mathcal{S}^2$  may be written

$$H^0(X) = \langle 1_X \rangle_{\mathbb{R}}, \quad H^1(X) = 0, \quad H^2(X) = \langle \alpha_1, \alpha_2 \rangle_{\mathbb{R}},$$

$$H^3(X) = 0, \quad H^4(X) = \langle \alpha_1 \cup \alpha_2 \rangle_{\mathbb{R}},$$

where  $\alpha_1 \cup \alpha_1 = 0$ ,  $\alpha_2 \cup \alpha_2 = 0$ , and  $\int_X \alpha_1 \cup \alpha_2 = 1$ .

[You may assume the Künneth Theorem, and a formula for  $H^k(\mathcal{S}^2)$ .]

Quote:  $H^0(\mathcal{S}^2) \cong H^2(\mathcal{S}^2) \cong \mathbb{R}$ ,  $H^1(\mathcal{S}^2) = 0$ .

Künneth Theorem:  $H^k(X \times Y) \cong \bigoplus_{i+j=k} H^i(X) \otimes H^j(Y)$ , where the  $H^i(X) \otimes H^j(Y)$  factor is the image of  $\pi_X^*(H^i(X)) \cup \pi_Y^*(H^j(Y))$ .

Write  $H^0(\mathcal{S}^2) = \langle 1 \rangle_{\mathbb{R}}$  and  $H^2(\mathcal{S}^2) = \langle \omega \rangle_{\mathbb{R}}$  with  $\int_{\mathcal{S}^2} \omega = 1$ . Write  $\pi_1, \pi_2 : \mathcal{S}^2 \times \mathcal{S}^2 \rightarrow \mathcal{S}^2$  for the projections to first and second factors.

Künneth says that  $H^0(X) = \langle \pi_1^*(1) \cup \pi_2^*(1) \rangle_{\mathbb{R}} = \langle 1 \rangle_{\mathbb{R}}$ ,  $H^1(X) = 0$ ,  $H^2(X) = \langle \pi_1^*(1) \cup \pi_2^*(\omega) \rangle_{\mathbb{R}} \oplus \langle \pi_1^*(\omega) \cup \pi_2^*(1) \rangle_{\mathbb{R}} = \langle \pi_2^*(\omega), \pi_1^*(\omega) \rangle_{\mathbb{R}}$ ,  $H^3(X) = 0$ , and  $H^4(X) = \langle \pi_1^*(\omega) \cup \pi_2^*(\omega) \rangle_{\mathbb{R}}$ .

Set  $\alpha_j = \pi_j^*(\omega)$ . Then  $H^2(X) = \langle \alpha_1, \alpha_2 \rangle_{\mathbb{R}}$ ,  $H^4(X) = \langle \alpha_1 \cup \alpha_2 \rangle_{\mathbb{R}}$  as we want. Also  $\alpha_1 \cup \alpha_1 = \pi_1^*(\omega) \cup \pi_1^*(\omega) = \pi_1^*(\omega \cup \omega) = 0$ , as  $\omega \cup \omega \in H^4(\mathcal{S}^2) = 0$ . Similarly  $\alpha_2 \cup \alpha_2 = 0$ . And

$$\int_X \alpha_1 \cup \alpha_2 = \int_{\mathcal{S}^2 \times \mathcal{S}^2} \pi_1^*(\omega) \cup \pi_2^*(\omega) = \left( \int_{\mathcal{S}^2} \omega \right) \cdot \left( \int_{\mathcal{S}^2} \omega \right) = 1 \cdot 1 = 1.$$

(d)[5 marks] The cohomology of the compact oriented 4-manifold  $Y = \mathbb{C}P^2$  may be written

$$H^0(Y) = \langle 1_Y \rangle_{\mathbb{R}}, \quad H^1(Y) = 0, \quad H^2(Y) = \langle \beta \rangle_{\mathbb{R}},$$

$$H^3(Y) = 0, \quad H^4(Y) = \langle \beta \cup \beta \rangle_{\mathbb{R}}, \quad \text{where} \quad \int_Y \beta \cup \beta = 1.$$

Show that any smooth map  $f : Y \rightarrow X$ , with  $X$  defined as in (c), has degree  $\deg f = 0$ .

Write  $f^*(\alpha_i) = a_i \beta$  for  $i = 1, 2$ . Then  $f^*(\alpha_i \cup \alpha_i) = a_i^2 \beta \cup \beta$ . But  $\alpha_i \cup \alpha_i = 0$  and  $\beta \cup \beta \neq 0$ , so  $a_i^2 = 0$ , and  $a_i = 0$ .

Hence  $f^*(\alpha_1 \cup \alpha_2) = a_1 a_2 \beta \cup \beta = 0$ . The commuting diagram

$$\begin{array}{ccc} H^4(X) = \langle \alpha_1 \cup \alpha_2 \rangle_{\mathbb{R}} & \xrightarrow{f^*} & H^4(Y) = \langle \beta^2 \rangle_{\mathbb{R}} \\ \cong \downarrow [\lambda] \mapsto \int_X \lambda & & [\lambda] \mapsto \int_Y \lambda \downarrow \cong \\ \mathbb{R} & \xrightarrow{\cdot \deg f} & \mathbb{R} \end{array}$$

now shows that  $\deg f = 0$ .