Numerical Solution of Partial Differential Equations

Endre Süli

Mathematical Institute
University of Oxford
2026

Lecture 1

Elements of function spaces

The accuracy of a numerical method for the approximate solution of PDEs depends on its ability to capture the important features of the analytical solution. One such feature is the smoothness of the solution, which depends on the smoothness of the data.

Elements of function spaces

The accuracy of a numerical method for the approximate solution of PDEs depends on its ability to capture the important features of the analytical solution. One such feature is the smoothness of the solution, which depends on the smoothness of the data.

Precise assumptions about the data and of the corresponding solution can be conveniently stated in terms of function spaces.

Elements of function spaces

The accuracy of a numerical method for the approximate solution of PDEs depends on its ability to capture the important features of the analytical solution. One such feature is the smoothness of the solution, which depends on the smoothness of the data.

Precise assumptions about the data and of the corresponding solution can be conveniently stated in terms of function spaces.

We present a brief overview of definitions and basic results form the theory of function spaces, focusing in particular on spaces of:

Elements of function spaces

The accuracy of a numerical method for the approximate solution of PDEs depends on its ability to capture the important features of the analytical solution. One such feature is the smoothness of the solution, which depends on the smoothness of the data.

Precise assumptions about the data and of the corresponding solution can be conveniently stated in terms of function spaces.

We present a brief overview of definitions and basic results form the theory of function spaces, focusing in particular on spaces of:

- Continuous functions;
- Integrable functions; and
- Sobolev spaces.

Spaces of continuous functions

$\mathbb{N}$ denotes the set of nonnegative integers.

An *n*-tuple $\alpha = (\alpha_1, \ldots, \alpha_n) \in \mathbb{N}^n$ is called a *multi-index*.
The nonnegative integer $|\alpha| := \alpha_1 + \cdots + \alpha_n$ is called the *length* of the multi-index $\alpha = (\alpha_1, \ldots, \alpha_n)$. We denote $(0, \ldots, 0)$ by $\mathbf{0}$; clearly $|\mathbf{0}| = 0$.

Spaces of continuous functions

$\mathbb{N}$ denotes the set of nonnegative integers.

An n-tuple $\alpha = (\alpha_1, \ldots, \alpha_n) \in \mathbb{N}^n$ is called a *multi-index*.
The nonnegative integer $|\alpha| := \alpha_1 + \cdots + \alpha_n$ is called the *length* of the multi-index $\alpha = (\alpha_1, \ldots, \alpha_n)$. We denote $(0, \ldots, 0)$ by $\mathbf{0}$; clearly $|\mathbf{0}| = 0$.

Let

$$D^\alpha := \left(\frac{\partial}{\partial x_1}\right)^{\alpha_1} \cdots \left(\frac{\partial}{\partial x_n}\right)^{\alpha_n} = \frac{\partial^{|\alpha|}}{\partial x_1^{\alpha_1} \cdots \partial x_n^{\alpha_n}}.$$

Spaces of continuous functions

$\mathbb{N}$ denotes the set of nonnegative integers.

An n-tuple $\alpha = (\alpha_1, \ldots, \alpha_n) \in \mathbb{N}^n$ is called a *multi-index*.
The nonnegative integer $|\alpha| := \alpha_1 + \cdots + \alpha_n$ is called the *length* of the multi-index $\alpha = (\alpha_1, \ldots, \alpha_n)$. We denote $(0, \ldots, 0)$ by $\mathbf{0}$; clearly $|\mathbf{0}| = 0$.

Let

$$D^\alpha := \left(\frac{\partial}{\partial x_1}\right)^{\alpha_1} \cdots \left(\frac{\partial}{\partial x_n}\right)^{\alpha_n} = \frac{\partial^{|\alpha|}}{\partial x_1^{\alpha_1} \cdots \partial x_n^{\alpha_n}}.$$

EXAMPLE. Suppose that $n = 3$ and $\alpha = (\alpha_1, \alpha_2, \alpha_3)$, $\alpha_j \in \mathbb{N}$, $j = 1, 2, 3$. Then, for u, a function of three variables x_1, x_2, x_3:

$$\sum_{|\alpha|=3} D^\alpha u = \frac{\partial^3 u}{\partial x_1^3} + \frac{\partial^3 u}{\partial x_1^2 \partial x_2} + \frac{\partial^3 u}{\partial x_1^2 \partial x_3} + \frac{\partial^3 u}{\partial x_1 \partial x_2^2} + \frac{\partial^3 u}{\partial x_1 \partial x_3^2}$$
$$+ \frac{\partial^3 u}{\partial x_3^2} + \frac{\partial^3 u}{\partial x_1 \partial x_2 \partial x_3} + \frac{\partial^3 u}{\partial x_2^2 \partial x_3} + \frac{\partial^3 u}{\partial x_2 \partial x_3^2} + \frac{\partial^3 u}{\partial x_3^3}.$$

We shall frequently write ∂_{x_j} instead of $\frac{\partial}{\partial x_j}$. ◇

Let Ω be an open set in $\mathbb{R}^n$, and let $k \in \mathbb{N}$.

Let Ω be an open set in $\mathbb{R}^n$, and let $k \in \mathbb{N}$.

We denote by $C^k(\Omega)$ the set of all continuous real-valued functions defined on Ω s.t. $D^\alpha u$ is continuous on Ω for all $\alpha = (\alpha_1, \ldots, \alpha_n)$ with $|\alpha| \leq k$.

Let Ω be an open set in $\mathbb{R}^n$, and let $k \in \mathbb{N}$.

We denote by $C^k(\Omega)$ the set of all continuous real-valued functions defined on Ω s.t. $D^\alpha u$ is continuous on Ω for all $\alpha = (\alpha_1, \ldots, \alpha_n)$ with $|\alpha| \leq k$.

Assuming that Ω is a *bounded* open set, $C^k(\overline{\Omega})$ will denote the set of all u in $C^k(\Omega)$ s.t. $D^\alpha u$ can be extended from Ω to a continuous function on $\overline{\Omega}$, the closure of the set Ω, for all $\alpha = (\alpha_1, \ldots, \alpha_n)$ with $|\alpha| \leq k$.

Let Ω be an open set in $\mathbb{R}^n$, and let $k \in \mathbb{N}$.

We denote by $C^k(\Omega)$ the set of all continuous real-valued functions defined on Ω s.t. $D^\alpha u$ is continuous on Ω for all $\alpha = (\alpha_1, \ldots, \alpha_n)$ with $|\alpha| \leq k$.

Assuming that Ω is a *bounded* open set, $C^k(\overline{\Omega})$ will denote the set of all u in $C^k(\Omega)$ s.t. $D^\alpha u$ can be extended from Ω to a continuous function on $\overline{\Omega}$, the closure of the set Ω, for all $\alpha = (\alpha_1, \ldots, \alpha_n)$ with $|\alpha| \leq k$.

The linear space $C^k(\overline{\Omega})$ can then be equipped with the norm

$$\|u\|_{C^k(\overline{\Omega})} := \sum_{|\alpha| \leq k} \sup_{x \in \Omega} \left| D^\alpha u(x) \right| .$$

Let Ω be an open set in $\mathbb{R}^n$, and let $k \in \mathbb{N}$.

We denote by $C^k(\Omega)$ the set of all continuous real-valued functions defined on Ω s.t. $D^\alpha u$ is continuous on Ω for all $\alpha = (\alpha_1, \ldots, \alpha_n)$ with $|\alpha| \leq k$.

Assuming that Ω is a *bounded* open set, $C^k(\overline{\Omega})$ will denote the set of all u in $C^k(\Omega)$ s.t. $D^\alpha u$ can be extended from Ω to a continuous function on $\overline{\Omega}$, the closure of the set Ω, for all $\alpha = (\alpha_1, \ldots, \alpha_n)$ with $|\alpha| \leq k$.

The linear space $C^k(\overline{\Omega})$ can then be equipped with the norm

$$\|u\|_{C^k(\overline{\Omega})} := \sum_{|\alpha| \leq k} \sup_{x \in \Omega} \left| D^\alpha u(x) \right| .$$

Note: When $k = 0$, we shall write $C(\overline{\Omega})$ instead of $C^0(\overline{\Omega})$.

The *support*, supp u, of a continuous function u on Ω is defined as the closure in Ω of the set

$$\{x \in \Omega : u(x) \neq 0\}.$$

In other words, supp u is the smallest closed subset of Ω such that $u = 0$ in $\Omega \backslash \text{supp } u$.

EXAMPLE. Let w be the function defined on $\mathbb{R}^n$ by

$$w(x) = \begin{cases} \mathrm{e}^{-\frac{1}{1-|x|^2}} & , \ |x| < 1, \\ 0, & \text{otherwise;} \end{cases}$$

here $|x| := (x_1^2 + \cdots + x_n^2)^{1/2}$ for $x \in \mathbb{R}^n$.

Clearly, supp w is the closed unit ball $\{x \in \mathbb{R}^n : |x| \leq 1\}$. $\diamond$

We denote by $C_0^k(\Omega)$ the set of all $u \in C^k(\Omega)$ such that supp $u \subset \Omega$ and supp u is bounded. Let

$$C_0^\infty(\Omega) = \bigcap_{k \geq 0} C_0^k(\Omega).$$

EXAMPLE.
The function w defined in the previous example belongs to $C_0^\infty(\mathbb{R}^n)$. $\diamond$

Spaces of integrable functions

Let p be a real number, $p \geq 1$; we denote by $L_p(\Omega)$ the set of all real-valued functions defined on Ω such that

$$\int_\Omega \left|u(x)\right|^p \, \mathrm{d}x < \infty.$$

Spaces of integrable functions

Let p be a real number, $p \geq 1$; we denote by $L_p(\Omega)$ the set of all real-valued functions defined on Ω such that

$$\int_\Omega \left|u(x)\right|^p \, \mathrm{d}x < \infty.$$

Functions which are equal almost everywhere (i.e., equal, except on a set of measure zero) on Ω are identified with each other.

A subset of $\mathbb{R}^n$ is said to be a *set of measure zero* if it can be contained in the union of countably many open balls of arbitrarily small total volume.

Spaces of integrable functions

Let p be a real number, $p \geq 1$; we denote by $L_p(\Omega)$ the set of all real-valued functions defined on Ω such that

$$\int_\Omega \left|u(x)\right|^p \, \mathrm{d}x < \infty.$$

Functions which are equal almost everywhere (i.e., equal, except on a set of measure zero) on Ω are identified with each other.

A subset of $\mathbb{R}^n$ is said to be a *set of measure zero* if it can be contained in the union of countably many open balls of arbitrarily small total volume.

$L_p(\Omega)$ is equipped with the norm

$$\|u\|_{L_p(\Omega)} := \left(\int_\Omega \left|u(x)\right|^p \, \mathrm{d}x \right)^{1/p}.$$

A particularly important case is $p = 2$; then,

$$\|u\|_{L_2(\Omega)} = \left(\int_\Omega \left| u(x) \right|^2 \, \mathrm{d}x \right)^{1/2}.$$

The space $L_2(\Omega)$ can be equipped with an inner product

$$(u, v) := \int_\Omega u(x) v(x) \, \mathrm{d}x.$$

Clearly $\|u\|_{L_2(\Omega)} = (u, u)^{1/2}$.

A particularly important case is $p = 2$; then,

$$\|u\|_{L_2(\Omega)} = \left(\int_\Omega \left| u(x) \right|^2 \, \mathrm{d}x \right)^{1/2}.$$

The space $L_2(\Omega)$ can be equipped with an inner product

$$(u, v) := \int_\Omega u(x) v(x) \, \mathrm{d}x.$$

Clearly $\|u\|_{L_2(\Omega)} = (u, u)^{1/2}$.

Lemma (The Cauchy–Schwarz inequality)

Let $u, v \in L_2(\Omega)$; then

$$\left| (u, v) \right| \leq \|u\|_{L_2(\Omega)} \|v\|_{L_2(\Omega)}.$$

Sobolev spaces

Suppose that u is locally integrable on Ω (i.e. $u \in L_1(\omega)$ for each bounded open set ω, with $\overline{\omega} \subset \Omega$).

Sobolev spaces

Suppose that u is locally integrable on Ω (i.e. $u \in L_1(\omega)$ for each bounded open set ω, with $\overline{\omega} \subset \Omega$). Suppose also that there exists a function w_α, locally integrable on Ω, and such that

$$\int_\Omega w_\alpha(x)\, v(x)\, \mathrm{d}x = (-1)^{|\alpha|} \int_\Omega u(x)\, D^\alpha v(x) \qquad \forall\, v \in C_0^\infty(\Omega).$$

Then w_α is called the *weak derivative* of u (of order $|\alpha| = \alpha_1 + \cdots + \alpha_n$) and we write $w_\alpha = D^\alpha u$.

Sobolev spaces

Suppose that u is locally integrable on Ω (i.e. $u \in L_1(\omega)$ for each bounded open set ω, with $\overline{\omega} \subset \Omega$). Suppose also that there exists a function w_α, locally integrable on Ω, and such that

$$\int_\Omega w_\alpha(x)\, v(x)\, \mathrm{d}x = (-1)^{|\alpha|} \int_\Omega u(x)\, D^\alpha v(x) \qquad \forall\, v \in C_0^\infty(\Omega).$$

Then w_α is called the *weak derivative* of u (of order $|\alpha| = \alpha_1 + \cdots + \alpha_n$) and we write $w_\alpha = D^\alpha u$.

Clearly, if u is a smooth function then its weak derivatives coincide with those in the classical (pointwise) sense.

EXAMPLE Let $\Omega = \mathbb{R}^1$, and let $u(x) = (1 - |x|)_+$. Here, for a real number y, $y_+ := \max\{y, 0\}$.

EXAMPLE Let $\Omega = \mathbb{R}^1$, and let $u(x) = (1 - |x|)_+$. Here, for a real number y, $y_+ := \max\{y, 0\}$. Clearly u is not differentiable at $x = 0, \pm 1$.

EXAMPLE Let $\Omega = \mathbb{R}^1$, and let $u(x) = (1 - |x|)_+$. Here, for a real number y, $y_+ := \max\{y, 0\}$. Clearly u is not differentiable at $x = 0, \pm 1$. However, because u is locally integrable on Ω it may still have a weak derivative.

EXAMPLE Let $\Omega = \mathbb{R}^1$, and let $u(x) = (1 - |x|)_+$. Here, for a real number y, $y_+ := \max\{y, 0\}$. Clearly u is not differentiable at $x = 0, \pm 1$. However, because u is locally integrable on Ω it may still have a weak derivative. Indeed, for any $v \in C_0^\infty(\Omega)$:

$$\begin{aligned}\int_{-\infty}^{+\infty} u(x)\, v'(x)\, \mathrm{d}x &= \int_{-\infty}^{+\infty} (1 - |x|)_+\, v'(x)\, \mathrm{d}x = \int_{-1}^{1} (1 - |x|)\, v'(x)\, \mathrm{d}x \\ &= \int_{-1}^{0} (1 + x)\, v'(x)\, \mathrm{d}x + \int_{0}^{1} (1 - x)\, v'(x)\, \mathrm{d}x \\ &= \int_{-1}^{0} (-1)\, v(x)\, \mathrm{d}x + \int_{0}^{1} (+1)\, v(x)\, \mathrm{d}x \\ &= -\int_{-\infty}^{+\infty} w(x)\, v(x)\, \mathrm{d}x,\end{aligned}$$

where

$$w(x) = \begin{cases} 0, & x < -1, \\ 1, & x \in (-1, 0), \\ -1, & x \in (0, 1), \\ 0, & x > 1. \end{cases}$$

Thus, $w = u' = Du$ $\diamond$

Let k be a nonnegative integer. We define (with D^α denoting a weak derivative of order $|\alpha|$)

$$H^k(\Omega) := \{u \in L_2(\Omega) : D^\alpha u \in L_2(\Omega), \quad |\alpha| \leq k\}.$$

$H^k(\Omega)$ is called a Sobolev space of order k; it is equipped with the (Sobolev) norm and semi-norm:

$$\|u\|_{H^k(\Omega)} := \left(\sum_{|\alpha| \leq k} \|D^\alpha u\|^2_{L_2(\Omega)} \right)^{1/2}, \qquad |u|_{H^k(\Omega)} := \left(\sum_{|\alpha| = k} \|D^\alpha u\|^2_{L_2(\Omega)} \right)^{1/2}$$

and the inner product

$$(u, v)_{H^k(\Omega)} := \sum_{|\alpha| \leq k} (D^\alpha u, D^\alpha v).$$

EXAMPLE

$$H^0(\Omega) = L_2(\Omega).$$

$$H^1(\Omega) := \left\{ u \in L_2(\Omega) : \partial_{x_j} u := \frac{\partial u}{\partial x_j} \in L_2(\Omega), \;\; j = 1, \ldots, n \right\},$$

$$\|u\|_{H^1(\Omega)} := \left\{ \|u\|^2_{L_2(\Omega)} + \sum_{j=1}^{n} \|\partial_{x_j} u\|^2_{L_2(\Omega)} \right\}^{1/2},$$

$$|u|_{H^1(\Omega)} := \left\{ \sum_{j=1}^{n} \|\partial_{x_j} u\|^2_{L_2(\Omega)} \right\}^{1/2}.$$

Similarly,

$$H^2(\Omega) := \left\{ u \in L_2(\Omega) : \partial_{x_j} u \in L_2(\Omega),\ \partial^2_{x_i x_j} u \in L_2(\Omega),\ i,j = 1,\dots,n \right\},$$

$$\|u\|_{H^2(\Omega)} := \left\{ \|u\|^2_{L_2(\Omega)} + \sum_{j=1}^{n} \|\partial_{x_j} u\|^2_{L_2(\Omega)} + \sum_{i,j=1}^{n} \|\partial^2_{x_i x_j} u\|^2_{L_2(\Omega)} \right\}^{1/2},$$

$$|u|_{H^2(\Omega)} := \left\{ \sum_{i,j=1}^{n} \|\partial^2_{x_i x_j} u\|^2_{L_2(\Omega)} \right\}^{1/2}.$$

We conclude with the following important result.

Lemma (Poincaré–Friedrichs inequality)

Suppose that Ω is a bounded open set in $\mathbb{R}^n$ (with a sufficiently smooth boundary $\partial\Omega$) and let $u \in H_0^1(\Omega)$; then, there exists a positive constant $c_\star(\Omega)$, independent of u, such that

$$\int_\Omega u^2(x)\,\mathrm{d}x \leq c_\star \sum_{i=1}^{n} \int_\Omega \left|\partial_{x_i} u(x)\right|^2 \,\mathrm{d}x. \tag{1}$$

PROOF. We shall prove this for the special case of a rectangular domain $\Omega = (a, b) \times (c, d)$ in $\mathbb{R}^2$. The proof for general Ω is analogous.

PROOF. We shall prove this for the special case of a rectangular domain $\Omega = (a, b) \times (c, d)$ in $\mathbb{R}^2$. The proof for general Ω is analogous. Evidently,

$$u(x, y) = u(a, y) + \int_a^x \partial_x u(\xi, y)\, \mathrm{d}\xi = \int_a^x \partial_x u(\xi, y)\, \mathrm{d}\xi, \qquad c < y < d.$$

PROOF. We shall prove this for the special case of a rectangular domain $\Omega = (a, b) \times (c, d)$ in $\mathbb{R}^2$. The proof for general Ω is analogous. Evidently,

$$u(x, y) = u(a, y) + \int_a^x \partial_x u(\xi, y)\, \mathrm{d}\xi = \int_a^x \partial_x u(\xi, y)\, \mathrm{d}\xi, \qquad c < y < d.$$

Thus, by the Cauchy–Schwarz inequality,

$$\begin{aligned}
\int_\Omega \left|u(x, y)\right|^2 \, \mathrm{d}x\, \mathrm{d}y &= \int_a^b \int_c^d \left| \int_a^x \partial_x u(\xi, y)\, \mathrm{d}\xi \right|^2 \, \mathrm{d}x\, \mathrm{d}y \\
&\leq \int_a^b \int_c^d (x - a) \left(\int_a^x \left|\partial_x u(\xi, y)\right|^2 \, \mathrm{d}\xi \right) \, \mathrm{d}x\, \mathrm{d}y \\
&\leq \int_a^b (x - a)\, \mathrm{d}x \left(\int_c^d \int_a^b \left|\partial_x u(\xi, y)\right|^2 \, \mathrm{d}\xi\, \mathrm{d}y \right) \\
&= \frac{1}{2}(b - a)^2 \int_\Omega \left|\partial_x u(x, y)\right|^2 \, \mathrm{d}x\, \mathrm{d}y.
\end{aligned}$$

Analogously,

$$\int_\Omega \left|u(x,y)\right|^2 \,\mathrm{d}x\,\mathrm{d}y \leq \frac{1}{2}(d-c)^2 \int_\Omega \left|\partial_y u(x,y)\right|^2 \,\mathrm{d}x\,\mathrm{d}y.$$

Analogously,

$$\int_\Omega \left|u(x,y)\right|^2 \,\mathrm{d}x\,\mathrm{d}y \le \frac{1}{2}(d-c)^2 \int_\Omega \left|\partial_y u(x,y)\right|^2 \,\mathrm{d}x\,\mathrm{d}y.$$

By adding the two inequalities, we obtain

$$\int_\Omega \left|u(x,y)\right|^2 \,\mathrm{d}x\,\mathrm{d}y \le c_\star \int_\Omega \left(\left|\partial_x u\right|^2 + \left|\partial_y u\right|^2\right) \,\mathrm{d}x\,\mathrm{d}y,$$

where $c_\star = \left(\frac{2}{(b-a)^2} + \frac{2}{(d-c)^2}\right)^{-1}$. □