Numerical Solution of Partial Differential Equations

Endre Süli

Mathematical Institute
University of Oxford
2026

Lecture 13

The implicit scheme: stability, consistency and convergence

For $M \geq 2$, we define $\Delta t := T/M$, and for $J \geq 2$ the spatial step is taken to be $\Delta x := (b-a)/J$. We let $x_j := a + j\Delta x$ for $j = 0, 1, \ldots, J$ and $t_m := m\Delta t$ for $m = 0, 1, \ldots, M$.

On the space-time mesh $\{(x_j, t_m) : 0 \leq j \leq J,\ 0 \leq m \leq M\}$ we consider the implicit finite difference scheme

$$
\begin{aligned}
\frac{U_j^{m+1} - 2U_j^m + U_j^{m-1}}{(\Delta t)^2} - c^2 \frac{U_{j+1}^{m+1} - 2U_j^{m+1} + U_{j-1}^{m+1}}{(\Delta x)^2} &= f(x_j, t_{m+1}) \quad \text{for} \begin{cases} j = 1, \ldots, J-1, \\ m = 1, \ldots, M-1, \end{cases} \\
U_j^0 &= u_0(x_j) \qquad \text{for } j = 0, 1, \ldots, J, \\
U_j^1 &= U_j^0 + \Delta t\, u_1(x_j) \quad \text{for } j = 1, 2, \ldots, J-1, \\
U_0^m = 0 \quad \text{and} \quad U_J^m &= 0 \qquad \text{for } m = 1, \ldots, M.
\end{aligned} \tag{1}
$$

Stability of the implicit scheme

Consider the inner products

$$(U, V) := \sum_{j=1}^{J-1} \Delta x \, U_j \, V_j,$$

$$(U, V\,] := \sum_{j=1}^{J} \Delta x \, U_j \, V_j,$$

and the associated norms, respectively, $\|\cdot\|$ and $\|\cdot]\|$, defined by

$$\|U\| := (U, U)^{\frac{1}{2}} \qquad \text{and} \qquad \|U\,]\| := (U, U\,]^{\frac{1}{2}}.$$

Note that for two mesh functions A and B defined on the computational mesh $\{x_j : j = 1, \ldots, J-1\}$ one has that

$$(A - B, A) = \frac{1}{2}(\|A\|^2 - \|B\|^2) + \frac{1}{2}\|A - B\|^2.$$

Thus, by taking $A = U^{m+1} - U^m$ and $B = U^m - U^{m-1}$, we have

$$\begin{aligned}&(U^{m+1} - 2U^m + U^{m-1}, U^{m+1} - U^m)\\ &= \frac{1}{2}(\|U^{m+1} - U^m\|^2 - \|U^m - U^{m-1}\|^2) + \frac{1}{2}\|U^{m+1} - 2U^m + U^{m-1}\|^2.\end{aligned}$$

Analogously, for two mesh functions A and B defined on the mesh $\{x_j : j = 1, \ldots, J\}$ we have that

$$(A - B, A] = \frac{1}{2}(\|A]\|^2 - \|B]\|^2) + \frac{1}{2}\|A - B]\|^2.$$

Hence, by summation by parts and taking $A = D_x^- U^{m+1}$ and $B = D_x^- U^m$

$$\begin{aligned}\left(-D_x^+ D_x^- U^{m+1}, U^{m+1} - U^m\right) &= (D_x^- U^{m+1}, D_x^-(U^{m+1} - U^m)\,] \\ &= (D_x^- U^{m+1} - D_x^- U^m, D_x^- U^{m+1}\,] \\ &= \frac{1}{2}(\|D_x^- U^{m+1}]\|^2 - \|D_x^- U^m\,]\|^2) \\ &\quad + \frac{1}{2}\|D_x^-(U^{m+1} - U^m)\,]\|^2.\end{aligned}$$

By taking the $(\cdot,\cdot)$ inner product of $(1)_1$ with $U^{m+1} - U^m$ and using the identities stated above we therefore obtain:

$$\begin{aligned}&\frac{1}{2}\left(\left\|\frac{U^{m+1} - U^m}{\Delta t}\right\|^2 - \left\|\frac{U^m - U^{m-1}}{\Delta t}\right\|^2\right) + \frac{1}{2}(\Delta t)^2\left\|\frac{U^{m+1} - 2U^m + U^{m-1}}{(\Delta t)^2}\right\|^2 \\ &\quad + \frac{c^2}{2}(\|D_x^- U^{m+1}]\|^2 - \|D_x^- U^m\,]\|^2) + \frac{c^2}{2}(\Delta t)^2\left\|D_x^-\left(\frac{U^{m+1} - U^m}{\Delta t}\right)\right]\right\|^2 \\ &\quad = (f(\cdot, t_{m+1}), U^{m+1} - U^m).\end{aligned} \tag{2}$$

In the special case when f is identically zero the equality (2) gives

$$\left\| \frac{U^{m+1} - U^m}{\Delta t} \right\|^2 + c^2 \| D_x^- U^{m+1}]|^2 \leq \left\| \frac{U^m - U^{m-1}}{\Delta t} \right\|^2 + c^2 \| D_x^- U^m]|^2. \tag{3}$$

Let us define:

$$\mathcal{M}^2(U^m) := \left\| \frac{U^{m+1} - U^m}{\Delta t} \right\|^2 + c^2 \| D_x^- U^{m+1}]|^2.$$

With this notation (3) becomes

$$\mathcal{M}^2(U^m) \leq \mathcal{M}^2(U^{m-1}), \qquad \text{for all } m = 1, \ldots, M-1,$$

and therefore

$$\mathcal{M}^2(U^m) \leq \mathcal{M}^2(U^0), \qquad \text{for all } m = 1, \ldots, M-1.$$

The mapping

$$U \mapsto \max_{m \in \{0,\ldots,M-1\}} [\mathcal{M}^2(U^m)]^{1/2}$$

is a norm on the linear space of mesh functions U defined on the space-time mesh $\{(x_j, t_m) : j = 0, 1, \ldots, J,\ m = 0, 1, \ldots, M\}$ such that $U_0^m = U_J^m = 0$ for all $m = 0, 1, \ldots, M$, called the discrete energy norm.

Thus we have shown that when f is identically zero the implicit scheme (1) is unconditionally stable in this norm, i.e.:

$$\max_{m \in \{0,\ldots,M-1\}} [\mathcal{M}^2(U^m)]^{1/2} \leq [\mathcal{M}^2(U^0)]^{1/2}.$$

We now return to the general case when f is not identically zero. Our starting point is the equality (2). By the Cauchy–Schwarz inequality,

$$
\begin{aligned}
(f(\cdot, t_{m+1}), U^{m+1} - U^m) &\le \|f(\cdot, t_{m+1})\| \, \|U^{m+1} - U^m\| \\
&= \sqrt{\Delta t \, T} \, \|f(\cdot, t_{m+1})\| \sqrt{\frac{\Delta t}{T}} \left\| \frac{U^{m+1} - U^m}{\Delta t} \right\| \\
&\le \frac{\Delta t \, T}{2} \, \|f(\cdot, t_{m+1})\|^2 + \frac{\Delta t}{2T} \left\| \frac{U^{m+1} - U^m}{\Delta t} \right\|^2 ,
\end{aligned}
\tag{4}
$$

where in the transition to the last line we used the elementary inequality

$$
\alpha\beta \le \frac{1}{2}\alpha^2 + \frac{1}{2}\beta^2, \qquad \text{for } \alpha, \beta \in \mathbb{R}.
$$

Substituting (4) into (2) we deduce that

$$\begin{aligned}\left(1-\frac{\Delta t}{T}\right)\left(\left\|\frac{U^{m+1}-U^{m}}{\Delta t}\right\|^{2}+c^{2}\|D_{x}^{-}U^{m+1}]\|^{2}\right)\\ \leq\left\|\frac{U^{m}-U^{m-1}}{\Delta t}\right\|^{2}+c^{2}\|D_{x}^{-}U^{m}]\|^{2}+\Delta t\ T\ \|f(\cdot,t_{m+1})\|^{2}.\end{aligned}\tag{5}$$

By recalling the definition of $\mathcal{M}^2(U^m)$ we can rewrite (5) in the following compact form:

$$\left(1-\frac{\Delta t}{T}\right)\mathcal{M}^{2}(U^{m})\leq\mathcal{M}^{2}(U^{m-1})+\Delta t\ T\ \|f(\cdot,t_{m+1})\|^{2}.$$

As, by assumption, $M \geq 2$, it follows that $\Delta t := T/M \leq T/2$, whereby $\Delta t/T \leq 1/2$. By noting that

$$1 - x \geq \frac{1}{1+2x} \qquad \forall\, x \in \left[0, \tfrac{1}{2}\right],$$

it follows with $x = \Delta t/T$ that, for $m = 1, 2, \ldots, M-1$,

$$\begin{aligned}
\mathcal{M}^2(U^m) &\leq \left(1 + \frac{2\,\Delta t}{T}\right) \mathcal{M}^2(U^{m-1}) + \Delta t\; T \left(1 + \frac{2\,\Delta t}{T}\right) \|f(\cdot, t_{m+1})\|^2 \\
&\leq \left(1 + \frac{2\,\Delta t}{T}\right) \mathcal{M}^2(U^{m-1}) + 2\,\Delta t\; T\, \|f(\cdot, t_{m+1})\|^2.
\end{aligned}$$

We need the following result, which is easily proved by induction.

Lemma

Suppose that $M \geq 2$ *is an integer,* $\{a_m\}_{m=0}^{M-1}$ *and* $\{b_m\}_{m=1}^{M-1}$ *are nonnegative real numbers,* $\alpha > 0$*, and*

$$a_m \leq \alpha\, a_{m-1} + b_m \qquad \text{for } m = 1, 2, \ldots, M-1.$$

Then,

$$a_m \leq \alpha^m a_0 + \sum_{k=1}^{m} \alpha^{m-k} b_k \qquad \text{for } m = 1, 2, \ldots, M-1.$$

We shall apply Lemma 1 with

$$a_m = \mathcal{M}^2(U^m), \quad b_m = 2\,\Delta t\; T\,\|f(\cdot, t_{m+1})\|^2, \quad \alpha = 1 + \frac{2\,\Delta t}{T}$$

to deduce that, for $m = 1, 2, \ldots, M-1$,

$$\mathcal{M}^2(U^m) \le \left(1 + \frac{2\,\Delta t}{T}\right)^m \mathcal{M}(U^0) + 2\,\Delta t\; T \sum_{k=1}^{m} \left(1 + \frac{2\,\Delta t}{T}\right)^{m-k} \|f(\cdot, t^{k+1})\|^2.$$

We note that

$$\left(1 + \frac{2\,\Delta t}{T}\right)^m \le \left(1 + \frac{2\,\Delta t}{T}\right)^M = \left(1 + \frac{2\,\Delta t}{T}\right)^{\frac{T}{\Delta t}} \le \mathrm{e}^2,$$

where the last inequality follows from the inequality

$$(1 + 2x)^{\frac{1}{x}} \le e^2 \quad \forall\, x \in \left(0, \tfrac{1}{2}\right],$$

with $x = \Delta t / T$.

Thus we deduce the following stability result for the implicit scheme (1).

Theorem

The implicit finite difference approximation (1) *of the initial-boundary-value problem, on a finite difference mesh of spacing* $\Delta x = (b-a)/J$ *with* $J \geq 2$ *in the* x*-direction and* $\Delta t = T/M$ *with* $M \geq 2$ *in the* t*-direction, is unconditionally stable in the sense that, for* $m = 1, \ldots, M-1$,

$$\mathcal{M}^2(U^m) \leq \mathrm{e}^2\, \mathcal{M}^2(U^0) + 2\,\mathrm{e}^2\, T \sum_{k=1}^{m} \Delta t\, \|f(\cdot, t_{k+1})\|^2 ,$$

independently of the choice of Δx *and* Δt.

Consistency of the implicit scheme

We define the consistency error of the scheme by

$$T_j^{m+1} := \frac{u_j^{m+1} - 2u_j^m + u_j^{m-1}}{(\Delta t)^2} - c^2 \frac{u_{j+1}^{m+1} - 2u_j^{m+1} + 2u_{j-1}^{m+1}}{(\Delta x)^2} - f(x_j, t_{m+1}),$$

for $m = 1, 2, \ldots, M-1$, and

$$T_j^1 := \frac{u_j^1 - u_j^0}{\Delta t} - u_1(x_j), \qquad j = 1, \ldots, J-1,$$

where $u_j^m := u(x_j, t_m)$, and recall that

$$f(x_j, t_{m+1}) = \frac{\partial^2 u}{\partial t^2}(x_j, t_{m+1}) - c^2 \frac{\partial^2 u}{\partial x^2}(x_j, t_{m+1}) \quad \text{and} \quad u_1(x_j) = \frac{\partial u}{\partial t}(x_j, 0).$$

Then, by Taylor series expanion with remainder term(s):

$$|T_j^{m+1}| \leq \frac{1}{12}c^2(\Delta x)^2 M_{4x} + \frac{5}{3}\Delta t M_{3t}, \qquad \begin{cases} j = 1, \ldots, J-1, \\ m = 1, \ldots, M-1, \end{cases} \tag{6}$$

where

$$M_{4x} := \max_{(x,t)\in[a,b]\times[0,T]} \left|\frac{\partial^4 u}{\partial x^4}(x,t)\right| \quad \text{and} \quad M_{3t} := \max_{(x,t)\in[a,b]\times[0,T]} \left|\frac{\partial^3 u}{\partial t^3}(x,t)\right|.$$

Similarly, again by Taylor series expansion with a remainder term:

$$|T_j^1| \leq \frac{1}{2}\Delta t\, M_{2t}, \quad j = 1, \ldots, J-1,$$

where

$$M_{2t} := \max_{(x,t)\in[a,b]\times[0,T]} \left|\frac{\partial^2 u}{\partial t^2}(x,t)\right|.$$

Convergence of the implicit scheme

We define the global error

$$e_j^m := u(x_j, t_m) - U_j^m, \qquad \begin{cases} j = 0, \ldots, J, \\ m = 0, \ldots, M. \end{cases}$$

It follows from the definitions of T_j^{m+1} and T_j^1 that

$$\frac{e_j^{m+1} - 2e_j^m + e_j^{m-1}}{(\Delta t)^2} - c^2 \frac{e_{j+1}^{m+1} - 2e_j^{m+1} + e_{j-1}^{m+1}}{(\Delta x)^2} = T_j^{m+1},$$

for $j = 1, \ldots, J-1$ and $m = 1, \ldots, M-1$, and

$$e_j^1 = e_j^0 + \Delta t \, T_j^1, \qquad j = 1, \ldots, J-1.$$

Furthermore, $e_j^0 = 0$ for $j = 0, 1, \ldots, J$, and $e_0^m = e_J^m = 0$ for $m = 1, \ldots, M$.

Hence, the global error e satisfies an analogous finite difference scheme as U, but with $f(x_j, t_{m+1})$ replaced by T_j^{m+1}, $U_j^0 = u_0(x_j)$ replaced by $e_j^0 = 0$, and $u_1(x_j)$ replaced by T_j^1.

Theorem 2 with U^m replaced by e^m, U^0 replaced by e^0 and $f(x_j, t_{k+1})$ replaced by T_j^{k+1} for $j = 1, \ldots, J-1$ and $k = 1, \ldots, M-1$, gives that

$$\mathcal{M}^2(e^m) \leq \mathrm{e}^2\, \mathcal{M}^2(e^0) + 2\,\mathrm{e}^2\, T \sum_{k=1}^{m} \Delta t \left\| T^{k+1} \right\|^2, \quad \text{for } m = 1, \ldots, M-1.$$

It remains to bound the terms on the r.h.s. of this inequality.

Because $m\Delta t \leq M\Delta t = T$ and $(J-1)\Delta x \leq b-a$, (6) implies that

$$\sum_{k=1}^{m} \Delta t \left\| T^{k+1} \right\|^2 \leq T \max_{1\leq k\leq m} \left\| T^{k+1} \right\|^2 = T \max_{1\leq k\leq m} \sum_{j=1}^{J-1} \Delta x \, |T_j^{k+1}|^2$$

$$\leq T(b-a) \left[\frac{1}{12} c^2 (\Delta x)^2 M_{4x} + \frac{5}{3} \Delta t \, M_{3t} \right]^2 .$$

Similarly,

$$\mathcal{M}^2(e^0) \leq (b-a) \left[\frac{1}{2} \Delta t \, M_{2t} \right]^2 + (b-a) \left[\frac{1}{2} (\Delta t)^2 M_{1x2t} \right]^2 .$$

Proof of the last inequality

$$\mathcal{M}^2(e^0) = \left\| \frac{e^1 - e^0}{\Delta t} \right\|^2 + \|D_x^- e^1]\|^2 = \|T^1\|^2 + \|D_x^- e^1]\|^2$$
$$\leq (b-a) \left[\frac{1}{2} \Delta t \, M_{2t} \right]^2 + \|D_x^- e^1]\|^2.$$

It remains to bound the term $\|D_x^- e^1]\|^2$:

$$D_x^- e_j^1 = D_x^- e_j^0 + \Delta t \, D_x^- T_j^1 = \Delta t \, D_x^- T_j^1 = \int_0^{\Delta t} (\Delta t - t) \, D_x^- \frac{\partial^2 u}{\partial t^2}(x_j, t) \, \mathrm{d}t$$
$$= \frac{1}{\Delta x} \int_0^{\Delta t} (\Delta t - t) \int_{x_{j-1}}^{x_j} \frac{\partial^3 u}{\partial x \, \partial t^2}(x, t) \, \mathrm{d}x \, \mathrm{d}t.$$

Thus,

$$|D_x^- e_j^1| \leq \frac{1}{2} (\Delta t)^2 \, M_{1x2t}, \qquad \text{where} \quad M_{1x2t} := \max_{(x,t) \in [a,b] \times [0,T]} \left| \frac{\partial^3 u}{\partial x \partial t^2} \right|,$$

and therefore

$$\|D_x^- e^1]\|^2 \leq (b-a) \left[\frac{1}{2} (\Delta t)^2 M_{1x2t} \right]^2.$$

Hence, finally,

$$\mathcal{M}^2(e^m) \leq \mathrm{e}^2(b-a)\left[\frac{1}{2}\Delta t\, M_{2t}\right]^2 + \mathrm{e}^2(b-a)\left[\frac{1}{2}(\Delta t)^2 M_{1x2t}\right]^2 + 2\mathrm{e}^2\, T^2(b-a)\left[\frac{1}{12}c^2(\Delta x)^2 M_{4x} + \frac{5}{3}\Delta t\, M_{3t}\right]^2$$

for $m = 1, \ldots, M-1$. Thus, provided that M_{2t}, M_{1x2t}, M_{4x} and M_{3t} are all finite, we have that

$$\max_{m\in\{1,\ldots,M-1\}} [\mathcal{M}^2(u^m - U^m)]^{\frac{1}{2}} = \mathcal{O}((\Delta x)^2 + \Delta t).$$

In summary then, the implicit scheme is unconditionally stable and is unconditionally convergent with order $\mathcal{O}((\Delta x)^2 + \Delta t)$.