Analytic Topology

Rolf Suabedissen

November 19, 2026

(November 2026) Various mistakes fixed and expanded the Stone duality section. Inevitably some mistakes will remain - consult a textbook or email me if something looks wrong.

This document contains the (important) definitions, statements and proof sketches for results which require a new idea. The other proofs should be straightforward.

Concepts and results which should be known from a previous course are typeset in a smaller size.

Throughout this document assume that X, Y, Z as well as $X_i, Y_i, Z_i, i \in I$ are topological spaces and I is some index set unless otherwise indicated.

1 Topological spaces, Bases, Subbases, Initial Topology, Products

1.1 Definitions

A topology on a set X is a collection τ of subsets of X containing the empty set, X that is closed under taking finite intersections and arbitrary unions.

A topological space is a pair (X, τ) such that τ is a topology on X.

In a topological space (X, τ), elements of X are called points, elements of τ are called open sets, complements of elements of τ are called closed sets and subsets of X that are closed and open are called clopen.

For a subset A of a topological space X, the closure of A, $\overline{A}$, is the smallest closed set containing A and the interior of A, $\operatorname{int}(A)$, is the largest open set contained in A.

A function $f\colon X \to Y$ is continuous if and only if preimages of Y-open sets under f are X-open.

If $A \subseteq X$, the subspace topology on A is $\{U \cap A\colon U \text{ open } \subseteq X\}$.

A basis for a topology τ on X is a collection $\mathcal{B} \subseteq \tau$ such that every open set is a union of a subcollection $\mathcal{B}'$ of $\mathcal{B}$. If a basis has been fixed, its elements are called basic open sets.

X is metrizable if and only if there is a metric d on X such that $\left\{B_\epsilon^d(x) \colon x \in X, \epsilon > 0\right\}$ is a basis for X.

A space is second countable if and only if it has a countable basis.

A subbasis for a topology τ on X is a collection $\mathcal{S} \subseteq \tau$ such that the set of finite intersections of elements of $\mathcal{S}$ is a basis for τ. If a subbasis has been fixed, its elements are called subbasic open sets.

Given a set X and a collection $\mathcal{F} = \{f_i \colon X \to Y_i \colon i \in I\}$ the initial topology with respect to $\mathcal{F}$ is the smallest (wrt $\subseteq$) topology on X such that each $f_i \in \mathcal{F}$ is continuous.

The Tychonoff product $\prod_{i \in I} X_i$ of topological spaces $X_i, i \in I$ is the topological space consisting of the Cartesian (set) product equipped with the initial topology with respect to the projections.

1.2 Results

Lemma 1.1 (Recall)**.** *1. If $A \subseteq X$, then $\overline{A}$ exists and equals*

$$\bigcap \{C \colon A \subseteq C \text{ closed } \subseteq X\} = \{x \in X \colon \forall \text{ open } U \ni x \; U \cap A \neq \emptyset\}.$$

2. The closure operator $A \mapsto \overline{A}$ satisfies $\overline{\emptyset} = \emptyset$, $\overline{\overline{A}} = \overline{A}$, $A \subseteq \overline{A}$, $\overline{A \cup B} = \overline{A} \cup \overline{B}$ and $\overline{\bigcap_i A_i} \subseteq \bigcap_i \overline{A_i}$. Dual results hold for the interior operator.

3. A function $f \colon X \to Y$ is continuous if and only if for every $A \subseteq X$, $f\left(\overline{A}\right) \subseteq \overline{f(A)}$.

4. If $B \subseteq A \subseteq X$ then $\overline{B}^A = \overline{B}^X \cap A$.

Theorem 1.2. *1. The set of topologies on a fixed set X is a complete lattice with respect to $\subseteq$, i.e. a partial order with arbitrary infima and suprema. The infimum of a collection $\tau_i, i \in I$ of topologies on X is $\bigcap_i \tau_i$. The greatest element of the complete lattice is the discrete topology $\mathcal{P}(X)$, the smallest element is the indiscrete topology $\{\emptyset, X\}$.*

2. A collection $\mathcal{B}$ of subsets of X is the basis for a (necessarily unique) topology $\tau = \{\bigcup \mathcal{B}' \colon \mathcal{B}' \subseteq \mathcal{B}\}$ on X if and only if $\bigcup \mathcal{B} = X$ and for every $B_1, B_2 \in \mathcal{B}$ there is $\mathcal{B}' \subseteq \mathcal{B}$ such that $B_1 \cap B_2 = \bigcup \mathcal{B}'$. Moreover, τ is the smallest topology on X containing $\mathcal{B}$.

3. Every collection $\mathcal{S}$ is a subbasis for a (necessarily unique) topology τ on X with basis $\{\bigcap \mathcal{F} \colon \mathcal{F} \text{ finite } \subseteq \mathcal{S}\}$. Moreover, τ is the smallest topology on X containing $\mathcal{S}$.

4. *If Y is a topological space with a fixed subbasis, a function $f\colon X \to Y$ is continuous if and only if preimages of subbasic open sets under f are open.*

5. *If X is a set, $Y_i, i \in I$ are topological spaces and $f_i\colon X \to Y_i$ are functions, the initial topology with respect to the f_i exists and has subbasis $\left\{{f_i}^{-1}(U) : i \in I, U \text{ open } \subseteq Y_i\right\}$. It is the unique topology on X such that for every topological space Z and every function $f\colon Z \to X$, f is continuous if and only if each $f_i \circ f$ is continuous.*

6. *The product topology on $\prod_i X_i$ has basis*

$$\left\{\prod_i U_i\colon U_i \text{ open } \subseteq X_i \text{ and } X_i = U_i \text{ except for finitely many } i\right\}.$$

7. ***Embedding Lemma:*** *If $f_i\colon X \to X_i$ are continuous maps such that for distinct $x, y \in X$ there is $i \in I$ with $f_i(x) \neq f_i(y)$ and such that $\left\{{f_i}^{-1}(U) : i \in I, U \text{ open } \subseteq X_i\right\}$ is a basis for X then the diagonal map $\Delta = \Delta_i f_i\colon X \to \prod_i X_i; x \mapsto (f_i(x))_i$ is a homeomorphic embedding.*

8. *Countable products of metrizable spaces are metrizable.*

1.3 Proofs

Most of the proofs are straightforward set arithmetic.

5. For uniqueness, suppose that τ_1, τ_2 are two topologies on X satisfying the condition. Then $\mathrm{id}_{1,1}\colon (X, \tau_1) \to (X, \tau_1)$ is continuous, so each $f_i\colon (X, \tau_1) \to Y_i = f_i \circ \mathrm{id}_{1,1}$ is continuous. Thus every $f_i \circ \mathrm{id}_{1,2}$ is continuous and hence $\mathrm{id}_{1,2}$ is continuous. By symmetry $\mathrm{id}_{2,1}$ is continuous and hence $\tau_1 = \tau_2$.

7. **Embedding Lemma:** The only non-trivial bit is to check that Δ is open onto its image. For this note that unions and images commute and hence it is sufficient to consider basic open sets of the form ${f_i}^{-1}(U)$. But $\Delta\left({f_i}^{-1}(U)\right) = {\pi_i}^{-1}(U) \cap \Delta(X)$.

8. Exercise Sheet.

2 Separation Properties

2.1 Definitions

X is T_0 if and only if for every distinct $x, y \in X$ there is open U that contains exactly one of x and y.

X is T_1 if and only if for every distinct $x, y \in X$ there is open U such that $x \in U \not\ni y$.

X is T_2 (Hausdorff) if and only if for every distinct $x, y \in X$ there are disjoint open $U \ni x, V \ni y$ (x and y are separated by open sets).

X is T_3 (regular) if and only if X is T_1 and for every $x \in X$ and every closed $C \not\ni x$ there are disjoint open $U \ni x, V \supseteq C$.

X is $T_{3.5}$ (Tychonoff) if and only if X is T_1 and for every $x \in X$ and every closed $C \not\ni x$ there is a continuous $f\colon X \to [0,1]$ such that $f(x) = 0$ and $f(C) \subseteq \{1\}$.

X is T_4 (normal) if and only if X is T_1 and for every disjoint closed C, D there are disjoint open $U \supseteq C, V \supseteq D$.

X is functionally normal if and only if X is T_1 and for every disjoint closed C, D there is a continuous function $f\colon X \to [0,1]$ such that $f(C) \subseteq \{0\}, f(D) \subseteq \{1\}$.

X is T_5 (hereditarily normal) if and only if every subspace of X is normal.

X is T_6 (perfectly normal) if and only if X is T_1 and for every closed subspace C of X there is a continuous $f\colon X \to [0,1]$ such that $C = f^{-1}(\{0\})$.

2.2 Results

Theorem 2.1. *1. X is T_1 if and only if every singleton is closed.*

2. *If a basis for X has been fixed then for $i \leq 2$, replacing 'open' in the definition of T_i by 'basic open' yields an equivalent property.*

3. *functionally normal $\implies$ $T_{3.5}$ $\implies$ T_3 $\implies$ T_2 $\implies$ T_1 $\implies$ T_0. (None of these reverse in general.)*

4. *X is Tychonoff if and only if X is (homeomorphic to) a subspace of a power of $[0,1]$.*

5. *For $i \leq 3.5$, products and subspaces of T_i-spaces are T_i.*

6. ***Urysohn's Lemma:** Functionally Normal $\iff$ T_4.*

7. *Subspaces of normal spaces need not be normal. Products (even squares) of normal spaces need not be normal.*

8. ***Urysohn's Metrization Theorem I:*** *If X is normal and second countable then X is metrizable.*

9. *Metric spaces are perfectly normal.*

10. $T_6 \implies T_5 \implies T_4$ *(**Not examinable as bookwork.**)*

11. *A normal space is perfectly normal if and only if every closed subset is a countable intersection of open subsets (a G_δ).(**Not examinable as bookwork.**)*

2.3 Proofs

General important ideas are:

- A, B disjoint is equivalent to $A \subseteq X \setminus B$ (and of course B is open if and only if $X \setminus B$ is closed).

- If $f\colon X \to [0,1]$ is continuous, then $f^{-1}([0,1/3)), f^{-1}((2/3,1])$ are disjoint open and $f^{-1}(0) = \bigcap_n f^{-1}([0,2^{-n}))$.

4. Apply the Embedding Lemma.

5. For productivity of Tychonoffness, let $x \in U_1 \times \cdots \times U_n \times \prod X_i$. For each $k = 1, \ldots, n$, find a continuous f_k which is 1 at $\pi_k(x)$ and 0 outside U_i and take the product of the $f_k \circ \pi_k$.

6. **Urysohn's Lemma:** Backwards direction: Well order the countable set $\mathbb{Q} \cap (0,1)$, set $\overline{U_0} = C$, $U_1 = X \setminus D$ and inductively construct open U_r such that $r < s \implies \overline{U_r} \subseteq U_s$.

   Now define $f\colon X \to [0,1]$ by $f(x) = \inf\{r \in \mathbb{Q} \cap (0,1)\colon x \in U_r\}$ (where $\inf \emptyset = 1$) clearly satisfying $f(C) \subseteq \{0\}$ and $f(D) \subseteq \{1\}$.

   Using the Approximation Property note that if $f(x) < \beta$ then we can find $r \in \mathbb{Q} \cap (0,1)$ with $x \in U_r$ and $f(U_r) \subseteq [0,\beta)$.

   Next note that for $g(x) = \inf\left\{r \in \mathbb{Q} \cap (0,1)\colon x \in \overline{U_r}\right\}$ we have $g(x) \leq f(x)$. If $g(x) < f(x)$ then take $r \in \mathbb{Q}$ with $g(x) < r < f(x)$ and $s < r$

such that $x \in \overline{U_s} \subseteq U_r$ by the Approximation property for g so that $f(x) \leq r$ a contradiction. Hence $f = g$.

Now if $\alpha < f(x) = g(x)$ then take $s \in \mathbb{Q}$ with $\alpha < s < g(x)$. By the nestedness of the U_q we have $x \in X \setminus \overline{U_s}$ and $g(X \setminus \overline{U_s}) \subseteq [s, 1] \subseteq (\alpha, 1]$.

Thus $x \in U_r \cap X \setminus \overline{U_s} \subseteq f^{-1}(\alpha, \beta)$ showing continuity of f.

7. Below you can replace $\aleph_1$ by any uncountable set.

   We let $Y_f = \aleph_1 \cup \{\star\}$ with topology $\mathcal{P}(\aleph_1) \cup \{Y_f \setminus C \colon C \text{ finite } \subseteq \aleph_1\}$. It is easy to check that this is normal (it is compact Hausdorff) and we let $X = \{0\} \cup \{2^{-n} \colon n \in \mathbb{N}\}$ (with its usual topology). Then $Y_f \times X$ is compact Hausdorff so normal. Consider the subspace $Y_f \times X \setminus \{(\star, 0)\}$. $C = \aleph_1 \times \{0\}$ and $D = \{\star\} \times \{2^{-n} \colon n \in \mathbb{N}\}$ are closed disjoint. If $U \supseteq D$, then for each $n \in \mathbb{N}$ there is a countable $C_n \subseteq \aleph_1$ such that $(\aleph_1 \setminus C_n) \times \{2^{-n}\} \subseteq U$. Pick $\alpha \in \aleph_1 \setminus \bigcup_n C_n$ (this is non-empty at $\bigcup_n C_n$ is countable) and note that $\{\alpha\} \times \{2^{-n} \colon n \in \mathbb{N}\} \subseteq U$. Thus $(\alpha, 0) \in \overline{U} \cap C$.

8. **Urysohn's Metrization Theorem I:** Let $\mathcal{B}$ be a countable basis and for each $(B, B') \in \mathcal{B}^2$ such that $\overline{B} \subseteq B'$ find a continuous $f \colon X \to [0, 1]$ such that $\overline{B} \subseteq f^{-1}(0), X \setminus B' \subseteq f^{-1}(1)$ and apply the Embedding Lemma to these (countably many) f.

9. If $C \subseteq X$ is closed then $d_C(x) = \inf\left\{\hat{d}(x, c) \colon c \in C\right\}$ is as required with $\hat{d}(x, c) = \min\{1, d(x, c)\}$ and $\inf \emptyset = 1$.

10. T_6 is hereditary. For normality, note that if C, D are disjoint closed and $C = f^{-1}(0)$, $D = g^{-1}(0)$ then $\frac{f}{f+g}$ is the required Urysohn function.

11. For the converse, let closed $C = \bigcap_n U_n$ and let $f_n : X \to [0, 2^{-n}]$ be continuous functions that are 0 on C and 2^{-n} outside U_n. Let $f = \sum_n f_n$.

3 Filters

3.1 Definitions

Suppose X is a set.

A filter $\mathcal{F}$ on X is a non-empty collection of subsets of X that does not contain $\emptyset$ and is closed under supersets and finite intersections.

A filter basis $\mathcal{B}$ for a filter $\mathcal{F}$ on X is a subcollection of $\mathcal{F}$ such that for every $F \in \mathcal{F}$ there is $\mathcal{B} \in \mathcal{B}$ with $B \subseteq F$.

Two collections $\mathcal{A}, \mathcal{B}$ of subsets of X mesh, written $\mathcal{A}\#\mathcal{B}$ if and only if for every $A \in \mathcal{A}, B \in \mathcal{B}$ we have $A \cap B \neq \emptyset$. We also write $\mathcal{A}\#\mathcal{B} = \{A \cap B \colon A \in \mathcal{A}, B \in \mathcal{B}\}$.

An ultrafilter on X is a filter on X that is maximal wrt $\subseteq$.

For $x \in X$, the principal filter at x is $\mathcal{P}_x = \{A \subseteq X \colon x \in A\}$.

If $f \colon X \to Y$ is a function and $\mathcal{F}$ is a filter on X then $f(F) := \{B \subseteq Y \colon f^{-1}(B) \in \mathcal{F}\}$.

Now assume that X is a topological space.

For $x \in X$, the neighbourhood filter at x is $\mathcal{N}_x = \{N \subseteq X \colon \exists \text{ open } U \ x \in U \subseteq N\}$.

If $\mathcal{F}$ is a filter on X, $\lim \mathcal{F} = \{x \in X \colon \mathcal{N}_x \subseteq \mathcal{F}\}$ and $\mathcal{F} \to x$ if and only if $x \in \lim \mathcal{F}$.

If $\mathcal{P}$ is a property of topological spaces then X is locally $\mathcal{P}$ if and only if every neighbourhood filter has a filter basis of sets that are $\mathcal{P}$ (with respect to the subspace topology).

3.2 Results I

Lemma 3.1. *Suppose X is a set.*

1. *A non-empty collection $\mathcal{B}$ of non-empty subsets of X is a filter basis for a (necessarily unique) filter $\mathcal{F}$ on X if and only if for every $B_1, B_2 \in \mathcal{B}$ there is $B_3 \in \mathcal{B}$ such that $B_3 \subseteq B_1 \cap B_2$.*

2. *If $\mathcal{C}$ is a family of subsets of X with the f.i.p. then $\{\bigcap \mathcal{F} \colon \mathcal{F} \text{ finite } \subseteq \mathcal{C}\}$ is a filter basis for the smallest filter containing $\mathcal{C}$.*

3. *A filter $\mathcal{U}$ on X is an ultrafilter if and only if for every $A \subseteq X$ exactly one of A and $X \setminus A$ belongs to $\mathcal{U}$ if and only if whenever $A \cup B \in \mathcal{U}$ at least one of A or B belongs to $\mathcal{U}$.*

4. ***Ultrafilter Extension Lemma:** Every filter can be extended to an ultrafilter.*

5. *If $f\colon X \to Y$ is a function and $\mathcal{F}$ a filter on X then $f(F)$ is a filter on Y with filter basis $\{f(F) : F \in \mathcal{F}\}$. Moreover, if $\mathcal{F}$ is an ultrafilter then so is $f(F)$.*

3.3 Proofs I

Most of this is easy (once comfortable with the notation) set arithmetic.

General important ideas are:

- A family of subsets of X with the f.i.p. can be extended to a filter.
- If two families $\mathcal{A}, \mathcal{B}$ mesh, their mesh $\mathcal{A}\#\mathcal{B}$ can be extended to a filter.

Specific Lemmas:

3. Suppose $\mathcal{U}$ is an ultrafilter. If $A \notin \mathcal{U}$, then $X \setminus A \# \mathcal{U}$, so by maximality $X \setminus A \in \mathcal{U}$. The converse is obvious. For the last if and only if: for the forward direction assume $A \notin \mathcal{U}$. Then $X \setminus A, A \cup B \in \mathcal{U}$, so $B \supseteq (X \setminus A) \cap (A \cup B) \in \mathcal{U}$. For the backwards direction note that $A \cup (X \setminus A) = X \in \mathcal{U}$.
4. **Ultrafilter Extension Lemma: Proof not examinable!** Note that the union of a chain of filters is a filter and apply Zorn's Lemma.

3.4 Results II

Theorem 3.2. *Suppose X is a topological space.*

1. *Suppose $A \subseteq X$ and $x \in X$. $x \in \overline{A}$ if and only if there is a filter $\mathcal{F} \ni A$ such that $\mathcal{F} \to x$ if and only if there is an ultrafilter $\mathcal{U} \ni A$ such that $\mathcal{U} \to x$.*
2. *$f\colon X \to Y$ is continuous if and only if for every $x \in X$ and filter $\mathcal{F} \to x$, $f(F) \to f(x)$ if and only if for every $x \in X$ and ultrafilter $\mathcal{U} \to x$, $f(U) \to f(x)$.*
3. *X is Hausdorff if and only if every filter converges to at most one point if and only if every ultrafilter converges to at most one point.*

4 Compactness, Compactifications, Local Compactness, Čech-completeness

4.1 Definitions

A topological space is compact if and only if every open cover has a finite subcover.

A topological space is Lindelöf if and only if every open cover has a countable subcover.

A Hausdorff compactification of a topological space X is a pair (h, Y) where Y is compact Hausdorff and $h\colon X \to Y$ is a dense homeomorphic embedding.

For a topological space X and two Hausdorff compactification (h_1, Y_1), (h_2, Y_2), we define $(h_2, Y_2) \leq (h_1, Y_1)$ if and only if there is a continuous $g\colon Y_1 \to Y_2$ such that $g \circ h_1 = h_2$. We define $(h_1, Y_1) \sim (h_2, Y_2)$ if and only if there is a homeomorphism $g\colon Y_1 \to Y_2$ such that $g \circ h_1 = h_2$.

For a topological space X and a Hausdorff compactification (h, Y) we say that (h, Y) satisfies the Stone-Čech-property with respect to continuous maps into compact Hausdorff spaces if and only if for every compact Hausdorff space Z and every continuous $f\colon X \to Z$ there is a continuous $F\colon Y \to Z$ such that $f = F \circ h$.

The Stone-Čech compactifiation of a topological space is (the unique, if it exists) Hausdorff compactification $(\beta, \beta X)$ of X satisfying the Stone-Čech property wrt continuous maps into compact Hausdorff spaces.

Recall that a space X is locally compact if and only if for every $x \in U$ open $\subseteq X$ there is compact K and open V with $x \in V \subseteq K \subseteq U$ or equivalently if the neighbourhood filter of every point has a filter basis consisting of compact sets.

The Alexandroff one-point compactification of a topological space X is (the unique, if it exists) Hausdorff compactification $(\omega, \omega X)$ of X such that $\omega X \setminus \omega(X)$ is a singleton.

A Tychonoff topological space is Čech-complete if and only if for every Hausdorff compactification (h, Y) of X, $Y \setminus h(X)$ is a countable union of closed subsets of Y (i.e. an F_σ).

4.2 Results

The key ideas are:

- Compactness properties are inherited by closed subsets.
- Diagonal maps!
- if $f\colon Y \to Z$ is continuous and X dense in Y then $f(Y) \subseteq \overline{f(X)}$.
- Compactness is preserved by images, closedness by pre-images.

Lemma 4.1 (Recall)**.** *1. A topological space is compact if and only if every family of closed sets with the finite intersection property has non-empty intersection.*

2. Every closed subset of a compact topological space is compact.

3. Every compact subset of a Hausdorff space is closed.

4. Every compact Hausdorff space is regular. Every compact regular space is normal.

5. If X is compact, Y is a topological space and $f\colon X \to Y$ is continuous then $f(X)$ is compact.

Theorem 4.2. *1. Every Lindelöf regular space is normal.*

2. Every second countable space is Lindelöf.

3. Every Lindelöf metric space is second countable.

4. ***Urysohn's Metrization Theorem II:*** *A compact Hausdorff space is metrizable if and only if it is second countable.*

5. X is compact if and only if every ultrafilter on X converges (to some point).

6. ***Tychonoff's Theorem:*** *Products of compact spaces are compact.*

7. X has a Hausdorff compactification if and only if it is Tychonoff.

8. $\sim$ is an equivalence relation on the Hausdorff compactifications of X. If (h_1, Y_1), (h_2, Y_2) are Hausdorff compactifications of X such that $(h_1, Y_1) \leq (h_2, Y_2) \leq (h_1, Y_1)$ then $(h_1, Y_1) \sim (h_2, Y_2)$ and thus $\leq$ induces a partial order on the equivalence classes of Hausdorff compactifications under $\sim$.

9. If (h_1, Y_1), (h_2, Y_2) are Hausdorff compactifications of X such that $(h_1, Y_1) \leq (h_2, Y_2)$ as witnessed by $g\colon Y_2 \to Y_1$ then $g(Y_2 \setminus h_2(X)) = Y_1 \setminus h_1(X)$.

10. *If X is Tychonoff, then the partial order of (equivalence classes of) Hausdorff compactifications has suprema of non-empty families. Moreover each equivalence class has a representative with cardinality at most $2^{2^{|X|}}$.*

11. *If X is Tychonoff, then X has a Stone-Čech compactification which is unique (up to equivalence) and is the greatest compactification of X.*

12. *A Hausdorff compactification of X satisfies the Stone-Čech property with respect to continuous maps into compact Hausdorff spaces if and only if it satisfies the Stone-Čech property with respect to continuous maps into $[0,1]$.*

13. *Open subsets of locally compact spaces are locally compact.*

14. *Compact Hausdorff spaces are locally compact.*

15. *If X is non-compact, locally compact, Hausdorff and $\infty \notin X$ then $\omega X = X \cup \{\infty\}$ with topology $\{U : U \text{ open } \subseteq X\} \cup \{\omega X \setminus K : K \text{ compact } \subseteq X\}$ and embedding $\omega : X \to \omega X; x \mapsto x$ is the unique one-point compactification of X.*

16. *If X is Tychonoff and non-compact then the following are equivalent:*

    (a) *X is locally compact.*

    (b) *X has a smallest Hausdorff compactification.*

    (c) *X has a one-point compactification.*

    (d) *$\beta X \setminus \beta(X)$ is closed.*

    (e) *For every Hausdorff compactification (h, Y) of X, $Y \setminus h(X)$ is closed.*

    (f) *For some Hausdorff compactification (h, Y) of X, $Y \setminus h(X)$ is closed.*

17. *If X is Tychonoff, the following are equivalent:*

    (a) *$\beta X \setminus \beta(X)$ is a F_σ.*

    (b) *X is Čech-complete.*

    (c) *For some Hausdorff compactification (h, Y) of X, $Y \setminus h(X)$ is a F_σ.*

4.3 Proofs

1. Exercise Sheet.

5. Suppose X is compact and $\mathcal{U}$ is an ultrafilter not converging to any $x \in X$. For each $x \in X$, choose open $U_x \ni x$ such that $U_x \notin \mathcal{U}$. Then $\{U_x \colon x \in X\}$ is an open cover with finite subcover $U_{x_1}, \ldots, U_{x_n}$. Thus one of $U_{x_i} \in \mathcal{U}$ (a contradiction).

   Now assume that X is not compact: let $\mathcal{C}$ be a family of closed subsets with the f.i.p. but empty intersection. Extend $\mathcal{C}$ to a filter and then to an ultrafilter $\mathcal{U}$. If $x \in X$ then $x \notin$ some C_x, so $x \in X \setminus C_x \in \mathcal{N}_x$ and hence $\mathcal{N}_x \not\subseteq \mathcal{U}$, i.e. $\mathcal{U} \not\to x$.

6. **Tychonoff's Theorem:** Let $\mathcal{U}$ be an ultrafilter on $\prod_i X_i$. For each $i \in I$, $\pi_i(\mathcal{U})$ is an ultrafilter on X_i, so converges to some x_i. Now check that $\mathcal{U} \to (x_i)_i$.

8. If $(h_1, Y_1) \leq (h_2, Y_2) \leq (h_1, Y_1)$ is witnessed by $g \colon Y_2 \to Y_1$ and $h \colon Y_1 \to Y_2$ respectively, note that $g \circ h$ and $h \circ g$ are the identity on $h_1(X)$ and $h_2(X)$ respectively. But $h_i(X)$ is dense in the Hausdorff Y_i, so $g \circ h$ and $h \circ g$ are the identity on Y_1 and Y_2 respectively. Hence g is a homeomorphism as required.

9. Wlog h_2 is the identity. Suppose $y \in Y_2 \setminus X$ and $x \in X$ with $g(y) = h_1(x)$. Let $\mathcal{F}$ be a filter on Y_2 containing X and converging to $y \in \overline{X}$. As Y_2 is Hausdorff $\mathcal{F} \not\to x$. Then $\mathcal{F}_X = \{F \cap X \colon F \in \mathcal{F}\}$ is a filter on X and $\mathcal{F}_X \not\to x$. As h_1 is a homeomorphism $X \to h_1(X)$, $h_1(\mathcal{F}_X) \not\to h_1(x) = g(y)$. But $h_1(\mathcal{F}_X) = g(\mathcal{F})$ and $g(\mathcal{F}) \to g(y)$ by continuity.

10. If (h_i, Y_i) are compactifications, then check that $(\Delta_i h_i, \overline{\Delta_i h_i(X)}^{\prod_i Y_i})$ is an upper bound. If (g, Z) is another upper bound witnessed by the $g_i \colon Z \to Y_i$, then $\Delta_i g_i \colon Z \to \prod_i Y_i$ is continuous and into $\overline{\Delta_i h_i(X)}$ since $\Delta_i g_i(Z) = \Delta_i g_i\left(\overline{g(X)}^Z\right) \subseteq \overline{\Delta_i g_i(g(X))} = \overline{\Delta_i h_i(X)}$.

    For the 'moreover' claim: suppose X is dense in Y and let $f \colon Y \to \mathcal{P}(\mathcal{P}(X)); y \mapsto \left\{A \subseteq X \colon y \in \overline{A}^Y\right\}$. As Y is Hausdorff, for $y \neq y'$ there is open $U \ni y$ with $y' \notin \overline{U}^Y = \overline{U \cap X}^Y$. Thus f is an injection.

11. First uniqueness up to $\sim$: suppose (h_1, Y_1), (h_2, Y_2) are Hausdorff compactifications of X satisfying the Stone-Čech property. Then $h_2 \colon X \to Y_2$ is a continuous map into a compact Hausdorff space, so there is $H_2 \colon Y_1 \to Y_2$ such that $H_2 \circ h_1 = h_2$, i.e. $(h_2, Y_2) \leq (h_1, Y_1)$. By symmetry $(h_1, Y_1) \leq (h_2, Y_2)$ and hence $(h_1, Y_1) \sim (h_2, Y_2)$.

    Now we show existence, by showing that the greatest Hausdorff compactification of X satisfies the Stone-Čech property: for each equivalence class, choose a representative, and let $(\beta, \beta X)$ be the supremum over these representatives (there are only set many). If $f \colon X \to Z$ is continuous into a compact Hausdorff Z, then $f\Delta\beta \colon X \to \overline{f\Delta\beta\left(X\right)}^{Z \times \beta X} = Y$ is an embedding by the Embedding Lemma and hence determines a Hausdorff compactification of X. As $(\beta, \beta X)$ is the greatest Hausdorff compactification of X, there is a continuous $g \colon \beta X \to Y$ with $g \circ \beta = f\Delta\beta$. Then $F = \pi_Z \circ g$ is as required.

    Note that the existence proof has a special case ($f = h \colon X \to Y$) that shows that any compactification with the Stone-Čech property must be the greatest compactification of X (up to equivalence).

12. It is enough to check that the Stone-Čech property for continuous $[0,1]$-valued maps implies the Stone-Čech property for continuous maps into compact Hausdorff spaces. To that end, note that every compact Hausdorff space is normal, so Tychonoff, so homeomorphic to a closed subspace C of $[0,1]^I$ (for some I). So assume (h, Y) satisfies the Stone-Čech property for continuous $[0,1]$-valued maps and wlog $X \subseteq Y, h = \mathrm{id}_X$. If $f \colon X \to C$ is continuous, then each $f_i = \pi_i \circ f$ extends to some $F_i \colon Y \to [0,1]$ and thus $\Delta_i f_i = f \colon X \to [0,1]^I$ extends to $\Delta = \Delta_i F_i \colon Y \to [0,1]^I$. But $\Delta\left(Y\right) = \Delta\left(\overline{X}\right) \subseteq \overline{\Delta\left(X\right)} \subseteq \overline{C} = C$. Hence Δ is as required.

16. Statements 1 and 3 are equivalent to 6. To see 4,5,6 are equivalent, we use that remainders map (on)to remainders: If $X \subseteq Y$ and Y compact Hausdorff with $Y \setminus X$ closed, then note that $\beta X \setminus \beta\left(X\right) = g^{-1}\left(Y \setminus X\right)$ where g witnesses $(\mathrm{id}, Y) \leq (\beta, \beta X)$ so that $\beta X \setminus \beta\left(X\right)$ is closed. If $\beta X \setminus \beta\left(X\right)$ is closed, it is compact and hence $Y \setminus X = g\left(\beta X \setminus \beta\left(X\right)\right)$ is compact so closed where $X \subseteq Y$, Y is compact Hausdorff and g witnesses $(\mathrm{id}_X, Y) \leq (\beta, \beta X)$ giving 4 implies 5. Finally note that X is Tychonoff, so 5 implies 6 as X has a compactification.

So assume one (hence all) of 1,3,4,5,6, let $(\omega, \omega X)$ be the one-point compactification of X and (h, Y) some Hausdorff compactification: we claim that $g\colon Y \to \omega X$ given by $g(h(x)) = \omega(x)$ and $g(y) = \star$ for $y \in Y \setminus h(X)$ is continuous: if C is closed so compact in ωX then either $C \subseteq X$ and hence $g^{-1}(C) = h(C)$ is compact so closed in Y or $C \ni \star$ and hence $g^{-1}(C) = Y \setminus h(X \setminus C)$ which is Y-closed since $X \setminus C$ is X-open and hence $h(C \cap X)$ is $h(X)$-open and then also Y-open as $h(X)$ is Y-open.

Finally assume 2 and that there is a smallest compactification (h, Y) that is not the one-point compactification: let $y_1, y_2 \in Y \setminus h(X)$ be distinct. Then $Y \setminus \{y_1, y_2\}$ is locally compact, Tychonoff and hence has a one-point compactification $Z = (Y \setminus \{y_1, y_2\}) \cup \{\star\}$. Since Y is a two-point compactification of $Y \setminus \{y_1, y_2\}$, there is a continuous $g\colon Y \to Z$ (which is the identity except that $g(y_1) = g(y_2) = \star$). But (h, Z) is also a Hausdorff compactification of X so that $(h, Y) \leq (h, Z)$. Thus the g above must be a homeomorphism, a contradiction to it not being injective.

17. Just like the equivalence of 4,5,6 in 16., noting that unions and images as well as pre-images commute.

5 Paracompactness, Bing's Metrization Theorem

5.1 Definitions

A family $\mathcal{A}$ of subsets of X is locally finite (resp. discrete) if and only if for every $x \in X$ there is open $U \ni x$ such that $\{A \in \mathcal{A}\colon U \cap A \neq \emptyset\}$ is finite (resp. empty or a singleton).

A family $\mathcal{A}$ of subsets of X is closure preserving if and only if for every $\mathcal{A}' \subseteq \mathcal{A}$, $\overline{\bigcup_{A \in \mathcal{A}'} A} = \bigcup_{A \in \mathcal{A}'} \overline{A}$.

A family $\mathcal{A}$ is a refinement of a family $\mathcal{B}$ (of subsets of X) if and only if for every $A \in \mathcal{A}$, there is $B \in \mathcal{B}$ such that $A \subseteq B$.

A topological space is paracompact if and only if every open cover $\mathcal{U}$ of X has a locally finite open refinement covering X.

The hedgehog of spininess κ is $H_\kappa = \{0\} \cup ((0, 1] \times \kappa)$ with metric d given by $d(0, (t, i)) = t$, $d((t, i), (s, i)) = |t - s|$, $d((t, i), (s, j)) = t + s$ for $i \neq j$.

5.2 Results

Theorem 5.1. *1. Locally finite families are closure preserving.*

2. *A paracompact regular space is normal.*

3. *For a regular space X the following are equivalent:*

   (a) *X is paracompact.*

   (b) *Every open cover $\mathcal{U}$ has a sequence $\mathcal{V}_n$ of locally finite, open refinements such that $\bigcup_n \mathcal{V}_n$ covers X (i.e. every open cover has a σ-locally finite open refinement covering X).*

   (c) *Every open cover $\mathcal{U}$ has a locally finite refinement covering X.*

   (d) *Every open cover $\mathcal{U}$ has a locally finite closed refinement covering X.*

4. ***Stone's Theorem:*** *Every metric space is paracompact. If X is a metric space and $\mathcal{U}$ is an open cover of X then there are refinements $\mathcal{V}_n, n \in \mathbb{N}$ of $\mathcal{U}$ such that each $\mathcal{V}_n$ is a discrete family, and $\bigcup_n \mathcal{V}_n$ covers X. I.e. Every open cover of X has an open, σ-discrete refinement covering X.*

5. ***Bing's Metrization Theorem:*** *A space is metrizable if and only if X is perfectly normal and has a σ-discrete basis if and only if X is homeomorphic to a subspace of a countable product of hedgehogs (of some spininess).*

5.3 Proofs

1. If $\mathcal{A}$ is locally finite and $\mathcal{A}' \subseteq \mathcal{A}$, $\mathcal{A}'$ is still locally finite. It is thus sufficient to show $\overline{\bigcup \mathcal{A}} = \bigcup_{A \in \mathcal{A}} \overline{A}$. $\supseteq$ is clear. For $\subseteq$, assume that $x \notin \bigcup_{A \in \mathcal{A}} \overline{A}$. Let $U \ni x$ be open such that U meets only finitely many elements of $\mathcal{A}$, say $A_1, \ldots, A_n$. For each $i = 1, \ldots, n$, $x \notin \overline{A_i}$, so choose open $V_i \ni x$ disjoint from A_i. Then $x \in U \cap \bigcap_i V_i$ and the RHS is open and disjoint from $\bigcup \mathcal{A}$ as required.

2. If C, D are disjoint closed, for each $c \in C$ choose open $U_c \ni c$ such that $\overline{U_c} \cap D = \emptyset$. Then $\{U_c \colon c \in C\} \cup \{X \setminus C\}$ is an open cover of X so has a locally finite open refinement $\mathcal{V}'$. Let $\mathcal{V} = \{V \in \mathcal{V}' \colon V \cap C \neq \emptyset\}$,

still a locally finite family which refines $\{U_c\colon c \in C\}$ and covers C. Since locally finite families are closure preserving we have $\overline{\bigcup_{V\in\mathcal{V}} V} = \bigcup_{V\in\mathcal{V}} \overline{V} \subseteq \bigcup_{c\in C} \overline{U_c}$ disjoint from D, so that $U = \bigcup \mathcal{V}$ is as required.

3. Exercise Sheet.

4. **Stone's Theorem:** Suppose $\mathcal{U}$ is an open cover of X. Well order $\mathcal{U}$ by some well-order $\leq$ (using Choice). For each $n \in \mathbb{N}$ and $U \in \mathcal{U}$, define

$$S_{U,n} = \left\{x \in X\colon B_{3/2^n}(x) \subseteq U \wedge \forall U' < U\ x \notin U'\right\}$$

and let

$$V_{U,n} = \bigcup_{x\in S_{U,n}} B_{1/2^n}(x),$$

an open subset of U. If $y \in V_{U,n}$ and $y' \in V_{U',n}$ with (wlog) $U < U'$ then there is $x \in S_{U,n}$ with $d(x,y) < 1/2^n$ and $x' \in S_{U',n}$ with $d(x',y') < 1/2^n$. But if $x' \in S_{U',n}$ then $x' \notin U$ so $d(x,x') \geq 3/2^n$ and hence $d(y,y') \geq d(x,x') - d(x,y) - d(x',y') \geq 1/2^n$. Thus each $B_{2^{-(n+1)}}(y), y \in X$ meets at most one $V_{U,n}, U \in \mathcal{U}$ and hence $\mathcal{V}_n = \{V_{U,n}\colon U \in \mathcal{U}\}$ is a discrete family. Clearly $V_{U,n} \subseteq U$. Finally $\bigcup_n \mathcal{V}_n$ covers X, since for $x \in X$, choose $U \in \mathcal{U}$ minimal such that $x \in U$ and as U is open, find $n \in \mathbb{N}$ with $B_{3/2^n}(x) \subseteq U$ giving $x \in V_{U,n}$.

Noting that metric spaces are regular and that σ-discrete implies σ-locally finite, we see that metric spaces are paracompact. (In fact, by defining $S_{U,n} = \left\{x \in X\colon B_{3/2^n}(x) \subseteq U \wedge \forall U' < U\ x \notin U' \wedge x \notin \bigcup_{U\in\mathcal{U},n'<n} V_{U,n'}\right\}$ you could directly obtain a locally finite open refinement.)

5. **Bing's Metrization Theorem:** $d_C(x) = \inf\{d(x,c)\colon c \in C\}$ witnesses perfect normality of metric spaces. For the σ-discrete base, apply Stone's Theorem to each $\{B_{2^{-n}}(x)\colon x \in X\}$ to obtain a σ-discrete open refinement $\mathcal{V}_n$ covering X and then note that $\bigcup_n \mathcal{V}_n$ is a σ-discrete basis of X.

Now assume that X is perfectly normal and has a σ-discrete basis $\mathcal{B} = \bigcup_n \mathcal{B}_n$ with each $\mathcal{B}_n$ discrete. Fix $n \in \mathbb{N}$. For each $B \in \mathcal{B}_n$, let $f_B\colon X \to [0,1]$ be continuous such that $f^{-1}(0) = X \setminus B$ and define $F_n\colon X \to H_{\mathcal{B}_n}$ by $F_n(x) = 0$ if $x \notin \bigcup \mathcal{B}_n$ and $F_n(x) = (f_B(x), B)$ if $x \in B \in \mathcal{B}_n$. This is well-defined since $\mathcal{B}_n$ is discrete (each x is in at most one B). It is continuous since each f_B is continuous and $\mathcal{B}_n$ is

discrete: for $x \in X$, choose open $U \ni x$ that meets at most one element of $\mathcal{B}_n$, say B. Then $F_n|_U = f_B$ (where we identify the co-domain $[0,1]$ of f_B with $\{0\} \cup (0,1] \times \{B\}$) is continuous on U, hence F_n is continuous at x. If U meets no element of $\mathcal{B}_n$ then F_n is constant with value 0 hence continuous on U. Note that $\left\{ {F_n}^{-1}(U) : U \text{ open } \subseteq H_{\mathcal{B}_n} \right\} \supseteq \mathcal{B}_n$. Thus $\{F_n : n \in \mathbb{N}\}$ satisfies the conditions of the Embedding Lemma and hence X is homeomorphic to a subspace of a countable product of hedgehogs.

Finally, a countable product of metric spaces is metrizable.

6 Connectedness, Zero-Dimensionality

6.1 Definitions

A disconnection of X is a partition of X into two non-empty closed-and-open (clopen) subsets. X is disconnected if and only if there is a disconnection of X.

X is connected if and only if it is not disconnected.

The component of a point $x \in X$ is the greatest connected subspace $C(x)$ of X containing x.

The quasicomponent of a point $x \in X$ is $Q(x) = \bigcap \{F \subseteq X : x \in F \text{ clopen}\}$.

X is totally disconnected if and only if every component is a singleton.

X is zero-dimensional if and only if X has a basis of clopen sets.

6.2 Results

Lemma 6.1 (Recall)**.** *1. X is connected if and only if every continuous function into the discrete two-point space is constant.*

2. Suppose $A, A_i \subseteq X, i \in I$ are connected. If for each $i \in I$, $A \cap A_i \neq \emptyset$ then $A \cup \bigcup_i A_i$ is connected.

3. The component of a point exists and equals $\bigcup \{C \subseteq X : x \in C \text{ connected}\}$.

Theorem 6.2. *1. Both the components and the quasicomponents of a space form a partition.*

2. For every $x \in X$, $C(x) \subseteq Q(x)$.

*3. **Sura-Bura Lemma:** If X is compact Hausdorff, then for every $x \in X$, $C(x) = Q(x)$.*

4. A totally disconnected compact Hausdorff space is zero-dimensional.

6.3 Proofs

3. **Sura-Bura Lemma:** Suppose that some quasicomponent $Q = Q(x)$ is disconnected, i.e. there are Q-clopen non-empty disjoint A, B such $Q = A \cup B$. As Q is closed (an intersection of closed sets), A, B are closed in X. As X is compact Hausdorff, it is normal, so there are disjoint open $U \supseteq A, V \supseteq B$. As X is compact and $\{U \cup V\} \cup \{X \setminus F : x \in F \text{ clopen} \subseteq X\}$ covers X, there are finitely many X-clopen $F_1, \ldots, F_n$ containing x such that $U \cup V, X \setminus F_1, \ldots, X \setminus F_n$ covers X so that $x \in F = \bigcap_i F_i \subseteq U \cup V$ and F is clopen (in X). Now $\overline{F \cap U} \subseteq \overline{F} \cap \overline{U} \subseteq F \cap (X \setminus V) \subseteq (F \cap (U \cup V)) \cap (X \setminus V) \subseteq F \cap U$. Hence $F \cap U$ is X-clopen and similarly $F \cap V$ is X-clopen. Wlog $x \in F \cap U$ and hence $Q \subseteq F \cap U$, contradicting $B \neq \emptyset$.

7 Stone Duality

7.1 Definitions

A Boolean Algebra is a partial order $(B, \leq)$ with greatest element $\mathbf{1}$, least element $\mathbf{0}$, binary infima $a \wedge b$, binary suprema $a \vee b$ that distribute over each other, i.e. $a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c)$ as well as $a \vee (b \wedge c) = (a \vee b) \wedge (a \vee c)$, and unary negation operator $\neg a$ such that $a \wedge \neg a = \mathbf{0}$, $a \vee \neg a = \mathbf{1}$, $\neg(a \wedge b) = (\neg a) \vee (\neg b)$, $\neg(a \vee b) = (\neg a) \wedge (\neg b)$.

Given two Boolean Algebras $(A, \leq_A), (B, \leq_B)$ a function $f : A \to B$ is a Boolean Algebra homomorphism if and only if it preserves all the above notions, i.e.

- $a \leq a' \Rightarrow f(a) \leq f(a')$ (here and below for the Boolean Algebra notions, the first is the one on A, the second the one on B)
- $f(\mathbf{1}) = \mathbf{1}$
- $f(\mathbf{0}) = \mathbf{0}$
- $f(a \wedge a') = f(a) \wedge f(a')$
- $f(a \vee a') = f(a) \vee f(a')$
- $f(\neg a) = \neg f(a)$

f is a Boolean Algebra isomorphism if and only if f is a Boolean Algebra homomorphism and is bijective.

For a topological space X, $\mathcal{B}_X = \{C \colon C \text{ clopen } \subseteq X\}$.

For a continuous map $f : X \to Y$, $\mathcal{B}_f : \mathcal{B}_Y \to \mathcal{B}_X; C \mapsto f^{-1}(C)$.

A filter $\mathcal{F}$ on a Boolean Algebra B is a subset of B such that

- $\mathbf{0} \notin \mathcal{F} \neq \emptyset$;
- $a \leq b, a \in \mathcal{F} \Rightarrow b \in \mathcal{F}$;
- $a, b \in \mathcal{F} \Rightarrow a \wedge b \in \mathcal{F}$;

An ultrafilter $\mathcal{U}$ on a Boolean Algebra B is a filter on B which is maximal wrt $\subseteq$.

For a Boolean Algebra B and $b \in B$, write $b^\star = \{\mathcal{U} \colon b \in \mathcal{U} \text{ ultrafilter on } B\}$.

For a Boolean Algebra B the Stone Space of B is $\mathcal{S}(B) = \{\mathcal{U} \colon \mathcal{U} \text{ ultrafilter on } B\}$ with topology τ_B generated by the basis $\{b^\star \colon b \in B\}$.

For a Boolean Algebra homomorphism $g : B \to A$ we define $\mathcal{S}(g) : \mathcal{S}(A) \to \mathcal{S}(B)$ by $\mathcal{S}(g)(\mathcal{U}) = \{b \in B \colon g(b) \in \mathcal{U}\}$.

A Stone space is a compact Hausdorff zero-dimensional topological space.

7.2 Results

Theorem 7.1. *Suppose B is a Boolean Algebra.*

1. *$\wedge$ and $\vee$ are commutative and associative.*
2. *If f is a Boolean Algebra isomorphism then f^{-1} is one as well.*
3. *If $\mathcal{F}$ is a filter on a Boolean Algebra then $\mathbf{1} \in \mathcal{F}$.*
4. *A filter $\mathcal{F}$ on a Boolean Algebra B is an ultrafilter if and only if for every $a \in B$ $a \in \mathcal{F}$ or $\neg a \in \mathcal{F}$.*
5. *A filter $\mathcal{F}$ on a Boolean Algebra B is an ultrafilter if and only if for every $a, b \in B$, $a \vee b \in \mathcal{F} \Rightarrow a \in \mathcal{F}$ or $b \in \mathcal{F}$.*
6. *Every filter on a Boolean Algebra can be extended to an ultrafilter.*
7. *Every non-empty collection of non-zero elements of B which is closed under finite minima can be extended to an ultrafilter.*

8. *If a, b are elements of a Boolean Algebra B then $(a \vee b)^\star = a^\star \cup b^\star$, $(a \wedge b)^\star = a^\star \cap b^\star$, $(\neg a)^\star = \mathcal{S}(B) \setminus a^\star$, $\mathbf{0}^\star = \emptyset$, $\mathbf{1}^\star = \mathcal{S}(B)$, $a \leq b \Rightarrow a^\star \subseteq b^\star$.*

9. *If a, b are distinct elements of a Boolean Algebra then $a^\star \neq b^\star$, so in particular $b^\star = \emptyset \Rightarrow b = \mathbf{0}$.*

10. *If $g : B \to A$ is a Boolean Algebra homomorphism and $\mathcal{U}$ is an ultrafilter on A then $\mathcal{S}(g)(\mathcal{U})$ is an ultrafilter on B.*

11. *$\mathcal{B}_X$ ordered by $\subseteq$ is a Boolean Algebra with $\mathbf{1} = X$, $\mathbf{0} = \emptyset$, $\vee = \cup$, $\wedge = \cap$, $\neg C = X \setminus C$.*

12. *If $f : X \to Y$ is a continuous map between topological spaces then $\mathcal{B}_f : \mathcal{B}_Y \to \mathcal{B}_X$ is a Boolean Algebra homomorphism.*

13. *If $f : X \to Y, g : Y \to Z$ are continuous maps between topological spaces then $\mathcal{B}_{g \circ f} = \mathcal{B}_f \circ \mathcal{B}_g$.*

14. *If B is a Boolean Algebra then τ_B is a compact Hausdorff zero-dimensional topology on $\mathcal{S}(B)$.*

15. *If $g : B \to A$ is a Boolean Algebra homomorphism then $\mathcal{S}(g)$ is continuous map.*

16. *If $g : B \to A, f : C \to B$ are Boolean Algebra homomorphisms then $\mathcal{S}(g \circ f) = \mathcal{S}(f) \circ \mathcal{S}(g)$.*

17. *If X is a Stone space then $\mathcal{S}(\mathcal{B}_X)$ is homeomorphic to X by the homeomorphism $h_X : X \to \mathcal{S}(\mathcal{B}_X); x \mapsto \{C \in \mathcal{B}_X : x \in C\}$.*

18. *If B is a Boolean Algebra then $\mathcal{B}_{\mathcal{S}(B)}$ is isomorphic to B by the isomorphism $h_B : B \to \mathcal{B}_{\mathcal{S}(B)}; b \mapsto b^\star$.*

19. *If $f : X \to Y$ is a continuous map between Stone spaces then $\mathcal{S}(\mathcal{B}_f) \circ h_X = h_Y \circ f$.*

20. *If $g : B \to A$ is a Boolean Algebra homomorphism then $\mathcal{B}_{\mathcal{S}(g)} \circ h_B = h_A \circ g$.*

21. *(Non-examinable) $\mathcal{S}(.)$ and $\mathcal{B}_.$ witness that the categories of Boolean Algebras and Stone spaces are dually equivalent.*

7.3 Proofs

9. If $b \wedge \neg a = \mathbf{0}$ then $b = b \wedge \mathbf{1} = b \wedge (a \vee \neg a) = (b \wedge a) \vee (b \wedge \neg a) = b \wedge a = \inf\{a, b\}$ and thus $b \leq a$. Thus if $a \neq b$ then one of $b \wedge \neg a$ and or $a \wedge \neg b$ is non-zero and thus contained in an ultrafilter $\mathcal{U}$ on B. But then $\mathcal{U} \in b^\star \setminus a^\star$ or $\mathcal{U} \in a^\star \setminus b^\star$.

14. $\{b^\star \colon b \in B\}$ is closed under finite intersections so a basis for a topology τ on $\mathcal{S}(B)$ consisting of clopen sets $(\mathcal{S}(B) \setminus b^\star = (\neg b)^\star)$.

    τ is Hausdorff since if $\mathcal{U} \neq \mathcal{V}$ are ultrafilters on B then find $a \in \mathcal{U} \setminus \mathcal{V}$ (wlog). Then $\neg a \in \mathcal{V} \setminus \mathcal{U}$ and $\mathcal{U} \in a^\star, \mathcal{V} \in (\neg a)^\star$ are disjoint open sets.

    We show that τ is compact by checking the finite intersection property for basic closed sets, i.e. those of the form $b^\star$, $b \in B$. So suppose $\mathcal{C}$ is a collection of basic closed sets such that every finite intersection is non-empty and write $\mathcal{C} = \{b_i^\star \colon i \in I\}$. Then $\left\{\bigwedge_{j \in J} b_j \colon J \text{ finite } \subseteq I\right\}$ can be extended to an ultrafilter $\mathcal{U}$ and $\mathcal{U} \in \bigcap \mathcal{C}$.

15. $\mathcal{S}(g)^{-1}(b^\star) = (g(b))^\star$

17. h_X is continuous since $h_X^{-1}(C^\star) = C$. If $x \neq y$ then there is clopen U such that $x \in U \not\ni y$ and hence $U \in h_X(x) \setminus h_X(y)$ and thus h_X is injective. It is surjective since if $\mathcal{U}$ is an ultrafilter on $\mathcal{B}_X$ then interpreting it as a collection of closed subsets of X, it has the finite intersection property and thus there is $x \in \bigcap \mathcal{U}$ giving $h_X(x) = \mathcal{U}$. Thus h_X is a continuous bijection from a compact into a Hausdorff space and thus a homeomorphism.

18. Note that every clopen subset of $\mathcal{S}(B)$ is a $b^\star$: it is a union of $b^\star$s and is compact so a finite union of them, say $b_1^\star \cup \ldots \cup b_n^\star = (b_1 \vee \cdots \vee b_n)^\star$. Thus h_B is surjective. By 9. it is injective and by 8. it is a homomorphism.