

# Infinite Groups

Cornelia Druțu

University of Oxford

Part C course MT 2022, Oxford

**William Thurston:** “Mathematics is not about numbers, equations, computations, or algorithms: it is about understanding.”

At the entrance to Plato's Academy, an inscription over the door said:

**Let no one destitute of geometry enter here.**

The same is written at the entrance to the Maths Institute, as an engraving on the floor though.

## Material available on the webpage

<https://courses.maths.ox.ac.uk/course/view.php?id=718>

- **Lecture Notes**
- **Revision Notes:** collect material seen in courses from previous years, to be used as a reference only. Not examinable. Occasional reminders in lectures.
- **Hand out Notes:** expand on some notions introduced in the course, as further reading for students who wish to have a better understanding of the latter. Not examinable.
- **Mini-projects Broadening.** Please send me an email asap, and **by week 4 at the latest**, specifying your supervisor, the approximate topic of your DPhil and what project on the list you would like. Feel free to suggest other projects as well, as long as they are related to this course.

# Topics of the course

We study **countable infinite groups**.

**Methods of study:**

- Endow these groups with a metric, a geometry.
  - Easiest way to do it: using **Cayley graphs**.
  - Works for all groups **with a finite generating set**.
  - A recurrent theme: try to connect algebraic features to geometric features of Cayley graphs.
- Approximate these groups by finite groups.
  - For instance **larger and larger finite quotients**.
  - Works for **residually finite groups**.
- Design algorithms/construct Turing machines finding solutions to algebraic questions.  
Works mainly for (some) finitely presented groups (**groups that can be fully described to a computer via finite data**).

## Topics of the course 2

- Represent infinite groups as groups of matrices.  
The groups that can be thus represented are called **linear groups**.

### Classes of infinite groups:

- **“Small”**: Abelian finitely generated  $\subset$  Nilpotent finitely generated  $\subset$  Polycyclic  $\subset$  Solvable.
- **“Large”**: Free groups  $\subset$  Hyperbolic groups.  
Free groups  $\subset$  Amalgamated products (in the sense of J.P. Serre).

The families of “small” groups will be the topic of the **“Infinite Groups” course**.

The families of “large” groups will be the topic of the **“Geometric Group Theory” course** in Hilary Term.

# Constructions of groups, old and new

## Direct product

The standard approach: take  $H_1, H_2$ , define operation on  $H_1 \times H_2$ .

**Another approach:** given  $G$  and  $H_1, H_2$  subgroups of  $G$ , how to decide if  $G$  isomorphic to  $H_1 \times H_2$ ?

Three conditions:

- $H_1, H_2$  both normal subgroups.
- $H_1 \cap H_2 = \{1\}$ .
- $H_1 H_2 = G$ .

### Generalization: direct sum

Let  $X \neq \emptyset$ ,  $\mathcal{G} = \{G_x \mid x \in X\}$  a collection of groups.

Consider

$$\text{Map}_f(X, \mathcal{G}) := \left\{ f : X \rightarrow \prod_{x \in X} G_x \mid f(x) \in G_x, f(x) \neq 1_{G_x} \right.$$

for only finitely many  $x \in X$   $\left. \right\}$ .

The **direct sum**  $\bigoplus_{x \in X} G_x$  is  $\text{Map}_f(X, \mathcal{G})$ , endowed with the pointwise multiplication:

$$(f \cdot g)(x) = f(x) \cdot g(x), \forall x \in X.$$

When  $G_x = G$  for all  $x \in X$ , the direct sum is denoted either by  $\bigoplus_{x \in X} G$  or by  $G^{\oplus X}$ .

## Semidirect product

Given two groups  $N$  and  $H$  and a group homomorphism  $\varphi : H \rightarrow \text{Aut}(N)$ , one can define a new group  $G = N \rtimes_{\varphi} H$  called **semidirect product of  $N$  and  $H$  with respect to  $\varphi$** :

- As a set,  $N \rtimes_{\varphi} H$  is defined as the cartesian product  $N \times H$ .
- Binary operation  $*$  on  $G$  defined by

$$(n_1, h_1) * (n_2, h_2) = (n_1 \varphi(h_1)(n_2), h_1 h_2), \quad \forall n_1, n_2 \in N \text{ and } h_1, h_2 \in H.$$

- If  $\varphi$  is trivial (i.e. has as image  $\{\text{id}_N\}$ ) then  $N \rtimes_{\varphi} H$  is the direct product  $N \times H$ .



## Semi-direct product 2

Given a group  $G$  and two subgroups  $H, N$  how to know if  $G$  isomorphic to  $N \rtimes_{\varphi} H$  for some  $\varphi$ ?

Again three conditions:

- $N$  normal subgroup.
- $N \cap H = \{1\}$ .
- $NH = G$ .

If the above are satisfied,  $G$  isomorphic to  $N \rtimes_{\varphi} H$ , where  $\varphi(h) =$  conjugation by  $h$  of every element in  $N$ .

## A more general notion

An **exact sequence** is a sequence of groups and group homomorphisms

$$\dots G_{n-1} \xrightarrow{\varphi_{n-1}} G_n \xrightarrow{\varphi_n} G_{n+1} \dots$$

such that  $\text{Im } \varphi_{n-1} = \ker \varphi_n$  for every  $n$ .

A **short exact sequence** is an exact sequence of the form:

$$\{1\} \longrightarrow N \xrightarrow{\varphi} G \xrightarrow{\psi} H \longrightarrow \{1\}. \quad (1)$$

In other words,  $\varphi$  is an isomorphism from  $N$  to a normal subgroup  $N' \triangleleft G$  and  $\psi$  defines an isomorphism  $G/N' \simeq H$ .

If  $G$  is isomorphic to  $N \rtimes_{\varphi} H$  then we have a short exact sequence as above. The converse is in general not true.

## Semidirect product and short exact sequence

A short exact sequence **splits** if there exists a homomorphism  $\sigma : H \rightarrow G$  (called a **section**) such that

$$\psi \circ \sigma = \text{id}.$$

A split exact sequence determines a decomposition of  $G$  as a semidirect product  $\varphi(N) \rtimes \sigma(H)$ .

### Examples

- 1 The dihedral group  $D_{2n}$  is isomorphic to  $\mathbb{Z}_n \rtimes_{\varphi} \mathbb{Z}_2$ , where  $\varphi(1)(k) = n - k$ .
- 2 The infinite dihedral group  $D_{\infty}$  is isomorphic to  $\mathbb{Z} \rtimes_{\varphi} \mathbb{Z}_2$ , where  $\varphi(1)(k) = -k$ .
- 3 The permutation group  $S_n$  is the semidirect product of  $A_n$  and  $\mathbb{Z}_2 = \{\text{id}, (12)\}$ .

## Wreath product

Consider a direct sum  $\bigoplus_{x \in H} G$  with index set a group  $H$ .

There is a natural action of  $H$  on the direct sum:

$$\varphi : H \rightarrow \text{Aut} \left( \bigoplus_{h \in H} G \right), \quad \varphi(h)f(x) = f(h^{-1}x), \quad \forall x \in H.$$

Thus, we define the semidirect product

$$\left( \bigoplus_{h \in H} G \right) \rtimes_{\varphi} H. \quad (2)$$

The semidirect product (2) is called **the wreath product of  $G$  with  $H$** , and it is denoted by  $G \wr H$ .

The wreath product  $G = \mathbb{Z}_2 \wr \mathbb{Z}$  is called the **lamplighter group**. Its name comes from the way in which  $\varphi(1)$  acts.

The wreath product construction is a source of interesting examples of groups, in particular of solvable groups.