


# Infinite Groups

Cornelia Druțu

University of Oxford

Part C course MT 2022

# Graham Higman



## Theorem

Every finitely generated *recursively presented* group can be embedded as a subgroup of some finitely presented group.

# Finite presentation

## Remark

$G$  finitely presented *does not* imply  $H \leq G$  finitely presented or  $G/N$  finitely presented, for  $N \triangleleft G$ .

## Proposition

Let  $G$  be a group, and  $H \leq G$  such that  $|G : H|$  is finite. Then  $G$  is FP if and only if  $H$  is FP.

**Proof** Suppose  $G = \langle X \mid R \rangle$  with  $X$  and  $R$  finite.

We have an epimorphism  $\pi : F = F(X) \rightarrow G$  with  $K = \ker \pi = \langle\langle R \rangle\rangle$ .

Let  $E = \pi^{-1}(H)$ . Then  $|F : E| = |G : H|$  is finite, so  $E = F(Y)$  for some finite  $Y$ .

Since  $K \leq E$ , each  $r \in R$  satisfies  $r = s_r(Y)$  for some word  $s_r$  on  $Y$ . Put  $S = \{s_r(Y) \mid r \in R\}$ . Then  $\pi_1 = \pi|_E : E \rightarrow H$  is an epimorphism and

$$\ker \pi_1 = K = \langle\langle S \rangle\rangle = \langle S^F \rangle.$$

## FP of finite index subgroups continued

Say  $F = a_1E \cup \dots \cup a_nE$ .

Then  $S^F = (S^{a_1} \cup \dots \cup S^{a_n})^E$  and

$$\langle S^F \rangle = \langle (S^{a_1} \cup \dots \cup S^{a_n})^E \rangle = \langle\langle S^{a_1} \cup \dots \cup S^{a_n} \rangle\rangle_E$$

Thus  $\langle Y; S^{a_1} \cup \dots \cup S^{a_n} \rangle$  is a presentation for  $H$ .

**Conversely**, suppose that  $H$  is FP.

Let  $N \leq H$  be a normal subgroup of finite index in  $G$  (see **Revision notes**).

Then  $|H : N|$  is finite, so  $N$  is FP by the first part.

Also  $G/N$  is FP because finite. Therefore  $G$  is FP. □

## Brief incursion into residual finiteness

The idea when introducing this concept is to approximate an infinite group by its finite quotients.

So one needs to have enough finite quotients.

### Proposition

Let  $G$  be a group. The following are equivalent:

①

$$\bigcap_{i \in I} H_i = \{1\},$$

where  $\{H_i : i \in I\}$  is the set of all finite-index subgroups in  $G$ ;

② for every  $g \in G \setminus \{1\}$ , there exists a finite group  $\Phi$  and a homomorphism  $\varphi : G \rightarrow \Phi$ , such that  $\varphi(g) \neq 1$ .

### Definition

A group satisfying the above is called **residually finite**.

## Residual finiteness, equivalence

**Proof.** The key remark is that

$$\bigcap_{i \in I} H_i = \bigcap_{j \in J} N_j,$$

where  $\{N_j : j \in J\}$  is the set of all finite-index **normal** subgroups in  $G$ . This is because: **for every  $H \leq G$  of finite index there exists  $N \triangleleft G$  of finite index,  $N \leq H$ .**

**(1)  $\Rightarrow$  (2)**  $\forall g \neq 1, \exists N \triangleleft G$  of finite index,  $g \notin N$ . Take  $\varphi : G \rightarrow G/N$ .

**(2)  $\Rightarrow$  (1)**  $\forall g \neq 1, \exists \varphi : G \rightarrow F$  finite, such that  $g \notin \ker \varphi$ . Therefore  $g \notin \bigcap_{j \in J} N_j$  □

## Examples of RF groups

### Example

*The group  $\Gamma = GL(n, \mathbb{Z})$  is residually finite.*

Indeed, we take subgroups  $\Gamma(p) \leq \Gamma$ ,  $\Gamma(p) = \ker(\varphi_p)$ , where  $\varphi_p : \Gamma \rightarrow GL(n, \mathbb{Z}_p)$  is the reduction modulo  $p$ .

Assume  $g \in \Gamma$  is a non-trivial element.

If  $g$  has a non-zero off-diagonal entry  $g_{ij} \neq 0$ , then  $g_{ij} \not\equiv 0 \pmod{p}$ , whenever  $p > |g_{ij}|$ . Thus,  $\varphi_p(g) \neq 1$ .

If  $g \in \Gamma$  has only zero entries off-diagonal then it is a diagonal matrix with only  $\pm 1$  on the diagonal, and at least one entry  $-1$ . Then  $\varphi_3(g)$  has at least one 2 on the diagonal, hence  $\varphi_3(g) \neq 1$ .

Thus  $\Gamma$  is residually finite.

## A Theorem of Mal'cev. A Lemma of Selberg

### Theorem (A. I. Mal'cev)

Let  $\Gamma$  be a *finitely generated* subgroup of  $GL(n, R)$ , where  $R$  is a commutative ring with unity. Then  $\Gamma$  is residually finite.

Mal'cev's theorem is complemented by the following result:

### Theorem (Selberg's Lemma)

Let  $\Gamma$  be a finitely generated subgroup of  $GL(n, F)$ , where  $F$  is a field of characteristic zero. Then  $\Gamma$  contains a torsion-free subgroup of finite index.



# Properties of RF

## Proposition

- ①  $G, H$  residually finite (RF)  $\Rightarrow G \times H$  RF;
- ②  $G$  RF and  $H \leq G \Rightarrow H$  RF;
- ③  $H \leq G$  of finite index and  $H$  RF  $\Rightarrow G$  RF;
- ④  $H$  *finitely generated* RF and  $Q$  RF  $\Rightarrow H \rtimes Q$  RF.

## Remark

There exist short exact sequences

$$\{1\} \longrightarrow \mathbb{Z}_2 \xrightarrow{i} G \xrightarrow{\pi} Q \longrightarrow \{1\},$$

with  $Q$  finitely generated RF and  $G$  not RF (J. Millson 1979).

## Corollary

*The free group  $F_2$  of rank 2 is residually finite. Every free group of (at most) countable rank is residually finite.*

## Remark

*This in particular shows that  $G$  RF does not imply  $G/N$  RF, for  $N \triangleleft G$ .*

## Remark

*Given a short exact sequence*

$$\{1\} \longrightarrow H \xrightarrow{i} G \xrightarrow{\pi} F(X) \longrightarrow \{1\},$$

*with  $H$  finitely generated RF and  $X$  finite or countable,  $G$  is residually finite.*

## Back to polycyclic groups

### Proposition

*Polycyclic groups are finitely presented and residually finite.*

**Proof** by induction on the length  $\ell(G)$ .

For  $\ell(G) = 1$ ,  $G$  is cyclic.

Assume that the statement is true for polycyclic groups of length  $n$ , let  $G$  be polycyclic with  $\ell(G) = n + 1$ .

Let  $N_1$  be the first (sub)normal subgroup in a cyclic series of minimal length  $n + 1$ . Then  $N_1$  is polycyclic of length  $n$ , hence finitely presented (respectively residually finite) by the induction hypothesis.

$N_1$  is always finitely generated, because polycyclic.

## Induction proving polycyclic groups are FP and RF

We have the short exact sequence

$$\{1\} \longrightarrow N_1 \xrightarrow{i} G \xrightarrow{\pi} C \longrightarrow \{1\},$$

where  $C$  is cyclic.

This implies  $G$  finitely presented.

When  $C$  finite,  $N$  has finite index, hence  $G$  RF.

When  $C = \mathbb{Z}$ ,  $G = N_1 \rtimes \mathbb{Z}$ , hence RF.



## Normal poly- $C_\infty$ subgroup

### Proposition

*A polycyclic group contains a normal subgroup of finite index which is poly- $C_\infty$ .*

**Proof** By induction on the length  $\ell(G) = n$ .

For  $n = 1$  the group  $G$  is cyclic and the statement true.

Assume the assertion is true for  $n$  and consider a polycyclic group  $G$  with a cyclic series of length  $n + 1$ .

The induction hypothesis implies that  $N_1$  (the first group in the series) contains a normal subgroup  $S$  of finite index which is poly- $C_\infty$ .

Proposition 2.8, (2), in **Revision Notes** implies that  $S$  contains  $S_1$  characteristic subgroup of  $N_1$  of finite index.

Since  $N_1 \triangleleft G$ ,  $S_1$  is normal in  $G$ .

$S_1 \leq S \Rightarrow S_1$  is poly- $C_\infty$ .

If  $G/N_1$  is finite then  $S_1$  has finite index in  $G$ .

## Normal poly- $C_\infty$ subgroup 2

Assume  $G/N_1$  is infinite cyclic.

Then the group  $K = G/S_1$  contains the finite normal subgroup  $F = N_1/S_1$  such that  $K/F$  is isomorphic to  $\mathbb{Z}$ .

In other words, we have the short exact sequence

$$\{1\} \longrightarrow F \xrightarrow{\varphi} K \xrightarrow{\psi} \mathbb{Z} \longrightarrow \{1\}.$$

Then  $K$  is a semidirect product of  $F$  and an infinite cyclic subgroup  $\langle x \rangle$ . The conjugation by  $x$  defines an automorphism of  $F$  and since  $\text{Aut}(F)$  is finite, there exists  $r$  such that the conjugation by  $x^r$  is the identity on  $F$ . Hence  $F\langle x^r \rangle$  is a finite-index subgroup in  $K$  and it is a direct product of  $F$  and  $\langle x^r \rangle$ .

We conclude that  $\langle x^r \rangle$  is a finite index normal subgroup of  $K$ .

We have that  $\langle x^r \rangle = G_1/S_1$ , where  $G_1$  is a finite index normal subgroup in  $G$ , and  $G_1$  is poly- $C_\infty$  since  $S_1$  is poly- $C_\infty$ .  $\square$